

Programming Bugs: Microbiota and the Developmental Origins of Brain Health and Disease

Martin G. Codagnone, Simon Spichak, Siobhain M. O'Mahony, Olivia F. O'Leary, Gerard Clarke, Catherine Stanton, Timothy G. Dinan, and John F. Cryan

ABSTRACT

It has been nearly 30 years since Dr. David Barker first highlighted the importance of prenatal factors in contributing to the developmental origins of adult disease. This concept was later broadened to include postnatal events. It is clear that the interaction between genetic predisposition and early life environmental exposures is key in this regard. However, recent research has also identified another important factor in the microbiota—the trillions of microorganisms that inhabit key body niches, including the vagina and gastrointestinal tract. Because the composition of these maternal microbiome sites has been linked to maternal metabolism and is also vertically transmitted to offspring, changes in the maternal microbiota are poised to significantly affect the newborn. In fact, several lines of evidence show that the gut microbiota interacts with diet, drugs, and stress both prenatally and postnatally and that these exogenous factors could also affect the dynamic changes in the microbiota composition occurring during pregnancy. Animal models have shown great utility in illuminating how these disruptions result in behavioral and brain morphological phenotypes reminiscent of psychiatric disorders (anxiety, depression, schizophrenia, and autism spectrum disorders). Increasing evidence points to critical interactions among the microbiota, host genetics, and both the prenatal and postnatal environments to temporally program susceptibility to psychiatric disorders later in life. Sex-specific phenotypes may be programmed through the influence of the microbiota on the hypothalamic-pituitary-adrenal axis and neuroimmune system.

Keywords: Brain development, Early life, Gut-brain axis, Microbiota, Neuropsychiatry, Stress

<https://doi.org/10.1016/j.biopsych.2018.06.014>

BUILDING ON BARKER: THE FOUR HITS HYPOTHESIS FOR DEVELOPMENT OF DISEASE

Studies on the developmental origins of health and disease have gleaned mechanistic insights as well as risk factors that affect development. Using large cohort studies, Dr. David Barker showed that low birth weight was a risk factor for cardiac disease and ischemia (1–4). He postulated that factors affecting the fetus, prenatal and early postnatal environment, and genetic predisposition programmed health and disease during adulthood (1–4). Disruptions of any of these factors will function as a hit toward disease, and multiple hits may interact and increase the magnitude of impact of these factors. Current research in neuroscience, physiology, and immunology has now shown that the human microbiota, referring to the individual microorganisms, and the microbiome, their collective genomes (5–7), play an important role in health and disease; thus, the microbiota could act as a fourth hit. The gut and vaginal microbiota both are affected by, and can act on, these other factors in Barker's hypothesis, showing integrative functionality in the programming of health and disease (see Figure 1).

The vaginal and gut maternal microbiome has been a recent focus for understanding the health and disease in offspring. In fact, there is evidence of universal vertical microbiota transmission between mother and offspring in all animals (8). There is further interest in the study of the maternal microbiota to elucidate its role in mediating the impact of prenatal factors, such as diet and pharmacological agents, on the offspring. During early development, the infant microbiota may also be vulnerable to shaping from similar postnatal exposures. Thus, one of the emerging questions in developmental psychopathology research is how might the maternal vaginal and gut microbiota interact with prenatal and postnatal environment and host genetics to program brain health and disease later in life.

For the purposes of this review, the programming of brain health refers to physiological, transcriptional, or epigenetic changes induced by prenatal environment or exposure to compounds that lead to altering the homeostatic set point for neurodevelopment during the first 1000 days of human life or the animal model equivalent (9). With a focus on bacteria, we describe how pregnancy-related changes in the composition of the maternal microbiota as well as changes in the microbiota




Figure 1. Four hits programming for brain health and disease. In addition to the three hits posited by Barker (prenatal and postnatal environments and genetic predisposition), we propose that the microbiota acts as a fourth hit. The human microbiota is both influenced by and able to influence the other three factors. Together, these four hits can program for brain health and disease later in life.

during early postnatal life may interact with host genetics and prenatal and early life environment to affect brain health. During fetal development as well as early life, critical periods exist where both negative and positive interventions could alter the trajectory of neurodevelopment and affect brain health later in life (see [Figure 2](#)).




Figure 2. Overlap of critical periods in neurodevelopment with alterations in microbiota diversity. Maternal vaginal and intestinal microbiota diversity increases over the course of pregnancy concurrently with cell birth and migration as well as apoptosis. After birth, microbiota diversity has been measured to increase throughout infancy.

THE MICROBIOTA

The human gut microbiota, consisting of trillions of bacteria and weighing approximately 400 g (10), and their genomes have a profound impact on physiology and behavior. It is the most well-characterized mammalian microbiome and is involved in modulating intestinal barrier permeability as well as gut-brain axis signaling through the vagal nerve, inflammatory molecules, endocrine molecules, and microbial metabolites (11–14). The microbiota-gut-brain axis includes components of the central nervous system, the neuroendocrine and neuro-immune systems, and components of the autonomic and enteric nervous systems (15). However, many of the mechanisms underlying this axis remain unknown.

Other areas of the body that host a bacterial community include the skin, oral cavity, conjunctiva, lungs, biliary duct, and vagina and possibly the placenta. During pregnancy, the communities in many body areas change, but most focus has been placed on the remodeling that takes place in the gut and vagina of the mother. To assess these changes, diversity is used to determine the amount of different types of bacteria as well as their relative abundances within a community.

Human Microbiota During Pregnancy

Gut Microbiota. During pregnancy, the human gut microbiome undergoes dynamic compositional changes that accompany changes in metabolism (16–19). As gestation progresses, intestinal microbiota diversity, and specifically Proteobacteria and Actinobacteria, increase (17). As a result, the gut microbiota composition during the third trimester is significantly different from its composition at the beginning of pregnancy (17). Gut microbial diversity has also been shown to increase postpartum in the mother and vaginally delivered newborns (17,20,21). Interestingly, germ-free mice that received a transplantation of human microbiota from the third trimester of pregnancy displayed symptoms of metabolic syndrome—increased adiposity, inflammation, and reduced glucose tolerance (17–19). These effects may be mediated by

Table 1. Prenatal and Postnatal Impact of Drugs

Drug	Animal/Model	Time of Exposure	Effect	Reference
Proton Pump Inhibitors	Humans	Postnatal	↑ Incidence of <i>Clostridium difficile</i> and other infections Change in gut microbiota composition and diversity ↑ Bacteroidales order, <i>Streptococcus</i> genus ↓ Firmicutes	(42)
Metformin	Humans	Postnatal	Changes in gut microbiota composition and changes in some measures of diversity	(42)
Nonsteroidal Anti-inflammatory Drugs	Humans	Postnatal	Distinct changes in microbiota composition allowing for prediction of changes to different types of drugs involved in overlapping intestinal development pathways	(42)
Opioids	Humans	Postnatal	↑ Alpha diversity, association with compositional distance ↑ Peptostreptococcaceae, ↓ <i>Parasutterella</i> ↑ Energy production and carbohydrate metabolism	(42)
Statins	Humans	Postnatal	Association with beta diversity Species-level changes in composition	(42)
Antipsychotics	Humans	Postnatal	↓ Alpha diversity, altered beta diversity ↑ Lachnospiraceae, ↓ <i>Akkermansia</i>	(42)
Antipsychotics: Risperidone	Human children	Early postnatal	↑ Gut microbiota diversity ↑ Weight gain associated with <i>Clostridium</i> sp., <i>Lactobacillus</i> sp., <i>Ralstonia</i> sp., and Erysipelotrichaceae family	(147)
Antipsychotics: Olanzapine	Female rats	Postnatal	↑ Firmicutes/Bacteroidetes ratio, body weight ↓ Locomotor activity	(45)
	Mice	Postnatal	↑ Body weight Effects dependent on gut microbiota ↑ Firmicutes/Bacteroidetes ratio	(146)
Antiepileptics: Valproic Acid	Mice	Prenatal	↑ Firmicutes/Bacteroidetes ratio in offspring ↑ Cecal butyrate in male offspring Sex-specific impact	(44)
	Mice	Prenatal	↓ Sensitivity to painful stimuli, ↑ sensitivity to nonpainful stimuli in offspring ↓ Prepulse inhibition, social behavior, exploratory activity in offspring ↑ Locomotor and repetitive/stereotypic-like hyperactivity in offspring	(65)
	Rats	Prenatal	↓ Exploratory activity and social interaction in offspring ↑ Interneuronal space in offspring ↑ Neuroinflammation and neuronal disorganization in offspring	(66)
Antibiotics: Succinyl Sulfathiazole	Female rats	Prenatal	↓ Social interaction, exploration in elevated plus maze in offspring ↑ Anxiety in offspring Altered sensorimotor gating in offspring	(43)
Antibiotic Combination: Neomycin, Bacitracin, Pimaricin, and Acetic Acid	Female mice	Prenatal	↓ Exploratory behavior with some persistence into postnatal weeks 7 and 8	(48)
Antibiotics: Penicillin	Mice	Prenatal and early postnatal	↓ Anxiety-like behavior in male offspring ↓ Social behavior in offspring and social novelty ↑ Aggression, reduced social avoidance in offspring Differences in composition and diversity ↑ Brain cytokine expression ↓ Bacteroidetes, ↑ Firmicutes, Proteobacteria	(139)
Antibiotics: Various – Penicillin, Vancomycin, or Penicillin + Vancomycin + Chlortetracycline	Mice	Early postnatal	↑ Adiposity ↑ Firmicutes/Bacteroidetes ratio, Lachnospiraceae Shift in microbiota composition	(140)
Antibiotics: Vancomycin or Cocktail (Pimaricin, Bacitracin, and Neomycin)	Rats	Early postnatal	Visceral hypersensitivity in males ↓ Transient receptor potential in cation channel subfamily V member 1, alpha-2A adrenergic receptor, cholecystokinin B receptor Changes in gut microbiota composition	(141)

Table 1. Continued

Drug	Animal/Model	Time of Exposure	Effect	Reference
Antibiotics	Humans	Early postnatal	↓ Gut microbiota diversity noticeable after first year of life ↓ Gut microbiota stability Transient increase in antibiotic-resistant genes	(137)
	Humans	Early postnatal	↑ Risk of being overweight during later childhood in boys ↑ Adiposity during childhood	(138)
Antidepressants: Ketamine	Mice (chronic social defeat stress)	Postnatal	Attenuated social defeat stress-specific decrease in <i>Butyrivimonas</i> in susceptible mice Attenuated social defeat stress-specific increase in <i>Deltaproteobacteria</i> in susceptible mice	(54)
Antidepressants: Selective Serotonin Reuptake Inhibitors	Humans and rodents	Prenatal	Change in gastrointestinal function, birth weight, metabolism, motor activity/development, and weight later in life	(57)
	Mice	Early postnatal	↓ Exploratory behavior ↑ Anxiety in elevated plus maze ↑ Latency in novelty-suppressed feeding ↓ Shock avoidance	(144)
	Mice	Early postnatal	↓ Exploratory behavior ↑ Depressive behavior in forced swim test and sucrose preference test ↑ Latency in novelty-suppressed feeding ↓ Shock avoidance	(145)
Drugs of Abuse: Ethanol	Mice	Postnatal	↑ Gut permeability, bacterial growth in small intestine ↓ Operational taxonomic units of Firmicutes, ↑ Unknown bacteria, Verrucomicrobia, and Bacteroidetes in the cecum	(59)
	Rats	Postnatal	Changes in colonic microbiota composition (reduction in richness, evenness, Shannon diversity)	(60)
	Mice	Postnatal	↑ <i>Alistepes</i> genus, ↓ <i>Clostridium</i> and alpha diversity Similar changes previously associated with inflammation	(61)
Drugs of Abuse: Cocaine	Mice	Postnatal	Intestinal depletion of bacteria enhances behavioral responses of cocaine Short-chain fatty acids can modulate cocaine responses	(62)

The prenatal and postnatal impact of drugs on the microbiota, the gut, the brain and behavior from studies cited in this review are summarized.

key gut microbial metabolites such as short-chain fatty acids, which were previously shown to be associated with these metabolic changes during pregnancy (16). Because most changes in the gut microbiota occur in the second and third trimesters, perhaps microbiota modulation during these critical time periods can affect neurodevelopment.

Vaginal Microbiota. The human vaginal microbiome consists of multiple stable community state types that destabilize during pregnancy (22,23). It is most abundant in *Lactobacillus* spp. before pregnancy, but this species may be reduced during pregnancy (24,25). However, only some studies found concurrent reduction in overall bacterial diversity during pregnancy (26), while others did not (24,25). In addition, postpartum bacterial diversity was found to be increased (25). Notably, there are also differences in the composition of the vaginal microbiome during pregnancy and postpartum between different ethnic groups, suggesting an interaction between bacterial colonization and host genetics and environment (25). The infant gut microbiome most closely resembles the vaginal microbiome of the mother, which initially colonizes the infant, consistent with patterns of vertical transmission across the animal kingdom (8,27,28).

Pregnancy-Related Changes in the Microbiota: Preclinical Models

Intestinal Microbiota. The intestinal microbiota in mouse models also undergoes compositional changes during

pregnancy, affecting maternal metabolism. While taxa from the Firmicutes and Tenericutes phyla decrease in composition (29,30), there is an increase in the circulation of short-chain fatty acids produced by other members of the microbiota along with tissue-specific transcriptional changes in its receptor, free fatty acid receptor 2 (29). Interestingly, mice deficient in this receptor experienced hyperglycemia and decreased insulin secretion (29).

Vaginal Microbiota. There is also evidence of changes in the vaginal microbiota composition in mice during pregnancy. Diversity of the bacterial community actually increased during early gestation and then decreased early postpartum (31), a finding disparate with human studies (see Vaginal Microbiota subsection of Human Microbiota During Pregnancy section above). In addition, the gut microbiota of the offspring resembled the maternal vaginal microbiota during early postnatal life, and it more resembled that of the maternal gut microbiota at weaning (31).

HOST GENETICS AND THE MICROBIOME

To our knowledge, there are currently no investigations into the interactions between host genetics and the microbiome during pregnancy or into offspring during early postnatal development in preclinical models. There is conflicting evidence regarding the role of host genetics in human and mouse gut microbiota colonization and composition (32–34).

PRENATAL ENVIRONMENT AND THE MICROBIOTA

Although it is commonly thought that initial gut colonization occurs at birth during vaginal delivery, there is some evidence, albeit controversial, of a prenatal microbiome (35,36). Bacteria have been isolated from human meconium, stool that forms in the fetus before birth, and others have found morphological and sequencing evidence of a placental microbiome resembling the oral microbiome (37–40). Although a placental microbiome might not necessarily be transferred to the fetus in utero, it may be involved in fetal development. However, placental colonization might occur during labor due to ruptures in the placental barrier (35). Although bacteria have been isolated from the umbilical cord of infants born by umbilical caesarean section (C-section) (41) many of these studies have been criticized due to a lack of proper controls for contamination, evidence of viable bacteria rather than finding evidence of bacterial genes through sequencing, or inappropriate molecular approaches to detect the bacteria (36). The existence of germ-free mouse models is also further evidence against the prenatal microbiome (36). However, it is clear that a number of factors that can influence the maternal microbiome during pregnancy can either directly or indirectly affect the developing fetal brain.

Drugs and the Microbiota

More than a quarter of pharmaceutical drugs have been shown to affect the composition of the microbiota (42–45), which has been implicated in both the efficacy and side effects of these drugs (46). See Table 1 for more details on the impact of prescription drugs on the microbiota and behavior of offspring. However, there are few studies investigating the prenatal effect of drugs on the microbiota.

Antibiotics. Antibiotics are the most commonly prescribed drug during pregnancy (47) and reduce the diversity and bacterial load of the microbiome through bactericidal or bacteriostatic actions. Succinyl sulfathiazole administration during pregnancy in rats caused sex-specific impairments social behavior of the offspring as well as impairments in prepulse inhibition—an endophenotype of schizophrenia (43). A combination of nonabsorbable antibiotics, administered to mice during pregnancy, reduced exploratory behavior in their offspring (48). This is further validated by germ-free mouse models that display social impairments as well as abnormal microglial development (12,49,50). Microglia have recently been shown to play an important role in neurodevelopment (51). Accordingly, sex-specific aspects of microglial activation that persisted during adulthood in germ-free mice may contribute to the sex biases in psychiatric disorders later in life (50).

Antidepressants. Different classes of antidepressant drugs, including tricyclic antidepressants (52) and selective serotonin inhibitors (53) as well as ketamine (54), which may be used in the future as a new treatment, negatively impact bacterial growth. It is unclear whether their bacteriocidal/bacteriostatic actions affect their efficacy. Prenatal exposure to selective serotonin reuptake inhibitors induces anxiety-like and depressive-like behavior during adulthood (55) in rodents, although the implications of these results for humans are still being discussed (56). In addition,

evidence suggests that these drugs may affect birth weight and motor development in humans as well as in animal models (57); however, potential mechanisms are unknown. Owing to the prevalence of antidepressant administration during pregnancy (58), it is crucial to characterize the interaction among maternal mental health, antidepressant administration, and the offspring's microbiota and mental health.

Drugs of Abuse. The impact of several substances of abuse on the microbiome has not yet been studied. In rodents, ethanol exposure (59–61) and cocaine exposure (62) have shown effects on the gut microbiota composition. In humans, alcohol exposure during pregnancy leads to the development of fetal alcohol spectrum disorders (63). The severity of these effects on the brain may, however, supersede its impact from changes in the maternal microbiota. Future studies should identify whether different gut microbiota compositions are more susceptible to the effects of these substances during pregnancy as well as their impacts on the newborn.

Valproate/Valproic Acid. Valproate is an antiepileptic and mood stabilizer but is also a known teratogen; exposure during gestation results in a 4% risk in autism spectrum disorder (64) and has been able to produce autistic endophenotypes in prenatally exposed rodents (44,65,66). Valproic acid increased the Firmicutes/Bacteroidetes ratio and altered cecal butyrate in the gut microbiota of the male offspring (44). However, these changes likely involve its activity as a histone deacetylase inhibitor (67) rather than its antiepileptic mechanism [which may involve voltage-dependent sodium channels and the gamma-aminobutyric acid system; see (68)]. Other pharmaceutical drugs may also alter the gut microbiota through secondary mechanisms of action.

Maternal Diet and the Microbiome

Unhealthy Diet. Unhealthy diet in humans leads to obesity and poor cardiovascular health, but recent research has also shown its impact on neurocognitive development in both humans and rodents (69). Common rodent models of unhealthy diet include high-fat diet (36%–60% kcal from fat), Western diet (high fat and high sugar), and diets focusing on specific types of fats. Some of these studies have been criticized because improper controls are often selected and may introduce confounds that complicate the interpretation of admittedly intriguing results (70,71). A maternal high-fat diet in mice resulted in higher-fold increases in abundance of bacteria in the gut microbiome of the mother during pregnancy and differences in compositional changes compared with animals fed a control diet (30). In addition, both prenatal and adolescent exposure to a high-fat diet changed the gut metabolome and microbiota composition in mice, rats, and macaques (30,72–74). Prenatal Western diet has also been shown to alter the gut microbiota composition, leading to an increased Firmicutes/Bacteroidetes ratio as well as sex-specific differences in colonic gene expression (75). Although the impact of prenatal trans-fatty acid exposure on the microbiota has not been determined, it can alter hypothalamic function of prenatally exposed rodents during adulthood (76). Furthermore, unhealthy diets during pregnancy in mice also

Microbiota, Brain, and Pre- and Postnatal Programming

lead to sex-specific differences in gene expression (77,78), social deficits (73,78), altered hypothalamic stress response (79), and inflammation (79,80) in the offspring (see Table 2). This suggests that a greater emphasis should be placed on nutrition during pregnancy, although it is unclear whether these changes in maternal microbiota directly affect the stress response in human infants.

Omega-3 Polyunsaturated Fatty Acids. Several nutrients may play a positive role in neurodevelopment and microbiota maturation. A high-fat diet supplemented with omega-3 polyunsaturated fatty acids increased the diversity of microbiota and enriched *Bifidobacterium* at a species level (81). Omega-3 intake during pregnancy regulated the hypothalamic-pituitary-adrenal (HPA) axis activity, shifted the

maternal stress-induced gut microbiota composition to be more similar to an unstressed composition, and conferred resilience to stress later in life; in contrast, a deficit in omega-3 polyunsaturated fatty acids affected the metabolome, impaired communication and social behavior, and increased depressive-like behavior (82–85).

Prebiotics. Prebiotics promote the growth of beneficial bacteria and include indigestible fibers that are fermented by colonic bacteria to produce short-chain fatty acids and provide a health benefit, although their effects on neurodevelopment have not been well studied (86). Administration of the prebiotics galactooligosaccharide and inulin in mice reduced immune activation and intestinal permeability in offspring through gut microbiota modulation (87). The prenatal administration of

Table 2. Impact of Unhealthy Maternal or Adolescent Diet

Diet	Model	Treatment Start Time	Effect of Diet	Reference
Maternal High-Fat Diet (36% kcal From Fat)	Nonhuman primate	Before mating	↓ Spirochetes (<i>Treponema</i> spp. and <i>Sporobacter termitidis</i>), which are involved in carbohydrate metabolism ↓ <i>Campylobacter</i> in mother and offspring	(74)
Western Diet	Mice	4 weeks old	Compositional gut microbiota changes ↑ Firmicutes/Bacteroidetes ratio ↑ Fat when microbiota transplanted into germ-free mice	(149)
Maternal Western Diet	Mice	6 weeks before mating	↑ Firmicutes/Bacteroidetes ratio ↓ Sex-specific differences in colonic gene expression involved in overlapping intestinal development pathways	(75)
	Neonatal mice	2 weeks before mating	Toll-like receptor 2/4-mediated toxicity in neonate from breast milk	(80)
High-Fat Diet (45% kcal From Fat)	Male rats	4 weeks old	↑ Verrucomicrobia and Desulfovibrionaceae ↑ Energy production and carbohydrate metabolism	(72)
	Mouse	Postweaning	↑ Adrenal gland weight in females, ↓ adrenal gland weight in males ↑ Plasma corticosterone, glucocorticoid receptor levels in hypothalamus in females Sexually dimorphic effects in behavioral tests	(150)
	Male mice	Binge eating during adolescence	↑ Anxiety, cocaine self-administration during adulthood ↑ Corticosterone during withdrawal ↓ CB1, MOR expression in nucleus accumbens ↑ GHSR in ventral tegmental area	(152)
Maternal High-Fat Diet (60% kcal From Fat)	Male mice (7–12 weeks)	8 weeks before mating	↑ Social deficits mediated by microbiota composition changes ↓ Microbiota diversity in mother and offspring	(73)
	Mouse	12- to 14-week diet before mating	Sex-specific differences in gene expression and pathways	(77)
	Mouse	7 weeks before mating	↑ Nurr77 expression in males ↓ Exploration for novel object recognition in males	(78)
	Adult mice	Start of pregnancy	Abnormal hypothalamic gene expression No response to lipopolysaccharide Impaired poststress response	(79)
High-Fat Diet (60% kcal From Fat)	Male rats	Juvenile	↓ Spatial memory ↑ Response to lipopolysaccharide-induced cytokines in hippocampus	(151)
High Trans-fatty Acids	Rats	Gestational exposure	↑ Serum endotoxins, hypothalamic cytokines during adulthood ↑ Hypothalamic inflammation via NFκBp65 during adulthood	(76)

The effects of unhealthy prenatal and postnatal diets on the microbiota, gut, brain, and behavior from studies cited in this review are summarized. CB1, cannabinoid receptor 1; GHSR, growth hormone secretagogue receptor; MOR, mu-opioid receptor.

caprine milk oligosaccharide in mice has been shown to increase Bifidobacteria and butyric acid in the offspring colon (88). Interestingly, the addition of inulin to a mouse maternal high-fat diet abrogated the negative metabolic effects of the high-fat diet on offspring (89).

Probiotics. Probiotics are beneficial strains of bacteria that confer a health benefit to the host. Administration during pregnancy in humans can reduce the risk of atopy but not of other immunity-related diseases such as asthma (90,91); however, many current supplements lack proof of effectiveness (92). Although there is a lack of research on the prenatal impact of probiotics within our current scope, several studies have investigated the effects of postnatal administration of probiotics for improving psychiatric symptoms (i.e., anxiety or autism spectrum disorder-like; see Probiotics subsection of Postnatal Diet and the Microbiota section below). It is evident that more animal and human research must be conducted to determine the impact of prenatal probiotics on the maternal and offspring microbiota.

Maternal Stress

Maternal stress is modulated by the HPA axis and has been shown to affect this axis in the offspring. In offspring, maternal stress has increased serum levels of corticosterone, increased anxiety, increased social impairment, and altered the resilience of different strains of rodents (93–97). Maternal stress can also alter the gut and vaginal microbiota during pregnancy, decreasing diversity of maternal gut microbiota as well as dysregulating glucose metabolism in mice (31). Varied prenatal stress interrupted normal pregnancy-related compositional changes in the mouse vaginal microbiota and also altered the protein content in the vaginal mucosa, which may have contributed to altered abundance of Firmicutes and Bacteroidetes in their offspring's gut microbiota (98). After pregnant mice experienced restraint stress specifically, female offspring showed more anxiety-like behaviors as well as decreases in brain-derived neurotrophic factor in the placenta concurrent with later decreases of its expression in the adult amygdala (99).

In humans, ongoing maternal stress (inclusive of prenatal and postnatal stress) has been associated with mental health problems in adult offspring (100) as well as with influencing the development of the offspring microbiota over the first 110 days after birth (101). However, the mechanism by which the negative impact of maternal stress is transmitted to the offspring's microbiota is unknown, although immunoglobulin A-mediated immunity could be implicated (102).

Maternal Immune Activation

The maternal immune activation model is based on the controversial idea that maternal infections during pregnancy can affect psychiatric outcomes in the children (103). A rodent model of maternal immune activation commonly administers viral mimetic polyinosinic:polycytidylic acid or bacterial lipopolysaccharide to produce psychiatric endophenotypes in offspring. Specifically, the viral mimetic polyinosinic:polycytidylic acid administered at embryonic day 12.5 in mice altered the gut microbiota composition and increased gut leakiness by decreasing claudin expression while also elevating intestinal cytokine levels, including interleukin 6, in offspring (104). This model also elevated the

bacterial production of, 4-ethylphenylsulfate, which induced anxiety-like symptoms in wild-type mice (104). However, there is heterogeneity in the time of administration of the viral mimetic polyinosinic:polycytidylic acid, which can lead to the development of different biomarkers or behaviors common to different disorders, including schizophrenia (105,106) and autism spectrum disorder-like behaviors (107) in mice and inconsistent depression-like endophenotypes in rats and mice (108).

Although there exists no current research of its impact on the microbiota, the bacterial mimetic lipopolysaccharide can also induce behavioral phenotypes for anxiety, depression, and autism spectrum disorder in offspring (109,110); impairments in hippocampal development and neurogenesis (111,112); and increased postnatal inflammation (110). In mice, lipopolysaccharide-induced maternal inflammation caused placental damage and fetal intestinal injuries that persisted during adulthood (113); further studies should focus on the potential of a placental microbiome to mediate these interactions.

C-SECTION

There are now a large number of studies showing a marked effect of mode of delivery on the gut microbiome (20,27,114–126), although some studies have found less of an influence than others (127). Infants born via C-section had a gut microbiota more similar to the maternal skin microbiota than the vagina (20,27,114,115), delayed *Bacteroides* colonization (114,115), delayed *Lactobacillus* colonization (27), lower circulating chemokines (20), and a higher risk of vertical obesity transmission from their mother (126). Similar to humans, C-section mice initially had a significantly different gut microbiota from their mother (128). In addition, they had significant differences in the abundance of bacteria in the orders Clostridiales and Bacteroidales as well as lower systemic interleukin 10 expression (128). Although restoration of the microbiota in C-section infants using a vaginal swab has been piloted, the long-term consequences of such an intervention have yet to be studied (129). Alterations in the microbiota induced by C-section may play a functional role in predisposing such infants to a greater relative risk of allergy, asthma, obesity, and type 1 diabetes (126,130,131). The relative contribution of such disturbances to brain health is less clear, although epidemiology and animal studies are beginning to unlock some clear links (132–136).

POSTNATAL ENVIRONMENT AND THE MICROBIOTA

Postnatal Drugs and the Microbiota

Few studies have assessed the impact of postnatal drugs on the microbiota and behavior (see Table 1); this section summarizes the existing literature.

Antibiotics. Exposure to antibiotics within the first 3 years of life in humans decreased microbiome stability and diversity while it transiently increased transcription of antibiotic-resistant genes (137) and increased adiposity in male individuals during childhood (138). Moreover, neonatal exposure to antibiotics in rodents has been shown to alter the microbiota, brain inflammation, and behavior (139); contribute to obesity (140); and increase visceral pain receptors and sensitivity (141). More research must be conducted to elucidate the




Figure 3. Prenatal and postnatal factors contribute to sex-specific programming of health later in life. The gut and vaginal microbiome of the mother is altered by diet, drugs, infection, and stress. If birth occurs through vaginal delivery, the maternal vaginal microbiota colonizes the newborn's gut. If caesarean delivery occurs, the maternal skin microbiome colonizes the infant gut. Early in life, diet, drugs, and stress can also affect the infant microbiota composition. The microbiota is important for sex-specific microglial maturation, which is responsible for the neuroinflammatory response. The hypothalamic-pituitary-adrenal (HPA) axis not only is influenced by but also can influence the gut microbiota. Normally, it can attenuate the inflammatory response from the microglia. These four hits could affect this axis and affect brain health later in life. There are also other factors during adulthood that can affect brain health and disease (i.e., environment, disease, stress, and sex).

effects of common prescriptions of early-life antibiotics on brain health later in life.

Antidepressants. In humans, antidepressants have been shown to transfer through breast milk, and some can reach a clinically significant concentration in the infant's serum (142), although their effects on the infant have not yet been established (143). In rodents, early postnatal exposure to selective serotonin inhibitors alters their social behavior and anxiety-like and depressive-like behaviors (144,145). To our knowledge, there are no studies investigating the impact of early exposure to these drugs on the gut microbiota composition.

Antipsychotics. Similarly, the effects of antipsychotics during breastfeeding—and whether they are able to transfer to the infant in a clinically significant concentration—are currently unknown. However, antipsychotics have been shown to affect the microbiota in rats and adolescent children, leading to weight gain (45,146,147). It is unclear whether similar effects on the microbiota occur as a result of antipsychotic transfer through breast milk and whether this interaction affects infant neurodevelopment.

Postnatal Diet and the Microbiota

The stability and composition of the early postnatal gut microbiota community is strongly dependent on diet, evident in the microbiota differences between breastfed and formula-fed

infants (148). There are now more studies examining the impact of both healthy and unhealthy diets early in life.

Unhealthy Diet. An unhealthy diet commencing during early postnatal life also alters the microbiota composition (149) and results in different behavioral and inflammatory phenotypes (see Table 2). A high-fat diet after weaning and during adolescence altered the HPA axis in female rats and impaired hippocampal memory and increased hippocampal lipopolysaccharide-induced cytokine response in male rats (150,151). Meanwhile, high-fat diet bingeing during adolescence in mice increased anxiety and cocaine self-administration during adulthood (152).

Prebiotics. Few published studies have assessed the impact of early life prebiotics administration in humans on psychiatric outcomes but have shown effects in reducing the risk of atopy, an autoimmune disease (153). In juvenile rats, inulin administered for 2 weeks increased basal corticosterone levels and elevated corticosterone after an anxiety-inducing stimulus, but the animals showed less anxiety in behavioral tests (154).

Probiotics. In humans, probiotics may reduce the risk of depression (155) and autism (156,157). In rodents, *Lactobacillus* administration had effects similar to those of inulin, reducing anxiety through the HPA axis (154). Interestingly,

when combined with inulin, it did not affect the corticosterone levels and increased serotonin 1A receptor messenger RNA in the hippocampus, a receptor associated with anxiety and depression (154). Notably, *Lactobacillus rhamnosus* and *Lactobacillus helveticus* administration in stressed infant rats (postnatal days 2–14) had a protective effect on fear conditioning memory and relapse after extinction early in life (158). Some strains, such as *Bifidobacterium longum* and *Bifidobacterium breve*, can mediate anxiolytic and antidepressive behaviors in preclinical rodent models (159). In mice, *B. breve* combined with prebiotics early in life exerted protection from the negative metabolic effects of a Western diet (160), showing promise for coadministration of prebiotics and probiotics.

Postnatal Stress and the Microbiota

Early postnatal stress affects the HPA axis and contributes to the programming of brain health later in life (161). Different types of early postnatal stress (social isolation and maternal separation) alter the gut microbiota composition and metabolism in rats (162–164) and their inflammatory profiles (163,164). Social isolation also impaired memory and learning in rats (163). Finally, germ-free mice were more vulnerable to restraint stress—resulting in higher adrenocorticotrophic hormone and corticosterone in plasma (165,166), a reduction in glucocorticoid receptor messenger RNA, and an increased stress response (165). Remarkably, these effects were rescued with microbiota transplantation during adolescence but not during adulthood (165).

SEX-SPECIFIC PROGRAMMING OF PSYCHIATRIC DISORDERS LATER IN LIFE

Many psychiatric disorders differ among the sexes in terms of prevalence or onset—including autism (167), mood disorders (168), anxiety disorders (168), and schizophrenia (169). The underlying biological basis of these sex differences is still unknown. Because microbiota-related alterations have shown sex-specific effects after exposure to prenatal or postnatal environmental stimuli (44,75,99,147), they may play a role in the sex-specific programming of health later in life.

The gut microbiota and its metabolites influence the development of the microglia, and in its absence development is disrupted in a sex-specific manner (12,49,50). Abnormal microglial phenotypes were rescued with short-chain fatty acids produced by certain members of the microbiota (12). The microglia are involved in synaptic pruning and maturation during neurodevelopment [see review by Salter and Beggs (170)].

Other hits that disrupt the microbiota could prevent it from reaching a healthy homeostasis (171,172) and result in increased inflammation. In fact, many psychiatric disorders may involve the dysregulation of the immune system—including mood disorders (173–175), schizophrenia (176,177), and autism spectrum disorder (178,179). Alterations of the microbiota through stress and other prenatal or postnatal factors affected the programming of the HPA axis in a sex-specific manner (82,83,150,152).

Because early programming of the HPA axis and immune system can be affected in a sex-specific manner by changes in maternal/offspring microbiota composition, a microbiota hit could program health and disease later in life.

CONCLUSIONS

Several decades after Barker first proposed the role of prenatal factors in health and disease later in life, his ideas have continued to be investigated and expanded on. The microbiota community transmitted to the offspring may have plasticity—like other traits described in the Barker hypothesis. Dynamic changes in the maternal microbiota and early offspring microbiota would occur owing to the prenatal and postnatal environments, respectively, as well as host genetics. Although these changes may be beneficial for immediate survival, they may result in immediate and later physiological and behavioral consequences. The microbiota act as a fourth hit that interacts with the other factors to program for brain health and disease later in life. The microbiota is involved in mediating the effects of both postnatal and prenatal diets, drug exposure, and stress (see Figure 3).

This field has potential for the development of new bioactives and dietary/lifestyle interventions that would modulate the microbiota during pregnancy to reduce the risk of disease later in life. Moreover, microbiome interventions may also influence maternal mental health, as recently seen with a placebo-controlled trial of *L. rhamnosus* during pregnancy and lactation (180). However, there are few studies directly assessing the impact of changes to the microbiota composition early in life on brain health outcomes during adulthood. Further research should focus on identifying the interactions among these four hits as well as critical periods for microbiota modulation to support early neurodevelopment and brain health later in life. As this review shows, certain bacterial species are modified in response to developmental stimuli; however, future work is needed to understand whether there is any convergence to such findings and to understand their functional implications.

ACKNOWLEDGMENTS AND DISCLOSURES

The APC Microbiome Institute is funded by Science Foundation Ireland (SFI) through the Irish Government's National Development Plan. JFC, TGD, and CS are supported by SFI (Grant 12/RC/2273).

We acknowledge Dr. Caitlin S. Cowan for advice with figure design.

The APC Microbiome Institute has conducted research funded by many pharmaceutical and food companies. GC has been invited to a meeting organized by Janssen. TGD has been an invited speaker at meetings organized by Servier, Lundbeck, Janssen, and AstraZeneca and has received research funding from Mead Johnson, Cremo, Suntory Wellness, Nutricia, and 4D Pharma. JFC has been an invited speaker at meetings organized by Mead Johnson, Yakult, Alkermes, and Janssen and has received research funding from Mead Johnson, Cremo, Suntory Wellness, Nutricia, Dupont, and 4D Pharma. MGC, SS, SMO, OFO, and CS report no biomedical financial interests or potential conflicts of interest.

ARTICLE INFORMATION

From APC Microbiome Ireland (MGC, SS, SMO, OFO, GC, CS, TGD, JFC), Department of Anatomy and Neuroscience (MGC, SS, SMO, OFO, JFC), Department of Psychiatry and Neurobehavioural Science (GC, TGD), Irish Centre for Fetal and Neonatal Translational Research and Cork University Maternity Hospital (GC, CS), University College Cork, and Teagasc Food Research Centre (CS), Moorepark, Fermoy, Cork, Ireland.

MGC and SS contributed equally to this work.

Address correspondence to John F. Cryan, Ph.D., Professor and Chair, Department of Anatomy and Neuroscience, 386 Western Gateway Building, University College Cork, Cork, Ireland; E-mail: j.cryan@ucc.ie.

Received Mar 15, 2018; revised May 29, 2018; accepted Jun 18, 2018.

REFERENCES

- Barker DJ (1990): The fetal and infant origins of adult disease. *BMJ* 301:1111.
- Barker DJ, Winter PD, Osmond C, Margetts B, Simmonds SJ (1989): Weight in infancy and death from ischaemic heart disease. *Lancet* 2:577–580.
- Barker DJ, Gluckman PD, Godfrey KM, Harding JE, Owens JA, Robinson JS (1993): Fetal nutrition and cardiovascular disease in adult life. *Lancet* 341:938–941.
- Barker DJ (2004): The developmental origins of adult disease. *J Am Coll Nutr* 23:588S–595S.
- Lederberg J, McCray AT (2001): 'Ome sweet 'omics—A genealogical treasury of words. *The Scientist* 17:1874228.
- Dave M, Higgins PD, Middha S, Rioux KP (2012): The human gut microbiome: Current knowledge, challenges, and future directions. *Transl Res* 160:246–257.
- Peterson J, Garges S, Giovanni M, McInnes P, Wang L, Schloss JA, et al. (2009): The NIH Human Microbiome Project. *Genome Res* 19:2317–2323.
- Funkhouser LJ, Bordenstein SR (2013): Mom knows best: The universality of maternal microbial transmission. *PLoS Biol* 11:e1001631.
- Enos MK, Burton JP, Dols J, Buhulata S, Changanlucha J, Reid G (2013): Probiotics and nutrients for the first 1000 days of life in the developing world. *Benef Microbes* 4:3–16.
- Sender R, Fuchs S, Milo R (2016): Revised estimates for the number of human and bacteria cells in the body. *PLoS Biol* 14:e1002533.
- de Weerth C (2017): Do bacteria shape our development? Crosstalk between intestinal microbiota and HPA axis. *Neurosci Biobehav Rev* 83:458–471.
- Erny D, Hrabě de Angelis AL, Jaitin D, Wieghofer P, Staszewski O, David E, et al. (2015): Host microbiota constantly control maturation and function of microglia in the CNS. *Nat Neurosci* 18:965–977.
- Bravo JA, Forsythe P, Chew MV, Escaravage E, Savignac HM, Dinan TG, et al. (2011): Ingestion of *Lactobacillus* strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proc Natl Acad Sci U S A* 108:16050–16055.
- König J, Wells J, Cani PD, García-Ródenas CL, MacDonald T, Mercenier A, et al. (2016): Human intestinal barrier function in health and disease. *Clin Transl Gastroenterol* 7:e196.
- Dinan TG, Cryan JF (2012): Regulation of the stress response by the gut microbiota: Implications for psychoneuroendocrinology. *Psychoneuroendocrinology* 37:1369–1378.
- Priyadarshini M, Thomas A, Reisetter AC, Scholtens DM, Wolever TM, Josefson JL, et al. (2014): Maternal short-chain fatty acids are associated with metabolic parameters in mothers and newborns. *Transl Res* 164:153–157.
- Koren O, Goodrich JK, Cullender TC, Spor A, Laitinen K, Bäckhed HK, et al. (2012): Host remodeling of the gut microbiome and metabolic changes during pregnancy. *Cell* 150:470–480.
- Santacruz A, Collado MC, García-Valdés L, Segura MT, Martín-Lagos JA, Anjos T, et al. (2010): Gut microbiota composition is associated with body weight, weight gain, and biochemical parameters in pregnant women. *Br J Nutr* 104:83–92.
- Collado MC, Isolauri E, Laitinen K, Salminen S (2008): Distinct composition of gut microbiota during pregnancy in overweight and normal-weight women. *Am J Clin Nutr* 88:894–899.
- Jakobsson HE, Abrahamsson TR, Jenmalm MC, Harris K, Quince C, Jernberg C, et al. (2014): Decreased gut microbiota diversity, delayed Bacteroidetes colonisation and reduced Th1 responses in infants delivered by caesarean section. *Gut* 63:559–566.
- Kurokawa K, Itoh T, Kuwahara T, Oshima K, Toh H, Toyoda A, et al. (2007): Comparative metagenomics revealed commonly enriched gene sets in human gut microbiomes. *DNA Res* 14:169–181.
- Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Nikita L, et al. (2014): The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome* 2:4.
- Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, et al. (2011): Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci U S A* 108(Suppl 1):4680–4687.
- DiGiulio DB, Callahan BJ, McMurdie PJ, Costello EK, Lyell DJ, Robaczewska A, et al. (2015): Temporal and spatial variation of the human microbiota during pregnancy. *Proc Natl Acad Sci U S A* 112:11060–11065.
- MacIntyre DA, Chandiramani M, Lee YS, Kindinger L, Smith A, Angelopoulos N, et al. (2015): The vaginal microbiome during pregnancy and the postpartum period in a European population. *Sci Rep* 5:8988.
- Aagaard K, Riehle K, Ma J, Segata N, Mistretta TA, Coarfa C, et al. (2012): A metagenomic approach to characterization of the vaginal microbiome signature in pregnancy. *PLoS One* 7:e36466.
- Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, Fierer N, et al. (2010): Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci U S A* 107:11971–11975.
- Ley RE, Peterson DA, Gordon JI (2006): Ecological and evolutionary forces shaping microbial diversity in the human intestine. *Cell* 124:837–848.
- Fuller M, Priyadarshini M, Gibbons SM, Angueira AR, Brodsky M, Hayes MG, et al. (2015): The short-chain fatty acid receptor, FFA2, contributes to gestational glucose homeostasis. *Am J Physiol Endocrinol Metab* 309:E840–E851.
- Gohir W, Whelan FJ, Surette MG, Moore C, Schertzer JD, Sloboda DM (2015): Pregnancy-related changes in the maternal gut microbiota are dependent upon the mother's periconceptual diet. *Gut Microbes* 6:310–320.
- Jašarević E, Howard CD, Misić AM, Beiting DP, Bale TL (2017): Stress during pregnancy alters temporal and spatial dynamics of the maternal and offspring microbiome in a sex-specific manner. *Sci Rep* 7:44182.
- Goodrich JK, Davenport ER, Clark AG, Ley RE (2017): The relationship between the human genome and microbiome comes into view. *Annu Rev Genet* 51:413–433.
- Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, et al. (2018): Environment dominates over host genetics in shaping human gut microbiota. *Nature* 555:210–215.
- Chung H, Pamp SJ, Hill JA, Surana NK, Edelman SM, Troy EB, et al. (2012): Gut immune maturation depends on colonization with a host-specific microbiota. *Cell* 149:1578–1593.
- Willyard C (2018): Could baby's first bacteria take root before birth? *Nature* 553:264–266.
- Perez-Muñoz ME, Arrieta MC, Ramer-Tait AE, Walter J (2017): A critical assessment of the “sterile womb” and “in utero colonization” hypotheses: Implications for research on the pioneer infant microbiome. *Microbiome* 5:48.
- Jiménez E, Marín ML, Martín R, Odriozola JM, Olivares M, Xaus J, et al. (2008): Is meconium from healthy newborns actually sterile? *Res Microbiol* 159:187–193.
- Ardissone AN, de la Cruz DM, Davis-Richardson AG, Rechcigl KT, Li N, Drew JC, et al. (2014): Meconium microbiome analysis identifies bacteria correlated with premature birth. *PLoS One* 9:e90784.
- Stout MJ, Conlon B, Landeau M, Lee I, Bower C, Zhao Q, et al. (2013): Identification of intracellular bacteria in the basal plate of the human placenta in term and preterm gestations. *Am J Obstet Gynecol* 208:226.e1–226.e7.
- Aagaard K, Ma J, Antony KM, Ganu R, Petrosino J, Versalovic J (2014): The placenta harbors a unique microbiome. *Sci Transl Med* 6:237ra265.

41. Jiménez E, Fernández L, Marín ML, Martín R, Odriozola JM, Nueno-Palop C, *et al.* (2005): Isolation of commensal bacteria from umbilical cord blood of healthy neonates born by cesarean section. *Curr Microbiol* 51:270–274.
42. Le Bastard Q, Al-Ghalith GA, Grégoire M, Chapelet G, Javaudin F, Dailly E, *et al.* (2018): Systematic review: Human gut dysbiosis induced by non-antibiotic prescription medications. *Aliment Pharmacol Ther* 47:332–345.
43. Degroote S, Hunting DJ, Baccarelli AA, Takser L (2016): Maternal gut and fetal brain connection: Increased anxiety and reduced social interactions in Wistar rat offspring following peri-conceptual antibiotic exposure. *Prog Neuropsychopharmacol Biol Psychiatry* 71:76–82.
44. de Theije CG, Wopereis H, Ramadan M, van Eijndhoven T, Lambert J, Knol J, *et al.* (2014): Altered gut microbiota and activity in a murine model of autism spectrum disorders. *Brain Behav Immun* 37:197–206.
45. Davey KJ, Cotter PD, O'Sullivan O, Crispie F, Dinan TG, Cryan JF, *et al.* (2013): Antipsychotics and the gut microbiome: Olanzapine-induced metabolic dysfunction is attenuated by antibiotic administration in the rat. *Transl Psychiatry* 3:e309.
46. Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, *et al.* (2018): Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature* 555:623–628.
47. Kuperman AA, Koren O (2016): Antibiotic use during pregnancy: How bad is it? *BMC Med* 14:91.
48. Tochtiani S, Ikeno T, Ito T, Sakurai A, Yamauchi T, Matsuzaki H (2016): Administration of non-absorbable antibiotics to pregnant mice to perturb the maternal gut microbiota is associated with alterations in offspring behavior. *PLoS One* 11:e0138293.
49. Desbonnet L, Clarke G, Shanahan F, Dinan TG, Cryan JF (2014): Microbiota is essential for social development in the mouse. *Mol Psychiatry* 19:146–148.
50. Thion MS, Low D, Silvin A, Chen J, Grisel P, Schulte-Schrepping J, *et al.* (2018): Microbiome influences prenatal and adult microglia in a sex-specific manner. *Cell* 172:500–516.e516.
51. Biliomorja PM, Stevens B (2015): Microglia function during brain development: New insights from animal models. *Brain Res* 1617:7–17.
52. Csizsar K, Molnar J (1992): Mechanism of action of tricyclic drugs on *Escherichia coli* and *Yersinia enterocolitica* plasmid maintenance and replication. *Anticancer Res* 12:2267–2272.
53. Munoz-Bellido JL, Munoz-Criado S, Garcia-Rodriguez JA (2000): Antimicrobial activity of psychotropic drugs: Selective serotonin reuptake inhibitors. *Int J Antimicrob Agents* 14:177–180.
54. Yang C, Qu Y, Fujita Y, Ren Q, Ma M, Dong C, *et al.* (2017): Possible role of the gut microbiota-brain axis in the antidepressant effects of (R)-ketamine in a social defeat stress model. *Transl Psychiatry* 7:1294.
55. Homberg JR, Schubert D, Gaspar P (2010): New perspectives on the neurodevelopmental effects of SSRIs. *Trends Pharmacol Sci* 31:60–65.
56. Gur TL, Kim DR, Epperson CN (2013): Central nervous system effects of prenatal selective serotonin reuptake inhibitors: Sensing the signal through the noise. *Psychopharmacology (Berl)* 227:567–582.
57. Hutchison SM, Mässe LC, Pawluski JL, Oberlander TF (2018): Perinatal selective serotonin reuptake inhibitor (SSRI) effects on body weight at birth and beyond: A review of animal and human studies. *Reprod Toxicol* 77:109–121.
58. Koren G, Nordeng H (2012): Antidepressant use during pregnancy: The benefit–risk ratio. *Am J Obstet Gynecol* 207:157–163.
59. Yan AW, Fouts DE, Brandl J, Stärkel P, Torralba M, Schott E, *et al.* (2011): Enteric dysbiosis associated with a mouse model of alcoholic liver disease. *Hepatology* 53:96–105.
60. Mutlu E, Keshavarzian A, Engen P, Forsyth CB, Sikaroodi M, Gillevet P (2009): Intestinal dysbiosis: A possible mechanism of alcohol-induced endotoxemia and alcoholic steatohepatitis in rats. *Alcohol Clin Exp Res* 33:1836–1846.
61. Peterson VL, Jury NJ, Cabrera-Rubio R, Draper LA, Crispie F, Cotter PD, *et al.* (2017): Drunk bugs: Chronic vapour alcohol exposure induces marked changes in the gut microbiome in mice. *Behav Brain Res* 323:172–176.
62. Kiraly DD, Walker DM, Calipari ES, Labonte B, Issler O, Pena CJ, *et al.* (2016): Alterations of the host microbiome affect behavioral responses to cocaine. *Sci Rep* 6:35455.
63. Sokol RJ, Delaney-Black V, Nordstrom B (2003): Fetal alcohol spectrum disorder. *JAMA* 290:2996–2999.
64. Christensen J, Grønberg TK, Sørensen MJ, Schendel D, Parner ET, Pedersen LH, *et al.* (2013): Prenatal valproate exposure and risk of autism spectrum disorders and childhood autism. *JAMA* 309:1696–1703.
65. Schneider T, Przewłocki R (2005): Behavioral alterations in rats prenatally exposed to valproic acid: Animal model of autism. *Neuropharmacology* 30:80–89.
66. Codagnone MG, Podestá MF, Uccelli NA, Reinés A (2015): Differential local connectivity and neuroinflammation profiles in the medial prefrontal cortex and hippocampus in the valproic acid rat model of autism. *Dev Neurosci* 37:215–231.
67. Phiel CJ, Zhang F, Huang EY, Guenther MG, Lazar MA, Klein PS (2001): Histone deacetylase is a direct target of valproic acid, a potent anticonvulsant, mood stabilizer, and teratogen. *J Biol Chem* 276:36734–36741.
68. Löscher W (1999): Valproate: A reappraisal of its pharmacodynamic properties and mechanisms of action. *Prog Neurobiol* 58:31–59.
69. Monk C, Georgieff MK, Osterholm EA (2013): Research review: Maternal prenatal distress and poor nutrition—Mutually influencing risk factors affecting infant neurocognitive development. *J Child Psychol Psychiatry* 54:115–130.
70. Almeida-Suhett CP, Scott JM, Graham A, Chen Y, Deuster PA (2017): Control diet in a high-fat diet study in mice: Regular chow and purified low-fat diet have similar effects on phenotypic, metabolic, and behavioral outcomes [published online ahead of print Jul 19]. *Nutr Neurosci*.
71. Pellizzon MA, Ricci MR (2018): The common use of improper control diets in diet-induced metabolic disease research confounds data interpretation: The fiber factor. *Nutr Metab (Lond)* 15:3.
72. Oberbach A, Haange SB, Schlichting N, Heinrich M, Lehmann S, Till H, *et al.* (2017): Metabolic in vivo labeling highlights differences of metabolically active microbes from the mucosal gastrointestinal microbiome between high-fat and normal chow diet. *J Proteome Res* 16:1593–1604.
73. Buffington SA, Di Prisco GV, Auchtung TA, Ajami NJ, Petrosino JF, Costa-Mattioli M (2016): Microbial reconstitution reverses maternal diet-induced social and synaptic deficits in offspring. *Cell* 165:1762–1775.
74. Ma J, Prince AL, Bader D, Hu M, Ganu R, Baquero K, *et al.* (2014): High-fat maternal diet during pregnancy persistently alters the offspring microbiome in a primate model. *Nat Commun* 5:3889.
75. Steegenga WT, Mischke M, Lute C, Boekschoten MV, Lendvai A, Pruis MG, *et al.* (2017): Maternal exposure to a Western-style diet causes differences in intestinal microbiota composition and gene expression of suckling mouse pups. *Mol Nutr Food Res* Jan;61(1).
76. Pimentel GD, Lira FS, Rosa JC, Oliveira JL, Losinskas-Hachul AC, Souza GI, *et al.* (2012): Intake of trans fatty acids during gestation and lactation leads to hypothalamic inflammation via TLR4/NFκBp65 signaling in adult offspring. *J Nutr Biochem* 23:265–271.
77. Edlow AG, Guedj F, Pennings JL, Sverdlov D, Neri C, Bianchi DW (2016): Males are from Mars, and females are from Venus: Sex-specific fetal brain gene expression signatures in a mouse model of maternal diet-induced obesity. *Am J Obstet Gynecol* 214:623.e1–623.e10.
78. Graf AE, Lallier SW, Waidyaratne G, Thompson MD, Tipple TE, Hester ME, *et al.* (2016): Maternal high fat diet exposure is associated with increased hepcidin levels, decreased myelination, and neurobehavioral changes in male offspring. *Brain Behav Immun* 58:369–378.

79. Grissom NM, George R, Reyes TM (2017): The hypothalamic transcriptional response to stress is severely impaired in offspring exposed to adverse nutrition during gestation. *Neuroscience* 342:200–211.
80. Du Y, Yang M, Lee S, Behrendt CL, Hooper LV, Saghatelian A, *et al.* (2012): Maternal Western diet causes inflammatory milk and TLR2/4-dependent neonatal toxicity. *Genes Dev* 26:1306–1311.
81. Patterson E, O'Doherty RM, Murphy EF, Wall R, O'Sullivan O, Nilaweera K, *et al.* (2014): Impact of dietary fatty acids on metabolic activity and host intestinal microbiota composition in C57BL/6J mice. *Br J Nutr* 111:1905–1917.
82. Robertson RC, Seira Oriach C, Murphy K, Moloney GM, Cryan JF, Dinan TG, *et al.* (2017): Omega-3 polyunsaturated fatty acids critically regulate behaviour and gut microbiota development in adolescence and adulthood. *Brain Behav Immun* 59:21–37.
83. Weiser MJ, Wynalda K, Salem N, Butt CM (2015): Dietary DHA during development affects depression-like behaviors and biomarkers that emerge after puberty in adolescent rats. *J Lipid Res* 56:151–166.
84. Robertson RC, Seira Oriach C, Murphy K, Moloney GM, Cryan JF, Dinan TG, *et al.* (2017): Deficiency of essential dietary n-3 PUFA disrupts the caecal microbiome and metabolome in mice. *Br J Nutr* 118:959–970.
85. Pusceddu MM, El Aidy S, Crispie F, O'Sullivan O, Cotter P, Stanton C, *et al.* (2015): N-3 polyunsaturated fatty acids (PUFAs) reverse the impact of early-life stress on the gut microbiota. *PLoS One* 10:e0139721.
86. Gibson GR, Hutkins R, Sanders ME, Prescott SL, Reimer RA, Salminen SJ, *et al.* (2017): Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol* 14:491–502.
87. Bouchaud G, Castan L, Chesné J, Braza F, Aubert P, Neunlist M, *et al.* (2016): Maternal exposure to GOS/inulin mixture prevents food allergies and promotes tolerance in offspring in mice. *Allergy* 71:68–76.
88. Thum C, McNabb WC, Young W, Cookson AL, Roy NC (2016): Prenatal caprine milk oligosaccharide consumption affects the development of mice offspring. *Mol Nutr Food Res* 60:2076–2085.
89. Zou J, Chassaing B, Singh V, Pellizzon M, Ricci M, Fythe MD, *et al.* (2018): Fiber-mediated nourishment of gut microbiota protects against diet-induced obesity by restoring IL-22-mediated colonic health. *Cell Host Microbe* 23:41–53.e44.
90. Elazab N, Mendy A, Gasana J, Vieira ER, Quizon A, Forno E (2013): Probiotic administration in early life, atopy, and asthma: A meta-analysis of clinical trials. *Pediatrics* 132:e666–e676.
91. Azad MB, Coneys JG, Kozyrskyj AL, Field CJ, Ramsey CD, Becker AB, *et al.* (2013): Probiotic supplementation during pregnancy or infancy for the prevention of asthma and wheeze: Systematic review and meta-analysis. *BMJ* 347:f6471.
92. Reid G, Kirjaivanen P (2005): Taking probiotics during pregnancy. Are they useful therapy for mothers and newborns? *Can Fam Physician* 51:1477–1479.
93. Lee YA, Kim YJ, Goto Y (2016): Cognitive and affective alterations by prenatal and postnatal stress interaction. *Physiol Behav* 165:146–153.
94. Rana S, Pugh PC, Jackson N, Clinton SM, Kerman IA (2015): Inborn stress reactivity shapes adult behavioral consequences of early-life maternal separation stress. *Neurosci Lett* 584:146–150.
95. Bale TL (2015): Epigenetic and transgenerational reprogramming of brain development. *Nat Rev Neurosci* 16:332–344.
96. Hiroi R, Carbone DL, Zuloaga DG, Bimonte-Nelson HA, Handa RJ (2016): Sex-dependent programming effects of prenatal glucocorticoid treatment on the developing serotonin system and stress-related behaviors in adulthood. *Neuroscience* 320:43–56.
97. Golubeva AV, Crampton S, Desbonnet L, Edge D, O'Sullivan O, Lomasney KW, *et al.* (2015): Prenatal stress-induced alterations in major physiological systems correlate with gut microbiota composition in adulthood. *Psychoneuroendocrinology* 60:58–74.
98. Jašarević E, Howerton CL, Howard CD, Bale TL (2015): Alterations in the vaginal microbiome by maternal stress are associated with metabolic reprogramming of the offspring gut and brain. *Endocrinology* 156:3265–3276.
99. Gur TL, Shay L, Palkar AV, Fisher S, Varaljay VA, Dowd S, *et al.* (2017): Prenatal stress affects placental cytokines and neurotrophins, commensal microbes, and anxiety-like behavior in adult female offspring. *Brain Behav Immun* 64:50–58.
100. Betts KS, Williams GM, Najman JM, Alati R (2015): The relationship between maternal depressive, anxious, and stress symptoms during pregnancy and adult offspring behavioral and emotional problems. *Depress Anxiety* 32:82–90.
101. Zijlmans MA, Korpela K, Riksen-Walraven JM, de Vos WM, de Weerth C (2015): Maternal prenatal stress is associated with the infant intestinal microbiota. *Psychoneuroendocrinology* 53:233–245.
102. Kang LJ, Koleva PT, Field CJ, Giesbrecht GF, Wine E, Becker AB, *et al.* (2018): Maternal depressive symptoms linked to reduced fecal immunoglobulin A concentrations in infants. *Brain Behav Immun* 68:123–131.
103. Estes ML, McAllister AK (2016): Maternal immune activation: Implications for neuropsychiatric disorders. *Science* 353:772–777.
104. Hsiao EY, McBride SW, Hsien S, Sharon G, Hyde ER, McCue T, *et al.* (2013): Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopmental disorders. *Cell* 155:1451–1463.
105. Juckel G, Manitz MP, Brüne M, Friebe A, Heneka MT, Wolf RJ (2011): Microglial activation in a neuroinflammatory animal model of schizophrenia—A pilot study. *Schizophr Res* 131:96–100.
106. Li Q, Cheung C, Wei R, Hui ES, Feldon J, Meyer U, *et al.* (2009): Prenatal immune challenge is an environmental risk factor for brain and behavior change relevant to schizophrenia: Evidence from MRI in a mouse model. *PLoS One* 4:e6354.
107. Malkova NV, Yu CZ, Hsiao EY, Moore MJ, Patterson PH (2012): Maternal immune activation yields offspring displaying mouse versions of the three core symptoms of autism. *Brain Behav Immun* 26:607–616.
108. Ronovsky M, Berger S, Molz B, Berger A, Pollak DD (2016): Animal models of maternal immune activation in depression research. *Curr Neuropharmacol* 14:688–704.
109. Depino AM (2015): Early prenatal exposure to LPS results in anxiety- and depression-related behaviors in adulthood. *Neuroscience* 299:56–65.
110. Oskvig DB, Elkahloun AG, Johnson KR, Phillips TM, Herkenham M (2012): Maternal immune activation by LPS selectively alters specific gene expression profiles of interneuron migration and oxidative stress in the fetus without triggering a fetal immune response. *Brain Behav Immun* 26:623–634.
111. Escobar M, Crouzin N, Cavalier M, Quentin J, Roussel J, Lanté F, *et al.* (2011): Early, time-dependent disturbances of hippocampal synaptic transmission and plasticity after in utero immune challenge. *Biol Psychiatry* 70:992–999.
112. Romero E, Ali C, Molina-Holgado E, Castellano B, Guaza C, Borrell J (2007): Neurobehavioral and immunological consequences of prenatal immune activation in rats: Influence of antipsychotics. *Neuropsychopharmacology* 32:1791–1804.
113. Fricke EM, Elgin TG, Gong H, Reese J, Gibson-Corley KN, Weiss RM, *et al.* (2018): Lipopolysaccharide-induced maternal inflammation induces direct placental injury without alteration in placental blood flow and induces a secondary fetal intestinal injury that persists into adulthood. *Am J Reprod Immunol* 79:e12816.
114. Mueller NT, Bakacs E, Combellick J, Grigoryan Z, Dominguez-Bello MG (2015): The infant microbiome development: Mom matters. *Trends Mol Med* 21:109–117.

115. Biasucci G, Benenati B, Morelli L, Bessi E, Boehm G (2008): Cesarean delivery may affect the early biodiversity of intestinal bacteria. *J Nutr* 138:1796S–1800S.
116. Bäckhed F, Roswall J, Peng Y, Feng Q, Jia H, Kovatcheva-Datchary P, *et al.* (2015): Dynamics and stabilization of the human gut microbiome during the first year of life. *Cell Host Microbe* 17:690–703.
117. Madan JC, Hoen AG, Lundgren SN, Farzan SF, Cottingham KL, Morrison HG, *et al.* (2016): Association of cesarean delivery and formula supplementation with the intestinal microbiome of 6-week-old infants. *JAMA Pediatr* 170:212–219.
118. Biasucci G, Rubini M, Riboni S, Morelli L, Bessi E, Retetangos C (2010): Mode of delivery affects the bacterial community in the newborn gut. *Early Hum Dev* 86(Suppl 1):13–15.
119. Azad MB, Konya T, Maughan H, Guttman DS, Field CJ, Chari RS, *et al.* (2013): Gut microbiota of healthy Canadian infants: Profiles by mode of delivery and infant diet at 4 months. *CMAJ* 185:385–394.
120. Salminen S, Gibson GR, McCartney AL, Isolauri E (2004): Influence of mode of delivery on gut microbiota composition in seven year old children. *Gut* 53:1388–1389.
121. Brumbaugh DE, Arruda J, Robbins K, Ir D, Santorico SA, Robertson CE, *et al.* (2016): Mode of delivery determines neonatal pharyngeal bacterial composition and early intestinal colonization. *J Pediatr Gastroenterol Nutr* 63:320–328.
122. Dogra S, Sakwinska O, Soh SE, Ngom-Bru C, Brück WM, Berger B, *et al.* (2015): Dynamics of infant gut microbiota are influenced by delivery mode and gestational duration and are associated with subsequent adiposity. *MBio* 6:e02419–14.
123. Grześkowiak Ł, Sales Teixeira TF, Bigonha SM, Lobo G, Salminen S, Ferreira CL (2015): Gut *Bifidobacterium* microbiota in one-month-old Brazilian newborns. *Anaerobe* 35:54–58.
124. Martin R, Makino H, Cetinyurek Yavuz A, Ben-Amor K, Roelofs M, Ishikawa E, *et al.* (2016): Early-life events, including mode of delivery and type of feeding, siblings and gender, shape the developing gut microbiota. *PLoS One* 11:e0158498.
125. Hill CJ, Lynch DB, Murphy K, Ulaszewska M, Jeffery IB, O’Shea CA, *et al.* (2017): Evolution of gut microbiota composition from birth to 24 weeks in the INFANTMET cohort. *Microbiome* 5:4.
126. Tun HM, Bridgman SL, Chari R, Field CJ, Guttman DS, Becker AB, *et al.* (2018): Roles of birth mode and infant gut microbiota in inter-generational transmission of overweight and obesity from mother to offspring. *JAMA Pediatr* 172:368–377.
127. Chu DM, Ma J, Prince AL, Antony KM, Seferovic MD, Aagaard KM (2017): Maturation of the infant microbiome community structure and function across multiple body sites and in relation to mode of delivery. *Nat Med* 23:314–326.
128. Hansen CH, Andersen LS, Krych L, Metzendorf SB, Hasselby JP, Skov S, *et al.* (2014): Mode of delivery shapes gut colonization pattern and modulates regulatory immunity in mice. *J Immunol* 193:1213–1222.
129. Dominguez-Bello MG, De Jesus-Laboy KM, Shen N, Cox LM, Amir A, Gonzalez A, *et al.* (2016): Partial restoration of the microbiota of cesarean-born infants via vaginal microbial transfer. *Nat Med* 22:250–253.
130. Bager P, Wohlfahrt J, Westergaard T (2008): Cesarean delivery and risk of atopy and allergic disease: Meta-analyses. *Clin Exp Allergy* 38:634–642.
131. Cardwell CR, Stene LC, Joner G, Cineke O, Svensson J, Goldacre MJ, *et al.* (2008): Cesarean section is associated with an increased risk of childhood-onset type 1 diabetes mellitus: A meta-analysis of observational studies. *Diabetologia* 51:726–735.
132. O’Neill SM, Curran EA, Dalman C, Kenny LC, Kearney PM, Clarke G, *et al.* (2016): Birth by caesarean section and the risk of adult psychosis: A population-based cohort study. *Schizophr Bull* 42:633–641.
133. Fond G, Bulzacka E, Boyer L, Llorca PM, Godin O, Brunel L, *et al.* (2017): Birth by cesarean section and schizophrenia: Results from the multicenter FACE-SZ data-set. *Eur Arch Psychiatry Clin Neurosci* 267:587–594.
134. Moya-Pérez A, Luczynski P, Renes IB, Wang S, Borre Y, Anthony Ryan C, *et al.* (2017): Intervention strategies for cesarean section-induced alterations in the microbiota-gut-brain axis. *Nutr Rev* 75:225–240.
135. Martínez KA, Devlin JC, Lacher CR, Yin Y, Cai Y, Wang J, *et al.* (2017): Increased weight gain by C-section: Functional significance of the primordial microbiome. *Sci Adv* 3:eaa01874.
136. Curran EA, Kenny LC, Dalman C, Kearney PM, Cryan JF, Dinan TG, *et al.* (2017): Birth by caesarean section and school performance in Swedish adolescents—A population-based study. *BMC Pregnancy Childbirth* 17:121.
137. Yassour M, Vatanen T, Siljander H, Hämäläinen AM, Härkönen T, Ryhänen SJ, *et al.* (2016): Natural history of the infant gut microbiome and impact of antibiotic treatment on bacterial strain diversity and stability. *Sci Transl Med* 8:343ra381.
138. Azad MB, Bridgman SL, Becker AB, Kozyrskyj AL (2014): Infant antibiotic exposure and the development of childhood overweight and central adiposity. *Int J Obes (Lond)* 38:1290–1298.
139. Leclercq S, Mian FM, Staniszc AM, Bindels LB, Cambier E, Ben-Amram H, *et al.* (2017): Low-dose penicillin in early life induces long-term changes in murine gut microbiota, brain cytokines and behavior. *Nat Commun* 8:15062.
140. Cho I, Yamanishi S, Cox L, Methé BA, Zavadil J, Li K, *et al.* (2012): Antibiotics in early life alter the murine colonic microbiome and adiposity. *Nature* 488:621–626.
141. O’Mahony SM, Felice VD, Nally K, Savignac HM, Claesson MJ, Scully P, *et al.* (2014): Disturbance of the gut microbiota in early-life selectively affects visceral pain in adulthood without impacting cognitive or anxiety-related behaviors in male rats. *Neuroscience* 277:885–901.
142. Sachs HC, Drugs CO (2013): The transfer of drugs and therapeutics into human breast milk: An update on selected topics. *Pediatrics* 132:e796–e809.
143. Glover ME, Clinton SM (2016): Of rodents and humans: A comparative review of the neurobehavioral effects of early life SSRI exposure in preclinical and clinical research. *Int J Dev Neurosci* 51:50–72.
144. Ansorge MS, Zhou M, Lira A, Hen R, Gingrich JA (2004): Early-life blockade of the 5-HT transporter alters emotional behavior in adult mice. *Science* 306:879–881.
145. Rebello TJ, Yu Q, Goodfellow NM, Caffrey Cagliostro MK, Teissier A, Morelli E, *et al.* (2014): Postnatal day 2 to 11 constitutes a 5-HT-sensitive period impacting adult mPFC function. *J Neurosci* 34:12379–12393.
146. Morgan AP, Crowley JJ, Nonneman RJ, Quackenbush CR, Miller CN, Ryan AK, *et al.* (2014): The antipsychotic olanzapine interacts with the gut microbiome to cause weight gain in mouse. *PLoS One* 9:e115225.
147. Bahr SM, Tyler BC, Wooldridge N, Butcher BD, Burns TL, Teesch LM, *et al.* (2015): Use of the second-generation antipsychotic, risperidone, and secondary weight gain are associated with an altered gut microbiota in children. *Transl Psychiatry* 5:e652.
148. Penders J, Thijs C, Vink C, Stelma FF, Snijders B, Kummeling I, *et al.* (2006): Factors influencing the composition of the intestinal microbiota in early infancy. *Pediatrics* 118:511–521.
149. Turnbaugh PJ, Bäckhed F, Fulton L, Gordon JI (2008): Diet-induced obesity is linked to marked but reversible alterations in the mouse distal gut microbiome. *Cell Host Microbe* 3:213–223.
150. Boukouvalas G, Antoniou K, Papalexi E, Kitraki E (2008): Post weaning high fat feeding affects rats’ behavior and hypothalamic pituitary adrenal axis at the onset of puberty in a sexually dimorphic manner. *Neuroscience* 153:373–382.
151. Boitard C, Cavaroc A, Sauvart J, Aubert A, Castanon N, Layé S, *et al.* (2014): Impairment of hippocampal-dependent memory induced by juvenile high-fat diet intake is associated with enhanced hippocampal inflammation in rats. *Brain Behav Immun* 40:9–17.
152. Blanco-Gandia MC, Cantacorps L, Aracil-Fernández A, Montagud-Romero S, Aguilar MA, Manzanares J, *et al.* (2017): Effects of bingeing on fat during adolescence on the reinforcing effects of cocaine in adult male mice. *Neuropharmacology* 113:31–44.

Microbiota, Brain, and Pre- and Postnatal Programming

153. Grüber C, van Stuijvenberg M, Mosca F, Moro G, Chirico G, Braegger CP, *et al.* (2010): Reduced occurrence of early atopic dermatitis because of immunoactive prebiotics among low-atopy-risk infants. *J Allergy Clin Immunol* 126:791–797.
154. Barrera-Bugueño C, Realini O, Escobar-Luna J, Sotomayor-Zárate R, Gotteland M, Julio-Pieper M, *et al.* (2017): Anxiogenic effects of a *Lactobacillus*, inulin and the synbiotic on healthy juvenile rats. *Neuroscience* 359:18–29.
155. Huang R, Wang K, Hu J (2016): Effect of probiotics on depression: A systematic review and meta-analysis of randomized controlled trials. *Nutrients* 8:e483.
156. Pärtty A, Kalliomäki M, Wacklin P, Salminen S, Isolauri E (2015): A possible link between early probiotic intervention and the risk of neuropsychiatric disorders later in childhood: A randomized trial. *Pediatr Res* 77:823–828.
157. Gilbert JA, Krajmalnik-Brown R, Porazinska DL, Weiss SJ, Knight R (2013): Toward effective probiotics for autism and other neurodevelopmental disorders. *Cell* 155:1446–1448.
158. Cowan CS, Callaghan BL, Richardson R (2016): The effects of a probiotic formulation (*Lactobacillus rhamnosus* and *L. helveticus*) on developmental trajectories of emotional learning in stressed infant rats. *Transl Psychiatry* 6:e823.
159. Savignac HM, Kiely B, Dinan TG, Cryan JF (2014): Bifidobacteria exert strain-specific effects on stress-related behavior and physiology in BALB/c mice. *Neurogastroenterol Motil* 26:1615–1627.
160. Mischke M, Arora T, Tims S, Engels E, Sommer N, van Limpt K, *et al.* (2018): Specific synbiotics in early life protect against diet-induced obesity in adult mice. *Diabetes Obes Metab* 20:1408–1418.
161. Heim C, Nemeroff CB (2001): The role of childhood trauma in the neurobiology of mood and anxiety disorders: Preclinical and clinical studies. *Biol Psychiatry* 49:1023–1039.
162. Farshim P, Walton G, Chakrabarti B, Givens I, Saddy D, Kitchen I, *et al.* (2016): Maternal weaning modulates emotional behavior and regulates the gut-brain axis. *Sci Rep* 6:21958.
163. Doherty FD, O'Mahony SM, Peterson VL, O'Sullivan O, Crispie F, Cotter PD, *et al.* (2018): Post-weaning social isolation of rats leads to long-term disruption of the gut microbiota-immune-brain axis. *Brain Behav Immun* 68:261–273.
164. O'Mahony SM, Marchesi JR, Scully P, Codling C, Ceolho AM, Quigley EM, *et al.* (2009): Early life stress alters behavior, immunity, and microbiota in rats: Implications for irritable bowel syndrome and psychiatric illnesses. *Biol Psychiatry* 65:263–267.
165. Sudo N, Chida Y, Aiba Y, Sonoda J, Oyama N, Yu XN, *et al.* (2004): Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. *J Physiol* 558:263–275.
166. Clarke G, Grenham S, Scully P, Fitzgerald P, Moloney RD, Shanahan F, *et al.* (2013): The microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. *Mol Psychiatry* 18:666–673.
167. Yeargin-Allsopp M, Rice C, Karapurkar T, Doernberg N, Boyle C, Murphy C (2003): Prevalence of autism in a US metropolitan area. *JAMA* 289:49–55.
168. Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, *et al.* (2010): Lifetime prevalence of mental disorders in U.S. adolescents: Results from the National Comorbidity Survey Replication-Adolescent Supplement (NCS-A). *J Am Acad Child Adolesc Psychiatry* 49:980–989.
169. Aleman A, Kahn RS, Selten JP (2003): Sex differences in the risk of schizophrenia: Evidence from meta-analysis. *Arch Gen Psychiatry* 60:565–571.
170. Salter MW, Beggs S (2014): Sublime microglia: Expanding roles for the guardians of the CNS. *Cell* 158:15–24.
171. Kabat AM, Srinivasan N, Maloy KJ (2014): Modulation of immune development and function by intestinal microbiota. *Trends Immunol* 35:507–517.
172. Maynard CL, Elson CO, Hatton RD, Weaver CT (2012): Reciprocal interactions of the intestinal microbiota and immune system. *Nature* 489:231–241.
173. Capuron L, Miller AH (2011): Immune system to brain signaling: Neuropsychopharmacological implications. *Pharmacol Ther* 130:226–238.
174. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW (2008): From inflammation to sickness and depression: When the immune system subjugates the brain. *Nat Rev Neurosci* 9:46–56.
175. Leboyer M, Soreca I, Scott J, Frye M, Henry C, Tamouza R, *et al.* (2012): Can bipolar disorder be viewed as a multi-system inflammatory disease? *J Affect Disord* 141:1–10.
176. Laske C, Zank M, Klein R, Stransky E, Batra A, Buchkremer G, *et al.* (2008): Autoantibody reactivity in serum of patients with major depression, schizophrenia and healthy controls. *Psychiatry Res* 158:83–86.
177. Horváth S, Mirmics K (2014): Immune system disturbances in schizophrenia. *Biol Psychiatry* 75:316–323.
178. Li X, Chauhan A, Sheikh AM, Patil S, Chauhan V, Li XM, *et al.* (2009): Elevated immune response in the brain of autistic patients. *J Neuroimmunol* 207:111–116.
179. Rossignol DA, Frye RE (2012): A review of research trends in physiological abnormalities in autism spectrum disorders: Immune dysregulation, inflammation, oxidative stress, mitochondrial dysfunction and environmental toxicant exposures. *Mol Psychiatry* 17:389–401.
180. Slykerman RF, Hood F, Wickens K, Thompson JMD, Barthow C, Murphy R, *et al.* (2017): Effect of *Lactobacillus rhamnosus* HN001 in pregnancy on postpartum symptoms of depression and anxiety: A randomised double-blind placebo-controlled trial. *EBioMedicine* 24:159–165.