



Review article

Neural and cognitive correlates of stigma and social rejection in individuals with Serious Mental Illness (SMI): A systematic review of the literature

Dubreucq J.^{a,b,c,*}, Franck N.^{a,d,e,f}

^a Centre de Neurosciences Cognitive, UMR 5229, CNRS & Université Lyon 1, France

^b Centre Référent de Réhabilitation Psychosociale et de Remédiation Cognitive (C3R), Centre Hospitalier Alpes Isère, 1 place du Conseil National de la Résistance, Grenoble, 38400 Saint Martin d'Hères, France

^c Fondation FondaMental, Créteil, France

^d Réseau Handicap Psychique, Grenoble, France

^e Centre Référent Lyonnais de Réhabilitation Psychosociale et de Remédiation Cognitive (CL3R), Centre Hospitalier Le Vinatier, France

^f Centre Ressource de Réhabilitation Psychosociale et de Remédiation Cognitive, Centre Hospitalier Le Vinatier, France

ARTICLE INFO

Keywords:

Cognitive neuroscience

Self-stigma

Stigma

Social rejection

Severe mental illness

ABSTRACT

Stigma and self-stigma are major issues for people with Serious Mental Illness (SMI). This review's aims were to determine the cognitive and neural processes underlying the effects of stigma and social rejection in people with SMI.

A stepwise systematic literature review (PRISMA) was conducted by searching PubMed, Medline and Web of Science using the following keywords: “cyberball” OR “stereotype threat” OR “implicit association test” AND “mental illness”. The articles included met the following criteria: (a) reporting on social rejection, stigma or self-stigma (b) diagnosis of SMI (c) available data on the underlying mechanisms.

Our search on July 31st 2018 found in 955 articles on PubMed and 3,362 on Web of Science. Hypersensitivity to acute social rejection was found and associated with more self-related negative emotions and more dysfunctional emotion regulation strategies. People with SMI under-performed in cognitive tasks when confronted with stigma. Stigma resistance was described with its neural correlates. Psychiatric symptoms, duration of illness and baseline non-specific distress influenced the response to acute social rejection or stigma. The samples, methods, and reported outcomes were heterogeneous and difficult to compare. Future studies should investigate the associations between self-stigma and responses to acute and chronic social rejection.

1. Introduction

Stereotypes can be defined as common social knowledge shared by people from the same cultural background. They make it possible to rapidly identify and categorize the individuals with whom one interacts and contribute to understanding social situations and shaping behavioral responses (Krendl et al., 2006). Stigma occurs when stereotypes generate negative emotional responses and behaviors towards the members of a group – or towards oneself in the case of self-stigma (Corrigan and Watson, 2002). Self-stigma - or Internalized Stigma (IS) - is an identity transformation process wherein a person's previously held social identity (defined by social roles such as son, brother, sister, friend, employee or potential partner) is progressively replaced by a devalued and stigmatized view of oneself termed “illness identity” (Yanos et al., 2008). Stigma Resistance (SR) is defined as one's ability to deflect or challenge stigmatizing beliefs (Thoits, 2011).

According to the US Substance Abuse and Mental Health Services Administration, individuals with “Serious Mental Illness” (SMI) are “persons aged 18 or older who currently or at any time in the past year have had a diagnosable mental, behavioral, or emotional disorder (excluding developmental and substance use disorders) of sufficient duration to meet diagnostic criteria specified within DSM-IV that has resulted in serious functional impairment, which substantially interferes with or limits one or more major life activities such as maintaining interpersonal relationships, activities of daily living, self-care, employment, and recreation”. Serious Mental Illnesses (SMI; SAMSAH, 2014) include schizophrenia (SZ), bipolar disorder (BD), and borderline personality disorder (BPD), as well as Major Depressive Disorder (MDD) and Anxiety Disorders that cause severe and persistent functional impairments. SMI are frequent (1% for SZ; 1.5% for BD; 1.5% for BPD) and heavily stigmatized (Angermeyer et al., 2013; Ellison et al., 2013). People with SZ, BD or MDD are more likely to

* Corresponding author at: Centre Référent de Réhabilitation Psychosociale et de Remédiation Cognitive (C3R), Centre Hospitalier Alpes Isère, 1 place du Conseil National de la Résistance, Grenoble, 38400 Saint Martin d'Hères, France.

E-mail address: jdubreucq@ch-alpes-isere.fr (D. J.).

<https://doi.org/10.1016/j.psychres.2019.02.023>

Received 18 November 2018; Received in revised form 9 February 2019; Accepted 9 February 2019

Available online 11 February 2019

0165-1781/ © 2019 Elsevier B.V. All rights reserved.

experience discrimination in a range of aspects of their daily life (Thornicroft et al., 2009; Farrelly et al., 2014; Lasalvia et al., 2013). They frequently expect to be discriminated in social contexts even when they have no previous experience of discrimination (Thornicroft et al., 2009; Farrelly et al., 2014; Lasalvia et al., 2013). Internalized Stigma (IS) is frequent (41.7% of the 1229 participants with SZ and 21.7% of the 1182 participants with BD in the 2010 GAMIAN-Europe study had moderate to high levels of self-stigma; Brohan et al., 2010, 2011) and associated with multiple negative consequences: lower therapeutic adherence, self-efficacy, self-esteem, empowerment and hope and a greater intensity of depressive symptoms and suicidal ideation (Livingston and Boyd, 2010; Ellison et al., 2013). Stigma resistance concerns a wide range of people with SMI (49.2% of the 1229 participants with SZ and 49.7% of the 1182 participants with BD in the 2010 GAMIAN-Europe study had moderate to high levels of self-stigma; Brohan et al., 2010, 2011). It was associated with higher levels of metacognitive abilities, self-efficacy, self-esteem, empowerment, hope, personal recovery and subjective quality of life (Nabors, 2014; Firmin et al., 2016). Several psychosocial interventions (mainly combinations of psychoeducation and Cognitive Behavior Therapy (CBT)) have been designed to reduce IS and its impact on clinical and functional outcomes, with preliminary results on IS, insight, and self-efficacy. Their impact on functional brain connectivity remains unknown to date.

Public stigma, i.e. the relatively automatic cognitive and behavioral response in a social context cued by a characteristic related to a stigmatized group (Krendl et al., 2006), is the most documented area in the literature (Krendl et al., 2013; Reihl et al., 2015). The processes leading to negative or stigmatizing attitudes towards individuals with SMI were investigated and showed the role of the SZ diagnosis, persistent negative symptoms, and functional impairments (mostly on social skills) play in participants' emotional reactions when interacting with people they believed to have SZ (Penn et al., 2000; Heenan et al., 2014). More severe negative symptoms and poorer social skills were associated with higher perceived strangeness and desire for social distance when interacting with individuals with SZ (Penn et al., 2000).

In contrast with public stigma (or endorsement of negative beliefs and discriminating attitudes by the general population), personal stigma refers to the beliefs experiences and attitudes towards mental illness endorsed by the members of a stigmatized group, in this case people with SMI (Gerlinger et al., 2013). It includes perceived stigma (or one's beliefs and anticipations about the perception and attitudes of the general population towards persons with mental illness), experienced stigma (or the actual experiences of social discrimination lived by the persons with mental illness) and self-stigma. In a recent study anticipated social discrimination mediated the relationships between experienced stigma and self-stigma (Quinn et al., 2015), showing that these concepts may be interrelated. Three experimental paradigms were used in the literature to investigate the cognitive and neural mechanisms underlying personal stigma. Implicit association tests (Greenwald et al., 2009) and stereotype threat conditions directly addressed stigma or self-stigma and the cyberball game was used to investigate social exclusion, a stigma-related concept.

Implicit association tests (IAT) are commonly used to assess implicit self-associations, emotional reactions and attitudes towards a concept or a population, as they are less influenced than explicit measures by social desirability biases and have good psychometric properties (Greenwald et al., 2009). IAT assess the strength of association between concepts by measuring response latencies during computerized categorization tasks: the quicker the stimuli are classified, the more likely they are to match the respondent's associations with a target concept (Greenwald et al., 2009). IAT are particularly well-suited to investigating automatically activated reactions towards stigmatized individuals, as these reactions can influence outcomes, and because deliberately endorsed stereotypes and negative attitudes towards stigmatized individuals have become less acceptable (Greenwald et al.,

2009). IAT was used to assess automatic stereotyping and negative emotional reactions towards individuals with SMI in the general population (Peris et al., 2008; Heenan et al., 2014), in mental health professionals (Stull et al., 2013), and in participants diagnosed with SZ or BD (Teachman et al., 2006; Rusch et al., 2010a, 2011a).

The stereotype threat (ST) model is a popular experimental paradigm in social psychiatry that has been used in numerous studies concerning discriminated populations and minorities (Derks et al., 2008). The ST paradigm refers to situations where one individual is at risk of confirming by his/her actions his/her negative stereotypes about themselves or his/her social group (Derks et al., 2008). As an individual can suffer simultaneously from various forms of discrimination, Shapiro and Neuberg created a multi-threat framework. In this model, self and group concept threats imply endorsement by an individual of negative beliefs about oneself or about one's social group, respectively. Own and group reputation threats correspond to one's fear of being identified as possessing stigmatizing attributes and therefore reinforce negative beliefs about his/her own social group (Shapiro and Neuberg, 2007). Given the proximity of these concepts to stigma and self-stigma, the number of studies using ST in individuals with SMI is surprisingly limited (Henry et al., 2010; Moritz et al., 2018; Shapiro, 2011).

Social rejection models recreate stigma-related environments under experimental conditions. The most popular paradigm is the cyberball game, a ball-tossing game in which participants believe they are playing with fellow participants, but are in fact randomly assigned to inclusion, exclusion and sometimes over-inclusion conditions and play with computer-controlled players. Cyberball is considered to be an ecologically valid paradigm for evaluating emotional reactions following social inclusion or exclusion, and was used in numerous studies involving healthy individuals (Hartergerink et al., 2015). The neural correlates of acute social exclusion/inclusion in healthy controls (HC) are well known (Bolling et al., 2011; Moor et al., 2012; Kawamoto et al., 2012). Enhanced activation of the insula and of the dorsal and ventral anterior cingulate cortex (ACC) during social exclusion was associated with self-reported experience of rejection (Bolling et al., 2011; Moor et al., 2012). Enhanced activation of the medio-prefrontal cortex (mPFC), the precuneus and the posterior cingulate cortex during social exclusion were identified as a potential protective mechanism modulating the affective response to acute social rejection (Bolling et al., 2011; Moor et al., 2012; Kawamoto et al., 2012). Cyberball was used in individuals with SMI, mostly BPD (Lawrence et al., 2011; Staebler et al., 2011; Domsalla et al., 2014; De Panfilis, 2015; Brown et al., 2017; Euler et al., 2018; Ernst et al., 2018; Weinbrecht et al., 2018).

In summary, the clinical and functional correlates of stigma in individuals with SMI are well known and several strategies have been recently developed to reduce stigma and self-stigma (Griffiths, 2015; Wood, 2016). In contrast, there appears to be surprisingly little cognitive neuroscience research into mental illness stigma. Only a few studies directly addressed the cognitive and neural processes underlying self-stigma or stigma resistance in individuals with SMI (Rusch 2010a, 2010b, 2011; Raji et al., 2014; Moritz et al., 2018). These mechanisms remain therefore unclear as the data are heterogeneous and come from various sources and different experimental paradigms (some of which not even directly related to stigma or self-stigma). To our knowledge, no literature review has yet been conducted on this particular topic.

The present review has two objectives: (i) to review the cognitive and neural processes underlying the effects of stigma, discrimination and social rejection in individuals with SMI; (ii) to evaluate the strengths and limits of the current evidence in order to guide future research.

2. Methods

A stepwise systematic literature review (PRISMA guidelines) was conducted by searching PubMed, Medline and Web of Science for published, peer-reviewed papers using the following keywords:




Fig. 1. Review process (Prisma flow diagram).

“cyberball” OR “stereotype threat” OR “implicit association test” AND “mental illness”. The articles included in this review had to meet all the following criteria: (a) reporting on social rejection, stigma, or experiences of discrimination; (b) diagnosis of SMI (c) available data on the neural and cognitive mechanisms underlying social rejection, experiences of discrimination and stigma or self-stigma.

3. Results

Our search on July 31st 2018 found 955 articles on PubMed and 3362 on Web of Science. After manually removing all duplicates, 1209 articles were retained. Based on their titles and abstracts, 1153 papers were excluded for lack of relevance. Most of these articles focused on stigma-related experimental paradigms in healthy individuals or in other discriminated populations. Our search strategy yielded 57 full papers. After conducting a full-text analysis of all these papers and after excluding those which did not meet the inclusion criteria we ended up with 34 relevant papers (Fig. 1).

3.1. Neural and cognitive correlates of self-stigma

Nine (23.6%) of the 38 included papers reported directly on self-stigma. However, these studies were characterized by the heterogeneity of samples, methods, and reported outcomes. Two-hundred-fifty-four people with SMI and 148 healthy controls were included. Sample sizes ranged from 36 to 135 participants, mean age from 26 to 45 years, and the proportion of male participants ranged from 12.5% to 75%. Five

studies reported data on the participants’ ethnic origins with a proportion of Caucasian participants ranging from 34 to 77% (Teachman et al., 2006; Rusch, 2010a, 2010b, 2010c, 2011). Four studies investigated a specific psychiatric diagnosis (OCD; Moritz et al., 2018; SZ; Henry et al., 2010; Raji et al., 2014) whereas the other five included a broader range of diagnoses (SZ-spectrum disorders, BD, unipolar MDD and OCD). Seven studies only included outpatients. One only included inpatients and one did not report the patients’ statuses at inclusion. The mean duration of illness was reported in six studies and ranged from 49 months (Raji et al., 2014) to 15 years (Rusch, 2010a, 2010b, 2010c, 2011; Henry et al., 2010). The mean number of psychiatric hospitalizations was recorded in four studies (Rusch, 2010a, 2010b, 2010c, 2011) which all came from the same sample. Three studies reported data on patients’ current medication status (Henry et al., 2010; Raji et al., 2014; Moritz et al., 2018). Five studies used IAT (Teachman et al., 2006; Rusch 2010a, 2010b, 2010c; Raji et al., 2014), one used Lexical Decision Task (Wittenbrink et al., 2001; Rusch et al., 2011) and two ST (Henry et al., 2010; Moritz et al., 2018). Two studies included f-MRI data, one acquired during passive emotion viewing with visual stimuli from the Karolinska Directed Emotional Faces (KDEF) catalogue and one during a visuo-motor rating task with stigma-related and non-stigma related visual stimuli (statements about schizophrenia or common cold) (Raji et al., 2014). Results are shown on Table 1.

Two cognitive models of self-stigma and one of stigma resistance were described. Implicit self-stigma (defined as the combination of implicit negative attitudes towards mental illness and low implicit self-esteem; Rusch 2010c) was strongly correlated with explicit self-stigma

Table 1
Articles reporting directly on self stigma.

Author	Socio-demographic characteristics				Psychiatric history				Treatments	Experimental design	results	
	n	Age (mean)	Sex (%)	Ethnicity (%)	Education (mean)	Diagnosis	Current CM	Statut				Psychiatric history
Teachman et al. (2006)	35 SMI 36 HC	43	32% women	77% caucasian	-	59% psychotic disorders, 31% mood disorders; 31% PD, 13% AD	none	Outpatient	No	-	IAT	Similar implicit negative beliefs and attitudes towards SMI between persons with SMI and controls
Rusch et al. (2011a)	85 SMI 50 HC	45	32% women	34% caucasian, 58% afro-american	13.5 years	27% SZ, 26% schizoaffective disorders, 30% BD, 12% unipolar MDD	39% SUD	Outpatient	Yes Illness duration: 15 years Mean number of hospitalizations = 9	-	Lexical Decision Task	Reduced automatic stereotyping in persons with SMI compared to controls
Rusch et al. (2010a)	85 SMI 50 HC	45	32% women	34% caucasian	13.5	27% SZ, 26% schizoaffective disorders, 30% BD, 12% unipolar MDD	39% SUD	Outpatient	Yes Illness duration: 15 years Mean number of hospitalizations = 9	-	IAT	Endorsement of biogenetic causes (but not neurobiological) of mental illness is associated to increased implicit self-guilt and self reported fear towards people with SMI
Rusch et al. (2010b)	75 SMI	44	69% men	34% caucasian	13.5	24% SZ, 28% schizoaffective disorder, 36% BD, 9% unipolar MDD,	43% SUD	Outpatient	Yes Illness duration: 15 years Mean number of hospitalizations = 9	-	IAT	Implicit shame-related associations at baseline predict increased perceived legitimacy of discrimination at 6 months follow-up
Rusch et al. (2010c)	85 SMI	45	68% men	34% caucasian	13.5	27% SZ, 26% schizoaffective disorders, 30% BD, 12% unipolar MDD	39% SUD	Outpatient	Yes Illness duration: 15 years Mean number of hospitalizations = 9	-	IAT	Implicit self-stigma predicts lower quality of life independently of explicit self-stigma, depressive symptoms and psychiatric diagnosis
Henry et al. (2010)	30 SZ	38	43% men	-	13.4	70% SZ, 30% schizoaffective disorder	none	Outpatient	Yes -Mean illness duration = 15.8 years -Mean age of onset = 23.1 years	100% atypical AP; 343.2 CE	ST with conversation probe role play test	Lower social skills (higher impairment in initiating conversations and switching topics) under ST
Moritz et al. (2018)	50 OCD 43 (PC)	38 (ST), 43 (PC)	87.5% women (ST), 69.2% (PC)	-	-	100% OCD	not recorded	Outpatient	No	52% Antidepressants	ST	Lower executive functioning under ST
Lindner	38 SZ 42 HC	30	61% men (SZ) 67.5% (HC)	-	13.1	100% SZ	BDI 12.7 STAI 46.4	Inpatient	Yes Mean illness duration: 6.8 years	-	Passive emotion viewing during fMRI	Reduced insula activation to subtle expressions of disgust and positive correlation between insula activation and social loneliness in participants with SZ

(continued on next page)

Table 1 (continued)

Raij et al. (2014)	Socio-demographic characteristics			Psychiatric history			Psychiatric history	Statut	Treatments	Experimental design	results
	n	Age (mean)	Sex (%)	Ethnicity (%)	Education (mean)	Diagnosis					
20 SZ 16 HC	Patients 26 Controls 28	Patients 70% men Controls 75% men	-	13	100% SZ	25% MDD, 20% OCD, 10% SAD, 15% PD	Not recorded	Yes	100% AP (575 CE)	IAT Visuomotor rating task during fMRI	- Lower implicit negative associations in patients in comparison with controls -Negative correlation between implicit associations and rostral ventral medio- prefrontal cortex - Negative correlation between rostral ventral mPFC activation and right amygdala activation

AD: Anxiety Disorders; AP: Antipsychotics; BD: Bipolar Disorders; BDI: Beck Depression Inventory; CE: Comorbidity; CM: Comorbidities; fMRI: Functional Magnetic Resonance Imaging; HC : Healthy Controls ; IAT: Implicit Associations Test; MDD: Major Depressive Disorder; mPFC: Medio Prefrontal Cortex; OCD: Obsessive Compulsive Disorder; PC: Patient Controls; PD: Personality Disorder; SAD: Social Anxiety Disorder; SMI: Severe Mental Illness ; ST: Stereotype Threat; STAI: Spielberger Trait Anxiety Inventory; SUB: Substance Use Disorder; SZ: Schizophrenia.

and identified as a predictor of subjective quality of life (Rusch, 2010c). Social alienation (or high perceived social loneliness associated with low personality trait agreeableness) was associated with reduced insula activation in participants with SZ when processing covert expressions of disgust. According to the authors, this reflected a decreased sensitivity to subtle signals of social rejection that might protect individuals against social alienation. There was however also a negative association between insula responsiveness to subtle facial expressions of disgust and low agreeableness in the control group.

Stigma resistance (defined as low negative implicit associations between SZ and social inferiority) was associated with the enhanced activation of rostral mPFC and decreased activation of the amygdala (Raij et al., 2014). Acute social rejection, low stigma resistance, and ST conditions led to underperformance in cognitive or social cognitive tasks (Raij et al., 2014; Moritz et al., 2018) and in social skills (Henry et al., 2010) in people with SMI. The associations between self-stigma, cognitive performance and social functioning are however unclear as self-stigma did not influence cognitive performance under ST conditions in people with OCD (Moritz et al., 2018) and was not recorded in the study by Henry et al. (2010).

3.2. Neural and cognitive correlates of social rejection

The consequences of social rejection were investigated in 25 studies (65.7%), with varying samples, methods, and reported outcomes. Most of the 629 participants in the experimental groups had BPD (19 studies; n = 475; 75.5%), the other diagnoses represented were SZ (n = 55; 8.7%; Gradin et al., 2012; Engel et al., 2016), a high clinical risk of psychosis (CHR; n = 25; 3.9%), SAD (n = 23; 3.6%; Heeren et al., 2017), MDD (n = 15; 2.3%; Kumar et al., 2017) and suicide attempters (SA; n = 36; 5.7%; Olié et al., 2017). No studies were conducted specifically on people with BD. However, four studies recruited participants with type I or II BD in their experimental group (10 to 56% of the total sample; Domsalla et al., 2014; Renneberg et al., 2012; Bungert et al., 2015; Olié et al., 2017). All studies recruited healthy controls (HC). Seven studies had also patient controls (PC) with SAD (Gutz et al., 2015, 2016; Weinbrecht et al., 2018), AD, MDD (Ernst et al., 2018; Olié et al., 2017) or non-suicidal self-injury (Brown et al., 2017). Sample sizes ranged from 32 to 122 participants, mean age from 19 to 45 years and the proportion of female participants from 15.4% to 100%. Education level was reported in 16 studies and ranged from 10.61 to 15.7 years of education. Three studies reported data on the participants' ethnic origins with a proportion of Caucasian participants ranging from 49 to 80% (Ruocco et al., 2010; Gratz et al., 2013). Eleven studies were conducted on outpatients only, five on stabilized inpatients (admitted to receive specific psychotherapy), four had mixed samples and five did not report the participants' status at inclusion. Almost all studies recorded current psychiatric comorbidities. Eleven studies recorded lifetime comorbidities and one recorded the number of past depressive or manic episodes and the age of onset. Fourteen studies reported data on patients' medication with the proportion of participants taking medication at inclusion ranging from 0% to 92%. Most studies used cyberball. Two studies added an over-inclusion condition (i.e. a condition where the participant received the ball more often than the other players) to the conventional experimental paradigm (De Panfilis, 2015; Weinbrecht et al., 2018) and one a re-inclusion condition (Heeren et al., 2017). One study used implicit and explicit emotion processing tasks (Schienle et al., 2015) and one an emotion reappraisal task during f-MRI (Koenigsberg et al., 2009). Two studies included biological measures (oxytocin and cortisol dosages; Jobst et al., 2014,2016), ten studies f-MRI (Koenigsberg et al., 2009; Ruocco et al., 2010; Gradin et al., 2012; Domsalla et al., 2014; Bungert et al., 2015; Schienle et al., 2015; Kumar et al., 2017; Heeren et al., 2017; Olié et al., 2017; Brown et al., 2017), two EEG (Gutz et al., 2015; Weinbrecht et al., 2018) and two cognitive assessments (Go-No Go; Reading Mind in the Eyes Test). The results are presented in Table 2.

Table 2
Articles on social rejection in SMI.

	Socio-demographic characteristics				Psychiatric history			Experimental design	Results		
	n	Age (mean)	Sex (%)	Education (mean)	Diagnosis	Current CM	Statut			Psychiatric history	Treatments
Staebler et al. (2011)	35 BPD 33 HC	27.88 to 32.11	100% women	10.61	100% BPD	20% PTSD	Inpatients	Yes Past 54% MDD, 37% SUD	34.28% AD ; 31.4% AP 28.57% none	Cyberball	Women with BPD reported more feeling excluded when included and had less positive and increased negative emotions when excluded No difference in mood rating after social exclusion between young adults with BPD and healthy controls -More reported exclusion feelings during social inclusion and exclusion in BPD -More reported anger during social exclusion in BPD -Hypersensitivity to social rejection and heightened threat to all social needs in BPD -Emotion dysregulation mediated the relationship between BPD and response to the task -Increased self-focused negative emotions in both SAD and BPD -Increased hostile intent attributions in BPD Greater levels of negative reported emotions during social exclusion, inclusion but not during overinclusion in BPD Higher threat to social needs levels after social exclusion predicted lower therapeutic alliance in mentalization-based group therapy -Lower response inhibition in BPD + MDD and correlation with childhood trauma -No improvement of response inhibition during social exclusion in BPD + MDD Lower ability to identify correctly neutral faces after social exclusion in young adults with BPD features More reported feelings of exclusion and enhanced P3b during social inclusion in BPD -Enhanced P3 amplitude in SAD and BPD compared to controls in all conditions -Stronger self-reported ostracism
Lawrence et al. (2011)	30 BPD 22 HC	18.56 to 19.33	86 to 90% women	-	75% full BPD; 25% sub-threshold BPD	63% MDD, 46% PTSD; 73% AD	Outpatients	No	66.6% AD, 7.4% BZD, 3.5% AP	Cyberball	
Renneberg et al. (2012)	30 BPD 30 HC	28.8 BPD 29.03 HC	86.7% women	10.6	100% BPD	30% MDD, 3.3% BD; 60% SUD, 43.3% AD, 36.7% PTSD, 50% ED	Inpatients	Yes Past MDD 50%	60% AD, 23.3% AP, 10% MS, 20% none	Cyberball	
Gratz et al. (2013)	53 BPD 34 HC	-	72% women (BPD), 50% (HC)	-	100% BPD	none	Outpatients	No	not recorded	Cyberball	
Gutz et al. (2016)	25 BPD 25 SAD 25 HC	25 to 28	84 to 92% women	11.57	100% BPD	12% MDD, 8% SAD, 24% PTSD, 24% ED, 8% dysthymia	Inpatients	Yes Past 48% MDD	not recorded	Cyberball	
De Pauflis et al. (2015)	61 BPD 61 HC	40.2 BPD 37.6 HC	77% women (BPD) and 70.5% (HC)	14.8% college / university	100% BPD	36% MDD, 18% ED, 4.9% AD, 3.3% adjustment disorder, 44.3% other PD	Outpatients	No	not recorded	Cyberball	
Euler et al. (2018)	23 BPD 28 HC	27.9 BPD 25.7 HC	87% women (BPD), 85% (HC)	12.8	100% BPD	34.8% MDD, 8.7% AD, 26.1% SUD, 13% ED, 8.7% SUD	Inpatients	No	69.6% on medication	Cyberball	
Ernst	22 MDD 22 MDD + BPD 20 HC	26.73 to 30.5	100% women	25% college students	100% BPD	Mean BDI 37.4 MDD + BPD; 33 MDD	Not recorded	No	40% AD	Cyberball and Go-No Go task	
Savage	17 BPD 16 HC	19.65 BPD 19.41 HC	56.3% to 64.7% women	-	100% BPD	Mean BDI 21.0, STAI 95.47	Students	No	Not recorded	Cyberball and RMIE test	
Gutz et al. (2015)	25 BPD 25 SAD 25 HC	25 to 28	84 to 92% women	11.57	100% BPD	12% MDD, 8% SAD, 24% PTSD, 24% ED	Inpatients	Yes Past 48% MDD	-	Cyberball and EEG	
Weinbrecht et al. (2018)	29 BPD 28 SAD 28 HC	27.86 to 28.86	79 to 86% women	-	100% BPD	7% MDD, 55% AD, 4% SAD, 7% other PD	62% inpatients for BPD	Yes Past CM: 48% MDD	35% AD	Cyberball and EEG	

(continued on next page)

Table 2 (continued)

n	Socio-demographic characteristics			Psychiatric history			Statut	Psychiatric history	Treatments	Experimental design	Results
	Age (mean)	Sex (%)	Education (mean)	Diagnosis	Current CM	Psychiatric history					
Ruocco et al. (2010)	10 BPD 10 HC	100% women	13.2	100% BPD	30% MDD, 20% AD, 10% PTSD	No	Not recorded	80% none, 10% AP, 10% stimulants	fMRI during cyberball	during social exclusion / inclusion but not overinclusion in SAD and BPD compared to controls -Higher self-reported threats to social needs in all conditions in BPD Left hyperactivation of the mPFC in BPD during social exclusion - Higher reported exclusion feelings during social inclusion task and control condition in BPD - No changes depending on the interaction situation (exclusion/inclusion) in insula and precuneus responsiveness in BPD	
Domsalla et al. (2014)	20 BPD 20 HC	100% female	12.1	90% BPD, 10% BPD + type II BD	25% PTSD, 75% AD, 40% ED	Yes Past 90% MDD, 25% PTSD, 60% AD, 5% ED	Outpatients	-	Cyberball and fMRI	- Enhanced feelings of social exclusion in patients groups - Enhanced activation in ventral anterior cingulate cortex in patients groups - Enhanced activation in dorsolateral and mPFC during social inclusion in BPD - Reduced amygdala activation during pain after social inclusion in BPD - Association between higher rejection sensitivity and lower activation differences during physical pain processing in amygdala and insula after inclusion / exclusion in BPD - No differences in cortisol plasma concentrations between BPD and controls - Negative association between physical and emotional abuse and return of oxytocin levels at baseline after social exclusion - No differences in cortisol plasma concentrations between BPD and controls - Lower oxytocin concentrations in BPD patients with disorganized attachment representations compared to other BPD	
Brown et al. (2017)	15 BPD + NSSI 14 NSSI 31 HC	76.9% to 100% women	-	100% BPD	100% MDD, 50% PTSD, 14.2% AD, 7.1% ED	Yes Mean number of NSSI: 308.5 (BPD) and 128.8 (NSSI)	In and outpatients	85% AD, 7.1% MS	Cyberball f-MRI	- Enhanced feelings of social exclusion in patients groups - Enhanced activation in ventral anterior cingulate cortex in patients groups - Enhanced activation in dorsolateral and mPFC during social inclusion in BPD - Reduced amygdala activation during pain after social inclusion in BPD - Association between higher rejection sensitivity and lower activation differences during physical pain processing in amygdala and insula after inclusion / exclusion in BPD - No differences in cortisol plasma concentrations between BPD and controls - Negative association between physical and emotional abuse and return of oxytocin levels at baseline after social exclusion - No differences in cortisol plasma concentrations between BPD and controls - Lower oxytocin concentrations in BPD patients with disorganized attachment representations compared to other BPD	
Bungert et al. (2015)	20 BPD 20 HC	100% women	12.1	90% BPD, 10% BPD + type II BD	25% PTSD, 35% AD, 25% SAD, 5% ED	Yes Past 90% MDD, 25% PTSD, 75% AD, 35% SUD	Outpatients	-	Cyberball f-MRI	- Enhanced feelings of social exclusion in patients groups - Enhanced activation in ventral anterior cingulate cortex in patients groups - Enhanced activation in dorsolateral and mPFC during social inclusion in BPD - Reduced amygdala activation during pain after social inclusion in BPD - Association between higher rejection sensitivity and lower activation differences during physical pain processing in amygdala and insula after inclusion / exclusion in BPD - No differences in cortisol plasma concentrations between BPD and controls - Negative association between physical and emotional abuse and return of oxytocin levels at baseline after social exclusion - No differences in cortisol plasma concentrations between BPD and controls - Lower oxytocin concentrations in BPD patients with disorganized attachment representations compared to other BPD	
Jobst et al. (2014)	22 BPD 21 HC	100% women	-	100% BPD	-	No	Not recorded	68.1% AD ; 63.6% AP ; 36.3% MS	Cyberball and oxytocin and cortisol dosages	- Enhanced feelings of social exclusion in patients groups - Enhanced activation in ventral anterior cingulate cortex in patients groups - Enhanced activation in dorsolateral and mPFC during social inclusion in BPD - Reduced amygdala activation during pain after social inclusion in BPD - Association between higher rejection sensitivity and lower activation differences during physical pain processing in amygdala and insula after inclusion / exclusion in BPD - No differences in cortisol plasma concentrations between BPD and controls - Negative association between physical and emotional abuse and return of oxytocin levels at baseline after social exclusion - No differences in cortisol plasma concentrations between BPD and controls - Lower oxytocin concentrations in BPD patients with disorganized attachment representations compared to other BPD	
Jobst et al. (2016)	20 BPD 20 HC	100% women	11.20	100% BPD	Mean BDI 32.75 25% MDD 15% ED	No	70% outpatients	75% AD, 70% AP, 40% MS	Cyberball and oxytocin and cortisol dosages	- Enhanced feelings of social exclusion in patients groups - Enhanced activation in ventral anterior cingulate cortex in patients groups - Enhanced activation in dorsolateral and mPFC during social inclusion in BPD - Reduced amygdala activation during pain after social inclusion in BPD - Association between higher rejection sensitivity and lower activation differences during physical pain processing in amygdala and insula after inclusion / exclusion in BPD - No differences in cortisol plasma concentrations between BPD and controls - Negative association between physical and emotional abuse and return of oxytocin levels at baseline after social exclusion - No differences in cortisol plasma concentrations between BPD and controls - Lower oxytocin concentrations in BPD patients with disorganized attachment representations compared to other BPD	
Olié et al. (2017)			14 SA, 15 PC				Inpatients		Cyberball f-MRI	- Enhanced feelings of social exclusion in patients groups - Enhanced activation in ventral anterior cingulate cortex in patients groups - Enhanced activation in dorsolateral and mPFC during social inclusion in BPD - Reduced amygdala activation during pain after social inclusion in BPD - Association between higher rejection sensitivity and lower activation differences during physical pain processing in amygdala and insula after inclusion / exclusion in BPD - No differences in cortisol plasma concentrations between BPD and controls - Negative association between physical and emotional abuse and return of oxytocin levels at baseline after social exclusion - No differences in cortisol plasma concentrations between BPD and controls - Lower oxytocin concentrations in BPD patients with disorganized attachment representations compared to other BPD	

(continued on next page)

Table 2 (continued)

n	Socio-demographic characteristics			Psychiatric history			Statut	Psychiatric history	Treatments	Experimental design	Results
	Age (mean)	Sex (%)	Education (mean)	Diagnosis	Current CM	Diagnosis					
36 SA 41 PC 28 HC		100% women		56% BD in SA group; 36% in PC group	SA group , 39.2% AD, 16.8 SAD, 2.8% PTSD, 16.8% ED, PC group BD, 36% AD, 4.8% SAD, 12% ED;	- Mean number of depressive episodes: 3.5 - Mean number of manic episodes: 1 - Age of onset first SA: 22 - Past CM: 25.2% SUD	Yes	33.6% on medication (SA); 38.4% (PC)		-Higher reported social distress in patients groups -Decreased contrast in the left insula and supramarginal gyrus during the exclusion vs inclusion condition	
15 SZ 20 HC	41.23 SZ 40.87 HC	84.6% men (SZ) 77.7% (HC)	-	100% SZ	Mean BDI 17.43	No	Outpatients	Not recorded	Cyberball fMRI	Increased activation of the mPFC cortex during social exclusion in HC but not in SZ patients Association between positive symptoms severity and blunted response to exclusion in SZ More intense negative anticipations before cyberball in patients Less intense positive emotions during social inclusion in patients Stronger increase in paranoid beliefs after social exclusion in CHR mediated by lower levels of functional / higher levels of dysfunctional emotion regulation and by higher self-reported negative emotions -Higher reported feelings of social exclusion in patients -Higher activation within the left inferior frontal gyrus during re-inclusion phase. -Correlation between and feelings of exclusion and cerebral activation Greater amygdala, insula and ventrolateral prefrontal cortex activation to increasing social exclusion in patients	
40 SZ 40 HC	36.32 SZ 26.45 HC	50% women	11.1	77.5% SZ, 17.5% schizoaffective disorder	-	No	Inpatients	92.5% AP	Cyberball		
25 CHR 40 AD 40 HC	34.72 CHR, 40.3 HC, 42.2 AD	62.5 to 72% women	15.7	76% APS 8% GR 16% both	96% mood disorders; 36% AD, 16% PTSD, 8% SUD; 8% OCD	No	Outpatients	52% on medication, 16% AP	Cyberball		
23 SAD 23 HC	24.96 SAD 25.30 HC	100% women	13.3	100% SAD	Mean BDI : 15.7 ; STAI : 51.04 ; LSAS : 74.2	No	Inpatients	Not recorded	Cyberball fMRI		
15 MDD 17 HC	45.27 MDD 41.8 HC	60% women (MDD) 58% (HC)	-	Not recorded	Mean BDI: 22.9; STAI: 54.6	No	Outpatients	-	Cyberball F-MRI		

AD: Anxiety Disorders; AP: Antipsychotics; APS: Attenuated Positives Symptoms Syndrome; BD: Bipolar Disorders; BDI: Beck Depression Inventory; BPD: Borderline Personality Disorder; CHR: Clinical High Risk; CM: Comorbidities; ED: Eating Disorder; EEG: Electroencephalogram; GR: Genetic Risk And Deterioration Syndrome; HC : Healthy Controls; fMRI: Functional Magnetic Resonance Imaging; LSAS: Leibowitz Social Anxiety Scale; MDD: Major Depressive Disorder; mPFC: Medio Prefrontal Cortex; MS: Mood Stabilizers; NSSI: Non Suicidal Self-Injury; PC: Patient Controls; PD: Personality Disorder; PTSD: Post-Traumatic Stress Disorder; RMIE: Reading Mind In the Eyes test; SA: Suicide Attempters; SAD: Social Anxiety Disorder; STAI: Spielberger Trait Anxiety Inventory; SUD: Substance Use Disorder; SZ: Schizophrenia.

The relationships between acute social rejection and self-stigma are still largely unknown. However, acute social exclusion results in greater social distress, more negative emotions (i.e. self-blame, more internal attributions, decreased feeling of control) and less use of functional emotion regulation strategies in people with SMI (Staebler et al., 2011; Gratz et al., 2013; Olié et al., 2017). The response to social rejection was influenced by the psychiatric diagnosis and might be stigma-related. In people with SZ, the results are controversial. Some studies supported the hypothesis that reduced sensitivity to social rejection might be related to social cognition deficits (Gradin et al., 2012) and protect against social alienation. However, another study found similar negative emotions during social exclusion, less positive emotions during social inclusion and increased negative expectations before the task (Engel et al., 2016). According to the authors, the decrease in positive emotions during inclusion was related to social anhedonia and increased negative expectations before exclusion, due to higher anticipatory social anxiety (Engel et al., 2016). Hypersensitivity to social rejection and increased paranoid beliefs during social exclusion were found in people with CHR and were mediated by difficulties with emotion regulation and self-blaming. The potential role of self-stigma and of experienced or anticipated discrimination cannot be determined, as these variables were not recorded.

Almost all studies of BPD participants found a hypersensitivity to social rejection during both social exclusion and social inclusion. This hypersensitivity was associated in some studies with enhanced P3 amplitudes and with enhanced activation within the salience network (dorsal anterior cingulate cortex (dACC), dorsal medio prefrontal cortex (dmPFC) and precuneus), supporting the hypothesis of an altered modulation of the affective response to social rejection in people with BPD (Gutz et al., 2015; Weinbrecht et al., 2018; Ruocco et al., 2010; Domsalla et al., 2014; Brown et al., 2017). Participants with BPD showed an enhanced activation of the amygdala when processing disgusted faces, this activation being associated with self-disgust. Emotion regulation difficulties were also found in people with BPD with lower amygdala deactivation and lower recruitment of the brain areas involved in cognitive reappraisal in HC (Koenigsberg et al., 2009). Self-stigma, anticipated social discrimination and stigma resistance might therefore play a role in the response to social rejection in people with BPD, but their potential role remains unknown to date. Hypersensitivity to social rejection was also found in people with MDD and SAD, with increased difficulties recovering from social rejection in those with SAD. There were no direct correlations with stigma or self-stigma.

3.3. Predictors of the response to stigma and social rejection

Socio-demographic characteristics (age, sex, ethnicity, education level), current and past psychiatric comorbidities, current psychotropic medication, insight into illness and global functioning did not appear to influence the cognitive mechanisms underlying the effects of stigma or social rejection on people with SMI (Rusch, 2007; 2010a, 2010b, 2010c, 2011a, 2011b; Ruocco et al., 2010; Staebler et al., 2011; Lawrence et al., 2011; Gratz et al., 2013; Rajj et al., 2014; Brown et al., 2017; Ernst et al., 2018; Euler et al., 2018). Group identification, implicit shame-related self-associations and implicit self-guilt did however contribute to self-stigma and to a lesser ability to resist stigma (Rusch, 2010a, 2010b, 2010c). The endorsement of genetic causal attributions about mental illness increased implicit self-guilt in individuals with SMI (Rusch 2010a) which has implications when designing anti-stigma campaigns or psychoeducation interventions for people with SMI (Rusch, 2010a). The psychiatric diagnosis did not influence implicit stigma-related self-negative associations in people with SMI (Rusch 2010a, 2010b, 2010c, 2011a). It did however influence the response to social rejection (Rusch, 2007; Staebler et al., 2011; Renneberg et al., 2012; Gratz et al., 2013; Gutz et al., 2015, 2016; Domsalla et al., 2014; De Panfilis, 2015; Engel et al., 2016; Weinbrecht et al., 2018). This concurs with the literature on explicit self-stigma, which is more

frequent in people with SZ (Brohan et al., 2010, 2011) but affects clinical and psychosocial outcomes in all people with SMI (Livingston and Boyd, 2010). Psychiatric symptoms influenced self-perceptions as symptomatic BD participants had more implicit negative self-associations (Jabben et al., 2014). The number of BPD features influenced social-cognitive performance after social rejection in people with BPD and the severity of positive symptoms affected the response to social rejection in participants with SZ (Gradin et al., 2012). Illness duration affected negatively stigma resistance (Rajj et al., 2014). There were no significant differences between inpatients and outpatients with SZ in their response to social rejection (Engel et al., 2016). The participants' social networks did not influence the response to social exclusion in people with BPD (Gratz et al., 2013). Baseline non-specific distress was associated with elevated distress during social exclusion (Gratz et al., 2013) suggesting that chronic social rejection might affect the response to acute social rejection and contribute to social alienation (Gutz et al., 2016). A history of trauma influenced the biological and behavioural response to social rejection (Jobst et al., 2015, 2016; Ernst et al., 2018) and was associated with more shame-related self-associations in people with BPD (Rusch et al., 2011a). However, emotion processing when confronted with disgusted faces (Schienle et al., 2015) and when distancing from negative social cues (Koenigsberg et al., 2009) were not affected by a history of trauma. A history of suicidal acts influenced the response to social rejection in people with SMI (Olié et al., 2017).

4. Discussion

Overall, the results of the present review can be summarized as follows: Acute social rejection and its consequences on individuals with SMI were the most documented area (25 studies). Hypersensitivity to acute social rejection was found in almost all the studies including people with SMI and was associated with more self-related negative emotions and with the use of less functional and more dysfunctional emotion regulation strategies. Enhanced activation within the salience network was associated with hypersensitivity to acute social rejection. Hypersensitivity to social rejection resulted in an increase in paranoid beliefs in people with CHR. Reduced sensitivity to acute social rejection and reduced activation within the salience network were, however, found in people with SZ and could be related to social cognition deficits or act as a protection mechanism against social alienation. Individuals with SMI under-performed in cognitive and social cognitive tasks when placed in stigma-related environments. Stigma resistance was associated with enhanced activation of the rostro-ventral medio prefrontal cortex (rvmPFC) and decreased activation of the amygdala in people with SZ. Psychiatric symptoms, illness duration and baseline non-specific distress influenced the response to acute social rejection or stigma. The potential role of chronic social discrimination and its effects on people with SMI should be further investigated.

The existing body of research on the cognitive neuroscience of stigma in individuals with SMI has several limitations. Only a few studies directly addressed the consequences of self-stigma on people with SMI. The samples, methods, experimental paradigms and reported outcomes used in the studies were heterogeneous and therefore difficult to compare. The results mostly came from individual studies that have not yet been replicated. The cognitive and neural correlates of personal stigma remain largely unknown and should therefore be further investigated. Neuroimaging studies may add to the knowledge about the neural mechanisms underlying personal stigma and stigma resistance. Personal stigma is however a complex construct difficult to model precisely in cognitive neuroscience. Studying the effects of experienced and anticipated social discrimination on brain functional connectivity and biological inflammatory response in persons with SMI might improve the understanding of the mechanisms underlying personal stigma but future research will be needed.

4.1. Relationships between acute social discrimination and self-stigma

The relationships between acute social rejection and self-stigma are unclear, as most studies did not record self-stigma. Implicit and explicit self-stigma predicted subjective quality of life (Rusch, 2010c; Livingston and Boyd, 2010). As expected, individuals with SMI underperformed in cognitive and social cognitive tasks under stigma-related conditions (stereotype threat or acute social rejection). This supports previous findings that explicit self-stigma is associated with lower cognitive, social and vocational functioning (Yanos et al., 2010; Hill and Startup, 2013). Individuals with SMI showed higher rejection expectancy and anticipatory anxiety before social exclusion. They had also more self-related negative emotions during social exclusion and an increased tendency to use avoidant strategies after being excluded when compared with HC (Gratz et al., 2013; Gutz et al., 2015,2016; Engel et al., 2016). This may back up previous research where anticipated social discrimination mediated the relationship between experienced discrimination and self-stigma in individuals with SMI (Quinn et al., 2015), and where self-stigma predicted social anxiety in people with SZ (Lysaker et al., 2010) and was associated with avoidant coping strategies (Vauth et al., 2007). Anticipated social discrimination and self-stigma may influence the response to social rejection and their potential role should therefore be investigated further. In people with CHR, social exclusion was associated with hypersensitivity to acute social rejection and increased paranoid beliefs that were mediated by emotion regulation difficulties and self-blaming. This might be related to self-stigma, associated with hypersensitivity to social rejection and paranoid delusions in a cognitive model of self-stigma in psychosis (Wood et al., 2017). Two studies found reduced sensitivity to social rejection and reduced activation within the salience network in people with SZ (Gradin et al., 2012). This might relate to social cognition deficits (Gradin et al., 2012) and protect against social alienation. It might, however, also more generally reflect the salience network dysfunction described in people with SZ (Palaniyappan et al., 2012). The “insight paradox” described by P. Lysaker (association between depression and good insight) and its relationships with social cognition and emotion regulation abilities might also explain the differences observed in people with SZ (Lysaker et al., 2007,2013). The presence (or the absence) of metacognitive deficits impeding one's abilities to deflect negative beliefs about oneself can also be a potential explaining factor (Nabors et al., 2014). The relationships between social cognition, metacognitive abilities, self-stigma and the response to acute social rejection should therefore be further investigated.

Although very few studies reported on self-stigma in BPD, there is some evidence supporting its potential role in moderating the response to acute social rejection. Participants with BPD showed an enhanced activation of the amygdala when processing disgusted faces, this activation being associated with self-disgust. As implicit self-disgust contributed to self-stigma by increasing the perceived legitimacy of discrimination and by decreasing the ability to resist stigma (Rusch, 2010b), this hypersensitivity to social rejection in people with BPD might be stigma-related. Emotion regulation difficulties in people with BPD were associated with lower deactivation of the amygdala and lower recruitment of the brain areas involved in cognitive reappraisal in HC (Koenigsberg et al., 2009). Although affect dysregulation is a key feature of BPD, it has been suggested that this mechanism might also be related to the internalization of psychiatric disorders or early-life maltreatment (Gunderson et al., 2018). In contrast stigma resistance was associated with preserved cognitive performance in people with SZ. The f-MRI showed SR to be associated with enhanced activation of the rostro-ventral medio prefrontal cortex (rvmPFC) and decreased amygdala activation in people with SZ after controlling for global functioning, insight and depressive symptoms (Raij et al., 2014). As the rvmPFC was identified as a regulator of the affective response to social exclusion (Eisenberger, 2004) this could be one of the mechanisms behind stigma resistance (Raij et al., 2014). Self-stigma, anticipated

social discrimination, stigma resistance, and social cognition and emotion regulation ability might influence the response to acute social rejection and their potential role should be investigated in future studies.

4.2. Implications for future research

Several predictors of the response to stigma or acute social rejection were identified. Self-stigma was influenced by group identification, implicit shame-related self-associations and implicit self-guilt, but not by psychiatric diagnosis, psychiatric comorbidities or socio-demographic characteristics. The association between implicit shame-related self-associations and self-stigma is also found in a previous study where explicit shame-proneness was identified as a mediator between insight and self-stigma (Hasson-Ohayon et al., 2012). Stigma resistance was negatively affected by illness duration in accordance with previous results (Sibitz et al., 2011a). The response to social rejection was not influenced by the participants' status or social network in people with SZ and BPD (Gratz et al., 2013; Engel et al., 2016). Further investigation is however needed, as inpatient status and reduced social network were associated with decreased stigma resistance in people with SZ (Sibitz et al., 2011a,2011b). The role of the psychiatric diagnosis in moderating the response to acute social rejection remains unclear, as all the studies were diagnosis-specific. The severity of the psychiatric symptoms influenced the response to acute social rejection. The number of BPD features influenced social-cognitive performance after social rejection in people with BPD and the severity of positive symptoms affected the response to social rejection in participants with SZ (Gradin et al., 2012). This might be stigma-related as self-blaming, social threat processing dysfunctions and self-stigma can underpin paranoid beliefs in people with psychosis (Corlett et al., 2010; Wood et al., 2017) and explicit self-stigma is associated with increased severity of positive and depressive symptoms (Livingston and Boyd, 2010; Cerit et al., 2012).

Baseline non-specific distress was associated with elevated distress during social exclusion (Gratz et al., 2013) suggesting that chronic social rejection might affect the response to acute social rejection and contribute to the development of social alienation (Gutz, 2016). However, research on chronic social rejection is extremely limited in HC and non-existent in people with SMI. Chronic social rejection was investigated in people without SMI, but confronted with prolonged ostracism (social rejection > 3-months; Williams, 2009; Riva et al., 2014). In contrast to acute social rejection, chronic social rejection was associated with feelings of resignation, helplessness, unworthiness and low self-worth and ultimately with social alienation and depression (Williams, 2009; Riva et al., 2014). An increased sensitivity to acute social exclusion was also found in people suffering from chronic social rejection (Williams, 2009; Riva et al., 2014). In people at risk of being ostracized, enhanced spontaneous amygdala resting-state activity and increased connectivity with the salience network were associated with a greater exposure to discrimination, independently of ethnicity, sex or current psychiatric comorbidities. Elevated pro-inflammatory cytokines (IL-6 and TNF-alpha) were associated with enhanced dorsal ACC and anterior insula activation after acute social rejection in HC (Slavich et al., 2010). Exposure to everyday social stressors and to social rejection was associated in HC with elevated inflammatory activity and with MDD (Slavich et al., 2010, 2014; Beatty et al., 2014; Van Dyke, 2017). Elevated CRP levels were found in people with SZ, BD or MDD (Miller et al., 2014; Horsdal et al., 2017) and associated with more severe depressive symptoms in people with SZ (Faugere et al., 2018). However, the correlations between inflammatory activity, experiences of discrimination and self-stigma in people with SMI have not been investigated to date. Future research should therefore investigate the potential role of current, past and anticipated social discrimination in the modulation of biological and neural responses to stigma or acute social rejection in people with SMI.

In conclusion, the studies conducted in the last decade have improved our understanding of the cognitive and neural correlates of stigma, self-stigma and acute social rejection in people with SMI. Several potential mechanisms and predictors have been identified and should be further documented. Future research should investigate the associations between self-stigma, performance under ST conditions and the responses to acute and chronic social rejection in people with SMI. Chronic social rejection in people with mental illness and its cognitive, biological and neural correlates should also be further investigated.

Acknowledgement

The authors would like to thank Mrs. Kim Barrett for proofreading the manuscript.

Role of the funding source

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interests

None.

References

- Angermeyer, M.C., Millier, A., Rémuzat, C., Refai, T., Toumi, M., 2013. Attitudes and beliefs of the French public about schizophrenia and major depression: results from a vignette-based population survey. *BMC Psychiatry* 13, 313. <https://doi.org/10.1186/1471-244X-13-313>.
- Beatty, D.L., Matthews, K.A., Bromberger, J.T., Brown, C., 2014. Everyday discrimination prospectively predicts inflammation across 7-Years in racially diverse midlife women: study of women's health across the nation. *J. Soc. Issues* 70 (2), 298–314.
- Bolling, D.Z., Pitskel, N.B., Deen, B., Crowley, M.J., McPartland, J.C., Kaiser, M.D., Wyk, B.C., Wu, J., Mayes, L.C., Pelphrey, K.A., 2011. Enhanced neural responses to rule violation in children with autism: a comparison to social exclusion. *Dev. Cogn. Neurosci.* 1 (3), 280–294. <https://doi.org/10.1016/j.dcn.2011.02.002>.
- Brohan, E., Elgie, R., Sartorius, N., Thornicroft, G., GAMIAN-Europe Study Group., 2010. Self-stigma, empowerment and perceived discrimination among people with schizophrenia in 14 European countries: the GAMIAN-Europe study. *Schizophr. Res.* 122 (1–3), 232–238. <https://doi.org/10.1016/j.schres.2010.02.1065>.
- Brohan, E., Gauci, D., Sartorius, N., Thornicroft, G., GAMIAN-Europe Study Group., 2011. Self-stigma, empowerment and perceived discrimination among people with bipolar disorder or depression in 13 European countries: the GAMIAN-Europe study. *J. Affect. Disord.* 129 (1–3), 56–63. <https://doi.org/10.1016/j.jad.2010.09.001>.
- Brown, R.C., Plener, P.L., Groen, G., Neff, D., Bonenberger, M., Abler, B., 2017. Differential neural processing of social exclusion and inclusion in adolescents with non-suicidal self-injury and young adults with borderline personality disorder. *Front. Psychiatry* 8, 267. <https://doi.org/10.3389/fpsyg.2017.00267>.
- Bungert, M., Koppe, G., Niedtfield, I., Vollstädt-Klein, S., Schmahl, C., Lis, S., Bohus, M., 2015. Pain processing after social exclusion and its relation to rejection sensitivity in borderline personality disorder. *PLoS One* 10 (8), e0133693. <https://doi.org/10.1371/journal.pone.0133693>.
- Cerit, C., Filizer, A., Tural, Ü., Tufan, A.E., 2012. Stigma: a core factor on predicting functionality in bipolar disorder. *Compr. Psychiatry* 53 (5), 484–489. <https://doi.org/10.1016/j.comppsy.2011.08.010>. Epub 2011 Oct 27.
- Corlett, P.R., Taylor, J.R., Wang, X.J., Fletcher, P.C., Krystal, J.H., 2010. Toward a neurobiology of delusions. *Prog. Neurobiol.* 92, 345–369.
- Corrigan, P.W., Watson, A.C., 2002. Understanding the impact of stigma on people with mental illness. *World Psychiatry* 1 (1), 16–20.
- De Panfilis, C., Riva, P., Preti, E., Cabrino, C., Marchesi, C., 2015. When social inclusion is not enough: implicit expectations of extreme inclusion in borderline personality disorder. *Personal. Disord. Theory Res. Treat.* 6 (4), 301–309. <https://doi.org/10.1177/1368430207088036>.
- Derks, B., Inzlicht, M., Kang, S., 2008. The neuroscience of stigma and stereotype threat. *Group Process. Intergroup Relat.* 11 (2), 163–181. <https://doi.org/10.1177/1368430207088036>.
- Domsalla, M., Koppe, G., Niedtfield, I., Vollstädt-Klein, S., Lis, S., 2014. Cerebral processing of social rejection in patients with borderline personality disorder. *Scan* 9, 1789–1797. <https://doi.org/10.1093/scan/nst176>.
- Eisenberger, N.I., Lieberman, M.D., 2004. Why rejection hurts: a common neural alarm system for physical and social pain. *Trends Cogn. Sci.* 8 (7), 294–300. <https://doi.org/10.1016/j.tics.2004.05.010>.
- Ellison, N., Mason, O., Scior, K., 2013. Bipolar disorder and stigma: a systematic review of the literature. *J. Affect. Disord.* 151 (3), 805–820. <https://doi.org/10.1016/j.jad.2013.08.014>.
- Engel, M., Fritzsche, A., Lincoln, T.M., 2016. Anticipation and experience of emotions in patients with schizophrenia and negative symptoms. An experimental study in a social context. *Schizophr. Res.* 170 (1), 191–197. <https://doi.org/10.1016/j.schres.2015.11.028>.
- Ernst, M., Mohr, H.M., Schhött, M., Rickmeyer, C., Fischmann, T., Leuzinger-Bohleber, M., Weiß, H., Grabhorn, R., 2018. The effects of social exclusion on response inhibition in borderline personality disorder and major depression. *Psychiatry Res.* 262, 333–339. <https://doi.org/10.1016/j.psychres.2017.03.034>.
- Euler, S., Wrege, J., Busmann, M., Lindenmeyer, H.J., Sollberger, D., Lang, U.E., Gaab, J., Walter, M., 2018. Exclusion-proneness in borderline personality disorder impairs alliance in mentalization-based group therapy. *Front. Psychol.* 9, 824. <https://doi.org/10.3389/fpsyg.2018.00824>.
- Farrelly, S., Clement, S., Gabbidon, J., Jeffery, D., Dockery, L., Lassman, F., Brohan, E., Henderson, R.C., Williams, P., Howard, L.M., Thornicroft, G., MIRIAD study group., 2014. Anticipated and experienced discrimination amongst people with schizophrenia, bipolar disorder and major depressive disorder: a cross sectional study. *BMC Psychiatry* 14, 157. <https://doi.org/10.1186/1471-244X-14-157>.
- Faugere, M., Micoulaud-Franchi, J.A., Faget-Agius, C., Lançon, C., Cermolacce, M., Richieri, R., 2018. High C-reactive protein levels are associated with depressive symptoms in schizophrenia. *J. Affect. Disord.* 225, 671–675. <https://doi.org/10.1016/j.jad.2017.09.004>.
- Firmin, R.L., Luther, L., Lysaker, P.H., Minor, K.S., Salyers, M.P., 2016. Stigma resistance is positively associated with psychiatric and psychosocial outcomes: a meta-analysis. *Schizophr. Res.* 175 (1–3), 118–128. <https://doi.org/10.1016/j.schres.2016.03.008>.
- Gerlinger, G., Hauser, M., De Hert, M., Lacluyse, K., Wampers, M., Correll, C.U., 2013. Personal stigma in schizophrenia spectrum disorders: a systematic review of prevalence rates, correlates, impact and interventions. *World Psychiatry* 12 (2), 155–164. <https://doi.org/10.1002/wps.20040>.
- Gradin, V.B., Waiter, G., Kumar, P., Stickle, C., Milders, M., Matthews, K., Reid, I., Hall, J., Steele, J.D., 2012. Abnormal neural responses to social exclusion in schizophrenia. *PLoS One* 7 (8), e42608. <https://doi.org/10.1371/journal.pone.0042608>.
- Gratz, K.L., Dixon-Gordon, K.L., Breetz, A., Tull, M., 2013. A laboratory-based examination of responses to social rejection in borderline personality disorder: the mediating role of emotion dysregulation. *J. Pers. Disord.* 27 (3), 426. <https://doi.org/10.1521/pe.2013.27.2.157>.
- Greenwald, A.G., Poehlman, T.A., Uhlmann, E.L., Banaji, M.R., 2009. Understanding and using the Implicit Association Test: III. Meta-analysis of predictive validity. *J. Pers. Soc. Psychol.* 97 (1), 17–41. <https://doi.org/10.1037/a0015575>.
- Gunderson, J.G., Herpertz, S.C., Skodol, A.E., Torgersen, S., Zanarini, M.C., 2018. Borderline personality disorder. *Nat. Rev. Dis. Primers* 4, 18029. <https://doi.org/10.1038/nrdp.2018.29>.
- Gutz, L., Renneberg, B., Roepke, S., Niedeggen, M., 2015. Neural processing of social participation in borderline personality disorder and social anxiety disorder. *J. Abnorm. Psychol.* 124 (2), 421–431. <https://doi.org/10.1037/a0038614>.
- Gutz, L., Roepke, S., Renneberg, B., 2016. Cognitive and affective processing of social exclusion in borderline personality disorder and social anxiety disorder. *Behav. Ther.* 87, 70–75. <https://doi.org/10.1016/j.brat.2016.08.020>.
- Hartergrink, C.H., van Beest, I., Wicherts, J.M., Williams, K.D., 2015. The ordinal effects of ostracism: a meta-analysis of 120 Cyberball studies. *PLoS One* 10 (5), e0127002. <https://doi.org/10.1371/journal.pone.0127002>.
- Hasson-Ohayon, I., Ehrlich-Ben Or, S., Vahab, K., Amiaz, R., Weiser, M., Roe, D., 2012. Insight into mental illness and self-stigma: the mediating role of shame proneness. *Psychiatry Res.* 200 (2–3), 802–806. <https://doi.org/10.1016/j.psychres.2012.07.038>.
- Heenan, A., Best, M.W., Ouellette, S.J., Meiklejohn, E., Troje, N.F., Bowie, C.R., 2014. Assessing threat responses towards the symptoms and diagnosis of schizophrenia using visual perceptual biases. *Schizophr. Res.* 159 (1), 238–242. <https://doi.org/10.1016/j.schres.2014.07.024>.
- Heeren, A., Dricot, L., Billieux, J., Philippot, P., Grynberg, D., de Timary, P., Maurage, P., 2017. Correlates of social exclusion in social anxiety disorder: an fMRI study. *Sci. Rep.* 7 (1), 260. <https://doi.org/10.1038/s41598-017-00310-9>.
- Henry, J.D., von Hippel, C., Shapiro, L., 2010. Stereotype threat contributes to social difficulties in people with schizophrenia. *Br. J. Clin. Psychol.* 49 (1), 31–41. <https://doi.org/10.1348/014466509x421963>.
- Hill, K., Startup, M., 2013. The relationship between internalized stigma, negative symptoms and social functioning in schizophrenia: the mediating role of self-efficacy. *Psychiatry Res.* 206 (2–3), 151–157. <https://doi.org/10.1016/j.psychres.2012.09.056>.
- Horsdal, H.T., Köhler-Forsberg, O., Benros, M.E., Gasse, C., 2017. C-reactive protein and white blood cell levels in schizophrenia, bipolar disorders and depression - associations with mortality and psychiatric outcomes: a population-based study. *Eur. Psychiatry* 44, 164–172. <https://doi.org/10.1016/j.eurpsy.2017.04.012>.
- Jabben, N., de Jong, P.J., Kupka, R.W., Gashouwer, K.A., Nolen, W.A., Penninx, B.W., 2014. Implicit and explicit self-associations in bipolar disorder: a comparison with healthy controls and unipolar depressive disorder. *Psychiatry Res.* 215 (2), 329–334. <https://doi.org/10.1016/j.psychres.2013.11.030>.
- Jobst, A., Albert, A., Bauriedl-Schmidt, C., Mauer, M.C., Renneberg, B., Buchheim, A., Sabass, L., Falkai, P., Zill, P., Padberg, F., 2014. Social exclusion leads to divergent changes of oxytocin levels in borderline patients and healthy subjects. *Psychother. Psychosom.* 83 (4), 252–254. <https://doi.org/10.1159/000358526>.
- Jobst, A., Sabass, L., Palagyi, A., Bauriedl-Schmidt, C., Mauer, M.C., Sarubin, N., Buchheim, A., Renneberg, B., Falkai, P., Zill, P., Padberg, F., 2015. Effects of social exclusion on emotions and oxytocin and cortisol levels in patients with chronic depression. *J. Psychiatr. Res.* 60, 170–177. <https://doi.org/10.1016/j.jpsychires.2014.11.001>.
- Jobst, A., Padberg, F., Mauer, M.C., Daltrozio, T., Bauriedl-Schmidt, C., Sabass, L., Sarubin, N., Falkai, P., Renneberg, B., Zill, P., Gander, M., Buchheim, A., 2016. Lower Oxytocin Plasma Levels in Borderline Patients with Unresolved Attachment

- Representations. *Front. Hum. Neurosci.* 10, 125. <https://doi.org/10.3389/fnhum.2016.00125>.
- Kawamoto, T., Onoda, K., Nakashima, K., Nittono, H., Yamaguchi, S., Ura, M., 2012. Is dorsal anterior cingulate cortex activation in response to social exclusion due to expectancy violation? An fMRI study. *Front. Evol. Neurosci.* 4, 11. <https://doi.org/10.3389/fnevo.2012.00011>.
- Koenigsberg, H.W., Fan, J., Ochsner, K.N., Lui, X., Guise, K.G., Pizzarello, S., Dorantes, C., Guerrerri, S., Tecuta, L., Goodman, M., New, A., Siever, L.J., 2009. Neural correlates of the use of psychological distancing to regulate responses to negative social cues: a study of patients with borderline personality disorder. *Biol. Psychiatry* 66 (9), 854–863. <https://doi.org/10.1016/j.biopsych.2009.06.010>.
- Krendl, A.C., Macrae, C.N., Kelley, W.M., Fugelsang, J.A., Heatherton, T.F., 2006. The good, the bad, and the ugly: an fMRI investigation of the functional anatomical correlates of stigma. *Soc. Neurosci.* 1 (1), 5–15. <https://doi.org/10.1080/17470910600670579>.
- Krendl, A.C., Moran, J.M., Ambady, N., 2013. Does context matter in evaluations of stigmatized individuals? An fMRI study. *Soc. Cogn. Affect. Neurosci.* 8 (5), 602–608. <https://doi.org/10.1093/scan/nns037>.
- Kumar, P., Waiter, G.D., Dubois, M., Milders, M., Reid, I., Steele, J.D., 2017. Increased neural response to social rejection in major depression. *Depress. Anxiety* 34 (11), 1049–1056. <https://doi.org/10.1002/da.22665>.
- Lasalva, A., Zoppi, S., Van Bortel, T., Bonetto, C., Cristofalo, D., Wahlbeck, K., Bacle, S.V., Van Audenhove, C., van Weeghel, J., Reneses, B., Germanavicius, A., Economou, M., Lanfredi, M., Ando, S., Sartorius, N., Lopez-Ibor, J.J., Thornicroft, G., ASPEN/INDIGO Study Group., 2013. Global pattern of experienced and anticipated discrimination reported by people with major depressive disorder: a cross-sectional survey. *Lancet* 381, 55–62.
- Lawrence, K.A., Chanan, A.M., Allen, J.S., 2011. The effect of ostracism upon mood in youth with Borderline Personality Disorder. *J. Personal. Disord.* 25 (5), 702–714. <https://doi.org/10.1521/pedi.2011.25.5.702>.
- Livingston, J.D., Boyd, J.E., 2010. Correlates and consequences of internalized stigma for people living with mental illness: a systematic review and meta-analysis. *Soc. Sci. Med.* 71 (12), 2150–2161. <https://doi.org/10.1016/j.socscimed.2010.09.030>.
- Lysaker, P.H., Roe, D., Yanos, P.T., 2007. Toward understanding the insight paradox: internalized stigma moderates the association between insight and social functioning, hope, and self-esteem among people with schizophrenia spectrum disorders. *Schizophr. Bull.* 33 (1), 192–199.
- Lysaker, P.H., Yanos, P.T., Outcalt, J., Roe, D., 2010. Association of stigma, self-esteem, and symptoms with concurrent and prospective assessment of social anxiety in schizophrenia. *Clin. Schizophr. Relat. Psychoses* 4 (1), 41–48. <https://doi.org/10.3371/CSRP.4.1.3>.
- Lysaker, P.H., Vohs, J., Hasson-Ohayon, I., Kukla, M., Wierwille, J., Dimaggio, G., 2013. Depression and insight in schizophrenia: comparisons of levels of deficits in social cognition and metacognition and internalized stigma across three profiles. *Schizophr. Res.* 148 (1–3), 18–23. <https://doi.org/10.1016/j.schres.2013.05.025>.
- Miller, B.J., Culpapper, N., Rapaport, M.H., 2014. C-reactive protein levels in schizophrenia: a review and meta-analysis. *Clin. Schizophr. Relat. Psychoses* 7 (4), 223–230. <https://doi.org/10.3371/CSRP.MICU.020813>.
- Moor, B.G., Güroğlu, B., Op de Macks, Z.A., Serge, A.R.B., Rombouts, S.A.R.B., Van der Molen, M.W., Crone, E.A., 2012. Social exclusion and punishment of excluders: neural correlates and developmental trajectories. *NeuroImage* 59 (2012), 708–717. <https://doi.org/10.1016/j.neuroimage.2011.07.02814>.
- Moritz, S., Spirandelli, K., Happach, I., Lion, D., Berna, F., 2018. Dysfunction by disclosure? Stereotype threat as a source of secondary neurocognitive malperformance in obsessive-compulsive disorder. *J. Int. Neuropsychol. Soc.* 24 (6), 584–592. <https://doi.org/10.1017/S1355617718000097>.
- Nabors, L.M., Yanos, P.T., Roe, D., Hasson-Ohayon, I., Leonhardt, B.L., Buck, K.D., Lysaker, P.H., 2014. Stereotype endorsement, metacognitive capacity, and self-esteem as predictors of stigma resistance in persons with schizophrenia. *Compr. Psychiatry* 55, 792–798. <https://doi.org/10.1016/j.comppsy.2014.01.011>.
- Olié, E., Jollant, F., Deverduin, J., de Champfleury, N.M., Cyprien, F., Le Bars, E., Mura, T., Bonafé, A., Courtret, P., 2017. The experience of social exclusion in women with a history of suicidal acts: a neuroimaging study. *Sci. Rep.* 7 (1), 89. <https://doi.org/10.1038/s41598-017-00211-x>.
- Palaniyappan, L., White, T.P., Liddle, P.F., 2012. The concept of salience network dysfunction in schizophrenia: from neuroimaging observations to therapeutic opportunities. *Curr. Top. Med. Chem.* 12 (21), 2324–2338.
- Penn, D.L., Kohlmaier, J.R., Corrigan, P.W., 2000. Interpersonal factors contributing to the stigma of schizophrenia: social skills, perceived attractiveness, and symptoms. *Schizophr. Res.* 45 (1–2), 37–45.
- Peris, T.S., Teachman, B.A., Nosek, B.A., 2008. Implicit and explicit stigma of mental illness: links to clinical care. *J. Nerv. Ment. Dis.* 196 (10), 752–760. <https://doi.org/10.1097/NMD.0b013e3181879df>.
- Quinn, D.M., Williams, M.K., Weisz, B.M., 2015. From discrimination to internalized mental illness stigma: the mediating roles of anticipated discrimination and anticipated stigma. *Psychiatr. Rehabil. J.* 38 (2), 103–108. <https://doi.org/10.1037/prj0000136>.
- Raij, T.T., Korkeila, J., Joutsenniemi, K., Saarni, S.I., Riekk, T.J., 2014. Association of stigma resistance with emotion regulation - functional magnetic resonance imaging and neuropsychological findings. *Compr. Psychiatry* 55 (3), 727–735. <https://doi.org/10.1016/j.comppsy.2013.10.010>.
- Reihl, K.M., Hurley, R.A., Taber, K.H., 2015. Neurobiology of implicit and explicit bias: implications for clinicians. *J. Neuropsychiatr. Clin. Neurosci.* 27 (4), A6–253. <https://doi.org/10.1176/appi.neuropsych.15080212>.
- Renneberg, B., Herm, K., Hahn, A., Staebler, K., Lammers, C.H., Roepke, S., 2012. Perception of social participation in borderline personality disorder. *Clin. Psychol. Psychother.* 19, 473–480. <https://doi.org/10.1002/cpp.772>.
- Riva, P., Wesselmann, E.D., Wirth, J.H., Carter-Sowell, A.R., Williams, K.D., 2014. When pain does not heal: the common antecedents and consequences of chronic social and physical pain. *Basic Appl. Soc. Psychol.* 36, 329–346. <https://doi.org/10.1080/01973533.2014.917975>.
- Ruocco, A.C., Medaglia, J.D., Tinker, J.R., Ayaz, H., Forman, E.M., Newman, C.F., Williams, J.M., Hillary, F.G., Platek, S.M., Onaral, B., Chute, D.L., 2010. Medial prefrontal cortex hyperactivation during social exclusion in borderline personality disorder. *Psychiatry Res. Neuroimag.* 181 (3), 233–236. <https://doi.org/10.1016/j.psychres.2009.12.001>.
- Rüsch, N., Lieb, K., Göttler, I., Hermann, C., Schramm, E., Richter, H., Jacob, G.A., Corrigan, P.W., Bohus, M., 2007. Shame and implicit self-concept in women with borderline personality disorder. *Am. J. Psychiatry* 164 (3), 500–508. <https://doi.org/10.1176/ajp.2007.164.3.500>.
- Rüsch, N., Todd, A.R., Bodenhausen, G.V., Olschewski, M., Corrigan, P.W., 2010a. Automatically activated shame reactions and perceived legitimacy of discrimination: a longitudinal study among people with mental illness. *J. Behav. Ther. Exp. Psychiatry* 41 (1), 60–63. <https://doi.org/10.1016/j.jbtep.2009.10.002>.
- Rüsch, N., Todd, A.R., Bodenhausen, G.V., Corrigan, P.W., 2010b. Biogenetic models of psychopathology, implicit guilt, and mental illness stigma. *Psychiatry Res.* 179 (3), 328–332. <https://doi.org/10.1016/j.psychres.2009.09.010>.
- Rüsch, N., Corrigan, P.W., Todd, A.R., Bodenhausen, G.V., 2010c. Implicit self-stigma in people with mental illness. *J. Nerv. Ment. Dis.* 198 (2), 150–153. <https://doi.org/10.1097/NMD.0b013e3181cc43b5>.
- Rüsch, N., Schulz, D., Valerius, G., Steil, R., Bohus, M., Schmahl, C., 2011a. Disgust and implicit self-concept in women with borderline personality disorder and posttraumatic stress disorder. *Eur. Arch. Psychiatry Clin. Neurosci.* 261 (5), 369–376. <https://doi.org/10.1007/s00406-010-0174-2>.
- Rüsch, N., Corrigan, P.W., Todd, A.R., Bodenhausen, G.V., 2011b. Automatic stereotyping against people with schizophrenia, schizoaffective and affective disorders. *Psychiatry Res.* 186 (1), 34–39. <https://doi.org/10.1016/j.psychres.2010.08.024>.
- Schienze, A., Wabnegger, A., Schöngassner, F., Leutgeb, V., 2015. Effects of personal space intrusion in affective contexts: an fMRI investigation with women suffering from borderline personality disorder. *Soc. Cogn. Affect. Neurosci.* 10 (10), 1424–1428. <https://doi.org/10.1093/scan/nsv034>.
- Shapiro, J.R., Neuberger, S.L., 2007. From stereotype threat to stereotype threats: implications of a multi-threat framework for causes, moderators, mediators, consequences, and interventions. *Pers. Soc. Psychol. Rev.* 11 (2), 107–130. <https://doi.org/10.1177/1088868306294790>.
- Shapiro, J.R., 2011. Different groups, different threats: a multi-threat approach to the experience of stereotype threats. *Pers. Soc. Psychol. Bull.* 37 (4), 464–480. <https://doi.org/10.1177/0146167211398140>.
- Sibitz, I., Amering, M., Unger, A., Seyringer, M.E., Bachmann, A., Schrank, B., Benesch, T., Schulze, B., Woppmann, A., 2011a. The impact of the social network, stigma and empowerment on the quality of life in patients with schizophrenia. *Eur. Psychiatry* 26 (1), 28–33. <https://doi.org/10.1016/j.eurpsy.2010.08.010>.
- Sibitz, I., Unger, A., Woppmann, A., Zidek, T., Amering, M., 2011b. Stigma resistance in patients with schizophrenia. *Schizophr. Bull.* 37 (2), 316–323. <https://doi.org/10.1093/schbul/sbp048>.
- Slavich, G.M., Way, B.M., Eisenberger, N.I., Taylor, S.E., 2010. Neural sensitivity to social rejection is associated with inflammatory responses to social stress. *Proc. Natl. Acad. Sci. USA.* 107 (33), 14817–14822. <https://doi.org/10.1073/pnas.1009164107>.
- Slavich, G.M., Irwin, M.R., 2014. From stress to inflammation and major depressive disorder: a social signal transduction theory of depression. *Psychol. Bull.* 140 (3), 774–815. <https://doi.org/10.1037/a0035302>.
- Staebler, K., Renneberg, B., Stopsack, M., Fiedler, P., Weiler, M., Roepke, S., 2011. Facial emotional expression in reaction to social exclusion in borderline personality disorder. *Psychol. Med.* 41 (9), 1929–1938. <https://doi.org/10.1017/S003291711000080>.
- Stull, L.G., McGrew, J.H., Salyers, M.P., Ashburn-Nardo, L., 2013. Implicit and explicit stigma of mental illness: attitudes in an evidence-based practice. *J. Nerv. Ment. Dis.* 201 (12), 1072–1079. <https://doi.org/10.1097/NMD.0000000000000056>.
- Substance Abuse and Mental Health Services Administration (SAMHSA), 2014. *Results from the 2013 National Survey on Drug Use and Health: mental Health Findings*. NSDUH Series H-49, HHS Publication No. (SMA) 14-4887. Substance Abuse and Mental Health Services Administration, Rockville, MD.
- Teachman, B.A., Wilson, J.G., Komarovskaya, I., 2006. Implicit and explicit stigma of mental illness in diagnosed and healthy samples. *J. Soc. Clin. Psychol.* 25 (1), 75–95.
- Thoits, P.A., 2011. Mechanisms linking social ties and support to physical and mental health. *J. Health. Soc. Behav.* 52 (2), 145–161. <https://doi.org/10.1177/0022146510395592>.
- Thornicroft, G., Brohan, E., Rose, D., Sartorius, N., Leese, M., INDIGO Study Group, 2009. Global pattern of experienced and anticipated discrimination against people with schizophrenia: a cross-sectional survey. *Lancet* 373 (9661), 408–415. [https://doi.org/10.1016/S0140-6736\(08\)61817-6](https://doi.org/10.1016/S0140-6736(08)61817-6).
- Van Dyke, M.E., Vaccarino, V., Dunbar, S.B., Pemu, P., Gibbons, G.H., Quyyumi, A.A., Lewis, T.T., 2017. Socioeconomic status discrimination and C-reactive protein in African-American and White adults. *Psychoneuroendocrinology* 82, 9–16. <https://doi.org/10.1016/j.psychneuen.2017.04.009>.
- Vauth, R., Kleim, B., Wirtz, M., Corrigan, P.W., 2007. Self-efficacy and empowerment as outcomes of self-stigmatizing and coping in schizophrenia. *Psychiatry Res.* 150 (1), 71–80.
- Weinbrecht, A., Niedeggen, M., Roepke, S., Renneberg, B., 2018. Feeling excluded no matter what? Bias in the processing of social participation in borderline personality disorder. *Neuroimage Clin.* 19, 343–350. <https://doi.org/10.1016/j.nicl.2018.04.031>.

- Williams, K.D., 2009. Ostracism: a temporal need-threat model. *Adv. Exp. Soc. Psychol.* 41, 275–314.
- Wittenbrink, B., Judd, C.M., Park, B., 2001. Evaluative versus conceptual judgments in automatic stereotyping and prejudice. *J. Exp. Soc. Psychol.* 37, 244–252.
- Wood, L., Byrne, R., Varese, F., Morrison, A.P., 2016. Psychosocial interventions for internalised stigma in people with a schizophrenia-spectrum diagnosis: a systematic narrative synthesis and meta-analysis. *Schizophr. Res.* 176 (2–3), 291–303. <https://doi.org/10.1016/j.schres.2016.05.001>.
- Wood, L., Byrne, R., Burke, E., Enache, G., Morrison, A.P., 2017. The impact of stigma on emotional distress and recovery from psychosis: the mediatory role of internalised shame and self-esteem. *Psychiatry Res.* 255, 94–100. <https://doi.org/10.1016/j.psychres.2017.05.016>.
- Yanos, P.T., Roe, D., Markus, K., Lysaker, P.H., 2008. Pathways between internalized stigma and outcomes related to recovery in schizophrenia spectrum disorders. *Psychiatr. Serv.* 59 (12), 1437–1442. <https://doi.org/10.1176/appi.ps.59.12.1437>.
- Yanos, P.T., Lysaker, P.H., Roe, D., 2010. Internalized stigma as a barrier to improvement in vocational functioning among people with schizophrenia-spectrum disorders. *Psychiatry Res.* 178 (1), 211–213. <https://doi.org/10.1016/j.psychres.2010.01.003>.