



# Mental health literacy for autism spectrum disorder and depression

Charlotte Godfrey Born<sup>a</sup>, Alastair McClelland<sup>b</sup>, Adrian Furnham<sup>c,\*</sup>

<sup>a</sup> Research Department of Clinical, Educational and Health Psychology, University College London, London WC1E 6BT, UK

<sup>b</sup> Research Department of Experimental Psychology, University College London, London WC1E 6BT, UK

<sup>c</sup> Norwegian Business School (BI), Nydølveien, Oslo, Norway



## ARTICLE INFO

### Keywords:

Mental Health Literacy  
Autistic Spectrum Disorder  
Depression  
Lay Recognition

## ABSTRACT

In this study, three hundred and sixty-eight participants answered a questionnaire consisting of three vignettes describing a person with depression, severe Autism Spectrum Disorder (ASD), and high-functioning Autism Spectrum Disorder/Asperger's Syndrome (AS). Each vignette was followed by thirteen questions concerning the participant's knowledge about diagnosis, treatment, and prognosis. Participants demonstrated good mental health literacy (MHL) for depression, but were poor at diagnosing ASD. Drug therapy was rated as the least effective treatment and 'talking to a family member or friend' was rated as the most effective for each disorder. The highest prognosis ratings were given to depression, and the lowest given to ASD. Participants demonstrated better MHL than expected, possibly due to the availability of information on the Internet and the increase in prevalence of mental illnesses such as depression and ASD among the general population.

## 1. Introduction

The term 'mental health literacy' (MHL) was first coined by Jorm et al. (1997) who defined it as the "knowledge and beliefs about mental disorders that aid the recognition, management or prevention of these disorders" (p. 182). There are probably now well over five hundred papers in this field, and the idea has become so popular that there are now papers concerned with very specific mental health "literacies" such as dementia literacy and schizophrenia literacy (see Furnham and Swami, 2018 for a review). It is argued that the conceptual models that lay individuals use to understand and explain mental health illnesses help shape their help-seeking behaviours for very specific psychiatric symptoms. In addition, such conceptual mental models are also believed to influence choice of treatment, compliance with treatment, and the stigmatising of individuals suffering from mental health disorders (Jorm, 2012, 2015).

The extensive literature suggests that the general public to have a poor understanding of mental health, which has impeded them in seeking and receiving treatment. The evidence also suggests that the general public typically favours psychosocial explanations over biological explanations in relation to the onset of a range of mental health disorders. Further, people tend to emphasise stressful circumstances in a person's life as the most likely cause of a range of mental health disorders (Furnham and Swami, 2018; Jorm, 2015).

The two mental disorders which have received most attention are

depression and schizophrenia. People tend to show a greater recognition of depression than of schizophrenia with around 70%–80% recognising depression and 20%–30% schizophrenia, independent of other participant attributes (e.g., male/female; urban/rural; age, country of study) (Sai and Furnham, 2013). Indeed, this is why many studies looking at MHL literacy in a particular disorder tend to include MHL depression as a "control group".

The current study compares the mental health literacy for Autism Spectrum Disorder (ASD), and high-functioning Autism Spectrum Disorder/Asperger's Syndrome (AS) to that of depression. Very little work has been done in this area (Furnham and Buck, 2003), but it is known that ASD is less common than depression and often misunderstood, which can allow distorted beliefs to develop (Huws and Jones, 2010). One example is the theory that ASD is caused by Thimerosal, a mercury-containing compound found in some childhood vaccines. However, irrefutable scientific evidence has found no link between Thimerosal and autism, and indeed, the number of cases of autism has actually increased since Thimerosal was removed from childhood vaccinations in 1999 (Smith, 2011).

This study aimed to examine lay peoples' ability to recognise the disorders listed above, accurately label them, recommend the appropriate treatments, and estimate the prognosis for the individual with the disorder in terms of symptoms, ability to form relationships and maintain a job. It was predicted that participants would show (H1) greater understanding of the diagnosis, treatment, and prognosis of

\* Corresponding author.

E-mail address: [a.furnham@ucl.ac.uk](mailto:a.furnham@ucl.ac.uk) (A. Furnham).

<https://doi.org/10.1016/j.psychres.2019.04.004>

Received 5 January 2019; Received in revised form 29 March 2019; Accepted 1 April 2019

Available online 17 April 2019

0165-1781/ © 2019 Elsevier B.V. All rights reserved.

depression, compared to ASD and AS. It was also predicted that (H2) more participants would identify severe ASD and depression as psychological disorders, but fewer would recognise the symptoms of high-functioning AS. It was predicted that (H3) women would show greater MHL (problem recognition and understanding) than men, as shown in previous research (Furnham et al., 2011; Wang et al., 2007). Finally, it was predicted there would be an (H4) effect of age on prognosis and treatment suggestions, with younger participants demonstrating better MHL (recommending professional treatment) than older participants. (Farrer et al., 2008; Fisher and Goldney, 2003).

## 2. Method

### 2.1. Participants

Three hundred and sixty-eight participants aged 18–73 years ( $M = 42.0$  years,  $SD = 15.3$  years) were included in the analysis (181 males and 187 females). All participants were volunteers from the UK, recruited through opportunity sampling using various online methods, including social media, and the network *Fulcrum*. The majority of the participants (57.6%) were paid employees, 12.2% were students, 12.5% were retired, and 17.4% were unemployed. Within the sample, 33.7% of participants said they had been diagnosed with some form of depression, and 1.4% with ASD. Thirty-two point three percent of subjects claimed to have a family member or friend who had been diagnosed with depression, and 25.3% with ASD. The majority of participants reported their ethnicity as ‘white’ (87.2%), 4.3% as ‘black/black British’ and 6.5% as ‘Asian/Asian British’.

### 2.2. Materials

Three different case vignettes were used in this research (examples of which can be found in [Appendix A](#)). One vignette described an individual with symptoms typical of someone with depression, the remaining two described individuals with differing severity of ASD: severe ASD and high-functioning AS. Each vignette included at least three of the most common and distinctive general symptoms necessary to be diagnosed with the disorder, and were based on an example vignette with the classical features of depression found in [Jorm et al. \(1997\)](#). The order of presentation of each vignette was randomised.

The gender and socio-economic status of the individual described in each vignette was also randomised, as factors such as age, gender, or income could have an effect on the participants’ impression of the disorder. In the study participants were first required to indicate if the individual in the vignette had a psychological disorder (‘yes’/‘no’). If the response was ‘yes’, they were asked to state what disorder they believed the individual to have (See [Appendix B](#)). Next, they were asked if they would recommend some form of treatment, and were required to rate the potential effectiveness of five different treatments on a seven-point scale, where one indicated ‘not effective’ and seven indicated ‘very effective’. This was to determine whether subjects thought treatment was necessary and also whether this should be mainly psychological (i.e., CBT), biological (i.e., drugs), or more to do with lifestyle choices (i.e., exercise, meditation, or talking to a family member or friend). Participants were then asked questions regarding the prognosis of the individual in the vignette including, ‘will their symptoms get better’, ‘do they have the potential to get a job?’, ‘do they have the potential to live alone?’, and ‘do they have the potential to develop close personal relationships?’. These four questions were each rated on a five-point scale ranging from ‘very unlikely’ to ‘very likely’. The penultimate question required a rating of the happiness of the individuals in the vignettes, which involved a seven-point scale ranging from ‘very unhappy’ to ‘very happy’.

**Table 1**

Frequencies of responses to presence of psychological disorder and treatment.

Disorder	Disorder present?		Treatment?	
	Yes	No	Yes	No
Autism	322 (87.5%)	46 (12.5%)	319 (86.7%)	49 (13.3%)
Asperger's	100 (27.2%)	268 (72.8%)	125 (34.0%)	243 (66.0%)
Depression	282 (76.6%)	86 (23.4%)	307 (83.4%)	61 (16.6%)

### 2.3. Procedure

Ethics committee permission was sought and received. Participants completed the entire experiment online. It took them approximately ten to fifteen minutes. All participants were required to give their consent, were assured of their anonymity, and at the end of the experimental session they were given debriefing information and thanked for their participation.

## 3. Results

The data for diagnosis, treatment, and prognosis were analysed separately.

### 3.1. Diagnosis

[Table 1](#) presents the frequencies of responses to the presence of a psychological disorder question. It can be seen that the majority of participants responded ‘yes’ for the ASD vignette, closely followed by the depression vignette. However, less than half of the participants recognised the presence of a psychological disorder in the AS vignette.

Cochran’s  $Q$  analysis revealed a significant difference between whether or not participants recognised the presence of a psychological disorder in the three case vignettes,  $Q(2, N = 368) = 313.46, p < .001$ . Additional analysis using the McNemar test showed that participants recognised a disorder in the ASD vignette significantly more often than in depression and AS vignettes,  $\chi^2(1, N = 368) = 16.18, p < .001$  and  $\chi^2(1, N = 368) = 210.52, p < .001$  respectively. Participants also recognised a disorder significantly more often in the depression vignette than in the AS vignette,  $\chi^2(1, N = 368) = 156.01, p < .001$ . These results confirm the hypothesis that autism ASD would be recognised as a disorder the least often. However, the prediction that depression would have the highest recognition rate as a mental health disorder was not confirmed.

Participants who responded ‘yes’ to the first question were asked to state the psychological disorder the individual may have. Of the participants who recognised the presence of a disorder in the Autism vignette (ASD), a content analysis of responses showed that 47.2% used the word ‘autism’ or ‘autistic’, 5.1% incorrectly used the word ‘Asperger’s’, and 1.2% mentioned ‘Autism Spectrum Disorder’. Of the participants who recognised the presence of a disorder in the Asperger’s vignette, 22.0% used the words ‘autism/autistic’, only 16.0% identified the disorder as ‘Asperger’s syndrome’, and only 2 participants (2.0%) stated the individual had some form of ASD. Seventy-six point six percent of participants recognised the presence of a disorder in the depression vignette. The content analysis revealed that 79.1% of these participants used the words ‘depression’ or ‘depressed’ to describe the individual. A small proportion of participants (3.1%) wrongly described the individual as having some form of anxiety, and a further 3.1% wrongly focussed on the weight gain described and suggested the presence of some form of eating disorder. A summary table of the content analysis can be found in [Appendix B](#).

### 3.2. Treatment

The frequency of answers to the treatment question for each

**Table 2**  
Mean ratings of types of therapy for different disorders.

Disorder	Young		Female		Middle aged		Female		Old		Female	
	Male	SD	M	SD	Male	SD	M	SD	Male	SD	M	SD
A. Effectiveness of drug therapy												
Autism	2.66	1.46	3.22	1.65	3.41	1.64	3.14	1.65	3.05	1.53	3.07	1.51
Asperger's	1.98	1.48	2.04	1.67	2.36	1.49	1.91	1.35	1.84	1.14	1.76	1.23
Depression	3.07	2.00	3.79	1.89	3.38	1.86	3.79	1.92	3.38	1.97	3.91	2.05
B. Effectiveness of psychological therapy												
Autism	4.92	1.53	4.88	1.39	4.55	1.59	4.74	1.70	4.38	1.37	4.45	1.63
Asperger's	2.89	1.98	3.03	1.95	3.41	1.93	3.47	1.94	3.53	1.94	3.16	1.94
Depression	4.28	1.73	4.92	1.69	4.18	2.09	5.02	1.68	4.56	1.72	4.84	1.77
C. Effectiveness of treatment exercise												
Autism	3.84	1.72	4.00	1.56	3.86	1.83	4.09	1.81	3.58	1.49	3.72	1.61
Asperger's	4.00	1.71	4.08	1.84	3.83	1.72	3.79	1.90	3.66	1.74	3.78	1.85
Depression	5.30	1.70	5.44	1.54	5.23	1.51	5.54	1.45	4.86	1.47	5.52	1.43
D. Effectiveness of meditation												
Autism	3.85	1.81	3.79	1.72	3.79	1.79	4.19	1.86	3.31	1.45	3.62	1.88
Asperger's	3.02	1.76	3.50	1.94	3.55	1.93	3.39	1.93	3.22	1.75	3.62	1.73
Depression	3.45	1.75	4.44	1.56	4.48	1.07	4.75	1.72	3.72	1.70	4.59	1.65
E. Effectiveness of talking to family member or friend												
Autism	4.49	1.67	4.24	1.74	4.30	1.82	4.21	1.86	3.98	1.67	3.84	1.90
Asperger's	4.79	1.69	4.63	1.82	4.57	1.81	4.37	1.88	4.16	1.64	4.33	1.79
Depression	5.38	1.66	5.53	1.27	5.34	1.48	5.25	1.68	4.88	1.54	5.41	1.88
F. Mean ratings of prognosis												
Autism	12.93	3.62	14.06	2.82	11.57	3.64	13.72	3.33	11.83	3.59	12.22	3.51
Asperger's	15.02	2.85	15.99	2.79	15.70	2.97	15.72	2.31	16.30	2.39	16.21	2.06
Depression	15.61	2.98	17.46	2.69	16.48	3.14	17.67	2.50	17.72	2.75	17.78	2.53
G. Mean ratings of prognosis												
Autism	3.62	1.24	3.93	1.24	3.86	1.35	2.93	1.03	3.52	1.05	3.59	1.11
Asperger's	4.74	1.14	4.33	1.35	4.41	1.32	4.74	1.14	4.80	0.95	4.41	1.12
Depression	3.25	1.47	2.92	1.32	3.05	1.38	3.04	1.51	2.89	1.27	2.88	1.44

disorder are displayed in Table 2. Most participants recommended treatment for autism and depression, whereas less than half responded in the affirmative for Asperger's. Cochran's Q analysis for this question was significant,  $Q(2, N = 368) = 295.43, p < .001$ . Further analysis using McNemar tests showed significantly more participants recommended treatment for Autism than for Asperger's,  $\chi^2(1, N = 368) = 180.82, p < .001$ . Similarly, significantly more participants recommended treatment for depression than for Asperger's,  $\chi^2(1, N = 368) = 160.59, p < .001$ . There was no significant difference between autism and depression,  $\chi^2(1, N = 369) = 1.73, p = .189$ .

### 3.3. Treatment drug therapy

Table 2a displays the results for the perceived effectiveness of drug therapy for each disorder. A  $3 \times 2 \times 3$  repeated measures ANOVA with disorder as the within subjects factor, and the age and gender of the individual described in the vignette participants as between-subjects factors revealed a highly significant difference between the vignettes, with respect to the effectiveness ratings given by participants,  $F(1.90, 688.24) = 123.38, p < .001, \eta_p^2 = 25.4\%$ . There was also a significant interaction between disorder and gender,  $F(1.90, 688.24) = 6.24, p = .002, \eta_p^2 = 1.7\%$ , indicating that drug therapy was thought to be more effective by females for the treatment of depression. Bonferroni-corrected post hoc analyses showed the mean score of effectiveness given to drug therapy as a potential treatment was significantly higher for ASD compared to AS ( $p < .001$ ), mean effectiveness was significantly higher for depression compared to both the AS ( $p < .001$ ), and ASD ( $p < .001$ ). There was no significant effect of either the age or gender of participants,  $F < 1$  and  $F(1, 362) = 1.78, p = .183$  respectively.

#### 3.3.1. Treatment psychological therapy

Table 2b shows participant effectiveness ratings of psychological therapy as a possible treatment were highest for autism, closely followed by depression. Participants thought psychological therapy would

be least effective as a possible treatment for Asperger's Syndrome. The results from a  $3 \times 2 \times 3$  repeated measures ANOVA showed a highly significant difference between disorders,  $F(2, 234) = 113.93, p < .001, \eta_p^2 = 23.9\%$ . There was also a significant interaction between disorder and gender,  $F(2, 724) = 5.09, p = .006, \eta_p^2 = 1.4\%$ , indicating that females believed psychological therapy was a more effective treatment for depression than did males. There was a significant interaction between disorder and age,  $F(4, 724) = 3.72, p = .005, \eta_p^2 = 2.0\%$ . For autism, the oldest participants thought psychological therapy less effective as a treatment than the youngest participants. Bonferroni-corrected post hoc analyses showed significantly higher ratings of effectiveness for autism ASD compared to AS ( $p < .001$ ), and significantly lower ratings of effectiveness for AS compared to depression ( $p < .001$ ). There was no significant difference between ratings for autism and depression ( $p > .05$ )

#### 3.3.2. Treatment exercise

It can be seen from Table 2c that participants rated exercise as an effective treatment for depression, whereas the effectiveness scores for Asperger's Syndrome and autism were much lower. A  $3 \times 2 \times 3$  repeated measures ANOVA revealed a highly significant difference between effectiveness ratings,  $F(2, 724) = 167.72, p < .001, \eta_p^2 = 31.7\%$ . There was no significant effect of either the age or gender of participants, and no significant interactions were found ( $F < 1$ ). Bonferroni-corrected post-hoc analyses showed significantly higher effectiveness ratings in the depression condition compared to both the autism condition ( $p < .001$ ) and the Asperger's condition ( $p < .001$ ). There was no significant difference between effectiveness ratings of exercise as a possible treatment for autism and Asperger's syndrome ( $p > .05$ ).

#### 3.3.3. Treatment meditation

Table 2d displays the means and standard deviations of participant effectiveness ratings of meditation as a potential treatment. It can be seen that participant ratings of effectiveness were highest for depression, followed by autism, and lowest for AS. A  $3 \times 2 \times 3$  repeated

measures ANOVA revealed a highly significant main effect of disorder,  $F(2, 724) = 46.02, p < .001, \eta_p^2 = 11.3\%$ , and a significant effect of gender was also found,  $F(1, 362) = 6.46, p = .011, \eta_p^2 = 1.8\%$ . Overall, females viewed meditation as a more effective treatment than males. There was a significant interaction between disorder and age,  $F(4, 724) = 3.19, p = .013, \eta_p^2 = 1.7\%$ , and between disorder and gender and,  $F(2, 724) = 4.50, p = .011, \eta_p^2 = 1.2\%$ . Meditation was seen by they oldest age group to be a more effective treatment for depression than the youngest age group, and females thought it would be particularly effective in the treatment of depression. There was no significant main effect of age,  $F(2, 362) = 2.23, p = .111$ . Bonferroni-corrected post-hoc analyses showed significantly higher ratings of effectiveness for depression compared to both autism ( $p < .001$ ) and AS ( $p < .001$ ). Ratings of effectiveness were also significantly higher for autism compared to the AS ( $p < .001$ ).

### 3.3.4. Treatment talking to a family member/friend

Table 2e shows that participants rated this treatment relatively highly for all of the vignettes, with depression receiving the highest score followed by Asperger's syndrome which was closely followed by autism. A  $3 \times 2 \times 3$  repeated measures ANOVA comparing mean ratings of effectiveness of talking to family members or friends, found a highly significant effect of disorder,  $F(2, 724) = 78.21, p < .001, \eta_p^2 = 17.8\%$ . There was no significant main effects of either age or gender, and no significant interactions were found ( $F < 1$ ). Bonferroni-corrected post-hoc analyses found significantly higher ratings of effectiveness for depression when compared with both autism ( $p < .001$ ) and AS ( $p < .001$ ). Ratings of effectiveness were significantly higher for AS compared to autism ( $p = .004$ ).

### 3.4. Prognosis

Table 2f presents the ratings for prognosis, and an analysis of these data revealed a significant main effect of disorder,  $F(1.84, 665.10) = 284.96, p < .001, \eta_p^2 = 44.0\%$ , with the prognosis for depression clearly being much better than the other two disorders. There was also a significant main effect of gender,  $F(1, 362) = 15.32, p < .001, \eta_p^2 = 4.1\%$  (with a better prognosis predicted by females), and a significant interaction between disorder and gender,  $F(1.83, 665.10) = 3.30, p < .042, \eta_p^2 = 0.9\%$  (with a better prognosis predicted by females, but only for autism and depression). A significant interaction between disorder and age was also found,  $F(3.68, 665.10) = 9.99, p < .001, \eta_p^2 = 5.2\%$ . The youngest participants were more optimistic than the oldest participants about the prognosis for autism and depression but not Asperger's. There was no significant main effect of age ( $F < 1$ ). Bonferroni-corrected post hoc analyses showed that the mean prognosis score for depression ( $M = 17.12$ ) was significantly higher ( $p < .001$ ) than for both Asperger's syndrome AD ( $M = 15.82$ ) and autism ASD ( $M = 12.72$ ), and the mean prognosis score was significantly higher ( $p < .001$ ) for Asperger's syndrome compared to autism.

### 3.5. Happiness

The means and standard deviations for the happiness question are displayed in Table 2g, with a higher score indicates greater happiness. A  $3 \times 2 \times 3$  repeated measures ANOVA revealed a significant main effect of disorder,  $F(1.93, 699.46) = 186.70, p < .001, \eta_p^2 = 34.0\%$ , but no significant effect of either age or gender. No significant interactions were found. Bonferroni-corrected post hoc analyses indicated significantly higher ( $p < .001$ ) happiness ratings for Asperger's syndrome ( $M = 4.57$ ) compared to autism ( $M = 3.74$ ) and depression ( $M = 3.00$ ). Participants also gave significantly higher happiness ratings ( $p < .001$ ) for the individual with autism than the individual with depression.

## 4. Discussion

As predicted, participants demonstrated good MHL for depression. Contrary to predictions however, severe ASD was recognised as a disorder by most participants. This suggests that the symptoms of severe ASD are relatively easy to recognize as being abnormal, yet lay individuals do not have the information to label the disorder 'correctly'. Asperger's Syndrome (AS) was least well recognised as a disorder. The poor recognition rate of AS indicates a lack of understanding among lay individuals with respect to an autism spectrum with varying levels of severity. This is likely because ASD was only introduced as an umbrella term to include AS and other autism spectrum disorders in DSM V (American Psychiatric Association, 2013), and lay individuals are very unlikely to be familiar with current theories and new developments such as this.

Participants were best at identifying depression correctly, less good at identifying severe ASD 'correctly', and worst at identifying high functioning ASD 'correctly'. A notable number of participants claimed the individual in the AS vignette had 'autism'. Due to the fact that Asperger's Syndrome is encompassed by the term Autism Spectrum Disorder it is arguable that these participants showed an understanding of ASD, but with only 2 participants using the 'correct' term it is difficult to say for certain. ASD was 'incorrectly' labelled as social anxiety, depression, and attention deficit hyperactivity disorder (ADHD) on a number of occasions. However, considerable similarities have been found between children with ASD and ADHD (Rumpf et al., 2012), as well as 50% comorbidity of ASD with conditions including major depressive disorder and anxiety (Gjevik et al., 2011; Lugnegard et al., 2011). Therefore, lay individuals have some understanding of the symptoms associated with ASD and what they entail, even though many participants in this study failed to label the disorder 'correctly'.

These results are similar to those of Sai and Furnham (2013) where, using case vignettes, it was found that a more common disorder (depression) was identified correctly 72.7% of the time, and a less common disorder (schizophrenia) was identified correctly 46.4% of the time. This raises the question as to why the general population seem to be better at diagnosing depression. In this current study 66.0% of participants claimed to have either been diagnosed with depression, or have a close friend or family member diagnosed with depression, compared to only 26.7% for ASD. This suggests more experience with a disorder leads to better mental health literacy for that disorder. Indeed, lay individuals tend to receive most of their information about mental illness through personal experience, the experiences of people close to them, and the media. Lay individuals often 'incorrectly' identify other disorders such as schizophrenia (and in this study, ASD) as depression which suggests a tendency to over-diagnose depression (Farrer et al., 2008).

An overarching finding of this study is that the depression vignette generated higher scores than ASD for the treatment and prognosis questions. Participants thought treatment in any form would be most effective for depression, with exercise and talking to a family member or friend seen as the most effective. The only treatment rated more effective for another disorder (autism) was psychological therapy, and the difference in ratings for the two disorders was not significant. In terms of mental health literacy, these findings indicate lay individuals possess some knowledge regarding the effectiveness of different treatment methods. The results show participants correctly acknowledged that there are existing drug therapies found to be effective for depression, but there is not a specific 'cure' for autism, and drugs that are often prescribed for the disorder tend to target the disordered sleeping and eating that individuals with autism often suffer from (Arroll et al., 2009; Bschor and Kilarski, 2016; Leucht et al., 2012). This research also suggests that lay individuals are aware of the more extensive psychological and behavioural interventions available for ASD, such as behaviour modification (Sheinkopf and Siegel, 1998).

However, the fact that the treatments categorized as lifestyle

changes were rated more highly is possibly because people are reluctant to seek help from mental health professionals and often report a fear of being embarrassed or judged (Barney et al., 2006). There is evidence that personal attitudes are more important than the attitudes of others in predicting intention to seek professional help (Bayer and Peay, 1997).

Drug therapy was rated as the least effective for all disorders, and there is evidence in the literature that psychotherapy is superior to drug therapy for treating depression (Steinbrueck et al., 1983). However, this scepticism of pharmacological treatment may be unfounded. It is important to provide better, more accessible information in this area as the public will only endorse treatments they understand and have confidence in (Furnham and Buck, 2003). Some pharmacological therapies have been found to be very effective in the treatment of mental disorders, particularly when taken alongside a psychotherapy programme (Nemeroff et al., 2003).

With regards to ASD, participants showed less confidence in the efficacy of treatment and tended to give scores between three and five (i.e., towards the middle of the scale). These results could reflect the fact that there are still debates surrounding the best treatment for ASD due to the extreme variability of the disorder, and the fact that there is no universal consensus of the cause (Furnham and Buck, 2003).

Interestingly, the highest prognosis ratings were given to depression, despite the majority of participants claiming the individual with

high-functioning ASD did not have a psychological disorder. Because of the high prevalence of depression in the population, lay individuals are aware of the possibility of symptoms improving and for individuals suffering from depression to live a relatively normal life. The lower scores given to AS possibly reflect the fact that depression can be successfully treated. Participants did show an understanding that severe ASD does not have a good prognosis, suggesting good mental health literacy. However, these results could also be a consequence of participants using information from the vignettes to answer the question, rather than knowledge they already possessed. This is not necessarily a negative observation, as it demonstrates how lay individuals were able to use the information given to make informed decisions about the prognosis of an individual. For example, social communication problems were outlined in both the ASD vignettes, which could have affected the overall prognosis result

Age and gender did not have as much of an effect as was predicted. For the prognosis questions female participants tended to give higher ratings, suggesting they are more optimistic about the prognosis of mental disorders.

This study, like all others had limitations. It is always desirable to have a large and representative population and to know more about them. We could have had longer and more detailed vignettes and enquired about other mental illness. Nevertheless, we believe the study makes an interesting contribution to a growing literature.

## Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.psychres.2019.04.004](https://doi.org/10.1016/j.psychres.2019.04.004).

## Appendix A

### Examples of Vignettes used in the Study

Severe ASD vignette (language problems, doesn't seek relationships, sensitive to change, single focus)

Sally is 20 years old and has exhibited language difficulties since the age of 2. She has always been irritable, introverted, and does not seek social relationships with her family or peers. She is sensitive to unexpected change and often reacts aggressively by hitting her head or throwing things. This behaviour caused her to get bullied at school. Sally knows everything about trains, and has pictures of the London tube maps on her wall. Sally's father earns a lot of money as the vice-president of an insurance company, her mother stays at home to look after Sally.

High functioning ASD (Need for routine, single focus, lack of social skills)

Anna is 22 years old and works at her local library where she used to spend every weekend as a child. Anna has a degree in computer science from university. Anna used to get in trouble at school for not doing her homework, she was only interested in working on her computer. Anna does not have any friends. She only likes talking about computers and her peers don't enjoy this, so they don't spend much time with her. Anna does not look people in the eyes when she talks to them, and she often uses a monotone voice. Anna lives at home with her parents. Anna's father works in a bank, and her mother is a university lecturer. They have a 3-storey house in Hampstead.

Depression vignette (sad, lack of interesting in activities once enjoyed, change in eating habits, lack of concentration, tired)

Stephanie is 18 years old and has just started her first year at university. She has been feeling unusually sad for the last 4 weeks. Stephanie has always loved playing lacrosse, but recently she has been avoiding training and matches. Stephanie has gained quite a lot of weight because she is eating too much. Stephanie finds it hard to concentrate in her lectures and is not doing well at university because she keeps missing deadlines. Stephanie feels run down every day and finds it hard to get out of bed. Stephanie went to a private school and both her parents are doctors.

## Appendix B

Frequencies of response to vignettes identified as describing a disorder.

Description	Disorder Autism	AS	Depression
<i>Autism/autistic</i>	152	22	0
<i>Asperger's</i>	17	16	0
<i>Spectrum</i>	4	2	0
<i>Depression/depressed</i>	6	7	223
<i>OCD</i>	2	3	0
<i>Anxiety/social anxiety</i>	13	5	7
<i>ADHD/ADD</i>	9	6	0
<i>Mental health illness</i>	13	1	1
<i>Eating disorder</i>	0	0	7
<i>No label</i>	106	38	44

## References

- American Psychiatric Association, 2013. Diagnostic and statistical manual of mental disorders, 5th ed. American Psychiatric Publishing, Arlington, VA.
- Arroll, B., Elley, C.R., Fishman, T., Goodyear-Smith, F.A., Kenealy, T., Blashki, G., et al., 2009. Antidepressants versus placebo for depression in primary care. *Cochrane Database Syst. Rev.* <https://doi.org/10.1002/14651858.CD007954>. Issue 3, Art. No.: CD007954.
- Barney, L.J., Griffiths, K.M., Jorm, A.F., Christensen, H., 2006. Stigma about depression and its impact on help-seeking intentions. *Austral. NZ J. Psychiatry* 40 (1), 51–54.
- Bayer, J.K., Peay, M.Y., 1997. Predicting intentions to seek help from professional mental health services. *Austral. N. Zealand J. Psychiatry* 31 (4), 504–513.
- Bschor, T., Kilarski, L.L., 2016. Are antidepressants effective? A debate on their efficacy for the treatment of major depression in adults. *Expert Rev. Neurother.* 16 (4), 367–374.
- Farrer, L., Leach, L., Griffiths, K.M., Christensen, H., Jorm, A.F., 2008. Age differences in mental health literacy. *BMC Public Health* 8 (1), 125.
- Fisher, L.J., Goldney, R.D., 2003. Differences in community mental health literacy in older and younger Australians. *Int. J. Geriatr. Psychiatry* 18 (1), 33–40.
- Furnham, A., Buck, C., 2003. A comparison of lay-beliefs about autism and obsessive-compulsive disorder. *Int. J. Soc. Psychiatry* 49 (4), 287–307.
- Furnham, A., Cook, R., Martin, N., Batey, M., 2011. Mental health literacy among university students. *J. Public Mental Health* 10 (4), 198–210.
- Furnham, A., Swami, V., 2018. Mental health literacy: a review of what it is and why it matters. *Int. Perspect. Psychol.* 7 (4), 240–257.
- Gjevik, E., Eldevik, S., Fjæran-Granum, T., Sponheim, E., 2011. Kiddie-SADS reveals high rates of DSM-IV disorders in children and adolescents with autism spectrum disorders. *J. Autism Dev. Disord.* 41 (6), 761–769.
- Huws, J.C., Jones, R.S.P., 2010. ‘They just seem to live their lives in their own little world’: lay perceptions of autism. *Disabil. Soc.* 25 (3), 331–344.
- Jorm, A.F., 2012. Mental health literacy: empowering the community to take action for better mental health. *Am. Psychol.* 67 (3), 231–243.
- Jorm, A., 2015. Why do we need the concept of “Mental Health Literacy”. *Health Commun.* 30, 1166–1168.
- Jorm, A.F., Korten, A.E., Jacomb, P.A., Christensen, H., Rodgers, B., Pollitt, P., 1997. Public beliefs about causes and risk factors for depression and schizophrenia. *Soc. Psychiatry Psychiatr. Epidemiol.* 32 (3), 143–148.
- Leucht, S., Tardy, M., Komossa, K., Heres, S., Kissling, W., Salanti, G., Davis, J.M., 2012. Antipsychotic drugs versus placebo for relapse prevention in schizophrenia: a systematic review and meta-analysis. *Lancet* 379 (9831), 2063–2071.
- Lugnegård, T., Hallerbäck, M.U., Gillberg, C., 2011. Psychiatric comorbidity in young adults with a clinical diagnosis of Asperger syndrome. *Res. Dev. Disabil.* 32 (5), 1910–1917.
- Nemeroff, C.B., Heim, C.M., Thasett, M.E., Kleins, D.N., Rush, A.J., Schatzberg, A.F., et al., 2003. Differential responses to psychotherapy versus pharmacotherapy in patients with chronic forms of major depression and childhood trauma. *Proc. Natl. Acad. Sci.* 100 (24), 14293–14296.
- Rumpf, A.L., Kamp-Becker, I., Becker, K., Kauschke, C., 2012. Narrative competence and internal state language of children with Asperger Syndrome and ADHD. *Res. Dev. Disabil.* 33 (5), 1395–1407.
- Sai, G., Furnham, A., 2013. Identifying depression and schizophrenia using vignettes: a methodological note. *Psychiatry Res.* 210 (1), 357–362.
- Sheinkopf, S.J., Siegel, B., 1998. Home-based behavioral treatment of young children with autism. *J. Autism Dev. Disord.* 28 (1), 15–23.
- Smith, R.B., 2011. Childhood Vaccinations and Autism. In *Multilevel Modeling of Social Problems*. Springer, Dordrecht, pp. 431–450.
- Steinbrueck, S.M., Maxwell, S.E., Howard, G.S., 1983. A meta-analysis of psychotherapy and drug therapy in the treatment of unipolar depression with adults. *J. Consult. Clin. Psychol.* 51 (6), 856–870.
- Wang, P.S., Aguilar-Gaxiola, S., Alonso, J., Angermeyer, M.C., Borges, G., Bromet, E.J., et al., 2007. Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. *Lancet* 370 (9590), 841–850.