



## Case report

# The role of avoidance motivation in the relationship between reward sensitivity and depression symptoms in adolescents: An ERP study

Alissa J. Ellis\*, Giulia Salgari, David J. Miklowitz, Sandra K. Loo

Semel Institute of Neuroscience and Human Behavior, Department of Psychiatry, University of California-Los Angeles, 760 Westwood Plaza, Los Angeles, CA 90025, USA



## ARTICLE INFO

## Keywords:

Reward response  
Depression  
ERP  
Approach motivation

## ABSTRACT

Blunted neural responses to reward in an EEG paradigm (RewP) are associated with vulnerability to depression, but the pathways linking this biomarker to depressive symptoms are unclear. We examined whether the relationship between reward response (RewP mean amplitude and latency) and depression was in part explained by approach-motivated behaviors in adolescents with varying levels of depression. EEG was collected during a game rigged to provide win/loss trials. Longer RewP latency was associated with depression symptoms only when scores on a measure of avoidance motivation were included. These results suggest that treatments targeting avoidance may decrease vulnerability to depressive episodes.

## 1. Introduction

Anhedonia—decreased motivation and reactivity toward pleasurable stimuli—is a feature of numerous psychiatric disorders, including depression (Admon and Pizzagalli, 2015; Der-Avakian and Markou, 2012). Depression onset often occurs in adolescence, the peak period for the neural development of reward circuitry (Auerbach et al., 2014; Forbes, 2009; Thapar et al., 2012). Whereas adolescence is typically marked by heightened reward sensitivity (Spear, 2000), depressed adolescents have a difficult time experiencing reward and overcoming avoidance (Forbes, 2009). Disruptions of reward circuitry in the medial prefrontal cortex and subcortical areas such as the anterior cingulate cortex (ACC) and ventral striatum (Carlson et al., 2014; Gottfried et al., 2003; Knutson et al., 2003) may increase risk for depression (Auerbach et al., 2014; Bress et al., 2013; Holroyd and Umemoto, 2016; Pizzagalli et al., 2005).

Sensitivity to rewarding versus non-rewarding stimuli can be indexed by the reward positivity (RewP), an event-related potential (ERP) acquired with EEG. This neural response is positively correlated with activation of the ventral striatum and dorsal anterior cingulate (Becker et al., 2014; Foti et al., 2011b). The RewP is typically maximal in medial frontocentral channels and has a deflection that occurs approximately 250–550 ms post reward or loss feedback and thus is considered an index of reward response. Reward responsiveness, using the RewP, purportedly reflects individual differences in approach-motivated affect and behaviors (Angus et al., 2015; Proudfit, 2015; Threadgill and Gable, 2016). Approach-motivation is activated by

signals of reward, whereas avoidance-motivation is activated by signals of loss/failure (Carver and Harmon-Jones, 2009; Coan and Allen, 2004). Dysregulation of the approach/avoidance-motivation system may be a mechanism involved in the association between blunted reward responses and depression (Depue and Iacono, 1989; Harmon-Jones et al., 2010).

While there is clear evidence that the RewP is associated with approach-motivated states and traits, the relationship between avoidance-motivation and the RewP is less clear. Lower RewP is associated with negative emotional states (e.g., anxiety, depression), (Foti and Hajcak, 2009; Gu et al., 2010) increased depression vulnerability and more severe symptom course, (Bress et al., 2013; Foti et al., 2011a; Kujawa et al., 2014; Weinberg et al., 2015; Whitton et al., 2016) but no work exists on the RewP and avoidance-motivation.

Given that depression symptoms have been associated with the avoidance of prospective rewards (Winer and Salem, 2016), we probed whether individual differences in approach- and avoidance-motivation play a role in the relationship between blunted reward processing and depression in adolescents. We used neurophysiological measures of reward response (RewP) and self-reported approach- and avoidance-motivation (Carver and White, 1994) to understand the mechanisms through which reduced reward responsiveness is associated with depressive symptoms. We explored three hypotheses: (1) Reduced RewP mean amplitude and longer peak latency would be associated with depression (Proudfit, 2015); (2) blunted reward response would be associated with avoidance-motivated affect and behavior (Winer and Salem, 2016); (3) avoidance-motivation would have a role in the relationship between the

\* Corresponding author.

E-mail address: [aellis@mednet.ucla.edu](mailto:aellis@mednet.ucla.edu) (A.J. Ellis).

<https://doi.org/10.1016/j.psychres.2019.03.011>

Received 11 January 2019; Received in revised form 5 March 2019; Accepted 6 March 2019

Available online 08 March 2019

0165-1781/ © 2019 Elsevier B.V. All rights reserved.

**Table 1**  
Demographic information and sample characteristics.

| <b>Demographics (SD)</b>   |                |
|--|----------------|
| M/F (%)  | 45/55          |
| Age (years)  | 14.53 (2.09)   |
| <b>Racial categories (%)</b>   |                |
| American Indian/ Alaska native   | 4              |
| Asian  | 0              |
| Native Hawaiian or other Pacific Islander                                | 5              |
| Black or African American  | 9              |
| White  | 59             |
| Other  | 14             |
| Unknown or not reported  | 9              |
| <b>Mood disorders (n = 15 with formal mood disorder diagnosis)</b>       |                |
| Past episode of major depression   | 5              |
| Current major depression   | 4              |
| Bipolar I  | 2              |
| Bipolar II   | 2              |
| Bipolar- NOS   | 2              |
| <b>Psychiatric comorbidities (of those with mood disorder diagnosis)</b> |                |
| No comorbidities   | 2              |
| Anxiety  | 5              |
| Obsessive–compulsive disorder  | 2              |
| ADHD   | 4              |
| Oppositional defiant disorder  | 3              |
| <b>Group mean (SD) measures</b>  |                |
| DRS (depression symptoms)  | 27.30 (11.38)  |
| BAS-drive  | 10.65 (3.03)   |
| BAS-fun  | 12.00 (2.51)   |
| BAS-reward   | 17.45 (2.26)   |
| BIS  | 20.16 (3.99)   |
| RewP latency (ms)*   | –21.29 (33.71) |
| RewP mean amplitude (μV)*  | 3.18 (3.40)    |

DRS, depression rating scale; BAS, behavior activation system; BIS, behavior inhibition system.

\* The *difference* score derived from the RewP calculation (e.g., latency indicates a difference of 21.29 ms between the win-loss conditions).

RewP and depression.

## 2. Method

### 2.1. Sample

Adolescents between 12–17 years ( $N = 20$ ) were recruited from a mood clinic. A dimensional approach to depression resulted in varied symptom profiles of participants. The majority had a history of a mood disorder ( $n = 15$ ), which was not necessary for inclusion ( $n = 5$ ) (no mood disorder) and had additional comorbidities (see Table 1). Parents and youth provided consent prior to study procedures approved by the University of California-Los Angeles Medical Institutional Review Board. Exclusion criteria were: (1) use of psychoactive drugs; (2) psychosis, autism spectrum disorder, substance abuse; (3) serious neurological conditions or brain trauma; or (4) sensory impairments. Sample characteristics are in Table 1.

### 2.2. Procedure

Following diagnostic interviewing, adolescents completed questionnaires. They were fitted with EEG electrodes and completed the computer task while continuous EEG data were collected.

### 2.3. Diagnostic assessments

The *Kiddie-Schedule for Affective Disorders and Schizophrenia (K-SADS)* (Chambers et al., 1985) was administered to participants and parents by a doctoral level clinician during clinic intake or during study visit (non-clinic participants). Participants were interviewed using the *Depression Rating Scale (DRS)* (Geller et al., 2001) to assess dimensional levels of depression symptoms over the previous week. Summary scores

were derived from the parent and child responses to the DRS using best estimate consensus ratings.

### 2.4. Affective Posner task

This task involved 100 trials beginning with a fixation cross (500–5000 ms), followed by a cue in one of two boxes positioned side-by-side (200 ms), then a target within one of the boxes. Participants respond as quickly as possible as to which box the target was presented. Feedback was adjusted where 40% of correct trials were provided true feedback and monetary reward (“You are Quick! Win 25 cents”); the remaining 60% of correct responses resulted in loss feedback (“Too Slow! Lose 25 cents”) presented on the computer screen.

### 2.5. BIS/BAS scales

Participants use a 4-point Likert scale ranging from *very true for me* to *very false for me* to indicate how much they agree with self-statements associated with approach and avoidance behaviors (Carver and White, 1994). The BIS includes items covering the anticipation of or response to punishment (Cronbach's  $\alpha = 0.83$ ). The BAS subscales are Drive ( $\alpha = 0.79$ ), Fun-Seeking ( $\alpha = 0.72$ ) and Reward Responsiveness ( $\alpha = 0.62$ ).

### 2.6. Electroencephalography (EEG)

EEG utilized 40 Ag/AgCL surface electrodes using an extended international 10/20-location system (ElectroCap, Eaton, OH). MANSCAN (Sam Technology, San Francisco, CA) hardware/software was used to continuously record EEG at a rate of 256 samples per second and referenced to linked-ears. Impedance was  $<10$  K $\Omega$ . Data were processed within Matlab (Mathworks) utilizing EEGLAB functions (Delorme and Makeig, 2004). A high-pass filter at  $<1$  Hz and low pass filter at 58 Hz were used, followed by manual removal of bad channels. Data were segmented into 0.5 s bins, and any segment outside 5 standard deviations of overall data was removed up to a 10% maximum.

Independent Component Analysis (ICA) separated neural and artifact signals using parameters by binica in EEGLAB. ICs representing non-brain activity were removed. Data were epoched 2 s before and after each feedback event. An average of 75% of trials (range 69–94) were retained after cleaning. We calculated mean ERPs using as baseline the 200 ms interval prior to feedback. To calculate RewP, difference scores (win-loss) were calculated using the Fz electrode. Mean amplitude 250–350 ms after feedback (win-loss) and latency of this peak (win-loss) were used for the RewP calculation (see Figs. 1 and 2).

## 3. Data analyses

Analyses used IBM SPSS v24 software. The four steps outlined by Baron and Kenny (1986) tested direct effects and hypotheses regarding the role of approach/avoidance motivation. In step one, linear regression examined whether neural response to reward (RewP amplitude/latency) predicted depression symptoms (DRS total). In step two, linear regression models examined whether the RewP was associated with motivation-related emotions/behaviors (i.e., subscales of the BIS/BAS). Third, linear regression examined whether the approach/avoidance motivation was associated with depression symptoms, controlling for RewP. Finally, we examined whether the effect of the predictor (RewP) on the outcome variable (DRS total) remained significant, controlling for the mediator (BIS/BAS). Totals were calculated for the outcome variables. In addition to the change score used to quantify RewP (Reward – Loss), RewP-reward and RewP-loss scores were examined as independent predictors of depression. Age was excluded as a covariate; it was not associated with the RewP, our predictor variable.




Fig. 1. ERP image from the FZ electrode during win (You're quick, orange line) and loss (Too slow, blue line) trials. Mean amplitude and latency were extracted between 250–350 ms. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. Scalp distribution of the difference between win (You're quick) and loss (Too slow) trials during the 250–350 ms segment used to calculate the RewP amplitude and latency.

4. Results

4.1. Preliminary analyses

BIS total scores were positively correlated with DRS total scores ( $R^2 = 0.70, p < .01$ ). RewP mean amplitude was not associated with depression symptoms or BIS/BAS scores, so steps 1 and 2 outlined above were not supported to do the additional analyses using mean amplitude as a predictor. BAS subscales were unrelated to RewP latency; the BIS/RewP latency relationship is described below. Table 1 displays means and standard deviations for outcome variables.

4.2. The role of avoidance motivation

Direct effects of the RewP latency on depression symptoms ( $R^2 = 0.30, \beta = 0.55, t = 2.60, p = .02$ ) and withdrawal behaviors (BIS subscore) ( $R^2 = 0.26, \beta = 0.51, t = 2.34, p = .03$ ) were revealed, satisfying the first two steps in the model. To demonstrate step 3, we

examined whether BIS score was associated with DRS, when controlling for RewP latency. Results confirmed that BIS score remained significantly associated with depression ( $R^2 = 0.60, \beta = 0.63, t = 3.23, p < .01$ ). This model also confirmed the final step—when the BIS score was included, the relationship between RewP latency and depression was no longer significant ( $p = .25$ ) (see Fig. 3).

4.3. Follow-up analysis

The difference between win/loss trials cannot identify whether this difference is due to quicker response to loss, longer response to reward, or both. To clarify whether response to loss or reward was more strongly associated with depression, we examined individual associations between RewP-loss latency and depression and RewP-reward latency and depression. Loss-latency was not associated with depression. However, longer latency of reward was associated with greater depression ( $r(19) = 0.52, p = .02$ ), but not to BIS score ( $r(19) = 0.37, p = .13$ ).




Fig. 3. The role of self-reported avoidance motivation (Behavioral Inhibition Scale [BIS]) score on the relationship between the Reward Positivity (RewP) latency and depression symptoms. \*denotes the direct effect of the RewP on depressive symptoms, not including BIS in the model.

## 5. Discussion

Reduced reward response and anhedonia are negative prognostic indicators for the remission of adolescent depression (McMakin et al., 2012). The RewP is considered to be a neural correlate of reward responsiveness and is a predictor of depression vulnerability and course. Less is known, however, about the specific pathways that link dysfunctions in the brain's reward circuitry, indexed by the RewP, to depressive symptoms like anhedonia. This study suggests that avoidance-motivation can help to explain the relationship between reduced reward responsiveness and depression symptoms.

A longer RewP latency of onset was associated with depression and avoidance-motivation. This latency reflected a delayed onset of the RewP to rewards, rather than a quicker processing of loss (Mulligan and Hajcak, 2018) suggesting that adolescents who were less efficient at processing rewards had greater depression. These preliminary findings extend the current literature to consider including *both* intensity (amplitude) and processing rate (latency) as features of blunted reward (Proudfit, 2015). Results were inconsistent with work focusing on the association between the RewP amplitude and depression (Proudfit, 2015). Whether previous studies have failed to find an effect of RewP latency in mood disorders or if latency was not examined is not clear. Both should be reported to clarify whether these mechanisms play different roles in the course of depression.

While the RewP has been associated with approach-motivation (Bress and Hajcak, 2013; Lange et al., 2012), results are the first to highlight the potential import of behavioral *avoidance* of reward, in understanding the relationship between the RewP and depression. These findings suggest initial support for the theory of reward devaluation which posits that depressed individuals actively avoid positive information, not just experience blunted reward responses (Winer and Salem, 2016). Behavioral avoidance is thus a mechanistic target for intervention. These avoidance tendencies may have consequences for the adolescent that affect the clinical course depression symptoms (Auerbach et al., 2014; Lewinsohn et al., 1992).

Strengths included the use of multiple levels of assessment (electrophysiology, self-report, clinician rated indices) and the Affective Posner task. The latter allowed for the potential tapping of underlying fears that approaching prospective rewards may result in punishment, as suggested by the Reward Devaluation theory. This cross-sectional pilot study is limited by the small sample size which may be underpowered, resulting in false negatives. The small, heterogeneous sample also precludes examination of the role of concurrent psychopathology, such as anxiety, on these relationships (Dickson and MacLeod, 2004), does not consider developmental factors like pubertal status, and may not be representative of the broader population. Future research should also aim to use alternatives to the BIS that specifically distinguish between avoidance of negative/punishing situations versus avoidance of rewarding situations that might ultimately lead to punishment in order to more robustly support Reward Devaluation Theory. While in need of replication, they provide preliminary evidence that avoidance of reward is a potentially crucial and understudied mechanism in adolescent depression.

In summary, results highlight avoidance tendencies as a behavioral mechanism contributing to how neural response to rewards may affect depression symptoms and provide support for the use of behavioral activation (BA) (Dimidjian et al., 2011; Ekers et al., 2014) in depression treatment. BA focuses on avoidance *and* activates the BAS by engaging patients in rewarding activities, despite high BIS, and may thus be a key to improving symptoms (Ekers et al., 2008).

## Acknowledgments

Alissa Ellis, Giulia Salgari, David Miklowitz and Sandra Loo, Semel Institute of Neuroscience and Human Behavior, Department of Psychiatry, University of California-Los Angeles. The authors have no

disclosures to report. Correspondence concerning this article should be addressed to Alissa Ellis, Semel Institute of Neuroscience and Human Behavior, University of California-Los Angeles, 760 Westwood Plaza, Los Angeles, CA 90025, 58-225. This work was supported by a grant from the National Institutes of Health: K23 MH106785 (PI: Ellis).

## References

- Admon, R., Pizzagalli, D.A., 2015. Dysfunctional reward processing in depression. *Curr. Opin. Psychol.* 4, 114–118. <https://doi.org/10.1016/j.copsyc.2014.12.011>.
- Angus, D.J., Kemkes, K., Schutter, D.J.L.G., Harmon-Jones, E., 2015. Anger is associated with reward-related electrocortical activity: evidence from the reward positivity. *Psychophysiology* 52, 1271–1280. <https://doi.org/10.1111/psyp.12460>.
- Auerbach, R.P., Admon, R., Pizzagalli, D.A., 2014. Adolescent depression: stress and reward dysfunction. *Harv. Rev. Psychiatry* 22, 139–148. <https://doi.org/10.1097/HRP.000000000000034>.
- Becker, M.P.I., Nitsch, A.M., Miltner, W.H.R., Straube, T., 2014. A single-trial estimation of the feedback-related negativity and its relation to bold responses in a time-estimation task. *J. Neurosci.* 34, 3005. <https://doi.org/10.1523/JNEUROSCI.3684-13.2014>.
- Bress, J.N., Foti, D., Kotov, R., Klein, D.N., Hajcak, G., 2013. Blunted neural response to rewards prospectively predicts depression in adolescent girls. *Psychophysiology* 50, 74–81. <https://doi.org/10.1111/j.1469-8986.2012.01485.x>.
- Bress, J.N., Hajcak, G., 2013. Self-report and behavioral measures of reward sensitivity predict the feedback negativity. *Psychophysiology* 50, 610–616. <https://doi.org/10.1111/psyp.12053>.
- Carlson, J., Foti, D., Harmon-Jones, E., Proudfit, G., 2014. Midbrain volume predicts fMRI and ERP measures of reward reactivity. *Brain Struct. Funct.* 1–6. <https://doi.org/10.1007/s00429-014-0725-9>.
- Carver, C.S., Harmon-Jones, E., 2009. Anger is an approach-related affect: evidence and implications. *Psychol. Bull.* 135, 183–204. <https://doi.org/10.1037/a0013965>.
- Carver, C.S., White, T.L., 1994. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: the BIS/BAS scales. *J. Pers. Soc. Psychol.* 67, 319–333. <https://doi.org/10.1037/0022-3514.67.2.319>.
- Chambers, W.J., et al., 1985. The assessment of affective disorders in children and adolescents by semistructured interview: test-retest reliability of the schedule for affective disorders and schizophrenia for school-age children, present episode version. *Arch. Gen. Psychiatry* 42, 696–702. <https://doi.org/10.1001/archpsyc.1985.01790300064008>.
- Coan, J.A., Allen, J.J., 2004. Frontal EEG asymmetry as a moderator and mediator of emotion. *Front. EEG Asymmetry Emot. Psychopathol* 67, 7–50. <https://doi.org/10.1016/j.biopsycho.2004.03.002>.
- Delorme, A., Makeig, S., 2004. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J. Neurosci. Methods* 134, 9–21. <https://doi.org/10.1016/j.jneumeth.2003.10.009>.
- Depue, R.A., Iacono, W.G., 1989. Neurobehavioral aspects of affective disorders. *Annu. Rev. Psychol.* 40, pp. 457–492 *Annual Review of Psychology. Annual Reviews, Palo Alto, CA*.
- Der-Avakian, A., Markou, A., 2012. The neurobiology of anhedonia and other reward-related deficits. *Spec. Issue Neuropsychiatr. Disord.* 35, 68–77. <https://doi.org/10.1016/j.tins.2011.11.005>.
- Dickson, J.M., MacLeod, A.K., 2004. Approach and avoidance goals and plans: their relationship to anxiety and depression. *Cogn. Ther. Res.* 28, 415–432. <https://doi.org/10.1023/B:COTR.0000031809.20488.ee>.
- Dimidjian, S., Barrera Jr., M., Martell, C., Muñoz, R.F., Lewinsohn, P.M., 2011. The origins and current status of behavioral activation treatments for depression. *Annu. Rev. Clin. Psychol.* 7, 1–38. <https://doi.org/10.1146/annurev-clinpsy-032210.104535>.
- Ekers, D., Richards, D., Gilbody, S., 2008. A meta-analysis of randomized trials of behavioural treatment of depression. *Psychol. Med.* 38, 611–623. <https://doi.org/10.1017/S0033291707001614>.
- Ekers, D., Webster, L., Van Straten, A., Cuijpers, P., Richards, D., Gilbody, S., 2014. Behavioural activation for depression; an update of meta-analysis of effectiveness and sub group analysis. *Plos One* 9, e100100. <https://doi.org/10.1371/journal.pone.0100100>.
- Forbes, E.E., 2009. Where's the fun in that? Broadening the focus on reward function in depression. *Biol. Psychiatry* 66, 199–200. <https://doi.org/10.1016/j.biopsycho.2009.05.001>.
- Foti, D., Hajcak, G., 2009. Depression and reduced sensitivity to non-rewards versus rewards: evidence from event-related potentials. *Biol. Psychol.* 81, 1–8. <https://doi.org/10.1016/j.biopsycho.2008.12.004>.
- Foti, D., Kotov, R., Klein, D., Hajcak, G., 2011a. Abnormal neural sensitivity to monetary gains versus losses among adolescents at risk for depression. *J. Abnorm. Child Psychol.* 39, 913–924. <https://doi.org/10.1007/s10802-011-9503-9>.
- Foti, D., Weinberg, A., Dien, J., Hajcak, G., 2011b. Event-related potential activity in the basal ganglia differentiates rewards from nonrewards: temporoparietal principal components analysis and source localization of the feedback negativity. *Hum. Brain Mapp.* 32, 2207–2216. <https://doi.org/10.1002/hbm.21182>.
- Geller, B., Zimmerman, B., Williams, M., Bolhofner, K., Craney, J.L., Delbello, M.P., Soutullo, C., 2001. Reliability of the Washington University in St. Louis Kiddie schedule for affective disorders and schizophrenia (WASH-U-KSADS) mania and rapid cycling sections. *J. Am. Acad. Child Adolesc. Psychiatry* 40, 450–455. <https://doi.org/10.1097/00004583-200104000-00014>.
- Gottfried, J.A., O'Doherty, J., Dolan, R.J., 2003. Encoding predictive reward value in

- human amygdala and orbitofrontal cortex. *Science* 301, 1104–1107.
- Gu, R., Huang, Y.-X., Luo, Y.-J., 2010. Anxiety and feedback negativity. *Psychophysiology* 47, 961–967. <https://doi.org/10.1111/j.1469-8986.2010.00997.x>.
- Harmon-Jones, E., Gable, P.A., Peterson, C.K., 2010. The role of asymmetric frontal cortical activity in emotion-related phenomena: a review and update. *Biopsychol. Emot. Curr. Theor. Empir. Perspect.* 84, 451–462. <https://doi.org/10.1016/j.biopsycho.2009.08.010>.
- Holroyd, C.B., Umemoto, A., 2016. The research domain criteria framework: the case for anterior cingulate cortex. *Neurosci. Biobehav. Rev.* 71, 418–443. <https://doi.org/10.1016/j.neubiorev.2016.09.021>.
- Knutson, B., Fong, G.W., Bennett, S.M., Adams, C.M., Hommer, D., 2003. A region of mesial prefrontal cortex tracks monetarily rewarding outcomes: characterization with rapid event-related fMRI. *NeuroImage* 18, 263–272. [https://doi.org/10.1016/S1053-8119\(02\)00057-5](https://doi.org/10.1016/S1053-8119(02)00057-5).
- Kujawa, A., Proudfit, G.H., Klein, D.N., 2014. Neural reactivity to rewards and losses in offspring of mothers and fathers with histories of depressive and anxiety disorders. *J. Abnorm. Psychol.* 123, 287–297. <https://doi.org/10.1037/a0036285>.
- Lange, S., Leue, A., Beauducel, A., 2012. Behavioral approach and reward processing: results on feedback-related negativity and P3 component. *Biol. Psychol.* 89, 416–425. <https://doi.org/10.1016/j.biopsycho.2011.12.004>.
- Lewinsohn, P.M., Munoz, R.F., Youngren, M.A., Zeiss, A.M., 1992. *Control Your Depression*. Fireside/Simon and Schuster, New York, New York.
- McMakin, D.L., Olino, T.M., Porta, G., Dietz, L.J., Emslie, G., Clarke, G., Wagner, K.D., Asarnow, J.R., Ryan, N.D., Birmaher, B., Shamseddeen, W., Mayes, T., Kennard, B., Spirito, A., Keller, M., Lynch, F.L., Dickerson, J.F., Brent, D.A., 2012. Anhedonia predicts poorer recovery among youth with selective serotonin reuptake inhibitor treatment-resistant depression. *J. Am. Acad. Child Adolesc. Psychiatry* 51, 404–411. <https://doi.org/10.1016/j.jaac.2012.01.011>.
- Mulligan, E.M., Hajcak, G., 2018. The electrocortical response to rewarding and aversive feedback: the reward positivity does not reflect salience in simple gambling tasks. *Int. J. Psychophysiol.* <https://doi.org/10.1016/j.ijpsycho.2017.11.015>.
- Pizzagalli, D.A., Jahn, A.L., O'Shea, J.P., 2005. Toward an objective characterization of an anhedonic phenotype: a signal-detection approach. *Biol. Psychiatry* 57, 319–327. <https://doi.org/10.1016/j.biopsycho.2004.11.026>.
- Proudfit, G.H., 2015. The reward positivity: from basic research on reward to a biomarker for depression. *Psychophysiology* 52, 449–459. <https://doi.org/10.1111/psyp.12370>.
- Spear, L.P., 2000. The adolescent brain and age-related behavioral manifestations. *Neurosci. Biobehav. Rev.* 24, 417–463. [https://doi.org/10.1016/S0149-7634\(00\)00014-2](https://doi.org/10.1016/S0149-7634(00)00014-2).
- Thapar, A., Collishaw, S., Pine, D.S., Thapar, A.K., 2012. Depression in adolescence. *Lancet* 379, 1056–1067. [https://doi.org/10.1016/S0140-6736\(11\)60871-4](https://doi.org/10.1016/S0140-6736(11)60871-4).
- Threadgill, A.H., Gable, P.A., 2016. Approach-motivated pregoal states enhance the reward positivity. *Psychophysiology* 53, 733–738. <https://doi.org/10.1111/psyp.12611>.
- Weinberg, A., Liu, H., Hajcak, G., Shankman, S.A., 2015. Blunted neural response to rewards as a vulnerability factor for depression: results from a family study. *J. Abnorm. Psychol.* 124, 878–889. <https://doi.org/10.1037/abn0000081>.
- Whitton, A.E., Kakani, P., Foti, D., Van't Veer, A., Haile, A., Crowley, D.J., Pizzagalli, D.A., 2016. Blunted neural responses to reward in remitted major depression: a high-density event-related potential study. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* 1, 87–95. <https://doi.org/10.1016/j.bpsc.2015.09.007>.
- Winer, E.S., Salem, T., 2016. Reward devaluation: dot-probe meta-analytic evidence of avoidance of positive information in depressed persons. *Psychol. Bull.* 142, 18–78. <https://doi.org/10.1037/bul0000022>.