

Remote-Access Thyroidectomy: A Multi-Institutional North American Experience with Transaxillary, Robotic Facelift, and Transoral Endoscopic Vestibular Approaches

Jonathon O Russell, MD, Christopher R Razavi, MD, Meghan E Garstka, MD, Lena W Chen, BA, Elya Vasiliou, MD, Sang-Wook Kang, MD, Ralph P Tufano, MD, MBA, FACS, Emad Kandil, MD, FACS

- BACKGROUND:** Many remote-access approaches (RAAs) to the thyroid have been described to circumvent anterior neck scarring, including the transaxillary, robotic facelift, and transoral endoscopic vestibular approaches. These techniques have been popularized in Asia, but adoption has been slow in North America. We aimed to examine multi-institutional North American outcomes with RAA thyroidectomy in the context of these institutions' transcervical approach (TCA) outcomes.
- STUDY DESIGN:** Cases of lobectomy and total thyroidectomy via transaxillary, robotic facelift, and transoral endoscopic vestibular approaches were reviewed. Demographic characteristics, outcomes, and complications were compared with the same measures in patients undergoing lobectomy and total thyroidectomy via TCA by the primary RAA surgeons at each institution. Patients who underwent parathyroidectomy or other concomitant neck dissection procedures were excluded.
- RESULTS:** Two hundred and sixteen RAA thyroidectomies were attempted (92 transoral endoscopic vestibular approaches, 70 transaxillary, and 54 robotic facelift) and 410 TCA thyroidectomies were performed. There was no difference in mean index nodule sizes between RAA (2.8 ± 1.6 cm) and TCA (2.9 ± 1.9 cm) cohorts ($p = 0.72$). Median operative times for lobectomy were 146 minutes (range 60 to 343 minutes) and 90 minutes (range 25 to 247 minutes) for the RAA and TCA cohorts, respectively ($p < 0.0001$). Median operative time for total thyroidectomy was 170 minutes (range 100 to 398 minutes) vs 126.5 minutes (range 51 to 260 minutes) for the RAA and TCA cohorts, respectively ($p < 0.0001$). There was no difference in the rates of permanent recurrent laryngeal nerve injury between the RAA (0 of 216 [0%]) and TCA (0 of 410 [0%]) cohorts ($p = 0.99$).
- CONCLUSIONS:** Remote-access approach thyroidectomy can be performed in a select North American patient population with outcomes comparable with TCA. (J Am Coll Surg 2019;228:516–524. © 2019 Published by Elsevier Inc. on behalf of the American College of Surgeons.)

Surgical techniques and approaches have undergone significant refinement and reformation in an effort to reduce morbidity and mortality during the last 100 years. These include the introduction of laparoscopic and robotic

techniques that have allowed for more complex procedures through smaller incisions.^{1,2} In contrast, thyroid operations have remained largely stagnant, with the overwhelming majority of these procedures still being

Disclosure Information: Nothing to disclose.

Disclosures outside the scope of this work: Dr Tufano is a paid consultant to Hemostatix and Medtronic. All other authors have nothing to disclose.

Presented at the Southern Surgical Association 130th Annual Meeting, Palm Beach, FL, December 2018.

Received December 9, 2018; Accepted December 10, 2018.

From the Department of Otolaryngology-Head and Neck Surgery, Johns Hopkins University School of Medicine, Baltimore, MD (Russell, Razavi,

Chen, Vasiliou, Tufano) and Division of Endocrine Surgery, Department of Surgery, Tulane University Medical Center, New Orleans, LA (Garstka, Kang, Kandil).

Correspondence address: Jonathon O Russell, MD, Department of Otolaryngology-Head and Neck Surgery, The Johns Hopkins School of Medicine, Johns Hopkins Outpatient Center, 601 N Caroline St, 6th Floor, Baltimore, MD 21287. email: jrusse41@jhmi.edu

Abbreviations and Acronyms

RAA	= remote-access approach
RF	= robotic-assisted facelift approach
RLN	= recurrent laryngeal nerve
TA	= transaxillary approach
TCA	= transcervical approach
TOETVA	= transoral endoscopic thyroidectomy vestibular approach

performed via an anterior-cervical incision first described by Kocher.³ This is despite the finding that an anterior-cervical neck scar can negatively impact patient quality of life, regardless of the severity of the scar.^{4,5} Although one might assume this is due to the lack of other safe and feasible techniques, this is not the case, as several remote-access approaches (RAAs) to thyroid procedures have been described with good success.⁶⁻¹⁹

These RAA procedures have gained favor internationally, particularly in Asia, but the same cannot be said in North America.^{10,11} Some have postulated that this is due to differences in the patient populations, with the larger body habitus of the North American cohort often cited as a confounding factor.^{8,10,19} Others have suggested that the surgical indications in the Asian cohort are more amenable to the use of these RAA techniques.^{10,17} Given this divergent experience between continents and concerns about the generalizability of the robust Asian RAA outcomes, there is a need for multi-institutional North American data with these approaches. In addition, as outcomes with the transcervical approach (TCA) to the thyroid continue to be perfected, this RAA experience must be presented in the context of the TCA data, in an increasingly value-based North American healthcare market.²⁰ Here we present outcomes with 3 widely used RAA techniques—the robotic-assisted facelift (RF) approach, robotic-assisted transaxillary approach (TA), and the transoral endoscopic thyroidectomy vestibular approach (TOETVA)—from 2 high-volume North American thyroid centers in the context of their TCA outcomes.

METHODS

Cases of RF, TA, and TOETVA performed at the Johns Hopkins Medical Institution and Tulane University Medical Center were retrospectively reviewed from December 1, 2012, to June 29, 2018. Demographic characteristics, outcomes, and complications were collected. For the purposes of this study, only cases of thyroid lobectomy and total thyroidectomy were included for analysis. Additionally, cases where other concomitant procedures, such

as parathyroidectomy, central/lateral neck dissection, or Sistrunk operations, were performed were excluded to maintain homogeneity of the data. The overwhelming majority of these cases were completed by 2 high-volume thyroid surgeons (one from each institution) with fellowship training in head and neck endocrine surgery. As such, the RAA data were then compared with the TCA outcomes from these 2 surgeons between 2015 and 2018 at their respective institutions. In a similar fashion, only cases of thyroid lobectomy and total thyroidectomy were included for review, again with exclusion of those cases with other concomitant procedures.

All included patients met criteria for thyroid lobectomy or total thyroidectomy based on the current American Thyroid Association guidelines at the time of operation.^{21,22} Patients included in the RAA cohort were offered these procedures based on selection criteria described previously.^{19,23-26} In a broad sense, these included those with a thyroid lobe <10 cm in patients highly motivated to avoid an anterior-cervical scar. Cytopathologically benign and indeterminate nodules and patients with Graves' disease were considered for surgical candidacy. Similarly, select patients with T1 well-differentiated thyroid carcinoma were offered RAA. Of note, given inherent limitations accessing the contralateral lobe with the RF technique, patients opting for or requiring initial total thyroidectomy were not routinely offered RF. Operative technique for RF, TA, and TOETVA have been described previously in detail by our group and in the literature.^{12,17,19,23,27}

Non-parametric independent data between the cohorts were compared with the Mann-Whitney U test, and differences in means of parametric outcomes were compared using an unpaired *t*-test. Fisher's exact test was used to compare categorical data. Analysis was performed in Stata Statistical Software, release 15 (StataCorp) using an α of 0.05 for statistical significance. IRB approval was obtained from both Johns Hopkins University School of Medicine and Tulane University Medical Center.

RESULTS

Two-hundred and sixteen RAA (54 RF, 70 TA, and 92 TOETVA) cases were completed compared with 410 TCA cases in the reviewed periods at these institutions. An additional 33 remote-access cases were excluded from the RAA cohort based on the criteria described, that is, these cases included concomitant parathyroid or neck dissection procedures. All TOETVA cases and 20 RF cases were completed at Johns Hopkins Medical Institution, and the remaining RAA cases were completed at Tulane University Medical Center. Of the 216 RAA

Table 1. Demographic Characteristics of the Remote-Access Approach and Transcervical Approach Cohorts

Variable	Remote-access approach	Transcervical approach	p Value
Lobectomy, n/N (%)	169/216 (78)	240/410 (59)	<0.0001*
BMI, kg/m ² , mean ± SD	27.4 ± 5.9	31.7 ± 8.3	<0.0001*
Nodule size, cm, mean ± SD	2.8 ± 1.6	2.9 ± 1.9	0.72
Age, y, mean ± SD	43.9 ± 12.1	53.5 ± 14.7	<0.0001*
Women, n/N (%)	202/216 (94)	307/410 (75)	<0.0001*

*Statistically significant.

cases, 169 (78%) were lobectomies (21 completion thyroidectomies) and 47 (22%) were total thyroidectomies. In the TCA cohort, 240 (59%) were lobectomies (93 completion thyroidectomies) and 170 (41%) were total thyroidectomies. Women represented the majority in both cohorts, but there was a significantly higher proportion of women in the RAA cohort (94% vs 75%; $p < 0.0001$). Mean age was younger for the RAA cohort (43.9 ± 12.1 years vs 53.5 ± 14.7 years; $p < 0.0001$). Similarly, mean BMI of the RAA cohort was lower (27.4 ± 5.9 kg/m² compared with 31.7 ± 8.3 kg/m² for the TCA cohort; $p < 0.0001$). There was no difference in mean index nodule size between RAA and TCA cohorts (Table 1).

Two-hundred and fifteen of 216 (99.5%) RAA cases were completed via the intended approach. One TOETVA lobectomy was converted to TCA after superior pole bleeding was encountered that could not be controlled transorally. This case was completed without recurrent laryngeal nerve (RLN) injury, or other complication, in 123 total minutes with the patient discharged to home on the day of operation. This case was not included in calculation of median operative time for the RAA or TCA cohorts.

No patients in either the RAA or TCA cohorts had permanent (deficits lasting >3 months) RLN injury (0 of 216 vs 0 of 410; $p = 0.99$). There was also no significant difference in rate of temporary RLN injury between RAA (7 of 216 [3.2%]) and TCA (6 of 410 [1.5%]) cohorts ($p = 0.15$). All cases of temporary RLN injury had

evidence of recovery of true vocal fold function on flexible fiberoptic laryngoscopy within 3 months. Median time to recovery across all 13 cases was 18.5 ± 24.9 days (range 7 to 82 days). No patients had permanent hypoparathyroidism in the RAA (0 of 68 [0%]) or TCA (0 of 263 [0%]) cohorts ($p = 0.99$) (Table 2). Permanent hypoparathyroidism was defined in the at-risk population (total and completion thyroidectomies) based on biochemical parameters with a duration lasting >6 months.²⁸ On subgroup analysis of RAA techniques, there was no significant difference in rates of temporary or permanent RLN injury, and similarly, no significant differences in rates of permanent hypoparathyroidism (Table 3).

With regard to RAA technique-specific complications, there was no incidence of permanent (deficits lasting >3 months) mental nerve injury or CO₂ embolism in the TOETVA subgroup. Similarly, there was no incidence of brachial plexus injury in the TA subgroup. Notably, somatosensory evoked potential monitoring was used during all TA cases as a tool to prevent this complication as has been described previously.²⁹ One of 54 patients in the RF subgroup required sacrifice of a small branch of the great auricular nerve to facilitate dissection. Therefore, only 1 of 216 (0.5%) patients in the RAA cohort had a complication specific to their respective RAA technique. There were no postoperative neck infections in the RAA cohort, regardless of technique. There was no significant difference in rate of hematoma between the cohorts, with an incidence of 3 of 216 (1.4%) vs 1 of 410 (0.2%) in the RAA and TCA cohorts,

Table 2. Outcomes and Complications

Variable	Remote-access approach	Transcervical approach	p Value
Lobectomy OT, min, (median)	146 (43–343)	90 (25–247)	<0.0001*
Total thyroid OT, min, (median)	170 (100–398)	126.5 (51–260)	<0.0001*
Permanent RLN injury, n/N (%)	0/216 (0)	0/410 (0)	0.99
Temporary RLN injury, n/N (%)	7/216 (3.2)	6/410 (1.5)	0.15
Permanent hypoparathyroidism, [†] n/N (%)	0/68 (0)	0/263 (0)	0.99
Hematoma, n/N (%)	3/216 (1.4)	1/410 (0.2)	0.12
Seroma, n/N (%)	5/216 (2.3)	2/410 (0.5)	0.051

*Statistically significant.

[†]Calculated only in at-risk population of total thyroidectomies and completion thyroidectomies.

OT, operative time; RLN, recurrent laryngeal nerve.

Table 3. Subgroup Analysis of Remote-Access Techniques

Variable	Robotic facelift approach	Robotic-assisted transaxillary approach	Transoral endoscopic thyroidectomy vestibular approach
Lobectomy OT, min, median (range)	179 (114–293)	139 (85–226)	126 (60–343)
p Value*	<0.0001 [†]	<0.0001 [†]	0.30
Total thyroidectomy OT, min, median (range)	NA	164.5 (100–241)	172.5 (127–398)
p Value*	NA	NA	0.11
Permanent RLN injury, n/N (%)	0/54 (0)	0/70 (0)	0/92 (0)
p Value*	0.99	0.99	0.99
Temporary RLN injury, n/N (%)	2/54 (3.7)	1/70 (1.4)	4/92 (4.3)
p Value*	0.58	0.99	0.39
Permanent hypoparathyroidism, n/N (%)	0/8 (0)	0/30 (0)	0/30 (0)
p Value*	0.99	0.99	0.99
Hematoma, n/N (%)	3/54 (5.6)	0/70 (0)	0/92 (0)
p Value*	0.08	0.048 [†]	0.99
Seroma, n/N (%)	3/54 (5.6)	2/70 (2.9)	1/92 (1.1)
p Value*	0.65	0.14	0.58

*First value represents column 1 vs 2, second 1 vs 3, and third 2 vs 3.

[†]Statistically significant.

OT, operative time; RLN, recurrent laryngeal nerve.

respectively ($p = 0.12$) (Table 2). Though, all postoperative hematomas occurred in the RF subgroup, which reached statistical significance compared with TOETVA ($p = 0.048$), but not TA ($p = 0.08$) (Table 3). There was no statistically significant difference in rate of seromas with an incidence of 5 of 216 (2.3%) vs 2 of 410 (0.5%) in the RAA and TCA cohorts, respectively ($p = 0.051$). Similarly, there was no difference in seroma rate between any RAA subgroups, with 3 cases of seroma in the RF cohort, 2 cases in the TA cohort, and 1 case in the TOETVA cohort (Table 3). All seromas were managed with outpatient aspiration without need for a subsequent operative procedure.

Median operative time (incision to closure) for RAA lobectomies was 146 minutes (range 60 to 343 minutes), significantly longer than the TCA cohort, which had a median operative time of 90 minutes (range 25 to 247 minutes) ($p < 0.0001$). Similarly, median total thyroidectomy operative time in the RAA cohort was significantly longer than the TCA cohort; 170 minutes (range 100 to 398 minutes) vs 126.5 minutes (range 51 to 260 minutes) ($p < 0.0001$). When performing a subgroup analysis among RAA techniques, both TA (139 minutes; $p < 0.0001$) and TOETVA (126 minutes; $p < 0.0001$) were significantly faster than RF (179 minutes) in performing thyroid lobectomy. However, there was no significant difference in operative time between TA and TOETVA for lobectomy or total thyroidectomy. No analysis was performed comparing RF with the other RAA techniques for total thyroidectomy, given only 1 such case was performed via RF (Table 3). One-

hundred and nine of 169 (64%) RAA lobectomies were discharged on the day of operation, and 44 of 47 (94%) RAA total thyroidectomies were discharged to home on postoperative day 1 or sooner.

DISCUSSION

Although multiple international studies have demonstrated the safety and efficacy of RAA for thyroid operation, there has been a paucity of North American case volume and, therefore, literature.^{8,10,11,15,30-32} Early North American experiences with RAA were marked by complications not encountered with TCA.^{7,23,33} These were attributed, in part, to dissection planes, which were unfamiliar to most thyroid surgeons, but also to differing patient characteristics in the Asian cohort, such as BMI and index nodule size.

The RAA cohort had a significantly higher proportion of women and patients undergoing lobectomies compared with the TCA cohort. In addition, the RAA cohort was younger and BMI was lower than in the TCA cohort. These findings are not surprising, given that these RAA techniques are only offered to a highly selected patient population, though criteria have begun to expand.^{10,24,25} It is important to note that there was no significant difference in mean index nodule size between cohorts.

The largest study to date describing the South Korean TA experience by Ban and colleagues⁸ reported a mean nodule size of 0.66 cm. In contrast, the RAA cohort in our series had a mean nodule size of 2.8 cm, more than 4 times larger, and consistent with the typical North

American thyroid cohort. Additionally, mean BMI in our RAA cohort was 27.5 kg/m² compared with 22 kg/m² reported by Ban and colleagues. Despite this discrepancy, outcomes and complications were commensurate between our RAA cohort and this series mentioned. More importantly, outcomes and complications were not significantly different between our RAA and TCA cohorts. There was no difference in rate of permanent RLN injury, permanent hypoparathyroidism, or postoperative hematoma, and only 1 of 216 (0.4%) patients had a complication specific to the RAA technique. Notably, this complication was a deliberate sacrifice of a small branch of the great auricular nerve during flap elevation in the RA subgroup, as opposed to an unintentional iatrogenic injury. Additionally, there was also no significant difference in rate of temporary RLN injury between RAA and TCA cohorts. This finding might suggest the ability of these RAA techniques to perform fine dissection of the RLNs in addition to their identification and preservation.

Previous studies have proposed that characteristics inherent to TOETVA might allow for enhanced visualization and protection of the RLNs.³⁴ The TOETVA allows for visualization of the bilateral RLNs at their insertion points, a view familiar to experienced thyroid surgeons. In addition, the approach allows for a favorable angle of dissection along the nerve. These hypotheses have held true in our TOETVA cohort, with no permanent RLN injuries to date. Similarly, the robotic-assisted RAA techniques provide a magnified view of critical anatomy, and the wristed instrumentation facilitates safe dissection of the RLNs. This allowed for preservation of all RLNs in our cohort of robotic-assisted techniques. We believe these benefits with the RAA techniques are maximized only after the learning curves for these techniques have been met, and important to consider is that the learning curves for the robotic-assisted techniques, including RF and TA, are notably longer than TOETVA. Studies estimate a learning curve of 35 to 50 cases for robotic-assisted RAA techniques vs 7 to 11 cases for TOETVA.^{16,17,19,35-37}

There was a significant difference in operative time between the RAA and TCA cohorts. Consistent with earlier literature, RAA techniques were found to have longer operative times for both thyroid lobectomy and total thyroidectomy.^{10,11,16} Among the RAA techniques, RF was found to take significantly longer than both TA and TOETVA, and there was no significant difference in operative time between TA and TOETVA on subgroup analysis (Table 3).

In our series, we found that RF takes significantly longer than the other RAA techniques, only allowing for adequate ipsilateral lobe access, and with a higher rate of postoperative hematoma (3 of 54 [5.6%] vs 0 of

162 [0%]; $p = 0.015$). As such, both institutions have moved away from this technique in favor of TA and TOETVA, respectively. Importantly, these trends in outcomes seen with RF at our institutions might be related to our respective case volumes in relation to the learning curve for the procedure, as others have reported favorable outcomes with the technique.^{6,11,19}

Although the RAA outcomes are in the context of a significantly smaller BMI than the TCA cohort, the mean BMI of 27.4 kg/m² is on par with the US national average of 28.7 kg/m² as per a 2012 study.³⁸ This suggests that our RAA cohort is an appropriate reflection of the North American patient population with regard to body habitus. In addition, both TA and TOETVA were successfully used to perform thyroidectomy in patients with BMI >40 kg/m² without complication. Future studies are needed to examine how BMI might affect case complexity and therefore operative time and complication rate via these RAA techniques.

Moving forward, we might understand that differences in outcomes between North American and Asian series observed initially were due to the lower case volume and differing levels of progress on the learning curves of these procedures, rather than inherent differences in patient population. As RAA techniques gain favor in North America once again, particularly with the recent interest in TOETVA, it is important that we maintain a framework for educating surgeons and institutions on the learning curves and ideal methods for successfully performing these techniques.^{25,35}

Ultimately, although this study demonstrates that these procedures can be performed safely, their widespread adoption will be dependent on the value they can provide, particularly in our current healthcare climate. Although cost is a frequent concern with these RAA techniques, other factors, such as quality of life outcomes and patients' willingness to pay to avoid cervical incisions, must be considered in determining value.³⁹ We understand that, similar to TCA, cost can be minimized when these RAA cases are performed by high-volume surgeons and centers.⁴⁰ In addition, as experience continues to grow with these techniques—particularly TOETVA, given its relative infancy and the fact that it does not use a surgical robot—we may see an additional decrease in operative time and, subsequently, cost.³⁵

CONCLUSIONS

The RAA thyroidectomy is safe and feasible in a North American patient population when performed by high-volume head and neck endocrine surgeons, with no significant differences in rates of RLN injury or

hypoparathyroidism compared with TCA. Although there is an associated increase in operative time with these RAA techniques, they can ultimately provide healthcare value in the appropriately selected patient and with the appropriate high-volume surgical teams. Accordingly, RAA approaches, particularly TA and TOETVA, are reasonable surgical options for thyroidectomy in a select North American patient population wishing to avoid an anterior-cervical scar.

Author Contributions

Study conception and design: Russell, Kang, Tufano, Kandil

Acquisition of data: Razavi, Garstka, Chen, Vasiliou

Analysis and interpretation of data: Russell, Razavi, Garstka, Chen, Vasiliou, Kang, Tufano, Kandil

Drafting of manuscript: Russell, Razavi, Garstka, Chen, Vasiliou, Kang, Tufano, Kandil

Critical revision: Russell, Razavi, Garstka, Chen, Vasiliou, Kang, Tufano, Kandil

REFERENCES

- Wilson EB. The evolution of robotic general surgery. *Scand J Surg* 2009;98:125–129.
- Marino MV, Shabat G, Gulotta G, Komorowski AL. From illusion to reality: a brief history of robotic surgery. *Surg Innov* 2018;25:291–296.
- Halsted WS. IV. (I) The Excision of Both Lobes of the Thyroid Gland for the Cure of Graves's Disease. (II) The Preliminary Ligation of the Thyroid Arteries and of the Inferior in Preference to the Superior Artery. *Ann Surg* 1913;58:178–182.
- Choi Y, Lee JH, Kim YH, et al. Impact of postthyroidectomy scar on the quality of life of thyroid cancer patients. *Ann Dermatol* 2014;26:693–699.
- Best AR, Shipchandler TZ, Cordes SR. Midcervical scar satisfaction in thyroidectomy patients. *Laryngoscope* 2017;127:1247–1252.
- Terris DJ, Singer MC, Seybt MW. Robotic facelift thyroidectomy: II. Clinical feasibility and safety. *Laryngoscope* 2011;121:1636–1641.
- Kuppersmith RB, Holsinger FC. Robotic thyroid surgery: an initial experience with North American patients. *Laryngoscope* 2011;121:521–526.
- Ban EJ, Yoo JY, Kim WW, et al. Surgical complications after robotic thyroidectomy for thyroid carcinoma: a single center experience with 3,000 patients. *Surg Endosc* 2014;28:2555–2563.
- Clark JH, Kim HY, Richmon JD. Transoral robotic thyroid surgery. *Gland Surg* 2015;4:429–434.
- Berber E, Berner V, Fahey TJ 3rd, et al. American Thyroid Association Statement on Remote-Access Thyroid Surgery. *Thyroid* 2016;26:331–337.
- Kandil E, Hammad AY, Walvekar RR, et al. Robotic thyroidectomy versus nonrobotic approaches: a meta-analysis examining surgical outcomes. *Surg Innov* 2016;23:317–325.
- Russell JO, Clark J, Noureldine SI, et al. Transoral thyroidectomy and parathyroidectomy—A North American series of robotic and endoscopic transoral approaches to the central neck. *Oral Oncol* 2017;71:75–80.
- Anuwong A. Transoral endoscopic thyroidectomy vestibular approach: a series of the first 60 human cases. *World J Surg* 2016;40:491–497.
- Jitpratoom P, Ketwong K, Sasanakietkul T, Anuwong A. Transoral endoscopic thyroidectomy vestibular approach (TOETVA) for Graves' disease: a comparison of surgical results with open thyroidectomy. *Gland Surg* 2016;5:546–552.
- Kim HY, Chai YJ, Dionigi G, et al. Transoral robotic thyroidectomy: lessons learned from an initial consecutive series of 24 patients. *Surg Endosc* 2018;32:688–694.
- Anuwong A, Ketwong K, Jitpratoom P, et al. Safety and outcomes of the transoral endoscopic thyroidectomy vestibular approach. *JAMA Surg* 2018;153:21–27.
- Razavi CR, Khadem MGA, Fondong A, et al. Early outcomes in transoral vestibular thyroidectomy: robotic versus endoscopic techniques. *Head Neck* 2018;40:2246–2253.
- Shan L, Liu J. A systemic review of transoral thyroidectomy. *Surg Laparosc Endosc Percutan Tech* 2018;28:135–138.
- Russell JO, Razavi CR, Al Khadem MG, et al. Anterior cervical incision-sparing thyroidectomy: comparing retroauricular and transoral approaches. *Laryngoscope Investig Otolaryngol* 2018;3:409–414.
- Dhillon VK, Rettig E, Noureldine SI, et al. The incidence of vocal fold motion impairment after primary thyroid and parathyroid surgery for a single high-volume academic surgeon determined by pre- and immediate post-operative fiberoptic laryngoscopy. *Int J Surg* 2018;56:73–78.
- American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer, Cooper DS, Doherty GM, Haugen BR, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2009;19:1167–2214.
- Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid* 2015;26: thy.2015.0020–thy.2015.0020.
- Kandil EH, Noureldine SI, Yao L, Slakey DP. Robotic transaxillary thyroidectomy: an examination of the first one hundred cases. *J Am Coll Surg* 2012;214:558–564.
- Razavi CR, Russell JO. Indications and contraindications to transoral thyroidectomy. *Ann Thyroid* 2017;2[5].
- Russell JO, Anuwong A, Dionigi G, et al. Transoral thyroid and parathyroid surgery vestibular approach: a framework for assessment and safe exploration. *Thyroid* 2018;28:825–829.
- Garstka M, Mohsin K, Ali DB, et al. Well-differentiated thyroid cancer and robotic transaxillary surgery at a North American institution. *J Surg Res* 2018;228:170–178.
- Terris DJ, Singer MC, Seybt MW. Robotic facelift thyroidectomy: patient selection and technical considerations. *Surg Laparosc Endosc Percutan Tech* 2011;21:237–242.
- Orloff LA, Wiseman SM, Berner VJ, et al. American Thyroid Association Statement on Postoperative Hypoparathyroidism: Diagnosis, Prevention, and Management in Adults. *Thyroid* 2018;28:830–841.
- Davis SF, Abdel Khalek M, Giles J, et al. Detection and prevention of impending brachial plexus injury secondary to arm positioning using ulnar nerve somatosensory evoked potentials during transaxillary approach for thyroid lobectomy. *Am J Electroneurodiagnostic Technol* 2011;51:274–279.

30. Choi JY, Lee KE, Chung KW, et al. Endoscopic thyroidectomy via bilateral axillo-breast approach (BABA): review of 512 cases in a single institute. *Surg Endosc* 2012;26:948–955.
31. Dionigi G, Lavazza M, Bacuzzi A, et al. Transoral endoscopic thyroidectomy vestibular approach (TOETVA): from A to Z. *Surg Technol Int* 2017;30:103–112.
32. Arora A, Garas G, Sharma S, et al. Comparing transaxillary robotic thyroidectomy with conventional surgery in a UK population: a case control study. *Int J Surg* 2016;27:110–117.
33. Duke WS, Holsinger FC, Kandil E, et al. Remote access robotic facelift thyroidectomy: a multi-institutional experience. *World J Surg* 2017;41:116–121.
34. Razavi CR, Fondong A, Tufano RP, Russell JO. Central neck dissection via the transoral approach. *Ann Thyroid* 2017;2[5].
35. Razavi CR, Vasiliou E, Tufano RP, Russell JO. Learning curve for transoral endoscopic thyroid lobectomy. *Otolaryngol Head Neck Surg* 2018;159:625–629.
36. Lee J, Yun JH, Choi UJ, et al. Robotic versus endoscopic thyroidectomy for thyroid cancers: a multi-institutional analysis of early postoperative outcomes and surgical learning curves. *J Oncol* 2012;2012:734541.
37. Park JH, Lee J, Hakim NA, et al. Robotic thyroidectomy learning curve for beginning surgeons with little or no experience of endoscopic surgery. *Head Neck* 2015;37:1705–1711.
38. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999-2010. *JAMA* 2012;307:491–497.
39. Nellis JC, Ishii M, Russell JO, et al. Perception of neck scars after thyroid surgery: is there social value in avoiding a neck scar? AAFPRS Spring Meeting. National Harbor, MD 2018.
40. Al-Qurayshi Z, Robins R, Hauch A, et al. Association of surgeon volume with outcomes and cost savings following thyroidectomy. *JAMA Otolaryngol Head Neck Surg* 2016;142:32–39.

Discussion



DR WILLIAM B INABNET III (New York, NY): Russell and colleagues from Tulane and Johns Hopkins present a 2-institution report of their collective experience with remote access thyroid surgery. Surgeons often learn only 1 of these techniques, and few teams learn more than 2; the authors are to be commended for developing the expertise to offer 3 different approaches, an accomplishment that permits personalization of surgical recommendations. The authors report no major complications in both the remote-access approach and transcervical approach cohorts, which is a major accomplishment that all endocrine surgeons, including myself, should strive to meet.

In the title of your paper, the words “a multi-institutional North American experience” is somewhat misleading in that the study was really performed at 2 institutions. This phrase fails to acknowledge or recognize the innovative work with remote access thyroid surgery being done in Durham, Chicago, Dallas, San Francisco, New York, and elsewhere in the country. Consideration should be given to changing your title to reflect that this study is really a 2-institution study, not a multi-institutional study that is reflective of the North American experience.

The universally accepted definition of permanent vocal cord paresis is vocal cord immobility > 6 months. The abstract in the book reports a permanent recurrent laryngeal nerve injury in 3 of 201 patients in the remote access group, and 3 of 382 in the conventional group, yet your manuscript and your presentation report no permanent recurrent laryngeal nerve injuries in either group. Please explain the discrepancy in this very important outcomes measure.

In the remote access group, you report a difference in the percentage of patients undergoing total thyroidectomy, 22% in the remote access group vs 41% in the conventional group. Knowing that the robotic faceless approach is really only applicable for unilateral thyroidectomy, what happens if you exclude those patients? Do the numbers even out with regard to total thyroidectomy? In high-volume thyroid programs, thyroid cancer typically comprises 30% to 40% of patients. Many of these patients require central lymph node dissection, yet in your analysis you excluded these patients. Please explain why they were excluded.

Finally, you note that the majority of cases were performed by 2 right-handed surgeons, yet the transaxillary and facelift cases were performed with robotic assistance, which really sort of negates the influence of the surgeon’s dominant hand. As a left-handed surgeon, I teach my trainees to use their nondominant hand if the instrument/tissue alignment ergonomically favors such. Why do you place such an emphasis on the handedness of the surgeons, and do you think this in any way affects outcomes?

DR HERBERT CHEN (Birmingham, AL): Certainly, experienced thyroid surgeons such as Drs Russell, Kandil, and Tufano can perform these technically challenging approaches. But like many operations, showing that you can perform them does not necessarily mean you should perform them. In full disclosure, I do not perform any of these procedures listed here.

A permanent hypoparathyroidism rate of zero is impressive and certainly due to the skill of these surgeons. However, at least 1 of the patients was excluded from the analysis because that patient had biochemical hypoparathyroidism but less than 6 months of follow-up. I do think this patient and other patients who still have that complication but have not reached the full 6 months, should be included as a permanent complication because they are as permanent at the time of the analysis.

Although the complication rate is similar between the groups, the operative time is significantly longer for the remote access procedures compared with the conventional. This leads to more anesthesia and more operating room (OR) time. There is also extra equipment. Have the authors examined the total OR costs between the remote access and conventional thyroid surgery?

Most high-volume endocrine surgery centers perform outpatient thyroidectomy. At our institution, virtually all patients who have conventional thyroid lobectomy or total thyroidectomy go home the same day. What percent of your conventional thyroid lobectomies and total thyroidectomies vs the remote access go home the same day? What is the mean length of stay for all 4 groups? If the length of stay is longer for remote access, why? Lastly, on what percent of your practice do you perform remote