



Correspondence

How to Approach Vulnerable Adolescents and Young Adults With Psychogenic Nonsyncopal Collapse



To the editor:

Available online 7 August 2019

Psychogenic nonsyncopal collapse (PNSC) in adolescents and young adults is a common and important issue, so we thank Heyer¹ and *Pediatric Neurology* for bringing attention to this vulnerable population. We would like to add an idea of how to approach these individuals.

PNSC is the basis of most cases of syncope, often leading to unnecessary research, loss of time, and increased workload. Pana et al.² showed that approximately one-quarter of all referrals to specialty epilepsy clinics are nonepileptic patients, those with psychiatric symptoms, syncope (probably mostly PNSC). Although physicians can make the diagnosis, they often refer these patients to specialized centers due to their fear of overlooking organic causes.² However, an accurate diagnosis can be made by a detailed history, minimal investigations, or even only by psychosocial evaluation of patients. In accordance with the proposal of Beghi et al.,³ we may engage psychiatrists into the diagnostic process because psychiatric disorders such as post-traumatic stress disorder, depression, anxiety, and eating disorders may also be present in these individuals.¹

In our opinion, psychiatrists should be included in the psychosocial evaluation process, diagnosis, and treatment of this population to prevent time and financial loss.

<https://doi.org/10.1016/j.pediatrneurol.2019.07.014>

Response to Yatim and Kılıç. “How to Approach Vulnerable Adolescents and Young Adults With Psychogenic Nonsyncopal Collapse”



A very acute pain in any of the more sensible parts of the body, or violent affections of the mind, as terror, grief, anger, or disappointments, will sometimes so strongly affect the whole nervous system, as to bring on hysteric faintings, with convulsions, altho' the body be in every respect healthful and sound...

Robert Whytt, MD, FRS (1765)

References

1. Heyer GL. Youth with psychogenic non-syncopal collapse have more somatic and psychiatric symptoms and lower perceptions of peer relationships than youth with syncope. *Pediatr Neurol.* 2018;79:34–39.
2. Pana R, Labbé A, Dubeau F, Kobayashi E. Evaluation of the “non-epileptic” patient in a tertiary center epilepsy clinic. *Epilepsy Behav.* 2018;79:100–105.
3. Beghi M, Erba G, Cornaggia CM, et al. Engaging psychiatrists in the diagnosis of psychogenic nonepileptic seizures. What can they contribute? *Seizure.* 2017;52:182–187.

Aylin Yetim, MD

Division of Adolescent Medicine, Department of Pediatrics, Istanbul University Istanbul School of Medicine, Istanbul, Turkey

Ayşe Kılıç, MD, PhD

Division of Adolescent Medicine, Department of Pediatrics, Istanbul University Istanbul School of Medicine, Istanbul, Turkey

Division of Social Pediatrics, Department of Pediatrics, Istanbul University Istanbul School of Medicine, Istanbul, Turkey
E-mail address: aylin.yetim@istanbul.edu.tr

In his 1765 monograph,¹ Whytt recognized that fainting attacks appeared differently in different patients. Some would “lie as in a deep sleep,” whereas others were “affected with catchings and strong convulsions.” Certainly, he was describing a conversion disorder (functional neurological symptom disorder) among some patients, although his description of fainting with “...a giddiness, a noise in the ears, and a loss of sight...” probably represented vasovagal syncope. Distinguishing the various forms of transient loss of consciousness continues to pose a challenge for present-day clinicians.

Psychogenic nonsyncopal collapse (PNSC) is a conversion disorder that resembles syncope. The attacks do not have corresponding hypotension, bradycardia, or electroencephalography changes that would occur with syncope but, rather, have a psychological origin. In their letter to the editor, Yetim and Kılıç² suggest that most syncope cases represent PNSC and that an accurate PNSC diagnosis can be made by “a detailed history, minimal investigations or by psychosocial evaluation [only]...” I disagree.

Most patients who present with fainting have vasovagal syncope, not PNSC. Stress can trigger syncope. So can pain. Some patients with PNSC will have bizarre signs and symptoms that can only be a conversion disorder, but most simply appear to faint. A detailed history can help to identify fainting features that are

atypical of syncope. The Fainting Assessment Inventory³ can also be used as a screening tool to help distinguish patients with PNCS from those with syncope. But a definitive diagnosis requires capture of an attack while blood pressure and heart rate (and preferably electroencephalography) are monitored. This is analogous to a documented diagnosis of psychogenic nonepileptic seizures.⁴ Relying solely on history will lead to diagnostic errors, and incorrectly diagnosing a conversion disorder in a patient with anxiety and vasovagal syncope would be grossly inappropriate.

Centuries after Whytt's descriptions of "hysterical faintings" we do not know enough about PNCS to make the diagnosis from history alone, at least in most patients.

References

1. Whytt R. *Observations on the Nature, Causes, and Cure of Those Disorders Which Have Been Commonly Called Nervous, Hypochondriac, Or Hysterical: To Which Are Prefixed Some Remarks on the Sympathy of the Nerves*. Edinburgh: Printed for T. Becket, and P. Du Hondt, London; and J. Balfour, Edinburgh; 1765.
2. Yetim A, Kılıç A. How to approach vulnerable adolescents and young adults with psychogenic nonsyncopal collapse. *Pediatr Neurol*. 2019;101:89–90.
3. Heyer GL. The Fainting Assessment Inventory: a 10-item instrument developed to screen for psychogenic nonsyncopal collapse among youth referred for syncope. *J Nerv Ment Dis*. 2019;207:255–263.
4. LaFrance Jr WC, Baker GA, Duncan R, Goldstein LH, Reuber M. Minimum requirements for the diagnosis of psychogenic nonepileptic seizures: a staged approach: a report from the International League Against Epilepsy Nonepileptic Seizures Task Force. *Epilepsia*. 2013;54:2005–2018.

Geoffrey L. Heyer, MD

E-mail address: geoffreyheyer@yahoo.com.

Available online 14 August 2019

<https://doi.org/10.1016/j.pediatrneurol.2019.08.003>