

Review

Risk factors for metachronous contralateral breast cancer: A systematic review and meta-analysis



Delal Akdeniz ^{a, b, c}, Marjanka K. Schmidt ^{b, c}, Caroline M. Seynaeve ^a, Danielle McCool ^b, Daniele Giardiello ^{b, e}, Alexandra J. van den Broek ^c, Michael Hauptmann ^b, Ewout W. Steyerberg ^{d, e}, Maartje J. Hooning ^{a, *}

^a Department of Medical Oncology, Family Cancer Clinic, Erasmus MC Cancer Institute, Rotterdam, Netherlands

^b Division of Psychosocial Research and Epidemiology, Netherlands Cancer Institute – Antoni van Leeuwenhoek Hospital, Amsterdam, Netherlands

^c Division of Molecular Pathology, Netherlands Cancer Institute – Antoni van Leeuwenhoek Hospital, Amsterdam, Netherlands

^d Department of Public Health, Erasmus MC, Rotterdam, Netherlands

^e Department of Medical Statistics and Bioinformatics, Leiden University Medical Centre, Leiden, Netherlands

ARTICLE INFO

Article history:

Received 13 September 2018

Received in revised form

12 November 2018

Accepted 16 November 2018

Available online 6 December 2018

Keywords:

Contralateral breast cancer

Metachronous

Risk factors

Systematic review

Meta-analysis

ABSTRACT

Background: The risk of developing metachronous contralateral breast cancer (CBC) is a recurrent topic at the outpatient clinic. We aimed to provide CBC risk estimates of published patient, pathological, and primary breast cancer (PBC) treatment-related factors.

Methods: PubMed was searched for publications on factors associated with CBC risk. Meta-analyses were performed with grouping of studies by mutation status (i.e., *BRCA1*, *BRCA2*, *CHEK2* c.1100delC), familial cohorts, and general population-based cohorts.

Results: Sixty-eight papers satisfied our inclusion criteria. Strong associations with CBC were found for carrying a *BRCA1* (RR = 3.7; 95%CI:2.8–4.9), *BRCA2* (RR = 2.8; 95%CI:1.8–4.3) or *CHEK2* c.1100delC (RR = 2.7; 95%CI:2.0–3.7) mutation. In population-based cohorts, PBC family history (RR = 1.8; 95%CI:1.2–2.6), body mass index (BMI) ≥ 30 kg/m² (RR = 1.5; 95%CI:1.3–1.9), lobular PBC (RR = 1.4; 95%CI:1.1–1.8), estrogen receptor-negative PBC (RR = 1.5; 95%CI:1.0–2.3) and treatment with radiotherapy <40 years (RR = 1.4; 95%CI:1.1–1.7) was associated with increased CBC risk. Older age at PBC diagnosis (RR per decade = 0.93; 95%CI:0.88–0.98), and treatment with chemotherapy (RR = 0.7; 95%CI:0.6–0.8) or endocrine therapy (RR = 0.6; 95%CI:0.5–0.7) were associated with decreased CBC risk.

Conclusions: Mutation status, family history, and PBC treatment are key factors for CBC risk. Age at PBC diagnosis, BMI, lobular histology and hormone receptor status have weaker associations and should be considered in combination with key factors to accurately predict CBC risk.

© 2018 Elsevier Ltd. All rights reserved.

Contents

1. Introduction	2
2. Methods	2
2.1. Search strategy	2
2.2. Statistical analyses	2
2.3. Quality assessment	6
3. Results	7
3.1. Quality assessment	9
4. Discussion	9
4.1. Implications for future research	11

* Corresponding author. Department of Medical Oncology, Erasmus MC Cancer Institute, PO Box 2040, 3000 CA, Rotterdam, Netherlands.

E-mail address: m.hooning@erasmusmc.nl (M.J. Hooning).

4.2. Clinical implications	11
5. Conclusion	11
Funding	11
Acknowledgements	11
Supplementary data	11
References	11

1. Introduction

Due to an increasing incidence of primary breast cancer (PBC) and improved breast cancer (BC) surveillance and treatment methods, an increasing number of women who have survived BC are at risk of developing a contralateral breast cancer (CBC) [1]. The annual CBC risk is around 0.5% in the general BC population and up to 3% in *BRCA1/2* mutation carriers [2,3].

A risk-reducing contralateral mastectomy minimizes the risk of developing a subsequent CBC and may improve survival in patients considered to be at high risk, i.e. hereditary BC patients [4–6]. On the other hand, the percentage of patients opting for a risk-reducing contralateral mastectomy has rapidly increased over the last decades, suggesting that more relatively low-risk BC patients are also treated [7–9]. Fear and overestimation of risk may play a role in the decision-making of these low-risk patients [10,11].

For both high-risk and low-risk PBC patients, accurate CBC risk prediction is crucial and can be achieved by taking into account the effect of patient, pathological, and treatment-related characteristics. However, CBC risk prediction as used in clinical practice is currently only based on *BRCA1/2* mutation status, family history of BC and age at PBC [2,12,13]. The association of other factors with CBC risk is either lacking or conflicting. Combinations of these factors may improve decision-making regarding surveillance, primary and risk-reducing therapies, and may enable patient-tailored counselling in both high-risk and low-risk patients.

Therefore, we aimed to quantify the association of various patient, pathological, and treatment-related characteristics with metachronous CBC risk.

2. Methods

For this systematic review we published an online protocol at Prospero including details on study design (registration number: CRD42015014381, link: http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42015014381) and we followed the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines.

2.1. Search strategy

In collaboration with a research librarian (EdC, see acknowledgements) a search strategy was developed. One reviewer searched PubMed for publications on search terms for metachronous CBC in combination with various predefined patient characteristics (carriership of *BRCA1*, *BRCA2* and *CHEK2* c.1100delC mutations, family history of (bilateral) BC, mammographic density, factors at PBC diagnosis: age, BMI, menopausal status), PBC characteristics (TN(M)-stage, tumor grade, Estrogen (ER), Progesterone (PR) and HER2 neu receptor status, histological subtype), and PBC treatment-related characteristics (radiotherapy, chemotherapy, endocrine therapy, targeted therapy, risk-reducing salpingo-oophorectomy (RRSO)). We also searched for publications on second BC risk, in the knowledge that a majority (95%) of the second breast

cancers are contralateral events rather than ipsilateral breast tumors [14]. Details of the full strategy applied are provided in [Supplementary Table A.1](#).

Abstracts were screened using the following inclusion criteria: experimental and observational studies published in English, between January 1990 and July 4, 2016, investigating CBC risk in women who have no prior history of other invasive malignancies. We included papers only from 1990 onwards to have a long-term follow-up while also being able to investigate the effects of adjuvant treatment options (which were considered mainly from the late eighties onwards). Further, we excluded papers if the reported number of second BC events was less than twenty (arbitrary cut-off), and also if no relative risk (RR) estimates (hazard ratio or odds ratio or relative risk) for CBC risk were provided.

Relevant full-text publications were considered for inclusion and critically appraised, on methodology, and comparability of groups, subgroups and their reference groups. If papers reported on specific subgroups that were non-combinable with other subgroups, these papers were excluded for the meta-analysis. In addition, potential overlap in (part of) patients due to selection from the same registries/hospitals in the same period was solved by selecting the most relevant cohort (i.e. the factor of interest for the meta-analysis was specifically published on) and/or selecting the most recent cohort with the longest follow-up. From the included papers, study design characteristics and all the available univariable and multivariable risk estimates were extracted and entered in a Microsoft Access database by four reviewers (DA, MKS, AjvdB, MJH) using a specifically designed data entry form.

2.2. Statistical analyses

We investigated the effects of carrying vs. not carrying a *BRCA1*, *BRCA2* or *CHEK2* c.1100delC mutation on the risk of developing CBC. We also investigated the effects of the aforementioned patient, pathological, and treatment characteristics separately in five different groups: 1. *BRCA1* mutation carriers; 2. *BRCA2* mutation carriers; 3. *CHEK2* c.1100delC mutation carriers; 4. Familial BC patients, i.e. patients who tested negative for a *BRCA1/2* or *CHEK2* c.1100delC mutation; 5. Population-based cohorts, i.e. patients from hospitals or official registries representing the general population, that have not been selected on gene mutation carriership or a positive family history for BC.

Papers with only combined results for *BRCA1* and *BRCA2* mutation carriers were excluded from the analyses, as these two groups represent different entities with different characteristics and should be analyzed separately (*BRCA1* mutation carriers are younger at PBC diagnosis and present more often with a triple negative BC phenotype (ER, PR and HER2 receptor negative) than *BRCA2* mutation carriers) [2,15,16]. For the analyses on carriership of a genetic *BRCA1*, *BRCA2* or *CHEK2* c.1100delC mutation, we included studies where the reference group consisted of familial patients (i.e. patients from non-*BRCA1/2*, and/or *CHEK2*-negative BC families) and excluded papers that used a sporadic population as a reference group. After all, studies that compare mutation carriers

recruited from Clinical Genetic departments with BC patients from the general population easily lead to overestimations [17]. Since this is no issue in population-based studies with genetic test results generally available, these studies were included as well.

Since various ranges for age were used in the different papers, we estimated the overall effect of age using the method described by Greenland et al. [18], typically defined in the context of dose-response studies. The requirements needed for this method are the risk estimates from every age category, the corresponding confidence levels or the standard errors, and the number of cases and controls or person-time in case of incidence rate data. If these were not given, the continuous age effect was estimated by linearly regressing the category-specific log relative risks on an age value representative for each age category. Representative values were

the median age at PBC diagnosis calculated from female BC patients in the Netherlands Cancer Registry [19], with a 10-year CBC risk of 4% which is comparable with published results from studies from various western countries [20–22].

All types of relative risk estimates were log transformed and subsequently pooled for every factor of interest. The available univariable and multivariable estimates were analyzed separately (and reported as crude and adjusted analyses, respectively). If only subgroup estimates were available in a paper, we combined these estimates to generate an overall estimate. A random effects model was used to perform the meta-analyses [23]. We tested for heterogeneity using I^2 statistics and the p -value for heterogeneity using the Cochran's Q-statistic was reported.

To conduct the meta-analyses, we used Metan from the Stata




Fig. 1. PRISMA flow chart of papers on risk factors for contralateral breast cancer. Abbreviations CBC = Contralateral breast cancer; SNP = Single Nucleotide Polymorphism.

Table 1

Study characteristics of papers publishing on risk factors for contralateral breast cancer included in the systematic review.

First author, Year	Country/Continent ^a	Study Design ^a	Population	Factors ^a	Mean follow-up total group (years) ^{a,b}	Mean age total group (years) ^a	N Patients included	N CBCs ^c
van den Broek, 2016 [2]	Netherlands	Cohort	(non-) BRCA1/2	FamHis, Age, Ctx, DNA,	Md. 12.5	N/A	6294	578
Goss, 2016 [25], ^e Aalders, 2016 [26]	USA, Europe, Canada Netherlands	RCT Cohort	Unselected Unselected	EtX Age, TNM, Grade, ER,PR,HER2, His	Md. 6.3 5.0	Md. 65.1 59.0	1918 52 626	43 1534
Sisti, 2015 [27]	USA, Denmark, Canada	(Nested) Case-Control	Unselected	Meno	Md Cases 6.3 Controls: 5.5 (matched)	46.0 (matched)	3733	1521
Menes, 2015 [16], ^e Kiderlen, 2015 [28]	USA, Australia, Canada Netherlands	Cohort Cohort	BRCA1/2 Unselected	Ctx, EtX, Rtx, RRSO Age	8.9 Md. 7.2	41.0 Md. 74.9	800 2926	86 75
Drooger, 2015 [29], ^e Basu, 2015 [30], ^e	Netherlands United Kingdom	Cohort Cohort	BRCA1/2 BRCA1/2	Age, Ctx, EtX, Rtx, RRSO, DNA Age, Meno, RRSO, DNA	Md. 8.6 Md. 7.8	N/A NA	691 1011	161 202
Rasmussen, 2014 [20] Mellemkjaer, 2014 [31]	Denmark Denmark	Cohort Cohort	Unselected Unselected	Age EtX	Md. 5.6 N/A	N/A N/A	85 863 37 533	3120 124
Kriege, 2014 [32]	Netherlands	Cohort	(non-) CHEK2	DNA	Md. 7.2 Non-carriers: 7.2	N/A	3502	197
Gronwald, 2014 [33]	USA, Europe, Canada	(Nested) Case-Control	BRCA1/2	EtX	7.2	50.9 (matched)	1504	411
Calip, 2014 [34]	USA	Cohort	Unselected	BMI	Md. 6.3	Md. 63.0	4216	145 (ipsilateral: n.a.)
van de Water, 2013 [35] Valuckas, 2013 [36]	USA, Japan, Europe Lithuania	RCT Cohort	Unselected Unselected	Age Age, Meno, BMI, TNM, Ctx, EtX, Rtx,	Md. 5.1 Md HRtx 10.1 CRT 10.4	Md. 64.0 Md. 53.4	9766 832	83 48 (ipsilateral: n.a.)
Sandberg, 2013 [37]	Sweden	(Nested) Case-Control	Unselected	BD	Cases: 8.25 Controls: 8.25 (matched)	(matched)	422	211
Reiner, 2013 [12]	USA, Denmark	(Nested) Case-Control	Non-BRCA1/2	FamHis	(matched)	(matched)	1713	594
Phillips, 2013 [15]	USA, Australia, New Zealand, Europe, Canada	Cohort	BRCA1/2	EtX	Md. 6.6	N/A	2464	520
Pacelli, 2013 [38], ^e Metzger-Filho, 2013 [39], ^e	Italy Australia, New Zealand, Europe, India, South-America, Africa	Cohort RCT	Unselected Unselected	ER/PR/HER2 ER/PR/HER2	Md. 4.9 Md. 12.5	Md. 53.0 53.9	468 1951	24 75
Mavaddat, 2013 [40] Maskarinec, 2013 [41], ^f	United Kingdom USA	Cohort Cohort	BRCA1/2 Unselected	RRSO BD	Md. 2.6 12.9	Md. 39.5 63.3	988 607	61 71
Dellapasqua, 2013 [42] Courdi, 2013 [43]	Italy France	Cohort Cohort	Unselected Unselected	Age, TNM, ER/PR/HER2 Rtx	Md. 6.3 Md. 12.8	Md. 52.0 N/A	6971 1630	129 116
Bernstein, 2013 [44]	USA, Denmark	(Nested) Case-Control	BRCA1/2	Rtx, DNA	(matched)	(matched)	1802	603
Weischer, 2012 [45]	USA, Australia, Europe, Canada	Cohort	CHEK2	DNA	Md. 6.6	N/A	25 094	647 (ipsilateral: n.a.)
Vichapat, 2012 [46] Saltzman, 2012 [47], ^f	Sweden USA	Cohort (Nested) Case-Control	Unselected Unselected	Age, TNM, His, EtX ER, PR, HER2	Md. 6.7 (matched)	N/A (matched)	37 393 1988	894 482
Neta, 2012 [48], ^f Mavaddat, 2012 [49]	USA USA, Australia, Europe	Cohort Cohort	Unselected BRCA1/2	Rtx ER	10.0 N/A	N/A N/A	205 316 6893	6924 1022
Filleron, 2012 [50] Brooks, 2012 [51]	France USA, Denmark	RCT (Nested) Case-Control	Unselected Unselected	Age, TNM, Grade BMI	Md. 4.4 Md. 4.2	Md. 49.0 Md. 45.0 (matched)	2820 1510	58 511
Zhang, 2011 [22] Vichapat, 2011 [52]	Italy United Kingdom	Cohort Cohort	Unselected Unselected	Rtx FamHis, Age, Meno, TNM, Grade, ER, PR, HER2, His, Ctx, EtX, Rtx	8.0 N/A	54.7 N/A	5248 4366	261 315
Metcalfe, 2011 [53]	USA, Canada	Cohort	BRCA1/2	FamHis, Age, TNM, Grade, ER, Ctx, EtX, Rtx, RRSO, DNA	11.1	Md. 42.0	846	149
Majed, 2011 [54]	France	Cohort	Unselected	FamHis, Age, Meno, BMI, TNM, Grade, ER, PR, His, Ctx, EtX, Rtx	Md. 10.0	54.0	15 166	1370
Hackshaw, 2011 [55], ^e Bouchardy, 2011 [56], ^f	Europe, Asia Switzerland	RCT Cohort	Unselected Unselected	EtX FamHis, Age, ER, EtX	Md. 10.1 Md. 5.2	Md. 62.0 59.8	3449 4152	118 63
Rubino, 2010 [57], ^e Rondeau, 2010 [58]	France France	Cohort Cohort	Unselected Unselected	Age, TNM TNM, Grade, Ctx	Md. 10.6 Md. 12.7	56.0 Md. 57.0	6629 919	673 69

Table 1 (continued)

First author, Year	Country/Continent ^a	Study Design ^a	Population	Factors ^a	Mean follow-up total group (years) ^{a,b}	Mean age total group (years) ^a	N Patients included	N CBCs ^c
Reding, 2010 [59]	USA, Denmark	(Nested) Case-Control	(non-) BRCA1/2	Ctx, Etx	(matched)	(matched)	1579	181
Poynter, 2010 [60], ^e	USA, Denmark	(Nested) Case-Control	(non-) BRCA1/2	Meno	(matched)	(matched)	2103	181
Malone, 2010 [3]	USA, Denmark	(Nested) Case-Control	BRCA1/2	DNA	(matched)	Md. 46.0 (matched)	2103	705
Cuzick, 2010 [61], ^e Buist, 2010 [62]	Unknown USA	RCT Cohort	Unselected Unselected	Etx FamHis, Age, BD, TNM, Ctx, Etx,	Md. 10.0 N/A	Md. 72.0 N/A	6241 17 286	178 344 (ipsilateral: 54)
Berrington de Gonzalez, 2010 [63]	USA	Cohort	Unselected	Rtx	13.0	N/A	182 057	6491
Li, 2009 [64] A	USA	(Nested) Case-Control	Unselected	Etx	(matched)	(matched)	1094	367
Li, 2009 [65], ^d B	USA	(Nested) Case-Control	Unselected	BMI	(matched)	(matched)	1091	365
Graeser, 2009 [13], ^e Bertelsen, 2009 [66] Alkner, 2009 [67] Stovall, 2008 [68], ^d	Germany Denmark Sweden USA, Denmark	Cohort Cohort RCT (Nested) Case-Control	BRCA1 Unselected Unselected Unselected	DNA Age Age, Etx Rtx	N/A 8.4 Md. 14.0 Cases: 5.0 Controls: 5.0 (matched)	N/A N/A N/A 51.0 (matched)	2020 8737 564 1806	381 466 52 606
Schaapveld, 2008 [69] Mellemkjaer, 2008 [70]	Netherlands USA, Denmark	Cohort (Nested) Case-Control	Unselected (non-) CHEK2	Age, TNM, Ctx, Etx, Rtx Ctx, Rtx, DNA	Md. 5.8 5.0	N/A (matched)	45 229 2103	1477 708
Hoening, 2008 [71] Bertelsen, 2008 [72]	Netherlands USA, Denmark	Cohort (Nested) Case-Control	Unselected Unselected	FamHis, Age, Ctx, Rtx Ctx, Etx	Md. 13.8 (matched)	N/A Md. 46.0 (matched)	7221 1792	503 634
van der Leest, 2007 [73], ^e Trentham-Dietz, 2007 [74], ^f	Netherlands USA	Cohort Cohort	Unselected Unselected	Ctx, Etx FamHis, Meno, BMI	Md. 8.5 7.1	Md. 37.5 59.4	758 10 953	59 488 (ipsilateral: n.a.)
Schmidt, 2007 [75], ^{d,f}	Netherlands	Cohort	CHEK2	DNA	Md. 10.1	43.0	1479	124 (ipsilateral: 13)
Rutqvist, 2007 [76] Largent, 2007 [77], ^e	Sweden USA, Denmark	RCT (Nested) Case-Control	Unselected Unselected	Etx Meno (at CBC diagnosis)	Md. 18.0 Cases: 5.0 Controls: 5.0 (matched)	N/A 45.0 (matched)	2738 2107	170 708
Kirova, 2007 [78] Hemminki, 2007 [79] Broeks, 2007 [80]	France Sweden Netherlands	Cohort Cohort Case-Only	Unselected Unselected BRCA1/2, CHEK2	Rtx FamHis Rtx	Md. 10.5 N/A N/A	Md. 55.0 N/A N/A	16 705 102 176 247	1343 5495 247
Brekeldmans, 2007 [81], ^{e,f}	Netherlands	Cohort	(non-) BRCA1/2	DNA	Md BRCA1/2: 4.3 NonBRCA: 4.8 Sporadic: 5.1	N/A	498	53
Tilanus-Linthorst, 2006 [82], ^e Pierce, 2006 [83], ^e	Netherlands USA, Israel	Cohort Cohort	Non-BRCA1/2 BRCA1/2, Unselected	FamHis Age, TNM, Ctx, Etx, DNA	Md 6.1 Md BRCA1/2: 7.9 Unselected: 6.7	(matched) (matched)	654 605	51 48
Levi, 2006 [84] Gronwald, 2006 [85], ^d	Switzerland USA, Israel, Europe, Canada	Cohort (Nested) Case-Control	Unselected BRCA1/2	Rtx Etx	7.8 Etx: 5.7 No Etx: 7.4	N/A N/A	6119 1007	222 356
Dignam, 2006 [86] Brekeldmans, 2006 [87], ^{e,f}	USA Netherlands	Cohort Cohort	Unselected BRCA1	BMI DNA	N/A Md. 5.1	N/A Md. 39.0 (matched)	4077 669	242 75
Nordenskjold, 2005 [88], ^e Roychoudhuri, 2004 [89] McCaskill-Stevens, 2004 [90], ^d Coombes, 2004 [91], ^e Li, 2003 [92], ^f	Sweden United Kingdom USA, Australia, South-America, Ireland USA, Australia, Europe USA	RCT Cohort RCT RCT Cohort	Unselected Unselected Unselected Unselected Unselected	Etx Rtx Etx Etx Etx	Md. 10.6 N/A N/A N/A Md. 2.6 9.0	N/A N/A Md. 50! 64.3 37.7	4610 64 782 10 619 4742 1285	138 308 494 29 77

(continued on next page)

Table 1 (continued)

First author, Year	Country/Continent ^a	Study Design ^a	Population	Factors ^a	Mean follow-up total group (years) ^{a,b}	Mean age total group (years) ^a	N Patients included	N CBCs ^c
				FamHis, Age, BMI, TNM, ER, PR, His				
Gao, 2003 [21], ^d	USA	Cohort	Unselected	Age, Rtx, His	Rtx: 5.7 No Rtx: 6.8	61.0	134 501	5679
Dignam, 2003 [93]	USA, Unknown	RCT	Unselected	BMI	Md. 13.8	N/A	3385	193
Fisher, 2002 [94]	USA, Canada	RCT	Unselected	Etx	Md. 7.2	N/A	1000	27
Li, 2001 [95], ^d	USA	Cohort	Unselected	Etx	Etx: 3.9 No Etx: 4.2	N/A	8981	189
Fisher, 2001 [96]	USA, Unknown	RCT	Unselected	Etx	Md. 6.8	56.0	1172	37
Vahtinen, 2000 [97]	Sweden	Cohort	Unselected	FamHis, Age	6.2	N/A	72 092	2529
Narod, 2000 [98], ^d	USA, Europe, Canada	(Nested) Case-Control	BRCA1/2	Ctx, Etx, Rtx, RRSO	9.7	40.2 (matched)	593	209
Matsuyama, 2000 [99]	Japan	Cohort	Unselected	Etx	Md. 7.6 Etx: 7.6 No Etx: 8.1	51.0	6148	30
Robson, 1999 [100], ^e	USA	Cohort	BRCA1/2	DNA	Md. 10.3	N/A	305	42
Newcomb, 1999 [101]	USA	Cohort	Unselected	Etx	6.4	N/A	54 821	1730
Kollias, 1999 [102]	United Kingdom	Cohort	Unselected	FamHis, Age, His	Md. 9.0	Md. 54.0	3211	83
Broet, 1999 [103]	France	Cohort	Unselected	Ctx	7.9	56.0	6185	334
Early Breast Cancer Trialists' Collaborative, 1998 [104], ^d	USA, New Zealand, Europe, South-America, Africa, Asia	RCT	Unselected	Etx	2.7	N/A	32 422	839
Swedish Breast Cancer Cooperative Group, 1996 [105], ^e	Sweden	RCT	Unselected	Etx	Md. 5.5	N/A	3545	51
Cook, 1996 [106]	USA	(Nested) Case-Control	Unselected	FamHis, Meno, BMI, ER, PR, His, Ctx, Rtx,	(matched)	N/A	640	216
Cook, 1995 [107], ^d	USA	(Nested) Case-Control	Unselected	Etx	Md. 3.3 (matched)	(matched)	673	234
Broet, 1995 [108]	Europe	Cohort	Unselected	His, Ctx	Md. 6.7	55.5	4748	282
Healey, 1993 [109]	USA	Cohort	Unselected	Age, TNM, Ctx, Etx	Md. 7.9	53.0	1624	77
Storm, 1992 [110], ^f	Denmark	(Nested) Case-Control	Unselected	FamHis, Meno, BMI, Rtx	(matched)	51.0 (matched)	56 540	529
Boice, 1992 [111], ^f	USA	(Nested) Case-Control	Unselected	Rtx	(matched)	51.7 (matched)	1844	655
Bernstein, 1992 [112], ^f A	USA	Cohort	Unselected	Age, Meno, BMI, His, Ctx, Rtx,	4.3	44.3	4550	136
Bernstein, 1992 [113] B	USA	Cohort	Unselected	FamHis	N/A	N/A	4660	136
Baum, 1992 [114]	United Kingdom	RCT	Unselected	Etx	Md. 7.8	55.1	1912	21
Andersson, 1991 [115], ^d	Denmark	RCT	Unselected	TNM, Etx	Md. 7.9	N/A	3538	143

^a Abbreviations: Age = age at PBC diagnosis, BD = breast density, BMI = body mass index, Ctx = chemotherapy, DNA = BRCA1/BRCA2/CHEK2 c.1100delC DNA mutation, ER = Estrogen hormone receptor status, Etx = endocrine therapy, FamHis = family history, Grade = tumor grade, HER2 = HER2 receptor status, His = histology, Md = median, Meno = menopausal status, N/A = not available, PR = Progesterone hormone receptor status, RCT = Randomized controlled trial RRSO = risk-reducing salpingo-oophorectomy, Rtx = radiotherapy, TNM = TNM-stage, USA = United States of America.

^b Md. median if no mean was given; (matched): patients were matched on follow-up; if no mean/median follow-up for the total group was given, the mean/median follow-up for the exposed versus the reference group was given.

^c If the number of CBC events was not provided, this was calculated from the figure in the paper; (ipsilateral): ipsilateral second breast cancers were included in the analyses.

^d Studies not used for the analyses due to overlap in patients.

^e Studies not used for the analyses due to reporting on subgroups that could not be combined with another estimate for the meta-analyses.

^f Studies that included patients with metastatic disease at primary breast cancer diagnosis as well.

Statistical Software package (version 14.0). To assess the effects of age at PBC diagnosis, the dosresmeta package from R software (version 3.2.2) was used.

2.3. Quality assessment

We used the QUality In Prognostic Studies (QUIPS) tool for assessing the quality and bias in the included papers [24]. As suggested by the developers of this tool, we modified the domains to be applicable to the specific study questions in our systematic review (Supplementary Table A.2). We excluded one domain, which assessed outcome measurement, since this was performed similarly in all studies and in a following domain we already scored whether a definition for outcome was given.

Using the modified tool, two reviewers (DA, MJH) scored 11 items in five domains. Every item was assigned 0 points if bias was unlikely, 0.5 points if bias was possibly present and 1 point if bias was likely present. When in doubt, the reviewers discussed with the other authors to reach consensus. The distribution of points for potential bias following the QUIPS tool was inspected using a boxplot; the overall mean score was 1.8 points (range 0–5.5). Results were comparable for case-control (2.0), cohort studies (1.8) and randomized controlled trials (1.8). Papers that were classified as high-quality papers (i.e. on a scale of 0–11 a total bias score of <2 was assigned; Supplementary Table A.3), were analyzed separately using a random-effects model.




Fig. 2. Forest plot of the adjusted meta-analyses per patient, pathological and treatment-related characteristic on the risk of developing contralateral breast cancer in population-based cohorts; Abbreviations: BMI = body mass index per kg/m²; ER = Estrogen hormone receptor; PR = Progesterone hormone receptor; T1 = tumor size ≤2 cm; T2 = tumor size 2.1–5.0 cm; T3 = tumor size >5.0 cm; Total_N = number of papers used for the analysis. Age concerns the age at primary breast cancer diagnosis; family history concerns the family history of breast cancer; estimate is a relative risk estimate combining hazard ratios, odds ratios and relative risks; I² test for heterogeneity; p-value for heterogeneity: p < 0.05 considered significant; patient and pathological factors are assessed at primary breast cancer diagnosis; treatment-related characteristics concern primary breast cancer treatment.

3. Results

In total, 100 papers out of 1789 identified records fulfilled the inclusion criteria (Flow diagram, see Fig. 1) [2,3,12,13,15,16,20–22,25–115]; study characteristics are depicted in Table 1. Eligibility was validated for 10% of the titles and abstracts by a second reviewer. Subsequently, potential overlap in patients included in different papers was evaluated and we selected either the most relevant (i.e. on topic) or most recent paper (n = 11 excluded). In addition, we evaluated whether risk estimates in their given form were usable and/or combinable (n = 21 excluded).

Eventually, 68 papers were used for the meta-analyses and these included between 247 and 205 316 PBC patients and 21 and 6924 second BCs per study. Twenty studies used data from patients diagnosed in Northern America (USA/Canada) solely, versus 24 European studies. The risk estimates mainly concerned population-based cohorts; for the specific genetic groups of interest and the familial BC group the number of estimates was limited (Supplementary Table A.4).

In the summary estimates reported below, the adjusted

estimates are reported (Figs. 2–5). Crude estimates are only provided in the main paper if the number of multivariable estimates was insufficient to perform a meta-analysis. An overview of the results from the crude analyses can be found in Supplementary Figure (S Fig.) B.1. Study-specific estimates per factor and per group of interest are provided in S Figs. B.2–B.40.

Population-based cohorts: Patient characteristics (Fig. 2; S Figs. B.2–B.12).

For the analyses concerning patient characteristics we reviewed 30 papers. Having a positive family history of BC was associated with an increased risk of CBC, but heterogeneity was substantial (RR = 1.72; 95% CI: 1.15–2.57; I² 93.1%; S Fig. B.2). The study performed by Hemminki et al. [79] was the main outlier. They used a non-conventional method to determine CBC risk, by doubling the risk, leading to overestimation. Heterogeneity as well as the relative risk estimate decreased when ignoring this study (RR = 1.43; 95% CI: 1.22–1.68; I² 41.6%).

CBC risk appeared to be higher in first than in second degree relatives (RR = 1.54; 95% CI: 1.25–1.90 and RR = 1.17; 95% CI: 0.90–1.52, respectively; S Figs. B.3 and B.4). Heterogeneity was also




Fig. 3. Forest plot of the adjusted meta-analyses comparing carrying a *BRCA1*, *BRCA2* or *CHEK2* c.1100delC mutation with patients who did not have the genetic mutation on the risk of developing contralateral breast cancer; Abbreviations: Total_N = number of papers used for the analysis. Estimate is a relative risk estimate combining hazard ratios, odds ratios and relative risks; I²: test for heterogeneity; p-value for heterogeneity: $p < 0.05$ considered significant.

present in the meta-analysis concerning first degree relatives (I² 60.1% vs. 0% in second degree relatives). Excluding the results from Buist et al. [62], which was the main outlier in this analysis, resulted in a decrease in heterogeneity and small increase in CBC risk (RR = 1.61; 95% CI: 1.41–1.85; I² 15.3%).

Age at PBC diagnosis was associated with a 7% decrease in CBC risk per decade (RR = 0.93; 95% CI: 0.88–0.98, I² 86.9%; S Fig. B.5). Although heterogeneity between studies was substantial, the estimates from the individual papers did not seem to vary widely.

For mammographic breast density (S Figs. B.6 and B.7) and menopausal status (S Fig. B.8) no association with CBC risk was observed.

Being overweight or obese (BMI ≥ 25 kg/m²) or being obese (BMI ≥ 30 kg/m²) compared to having normal weight (BMI < 25 kg/m²), was associated with an increased risk of developing CBC (RR = 1.26; 95% CI: 1.10–1.44; I² 44.4% and RR = 1.54; 95% CI: 1.26–1.87; I² 0%, respectively; S Figs. B.9 and B.11).

Population-based cohorts: Pathological characteristics (Fig. 2; S Figs. B.13–B.19).

For the analyses concerning pathological characteristics we analyzed 15 papers. Having a PBC with a larger size was associated with increased CBC risk (tumor size > 2 cm vs. ≤ 2 cm; RR = 1.17; 95% CI: 1.03–1.34; I² 21.6%; S Fig. B.13). For nodal status and tumor grade no association with CBC was observed (S Fig. B.14 and B.15, respectively). Both negative ER and PR hormone receptor status (vs. positive) were associated with an increased risk of CBC as well (RR = 1.53; 95% CI: 1.04–2.26; S Fig. B.16; and RR = 1.23; 95% CI: 1.02–1.48; S Fig. B.17, respectively), although for ER status there was evidence of substantial heterogeneity (I² 67.3% vs. 0% for PR status). Excluding the outlying estimate reported by Filleron et al. [50] (possibly large effect size due to a small study population available for this factor), resulted in a decrease in heterogeneity and a non-

significant association between ER status and CBC risk (RR = 1.32; 95% CI: 0.99–1.76; I² 38.5%). For Her2 status no association with CBC risk was observed (S Fig. B.18).

Lobular morphology vs. ductal/non-lobular morphology was also associated with an increased risk of developing CBC, which in the forest plot was observed mainly in the older publications (RR = 1.43; 95% CI: 1.13–1.82; I² 42.5%; S Fig. B.19).

Population-based cohorts: Treatment-related characteristics (Fig. 2; S Figs. B.20–B.27).

Nine papers were included on treatment with adjuvant chemotherapy and 15 studies on adjuvant endocrine therapy; both factors were associated with a lower CBC risk (RR = 0.70; 95% CI: 0.62–0.79; I² 21.3%; S Fig. B.20 and RR = 0.61; 95% CI: 0.53–0.72; S Fig. B.21, respectively). Results for patients aged below and above 50 years at PBC diagnosis were similar (data not shown). Heterogeneity was high in the meta-analysis concerning endocrine therapy (I² 73.6%), but decreased substantially (I² 19.4%) when we selected papers including only patients with ER-positive tumors (RR = 0.57; 95% CI: 0.49–0.66; S Fig. B.22).

Treatment with radiotherapy (vs. no radiotherapy) was analyzed in 8 papers and associated with a modestly increased CBC risk when diagnosed at least five years after PBC (RR = 1.10; 95% CI: 1.05–1.15; I² 0%; S Fig. B.24). In patients aged below 40 years at PBC diagnosis this risk appeared to be higher, both for CBCs occurring any time after PBC and for CBCs occurring at least 5 years after PBC diagnosis (RR = 1.37; 95% CI: 1.13–1.66; I² 0% and RR = 1.34; 95% CI: 1.07–1.67; I² 0%, respectively; S Figs. B.26 and B.27). The association appeared to attenuate when the age cut-off was raised to 45 years at PBC diagnosis (RR = 1.22; 95% CI: 1.09–1.36, I² 0.0% and RR = 1.20; 95% CI: 1.06–1.35, I² 0.0%, respectively, data not shown).

Mutation carriers vs. patients from mutation-negative BC families (Fig. 3, S Figs. B.28–B.30).




Fig. 4. Forest plot of the overall adjusted meta-analyses per patient, pathological or treatment-related characteristic on the risk of developing contralateral breast cancer in *BRCA1* and *BRCA2* mutation carriers. Abbreviations: Total_N = number of papers used for the analysis. Age concerns the age at primary breast cancer diagnosis; family history concerns the family history for breast cancer; estimate is a relative risk estimate combining hazard ratios, odds ratios and relative risks; I²: test for heterogeneity; p-value for heterogeneity: $p < 0.05$ considered significant; treatment-related characteristics concerns primary breast cancer treatment.

The effect of mutation status on CBC risk was analyzed in 5 papers [2,3,32,45,70]. Carriership of a *BRCA1*, *BRCA2* or *CHEK2* c.1100delC mutation vs. non-carriership was associated with an increased risk of CBC (RR = 3.68; 95% CI: 2.76–4.89; I² 12.4%; RR = 2.75; 95% CI: 1.77–4.27; I² 20.8%; RR = 2.68, 95% CI: 1.96–3.65; I² 0%; S Figs. B.28–B.30; respectively).

BRCA1 and *BRCA2* mutation carriers (Fig. 4; S Figs. B.31–B.40).

Seven papers reported on risk factors in both *BRCA1* and *BRCA2* mutation carriers [3,15,33,40,53,59,80], and one in *BRCA1* mutation carriers only [2]. Although the number of papers was limited for *BRCA1* and *BRCA2* mutation carriers, effects of the meta-analyses pointed in the same direction as in the population based cohorts for family history of BC, age at PBC diagnosis and endocrine therapy. RRSO was associated with a decreased CBC risk in *BRCA1* mutation carriers (crude RR = 0.56; 95% CI: 0.32–0.99; I² 46.8%; S Fig. B.36).

3.1. Quality assessment

Results from the boxplot on the distribution of points for potential bias following the QUIPS tool are shown in S Figure B.41. We classified 46 out of 68 papers as being high quality which were subsequently used for the sensitivity analysis (S Table A.3).

Following the sensitivity analysis, heterogeneity became 0% in the meta-analysis concerning ER status and BMI (25–29.9 vs < 25 kg/m²) and decreased for age at PBC diagnosis (I² 58.4%). Further, a significant association between BMI and CBC risk was

observed (BMI 25–29.9 vs < 25 kg/m²: RR = 1.39; 95% CI: 1.14–1.69), but we no longer observed an association between T2 vs. T1/T0 PBC and CBC risk (Fig. 5). Concerning *BRCA1* and *BRCA2* mutation carriers, an insufficient number of papers remained to perform meta-analyses, especially due to evidence for selection bias.

Funnel plots were generated for the factors with multiple papers available (i.e. family history, age at PBC diagnosis, TNM-stage, treatment); we observed no evidence for publication bias (Supplementary Figures B.42–B.48).

4. Discussion

In this systematic review with meta-analyses, we aimed to quantify the association of several patient, pathological, and treatment-related characteristics and their influence on CBC risk. For the general BC population, confirming current clinical practice, we observed that carrying a *BRCA1*, *BRCA2* or *CHEK2* c.1100delC mutation comprises the strongest predictors for CBC risk. Family history of BC was also associated with increased CBC risk. In addition, a moderately increased risk was observed following lobular PBC, ER/PR negative PBC, radiotherapy for PBC (at young age) or having a high BMI at PBC diagnosis. Administration of adjuvant chemotherapy or endocrine therapy was associated with decreased CBC risk, as well as older age at PBC diagnosis, although to a lesser extent. For *BRCA1*, *BRCA2* and *CHEK2* c.1100delC mutation carriers,




Fig. 5. Forest plot of the overall adjusted meta-analyses per patient, pathological or treatment-related characteristic on the risk of developing contralateral breast cancer in population-based cohorts using only high-quality papers following the QUIPS bias scoring tool. Abbreviations: BMI = body mass index per kg/m²; ER = Estrogen hormone receptor; PR = Progesterone hormone receptor; Total_N = number of papers used for the analysis. Age concerns the age (years) at primary breast cancer diagnosis; family history concerns the family history of breast cancer estimate is a relative risk estimate combining hazard ratios, odds ratios and relative risks; I²: test for heterogeneity; p-value for heterogeneity: p < 0.05 considered significant; patient and pathological characteristics are assessed at primary breast cancer diagnosis; treatment-related characteristics concerns primary breast cancer treatment.

all estimates on risk factors went in the same direction. However, the number of papers was insufficient to draw strong conclusions. Most importantly, we confirmed the protective effect of adjuvant chemotherapy and endocrine therapy on CBC risk in population-based studies, as reported in large consortia such as the Early Breast Cancer Trialists' Collaborative Group [104,116]. In addition, the protective effect of adjuvant endocrine therapy was also found in *BRCA1/2* mutation carriers, specifically.

Radiotherapy for primary BC was associated with an increased risk of CBC, especially in patients irradiated at younger age (<40 years). This negative effect of radiotherapy is likely a consequence of scattered radiation dose in the contralateral breast [68]. In addition, in younger patients the cells are at higher risk of damage after radiotherapy due to a higher breast cell proliferation and increased DNA synthesis [117]. The late adverse effects of radiotherapy occur at least 10–12 years after PBC diagnosis, as has been

shown by Land et al. who studied atomic bomb survivors, and by Ronckers et al. who investigated the effects of x-rays for spine deformities [118,119]. Interestingly, we observed an increased risk of CBC already 5 years following radiotherapy for PBC.

We observed an increased CBC risk in patients with large tumors, and ER/PR negative PBC. Although we cannot deny these associations, both features are also associated with worse prognosis of BC, raising the question whether some CBCs were distant PBC metastases. Only recently it became possible to genetically distinguish a true CBC from recurrent disease. In the latter case, we might misclassify a malignant tumor in the contralateral breast as a new entity, while in fact we are dealing with recurrent disease (misclassification of outcome) [120–123]. This can lead to overestimation of CBC risk for these features. Furthermore, some studies did not rule out the ascertainment of CBCs in the presence of distant metastasis [47,56,112] or did not mention this.

Misclassification of outcome may then occur more often, especially when considering tumor features with high recurrence rate. We can thus not rule out that part of the CBCs were in fact recurrences.

We observed an increased association with CBC risk for lobular PBC, which is in line with some older studies [112,124]. In the papers published before 2000 lobular PBC appeared to be associated with a higher risk of CBC. The effect of lobular histology on CBC risk was less observed in the papers published after 2000, an era in which adjuvant systemic therapy was more widely given (S Fig. B.19). The latter phenomenon has also been reported for CBC in general [20]. In our opinion, this reflects the risk reducing effect of adjuvant systemic therapy, and is in line with our earlier mentioned results on the impact of systemic therapy for PBC on CBC risk.

Results from the QUIPS underscored the importance of interpreting the results of studies in *BRCA1/2* mutation carriers with caution because of several potential forms of bias. In particular, survival bias was observed, which was mainly due to the retrospective design with inclusion of only mutation carriers who were still alive at the time of genetic testing [125]. Additionally, selection bias played a role specifically in the papers published on factors associated with the DNA test result, such as RRSO. These studies showed a protective effect from RRSO in the meta-analyses, but were potentially biased and led to an overestimation of the protective effect.

Our study had some limitations. First, we used reported results rather than individual patient data for the meta-analyses. Nonetheless, for most factors we observed acceptable levels of heterogeneity, which make our results reliable. Second, we cannot completely exclude the possibility of some publication bias, although the funnel plots did not provide evidence for the factors where we had enough papers to inspect this. Last, we only included papers with relative risk estimates, excluding 89 papers which reported cumulative incidences or standardized incidence rates only. However, those papers presented univariable estimates (factors were sometimes only stratified for a potential effect modifier), while we preferred multivariable estimates since these results are potentially less biased.

4.1. Implications for future research

Results from our meta-analyses have provided information on multiple CBC risk factors that should be incorporated in a CBC risk prediction model, but have also identified several topics needing further attention. First, although we observed considerable bias according to the QUIPS tool in the studies on *BRCA1*, *BRCA2* or *CHEK2* c.1100delC mutation carriers, the effect of carrying either one of these mutations has the largest impact on CBC risk and remains, therefore, the most important factor in estimating CBC risk. We will need more data on the effects of other risk factors within these groups to provide more personalized CBC risk estimates. This also accounts for familial BC cohorts. Second, concerning treatment, the effects of various adjuvant chemotherapy regimens, various targeted therapies and the long-term effects of radiotherapy (in young patients) should be investigated more extensively. Third, the effects of breast density on CBC risk should be investigated in large and prospective studies to determine the effects of breast density at PBC diagnosis and changes in density over time, also in relation to adjuvant systemic treatment. Fourth, we propose to investigate SNPs and polygenic risk scores within one large international dataset. This will enable researchers to explore interaction between different SNPs (and between SNPs and other factors) and to further personalize CBC risk estimates. In general, large cohorts (i.e. multicenter/international studies) with individual patient data and sufficiently long follow-up of at least 10–15 years are needed to

accurately predict the risk of CBC.

4.2. Clinical implications

CBC risk is a growing concern in patients diagnosed with PBC, not only resulting in a psychological burden, but also determining survival in certain cases [126,127]. Risk-reducing mastectomy may be offered to those at high risk of developing CBC. On the other hand, overtreatment and exposing patients to side-effects of such radical surgery should be avoided as long as survival benefit has not been demonstrated. Especially in low-risk patients, where the number of patients opting for contralateral risk-reducing mastectomy is increasing, but no survival benefit has been observed, more thorough discussion on the individual CBC risk estimation considering various risk factors is important. This also includes discussing potential alternative risk-reducing options [128].

For example, extended endocrine treatment (beyond 5 years of initial/standard therapy) has been recently associated with a reduced risk of CBC as well [25]. In specific subgroups where the benefit from contralateral mastectomy is undecided, (extended) endocrine treatment as an alternative to reduce the risk of CBC may be advised. Nonetheless, the side-effects of (extended) endocrine treatment should also be considered.

For young PBC patients it is important to take into consideration the long-term side-effects of radiotherapy. Although local recurrence rates are decreased by more than 50% after radiotherapy in young PBC patients [129], CBC risk after radiotherapy is quite substantial in this group, and options to further reduce the scattered radiation dose towards the contralateral breast, as is done with more recent techniques, should thus be focused on.

Having a high BMI is one of the few modifiable risk factors that we have identified. Physicians should inform overweight patients about weight loss intervention programs that already have gained some success in BC patients [130,131].

5. Conclusion

Based on this review with meta-analyses, key prognostic factors for CBC risk are mutation status, family history of BC, and treatment for primary BC. Age at primary BC diagnosis, BMI, lobular histology and hormone receptor status of the primary BC have a weaker association and should be considered in combination with key factors to accurately predict CBC risk.

Funding

This work was supported by the Dutch Cancer Society/Alpe d'HuZes [A6C/6253].

Acknowledgements

We would like to thank Elda de Cuba (Netherlands Cancer Institute) for her efforts in designing our literature search strategy.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.breast.2018.11.005>.

References

- [1] Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Canc* 2015 Mar 01;136(5):E359–86.
- [2] van den Broek AJ, van 't Veer LJ, Hoening MJ, Cornelissen S, Broeks A, Rutgers EJ, et al. Impact of age at primary breast cancer on contralateral

- breast cancer risk in BRCA1/2 mutation carriers. *J Clin Oncol* 2016 Feb 10;34(5):409–18.
- [3] Malone KE, Begg CB, Haile RW, Borg A, Concannon P, Tellhed L, et al. Population-based study of the risk of second primary contralateral breast cancer associated with carrying a mutation in BRCA1 or BRCA2. *J Clin Oncol* 2010 May 10;28(14):2404–10.
 - [4] Dutch guideline (hereditary) breast cancer version 2.0. 2012. https://richtlijnendatabase.nl/en/richtlijn/breast_cancer/breast_cancer.html.
 - [5] Institute NC. Surgery to reduce the risk of breast cancer. 2013 August 12. 2013. <https://www.cancer.gov/types/breast/risk-reducing-surgery-fact-sheet#r1>.
 - [6] Heemskerk-Gerritsen BA, Rookus MA, Aalfs CM, Ausems MG, Collee JM, Jansen L, et al. Improved overall survival after contralateral risk-reducing mastectomy in BRCA1/2 mutation carriers with a history of unilateral breast cancer: a prospective analysis. *Int J Canc* 2015 Feb 01;136(3):668–77.
 - [7] Wong SM, Freedman RA, Sagara Y, Aydogan F, Barry WT, Golshan M. Growing use of contralateral prophylactic mastectomy despite no improvement in long-term survival for invasive breast cancer. *Ann Surg* 2017 Mar;265(3):581–9.
 - [8] Murphy JA, Milner TD, O'Donoghue JM. Contralateral risk-reducing mastectomy in sporadic breast cancer. *Lancet Oncol* 2013 Jun;14(7):e262–9.
 - [9] Tracy MS, Rosenberg SM, Dominici L, Partridge AH. Contralateral prophylactic mastectomy in women with breast cancer: trends, predictors, and areas for future research. *Breast Canc Res Treat* 2013 Aug;140(3):447–52.
 - [10] Rosenberg SM, Tracy MS, Meyer ME, Sepucha K, Gelber S, Hirshfield-Bartek J, et al. Perceptions, knowledge, and satisfaction with contralateral prophylactic mastectomy among young women with breast cancer: a cross-sectional survey. *Ann Intern Med* 2013 Sep 17;159(6):373–81.
 - [11] Abbott A, Rueth N, Pappas-Varco S, Kuntz K, Kerr E, Tuttle T. Perceptions of contralateral breast cancer: an overestimation of risk. *Ann Surg Oncol* 2011 Oct;18(11):3129–36.
 - [12] Reiner AS, John EM, Brooks JD, Lynch CF, Bernstein L, Mellemkjaer L, et al. Risk of asynchronous contralateral breast cancer in noncarriers of BRCA1 and BRCA2 mutations with a family history of breast cancer: a report from the Women's Environmental Cancer and Radiation Epidemiology Study. *J Clin Oncol* 2013 Feb 01;31(4):433–9.
 - [13] Graeser MK, Engel C, Rhiem K, Gadzicki D, Bick U, Kast K, et al. Contralateral breast cancer risk in BRCA1 and BRCA2 mutation carriers. *J Clin Oncol* 2009 Dec 10;27(35):5887–92.
 - [14] Witteveen A, Kwast AB, Sonke GS, MJ JJ, Siesling S. Survival after locoregional recurrence or second primary breast cancer: impact of the disease-free interval. *PLoS One* 2015;10(4). e0120832.
 - [15] Phillips KA, Milne RL, Rookus MA, Daly MB, Antoniou AC, Peock S, et al. Tamoxifen and risk of contralateral breast cancer for BRCA1 and BRCA2 mutation carriers. *J Clin Oncol* 2013 Sep 01;31(25):3091–9.
 - [16] Menes TS, Terry MB, Goldgar D, Andrulis IL, Knight JA, John EM, et al. Second primary breast cancer in BRCA1 and BRCA2 mutation carriers: 10-year cumulative incidence in the Breast Cancer Family Registry. *Breast Canc Res Treat* 2015 Jun;151(3):653–60.
 - [17] van den Broek AJ, Schmidt MK, van 't Veer LJ, Tollenaar RA, van Leeuwen FE. Worse breast cancer prognosis of BRCA1/BRCA2 mutation carriers: what's the evidence? A systematic review with meta-analysis. *PLoS One* 2015;10(3). e0120189.
 - [18] Greenland S, Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis. *Am J Epidemiol* 1992 Jun 01;135(11):1301–9.
 - [19] Netherlands Cancer Registry. www.iknl.nl.
 - [20] Rasmussen CB, Kjaer SK, Ejlersen B, Andersson M, Jensen MB, Christensen J, et al. Incidence of metachronous contralateral breast cancer in Denmark 1978–2009. *Int J Epidemiol* 2014 Dec;43(6):1855–64.
 - [21] Gao X, Fisher SG, Emami B. Risk of second primary cancer in the contralateral breast in women treated for early-stage breast cancer: a population-based study. *Int J Radiat Oncol Biol Phys* 2003 Jul 15;56(4):1038–45.
 - [22] Zhang W, Becciolini A, Biggeri A, Pacini P, Muirhead CR. Second malignancies in breast cancer patients following radiotherapy: a study in Florence, Italy. *Breast Cancer Res* 2011 Apr 04;13(2):R38.
 - [23] Borenstein M, Hedges LV, Higgins JP, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods* 2010 Apr;1(2):97–111.
 - [24] Hayden JA, van der Windt DA, Cartwright JL, Cote P, Bombardier C. Assessing bias in studies of prognostic factors. *Ann Intern Med* 2013 Feb 19;158(4):280–6.
 - [25] Goss PE, Ingle JN, Pritchard KI, Robert NJ, Muss H, Gralow J, et al. Extending aromatase-inhibitor adjuvant therapy to 10 years. *N Engl J Med* 2016 Jul 21;375(3):209–19.
 - [26] Aalders KC, van Bommel AC, van Dalen T, Sonke GS, van Diest PJ, Boersma LJ, et al. Contemporary risks of local and regional recurrence and contralateral breast cancer in patients treated for primary breast cancer. *Eur J Canc* 2016 Aug;63:118–26.
 - [27] Sisti JS, Bernstein JL, Lynch CF, Reiner AS, Mellemkjaer L, Brooks JD, et al. Reproductive factors, tumor estrogen receptor status and contralateral breast cancer risk: Results from the WECARE study. *SpringerPlus* 2015;4:825.
 - [28] Kiderlen M, van de Water W, Bastiaannet E, de Craen AJ, Westendorp RG, van de Velde CJ, et al. Survival and relapse free period of 2926 unselected older breast cancer patients: a FOCUS cohort study. *Cancer Epidemiol* 2015 Feb;39(1):42–7.
 - [29] Drooger J, Akdeniz D, Pignol JP, Koppert LB, McCool D, Seynaeve CM, et al. Adjuvant radiotherapy for primary breast cancer in BRCA1 and BRCA2 mutation carriers and risk of contralateral breast cancer with special attention to patients irradiated at younger age. *Breast Canc Res Treat* 2015 Nov;154(1):171–80.
 - [30] Basu NN, Ingham S, Hodson J, Lalloo F, Bulman M, Howell A, et al. Risk of contralateral breast cancer in BRCA1 and BRCA2 mutation carriers: a 30-year semi-prospective analysis. *Fam Cancer* 2015 Dec;14(4):531–8.
 - [31] Mellemkjaer L, Steding-Jessen M, Frederiksen K, Andersson M, Ejlersen B, Jensen MB, et al. Risk of contralateral breast cancer after tamoxifen use among Danish women. *Ann Epidemiol* 2014 Nov;24(11):843–8.
 - [32] Kriege M, Holte A, Jager A, Huijts PE, Berns EM, Sieuwerts AM, et al. Survival and contralateral breast cancer in CHEK2 1100delC breast cancer patients: impact of adjuvant chemotherapy. *Br J Canc* 2014 Aug 26;111(5):1004–13.
 - [33] Gronwald J, Robidoux A, Kim-Sing C, Tung N, Lynch HT, Foulkes WD, et al. Duration of tamoxifen use and the risk of contralateral breast cancer in BRCA1 and BRCA2 mutation carriers. *Breast Canc Res Treat* 2014 Jul;146(2):421–7.
 - [34] Calip GS, Malone KE, Gralow JR, Stergachis A, Hubbard RA, Boudreau DM. Metabolic syndrome and outcomes following early-stage breast cancer. *Breast Canc Res Treat* 2014 Nov;148(2):363–77.
 - [35] van de Water W, Seynaeve C, Bastiaannet E, Markopoulos C, Jones SE, Rea D, et al. Elderly postmenopausal patients with breast cancer are at increased risk for distant recurrence: a tamoxifen exemestane adjuvant multinational study analysis. *Oncol* 2013;18(1):8–13.
 - [36] Valuckas KP, Atkocius V, Kuzmickiene I, Aleknavicius E, Liukpetryte S, Ostapenko V. Second malignancies following conventional or combined (2/5)2Cf neutron brachytherapy with external beam radiotherapy for breast cancer. *J Radiat Res* 2013 Sep;54(5):872–9.
 - [37] Sandberg ME, Li J, Hall P, Hartman M, dos-Santos-Silva I, Humphreys K, et al. Change of mammographic density predicts the risk of contralateral breast cancer—a case-control study. *Breast Cancer Res* 2013;15(4):R57.
 - [38] Pacelli R, Conson M, Cella L, Liuzzi R, Troncone G, Iorio V, et al. Radiation therapy following surgery for localized breast cancer: outcome prediction by classical prognostic factors and approximated genetic subtypes. *J Radiat Res* 2013 Mar 01;54(2):292–8.
 - [39] Metzger-Filho O, Sun Z, Viale G, Price KN, Crivellari D, Snyder RD, et al. Patterns of Recurrence and outcome according to breast cancer subtypes in lymph node-negative disease: Results from international breast cancer study group trials VIII and IX. *J Clin Oncol* 2013 Sep 01;31(25):3083–90.
 - [40] Mavaddat N, Peock S, Frost D, Ellis S, Platte R, Fineberg E, et al. Cancer risks for BRCA1 and BRCA2 mutation carriers: Results from prospective analysis of EMBRACE. *J Natl Cancer Inst* 2013 Jun 05;105(11):812–22.
 - [41] Maskarinec G, Pagano IS, Little MA, Conroy SM, Park SY, Kolonel LN. Mammographic density as a predictor of breast cancer survival: the Multi-ethnic Cohort. *Breast Cancer Res* 2013 Jan 22;15(1):R7.
 - [42] Dellapasqua S, Maisonneuve P, Viale G, Pruneri G, Mazzarol G, Ghislini R, et al. Immunohistochemically defined subtypes and outcome of apocrine breast cancer. *Clin Breast Canc* 2013 Apr;13(2):95–102.
 - [43] Courdi A, Chamorey E, Ferrero JM, Hannoun-Levi JM. Influence of internal mammary node irradiation on long-term outcome and contralateral breast cancer incidence in node-negative breast cancer patients. *Radiother Oncol* 2013 Aug;108(2):259–65.
 - [44] Bernstein JL, Thomas DC, Shore RE, Robson M, Boice Jr JD, Stovall M, et al. Contralateral breast cancer after radiotherapy among BRCA1 and BRCA2 mutation carriers: a WECARE study report. *Eur J Canc* 2013 Sep;49(14):2979–85.
 - [45] Weischer M, Nordestgaard BG, Pharoah P, Bolla MK, Nevanlinna H, Van't Veer LJ, et al. CHEK2*1100delC heterozygosity in women with breast cancer associated with early death, breast cancer-specific death, and increased risk of a second breast cancer. *J Clin Oncol* 2012 Dec 10;30(35):4308–16.
 - [46] Vichapat V, Garmo H, Holmqvist M, Liljegren G, Warnberg F, Lambe M, et al. Tumor stage affects risk and prognosis of contralateral breast cancer: Results from a large Swedish-population-based study. *J Clin Oncol* 2012 Oct 01;30(28):3478–85.
 - [47] Saltzman BS, Malone KE, McDougall JA, Daling JR, Li CI. Estrogen receptor, progesterone receptor, and HER2-neu expression in first primary breast cancers and risk of second primary contralateral breast cancer. *Breast Canc Res Treat* 2012 Oct;135(3):849–55.
 - [48] Neta G, Anderson WF, Gilbert E, Berrington A. Variation in the risk of radiation-related contralateral breast cancer by histology and estrogen receptor expression in SEER. *Breast Canc Res Treat* 2012 Feb;131(3):1021–7.
 - [49] Mavaddat N, Barrowdale D, Andrulis IL, Domchek SM, Eccles D, Nevanlinna H, et al. Pathology of breast and ovarian cancers among BRCA1 and BRCA2 mutation carriers: Results from the Consortium of Investigators of Modifiers of BRCA1/2 (CIMBA). *Cancer Epidemiol Biomark Prev* 2012 Jan;21(1):134–47.
 - [50] Filleron T, Kramar A, Dalenc F, Spielmann M, Fumoleau P, Kerbrat P, et al. Failure event types and prognostic factors after node-positive breast cancer in patients treated by adjuvant chemotherapy: impact on follow-up. *Bull Cancer* 2012 Jun;99(6):E64–74.
 - [51] Brooks JD, John EM, Mellemkjaer L, Reiner AS, Malone KE, Lynch CF, et al.

- Body mass index and risk of second primary breast cancer: the WECARE Study. *Breast Canc Res Treat* 2012 Jan;131(2):571–80.
- [52] Vichapat V, Gillett C, Fentiman IS, Tutt A, Holmberg L, Lichtenborg M. Risk factors for metachronous contralateral breast cancer suggest two aetiological pathways. *Eur J Canc* 2011 Sep;47(13):1919–27.
- [53] Metcalfe K, Gershman S, Lynch HT, Ghadirian P, Tung N, Kim-Sing C, et al. Predictors of contralateral breast cancer in BRCA1 and BRCA2 mutation carriers. *Br J Canc* 2011 Apr 26;104(9):1384–92.
- [54] Majed B, Dozol A, Ribassin-Majed L, Senouci K, Asselain B. Increased risk of contralateral breast cancers among overweight and obese women: a time-dependent association. *Breast Canc Res Treat* 2011 Apr;126(3):729–38.
- [55] Hackshaw A, Roughton M, Forsyth S, Monson K, Reczko K, Sainsbury R, et al. Long-term benefits of 5 years of tamoxifen: 10-year follow-up of a large randomized trial in women at least 50 years of age with early breast cancer. *J Clin Oncol* 2011 May 01;29(13):1657–63.
- [56] Bouchardy C, Benhamou S, Fioretta G, Verkooyen HM, Chappuis PO, Neyroud-Caspar I, et al. Risk of second breast cancer according to estrogen receptor status and family history. *Breast Canc Res Treat* 2011 May;127(1):233–41.
- [57] Rubino C, Arriagada R, Delaloge S, Le MG. Relation of risk of contralateral breast cancer to the interval since the first primary tumour. *Br J Canc* 2010 Jan 05;102(1):213–9.
- [58] Rondeau V, Mathoulin-Pelissier S, Tanneau L, Sasso AJ, Macgrogan G, Debled M. Separate and combined analysis of successive dependent outcomes after breast-conservation surgery: recurrence, metastases, second cancer and death. *BMC Canc* 2010 Dec 31;10:697.
- [59] Reding KW, Bernstein JL, Langholz BM, Bernstein L, Haile RW, Begg CB, et al. Adjuvant systemic therapy for breast cancer in BRCA1/BRCA2 mutation carriers in a population-based study of risk of contralateral breast cancer. *Breast Canc Res Treat* 2010 Sep;123(2):491–8.
- [60] Poynter JN, Langholz B, Largent J, Mellemkjaer L, Bernstein L, Malone KE, et al. Reproductive factors and risk of contralateral breast cancer by BRCA1 and BRCA2 mutation status: Results from the WECARE study. *Cancer Causes Control* 2010 Jun;21(6):839–46.
- [61] Cuzick J, Sestak I, Baum M, Buzdar A, Howell A, Dowsett M, et al. Effect of anastrozole and tamoxifen as adjuvant treatment for early-stage breast cancer: 10-year analysis of the ATAC trial. *Lancet Oncol* 2010 Dec;11(12):1135–41.
- [62] Buist DS, Abraham LA, Barlow WE, Krishnaraj A, Holdridge RC, Sickles EA, et al. Diagnosis of second breast cancer events after initial diagnosis of early stage breast cancer. *Breast Canc Res Treat* 2010 Dec;124(3):863–73.
- [63] Berrington de Gonzalez A, Curtis RE, Gilbert E, Berg CD, Smith SA, Stovall M, et al. Second solid cancers after radiotherapy for breast cancer in SEER cancer registries. *Br J Canc* 2010 Jan 05;102(1):220–6.
- [64] Li CI, Daling JR, Porter PL, Tang MT, Malone KE. Adjuvant hormonal therapy for breast cancer and risk of hormone receptor-specific subtypes of contralateral breast cancer. *Cancer Res* 2009 Sep 01;69(17):6865–70.
- [65] Li CI, Daling JR, Porter PL, Tang MT, Malone KE. Relationship between potentially modifiable lifestyle factors and risk of second primary contralateral breast cancer among women diagnosed with estrogen receptor-positive invasive breast cancer. *J Clin Oncol* 2009 Nov 10;27(32):5312–8.
- [66] Bertelsen L, Mellemkjaer L, Christensen J, Rawal R, Olsen JH. Age-specific incidence of breast cancer in breast cancer survivors and their first-degree relatives. *Epidemiology* 2009 Mar;20(2):175–80.
- [67] Alkner S, Bendahl PO, Ferno M, Nordenskjold B, Ryden L, South S, et al. Tamoxifen reduces the risk of contralateral breast cancer in premenopausal women: Results from a controlled randomised trial. *Eur J Canc* 2009 Sep;45(14):2496–502.
- [68] Stovall M, Smith SA, Langholz BM, Boice Jr JD, Shore RE, Andersson M, et al. Dose to the contralateral breast from radiotherapy and risk of second primary breast cancer in the WECARE study. *Int J Radiat Oncol Biol Phys* 2008 Nov 15;72(4):1021–30.
- [69] Schaapveld M, Visser O, Louwman WJ, Willemse PH, de Vries EG, van der Graaf WT, et al. The impact of adjuvant therapy on contralateral breast cancer risk and the prognostic significance of contralateral breast cancer: a population based study in The Netherlands. *Breast Canc Res Treat* 2008 Jul;110(1):189–97.
- [70] Mellemkjaer L, Dahl C, Olsen JH, Bertelsen L, Guldberg P, Christensen J, et al. Risk for contralateral breast cancer among carriers of the CHEK2*1100delC mutation in the WECARE Study. *Br J Canc* 2008 Feb 26;98(4):728–33.
- [71] Hoening MJ, Aleman BM, Hauptmann M, Baaijens MH, Klijn JG, Noyon R, et al. Roles of radiotherapy and chemotherapy in the development of contralateral breast cancer. *J Clin Oncol* 2008 Dec 01;26(34):5561–8.
- [72] Bertelsen L, Bernstein L, Olsen JH, Mellemkjaer L, Haile RW, Lynch CF, et al. Effect of systemic adjuvant treatment on risk for contralateral breast cancer in the Women's Environment, Cancer and Radiation Epidemiology Study. *J Natl Cancer Inst* 2008 Jan 02;100(1):32–40.
- [73] van der Leest M, Evers L, van der Sangen MJ, Poortmans PM, van de Poll-Franse LV, Vulto AJ, et al. The safety of breast-conserving therapy in patients with breast cancer aged < or = 40 years. *Cancer* 2007 May 15;109(10):1957–64.
- [74] Trentham-Dietz A, Newcomb PA, Nichols HB, Hampton JM. Breast cancer risk factors and second primary malignancies among women with breast cancer. *Breast Canc Res Treat* 2007 Oct;105(2):195–207.
- [75] Schmidt MK, Tollenaar RA, de Kemp SR, Broeks A, Cornelisse CJ, Smit VT, et al. Breast cancer survival and tumor characteristics in premenopausal women carrying the CHEK2*1100delC germline mutation. *J Clin Oncol* 2007 Jan 01;25(1):64–9.
- [76] Rutqvist LE, Johansson H, Stockholm Breast Cancer Study G. Long-term follow-up of the randomized Stockholm trial on adjuvant tamoxifen among postmenopausal patients with early stage breast cancer. *Acta Oncol* 2007;46(2):133–45.
- [77] Largent JA, Capanu M, Bernstein L, Langholz B, Mellemkjaer L, Malone KE, et al. Reproductive history and risk of second primary breast cancer: the WECARE study. *Cancer Epidemiol Biomark Prev* 2007 May;16(5):906–11.
- [78] Kirova YM, Gambotti L, De Rycke Y, Vilcoq JR, Asselain B, Fourquet A. Risk of second malignancies after adjuvant radiotherapy for breast cancer: a large-scale, single-institution review. *Int J Radiat Oncol Biol Phys* 2007 Jun 01;68(2):359–63.
- [79] Hemminki K, Ji J, Forsti A. Risks for familial and contralateral breast cancer interact multiplicatively and cause a high risk. *Cancer Res* 2007 Feb 01;67(3):868–70.
- [80] Broeks A, Braaf LM, Huseinovic A, Nooijen A, Urbanus J, Hogervorst FB, et al. Identification of women with an increased risk of developing radiation-induced breast cancer: a case only study. *Breast Cancer Res* 2007;9(2):R26.
- [81] Brekelmans CT, Tilanus-Linthorst MM, Seynaeve C, vd Ouweland A, Menke-Pluymers MB, Bartels CC, et al. Tumour characteristics, survival and prognostic factors of hereditary breast cancer from BRCA2-, BRCA1- and non-BRCA1/2 families as compared to sporadic breast cancer cases. *Eur J Canc* 2007 Mar;43(5):867–76.
- [82] Tilanus-Linthorst MM, Bartels KC, Alves C, Bakri B, Crepin E, van den Ouweland A, et al. Selection bias influences reported contralateral breast cancer incidence and survival in high risk non-BRCA1/2 patients. *Breast Canc Res Treat* 2006 Jan;95(2):117–23.
- [83] Pierce LJ, Levin AM, Rebbeck TR, Ben-David MA, Friedman E, Solin LJ, et al. Ten-year multi-institutional results of breast-conserving surgery and radiotherapy in BRCA1/2-associated stage I/II breast cancer. *J Clin Oncol* 2006 Jun 01;24(16):2437–43.
- [84] Levi F, Randimbison L, Te VC, La Vecchia C. Cancer risk after radiotherapy for breast cancer. *Br J Canc* 2006 Aug 07;95(3):390–2.
- [85] Gronwald J, Tung N, Foulkes WD, Offit K, Gershoni R, Daly M, et al. Tamoxifen and contralateral breast cancer in BRCA1 and BRCA2 carriers: an update. *Int J Canc* 2006 May 01;118(9):2281–4.
- [86] Dignam JJ, Wieand K, Johnson KA, Raich P, Anderson SJ, Somkin C, et al. Effects of obesity and race on prognosis in lymph node-negative, estrogen receptor-negative breast cancer. *Breast Canc Res Treat* 2006 Jun;97(3):245–54.
- [87] Brekelmans CT, Seynaeve C, Menke-Pluymers M, Bruggenwirth HT, Tilanus-Linthorst MM, Bartels CC, et al. Survival and prognostic factors in BRCA1-associated breast cancer. *Ann Oncol* 2006 Mar;17(3):391–400.
- [88] Nordenskjold B, Rosell J, Rutqvist LE, Malmstrom PO, Bergh J, Bengtsson NO, et al. Coronary heart disease mortality after 5 years of adjuvant tamoxifen therapy: Results from a randomized trial. *J Natl Cancer Inst* 2005 Nov 02;97(21):1609–10.
- [89] Roychoudhuri R, Evans H, Robinson D, Moller H. Radiation-induced malignancies following radiotherapy for breast cancer. *Br J Canc* 2004 Aug 31;91(5):868–72.
- [90] McCaskill-Stevens W, Wilson J, Bryant J, Mamounas E, Garvey L, James J, et al. Contralateral breast cancer and thromboembolic events in African American women treated with tamoxifen. *J Natl Cancer Inst* 2004 Dec 01;96(23):1762–9.
- [91] Coombes RC, Hall E, Gibson LJ, Paridaens R, Jassem J, Delozier T, et al. A randomized trial of exemestane after two to three years of tamoxifen therapy in postmenopausal women with primary breast cancer. *N Engl J Med* 2004 Mar 11;350(11):1081–92.
- [92] Li CI, Malone KE, Porter PL, Daling JR. Epidemiologic and molecular risk factors for contralateral breast cancer among young women. *Br J Canc* 2003 Aug 04;89(3):513–8.
- [93] Dignam JJ, Wieand K, Johnson KA, Fisher B, Xu L, Mamounas EP. Obesity, tamoxifen use, and outcomes in women with estrogen receptor-positive early-stage breast cancer. *J Natl Cancer Inst* 2003 Oct 01;95(19):1467–76.
- [94] Fisher B, Bryant J, Dignam JJ, Wickerham DL, Mamounas EP, Fisher ER, et al. Tamoxifen, radiation therapy, or both for prevention of ipsilateral breast tumor recurrence after lumpectomy in women with invasive breast cancers of one centimeter or less. *J Clin Oncol* 2002 Oct 15;20(20):4141–9.
- [95] Li CI, Malone KE, Weiss NS, Daling JR. Tamoxifen therapy for primary breast cancer and risk of contralateral breast cancer. *J Natl Cancer Inst* 2001 Jul 04;93(13):1008–13.
- [96] Fisher B, Dignam J, Bryant J, Wolmark N. Five versus more than five years of tamoxifen for lymph node-negative breast cancer: updated findings from the National Surgical Adjuvant Breast and Bowel Project B-14 randomized trial. *J Natl Cancer Inst* 2001 May 02;93(9):684–90.
- [97] Vaittinen P, Hemminki K. Risk factors and age-incidence relationships for contralateral breast cancer. *Int J Canc* 2000 Dec 15;88(6):998–1002.
- [98] Narod SA, Brunet JS, Ghadirian P, Robson M, Heimdal K, Neuhausen SL, et al. Tamoxifen and risk of contralateral breast cancer in BRCA1 and BRCA2 mutation carriers: a case-control study. Hereditary Breast Cancer Clinical Study Group. *Lancet* 2000 Dec 02;356(9245):1876–81.
- [99] Matsuyama Y, Tominaga T, Nomura Y, Koyama H, Kimura M, Sano M, et al. Second cancers after adjuvant tamoxifen therapy for breast cancer in Japan.

- Ann Oncol 2000 Dec;11(12):1537–43.
- [100] Robson M, Levin D, Federici M, Satagopan J, Bogolminy F, Heerdt A, et al. Breast conservation therapy for invasive breast cancer in Ashkenazi women with BRCA gene founder mutations. *J Natl Cancer Inst* 1999 Dec 15;91(24):2112–7.
 - [101] Newcomb PA, Solomon C, White E. Tamoxifen and risk of large bowel cancer in women with breast cancer. *Breast Canc Res Treat* 1999 Feb;53(3):271–7.
 - [102] Kollias J, Ellis IO, Elston CW, Blamey RW. Clinical and histological predictors of contralateral breast cancer. *Eur J Surg Oncol* 1999 Dec;25(6):584–9.
 - [103] Broet P, de la Rochefordiere A, Scholl SM, Fourquet A, De Rycke Y, Pouillart P, et al. Analyzing prognostic factors in breast cancer using a multistate model. *Breast Canc Res Treat* 1999 Mar;54(1):83–9.
 - [104] Early Breast Cancer Trialists' Collaborative G. Tamoxifen for early breast cancer: an overview of the randomised trials. *Early Breast Cancer Trialists' Collaborative Group. Lancet* 1998 May 16;351(9114):1451–67.
 - [105] Swedish Breast Cancer Cooperative Group. Randomized trial of two versus five years of adjuvant tamoxifen for postmenopausal early stage breast cancer. *J Natl Cancer Inst* 1996 Nov 06;88(21):1543–9.
 - [106] Cook LS, White E, Schwartz SM, McKnight B, Daling JR, Weiss NS. A population-based study of contralateral breast cancer following a first primary breast cancer (Washington, United States). *Cancer Causes Control* 1996 May;7(3):382–90.
 - [107] Cook LS, Weiss NS, Schwartz SM, White E, McKnight B, Moore DE, et al. Population-based study of tamoxifen therapy and subsequent ovarian, endometrial, and breast cancers. *J Natl Cancer Inst* 1995 Sep 20;87(18):1359–64.
 - [108] Broet P, de la Rochefordiere A, Scholl SM, Fourquet A, Mosseri V, Durand JC, et al. Contralateral breast cancer: annual incidence and risk parameters. *J Clin Oncol* 1995 Jul;13(7):1578–83.
 - [109] Healey EA, Cook EF, Orav EJ, Schnitt SJ, Connolly JL, Harris JR. Contralateral breast cancer: clinical characteristics and impact on prognosis. *J Clin Oncol* 1993 Aug;11(8):1545–52.
 - [110] Storm HH, Andersson M, Boice Jr JD, Blettner M, Stovall M, Mouridsen HT, et al. Adjuvant radiotherapy and risk of contralateral breast cancer. *J Natl Cancer Inst* 1992 Aug 19;84(16):1245–50.
 - [111] Boice Jr JD, Harvey EB, Blettner M, Stovall M, Flannery JT. Cancer in the contralateral breast after radiotherapy for breast cancer. *N Engl J Med* 1992 Mar 19;326(12):781–5.
 - [112] Bernstein JL, Thompson WD, Risch N, Holford TR. Risk factors predicting the incidence of second primary breast cancer among women diagnosed with a first primary breast cancer. *Am J Epidemiol* 1992 Oct 15;136(8):925–36.
 - [113] Bernstein JL, Thompson WD, Risch N, Holford TR. The genetic epidemiology of second primary breast cancer. *Am J Epidemiol* 1992 Oct 15;136(8):937–48.
 - [114] Baum M, Houghton J, Riley D. Results of the cancer research campaign adjuvant trial for perioperative cyclophosphamide and long-term tamoxifen in early breast cancer reported at the tenth year of follow-up. *Cancer research campaign breast cancer trials group. Acta Oncol* 1992;31(2):251–7.
 - [115] Andersson M, Storm HH, Mouridsen HT. Incidence of new primary cancers after adjuvant tamoxifen therapy and radiotherapy for early breast cancer. *J Natl Cancer Inst* 1991 Jul 17;83(14):1013–7.
 - [116] Early Breast Cancer Trialists' Collaborative Group, Clarke M, Coates AS, Darby SC, Davies C, Gelber RD, et al. Adjuvant chemotherapy in oestrogen-receptor-poor breast cancer: patient-level meta-analysis of randomised trials. *Lancet* 2008 Jan 05;371(9606):29–40.
 - [117] Drooger JC, Hoening MJ, Seynaeve CM, Baaijens MH, Obdeijn IM, Sleijfer S, et al. Diagnostic and therapeutic ionizing radiation and the risk of a first and second primary breast cancer, with special attention for BRCA1 and BRCA2 mutation carriers: a critical review of the literature. *Cancer Treat Rev* 2015 Feb;41(2):187–96.
 - [118] Land CE. Studies of cancer and radiation dose among atomic bomb survivors. The example of breast cancer. *J Am Med Assoc* 1995 Aug 02;274(5):402–7.
 - [119] Ronckers CM, Doody MM, Lonstein JE, Stovall M, Land CE. Multiple diagnostic X-rays for spine deformities and risk of breast cancer. *Cancer Epidemiol Biomark Prev* 2008 Mar;17(3):605–13.
 - [120] Alkner S, Tang MH, Brueffer C, Dahlgren M, Chen Y, Olsson E, et al. Contralateral breast cancer can represent a metastatic spread of the first primary tumor: determination of clonal relationship between contralateral breast cancers using next-generation whole genome sequencing. *Breast Cancer Res* 2015 Aug 05;17:102.
 - [121] Begg CB, Ostrovskaya I, Geyer FC, Papanastasiou AD, Ng CK, Sakr R, et al. Contralateral breast cancers: independent cancers or metastases? *Int J Cancer* 2018 Jan 15;142(2):347–56.
 - [122] Song F, Li X, Song F, Zhao Y, Li H, Zheng H, et al. Comparative genomic analysis reveals bilateral breast cancers are genetically independent. *Oncotarget* 2015 Oct 13;6(31):31820–9.
 - [123] Klevebring D, Lindberg J, Rockberg J, Hilliges C, Hall P, Sandberg M, et al. Exome sequencing of contralateral breast cancer identifies metastatic disease. *Breast Canc Res Treat* 2015 Jun;151(2):319–24.
 - [124] Horn PL, Thompson WD. Risk of contralateral breast cancer. Associations with histologic, clinical, and therapeutic factors. *Cancer* 1988 Jul 15;62(2):412–24.
 - [125] Klaren HM, van't Veer LJ, van Leeuwen FE, Rookus MA. Potential for bias in studies on efficacy of prophylactic surgery for BRCA1 and BRCA2 mutation. *J Natl Cancer Inst* 2003 Jul 02;95(13):941–7.
 - [126] Font-Gonzalez A, Liu L, Voogd AC, Schmidt MK, Roukema JA, Coebergh JW, et al. Inferior survival for young patients with contralateral compared to unilateral breast cancer: a nationwide population-based study in The Netherlands. *Breast Canc Res Treat* 2013 Jun;139(3):811–9.
 - [127] Vichapat V, Garmo H, Holmberg L, Fentiman IS, Tutt A, Gillett C, et al. Prognosis of metachronous contralateral breast cancer: importance of stage, age and interval time between the two diagnoses. *Breast Canc Res Treat* 2011 Nov;130(2):609–18.
 - [128] Fayanju OM, Stoll CR, Fowler S, Colditz GA, Margenthaler JA. Contralateral prophylactic mastectomy after unilateral breast cancer: a systematic review and meta-analysis. *Ann Surg* 2014 Dec;260(6):1000–10.
 - [129] Group EBCTC, Darby S, McGale P, Correa C, Taylor C, Arriagada R, et al. Effect of radiotherapy after breast-conserving surgery on 10-year recurrence and 15-year breast cancer death: meta-analysis of individual patient data for 10,801 women in 17 randomised trials. *Lancet* 2011 Nov 12;378(9804):1707–16.
 - [130] Reeves MM, Terranova CO, Eakin EG, Demark-Wahnefried W. Weight loss intervention trials in women with breast cancer: a systematic review. *Obes Rev* 2014 Sep;15(9):749–68.
 - [131] Playdon M, Thomas G, Sanft T, Harrigan M, Ligibel J, Irwin M. Weight loss intervention for breast cancer survivors: a systematic review. *Curr Breast Cancer Rep* 2013 Sep;5(3):222–46.