

GYNECOLOGY

Risk factors for surgical intervention of early medical abortion



Amani Meaidi, MD; Sarah Friedrich, PhD; Thomas Alexander Gerds, MS; Oejvind Lidegaard, DMSc

BACKGROUND: By being noninvasive, medical termination of pregnancy has increased worldwide access to abortion and improved safety of unsafe abortion. However, secondary surgical intervention is the most frequent complication to medical abortion.

OBJECTIVE: We aimed to identify and quantify risk factors for surgical intervention in women undergoing medically induced termination of pregnancy before 9 completed weeks of gestation.

STUDY DESIGN: We conducted a nationwide cohort study, including all pregnancies terminated before 63 gestational days in women aged 15–49 years during the period 2005–2015. Induction regimen was 200 mg mifepristone followed 24–48 hours later by 0.8 mg vaginal misoprostol. All included pregnancies were followed up for 8 weeks from mifepristone administration. Data were retrieved from national health registers. Multiple logistic regression provided adjusted odds ratios of surgical intervention with 95% confidence intervals. The discriminative ability of the risk factors in identifying surgical intervention was assessed by cross-validated area under the receiver operating characteristic curve.

RESULTS: Of 86,437 early medical abortions, 5320 (6.2%) underwent a surgical intervention within 8 weeks after induction. The proportion of surgical interventions increased from 3.5% in the 5th to 6th gestational week to 10.3% in week 9, odds ratio, 3.2 (95% confidence interval, 2.9–3.6). Compared with women aged 15–19 years, the risk of surgical intervention increased with increasing maternal age until the age of 30–34 years, odds ratio, 1.7 (95% confidence interval, 1.5–1.9), where

after the risk decreased to an odds ratio for age group 40–49 of 1.2 (95% confidence interval, 1.0–1.4). Compared with nulliparous women, a history of only vaginal deliveries with spontaneous delivery of placenta implied an odds ratio of 1.1 (95% confidence interval, 1.0–1.2), women with a history of at least 1 cesarean delivery, an odds ratio of 1.5 (95% confidence interval, 1.3–1.6), and women having experienced a manual removal of placenta after a vaginal birth, an odds ratio of 2.0 (95% confidence interval, 1.7–2.4). Previous medically induced abortion decreased the risk of surgical intervention, odds ratio 0.84 (95% confidence interval, 0.78–0.91), whereas previous early (before 56 days of gestation) surgically induced abortion implied a 53% (95% confidence interval, 1.4–1.7) increased risk of surgical intervention. Previous surgical abortion after 55 days of gestation increased the risk by 17% (95% confidence interval, 1.1–1.3). The area under the receiver operating characteristic curve of the model including all quantified risk factors was 63% (95% confidence interval, 62–64%).

CONCLUSION: Gestational age, maternal age, previous deliveries, and history of medically and surgically induced abortions all had a significant influence on the risk of surgical intervention of early medical abortion. However, inclusion of all quantified risk factors still left most interventions unpredictable.

Key words: Cesarean delivery, complication, gestational age, induced abortion, maternal age, medical abortion, retained placenta, surgical abortion, uterine vacuum aspiration, vaginal delivery

Medical termination of pregnancy before 9 gestational weeks (early medical abortion) is a recognized procedure increasingly used worldwide.^{1,2} Being noninvasive, early medical abortion improves safety and minimizes the infrastructural demands for the handling of terminations of pregnancies.²

Secondary surgical intervention is, however, the most frequent, clinically significant complication to early medical abortion and for the approximately 5% who experience this intervention, the otherwise obvious administrative,

economical, and health-related advantages are challenged.

Despite the high and increasing use of early medical abortion worldwide, not much research has been performed on possible risk factors for secondary surgical intervention.² Evidence on both acknowledged risk factors for surgical intervention, such as gestational age at time of the induction, as well as suggested risk factors, such as high maternal age, previous deliveries, and history of induced abortion, is sparse, inconsistent, and often based on outdated medical regimes.^{3–8} To our knowledge, the predictive value of each of these recognized or possible risk factors has not been sufficiently evaluated.

Since 1997, early medical abortion with mifepristone and misoprostol has been available for all women with an unwanted pregnancy in Denmark. Apart

from the first years after the introduction of the procedure, where 600 mg mifepristone was followed by 0.4 mg vaginal misoprostol, the typical regimen has been 200 mg mifepristone followed 24–48 hours later by 0.8 mg vaginally administered misoprostol.^{9,10} All legally induced abortions are registered in the Danish Register of Legally Induced Abortions (“abortion register”).¹¹

Considering the health and socioeconomic advantages of preventing the most frequent complication to one of the most commonly executed procedures within gynecology and the opportunities provided by the Danish registers, we followed a Danish nationwide cohort of early medical abortions with the aim of determining how gestational age, maternal age, previous deliveries, and history of induced abortion influence the risk of surgical intervention and to

Cite this article as: Meaidi A, Friedrich S, Gerds TA, et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019;220:478.e1-15.

0002-9378/\$36.00

© 2019 Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.ajog.2019.02.014>

AJOG at a Glance

Why was this study conducted?

Being noninvasive, early medical termination of unwanted pregnancy has increased worldwide access to induced abortion and improved safety of unsafe abortion. Thus, avoidance of secondary surgical intervention has important logistical and health-related implications worldwide. We therefore aimed to identify and quantify risk factors for surgical intervention of early first-trimester medical abortions.

Key findings

Gestational age, maternal age, previous deliveries, and history of induced abortion were all found to independently influence the risk of surgical intervention of early medically induced abortion.

What does this add to what is known?

This is the first nationwide study following a cohort of non-trial, real-life early medically induced abortions, identifying and quantifying risk factors for surgical intervention. The study both confirmed previous findings as well as assessed and quantified new risk factors.

estimate the predictive performance of a model including these factors.

Materials and Methods**Early medical abortion in Denmark**

Since 2005, early medical abortions have been induced at home, and the typical follow-up strategy to ensure completion of the abortion has been a halving of serum human chorionic gonadotropin 1 week after the mifepristone administration.¹⁰ If the follow-up serum human chorionic gonadotropin has been reduced by less than 50%, women undergo transvaginal ultrasound examination. Surgical intervention has been offered when ultrasound has shown a persistent gestational sac or viable pregnancy. Otherwise, the decision to surgically intervene has been a clinical estimate made by the gynecologist on duty.¹⁰

Study population

We included all medical abortions induced in Denmark at a gestational age of less than 63 days during the period 2005–2015 among women aged 15–49 years, using the specific diagnostic and treatment codes by which medical abortions are registered in the abortion register (Supplemental Table 1).¹¹ The abortions were induced with 200 mg mifepristone followed 24–48 hours later

by 0.8 mg vaginally administered misoprostol.¹⁰

Study design

We followed all included pregnancies for 8 weeks from mifepristone administration. Women receiving a surgical intervention (either uterine vacuum aspiration or a hysteroscopic excision of anticipated retained tissue) to complete the abortion additionally are given specific surgical codes at the time of surgical intervention in The Danish National Patient Register.¹² We defined a medical abortion as being surgically intervened, if one of these surgical codes was given during the 8 weeks of follow-up (Supplemental Table 1).

Information on gestational and maternal age at first medical administration was extracted from the abortion register.¹¹ Data on history of induced abortion were achieved from the same register. We distinguished between previous medically induced abortion, previous surgically induced abortion induced before a gestational age of 56 days, and previous surgically induced abortion induced at a gestational age of ≥ 56 days (Supplemental Table 1).

The Danish Medical Birth Register and The Danish National Patient Register provided data on previous vaginal delivery, previous cesarean delivery, and

previous manual removal of placenta (Supplemental Table 1).^{12,13} A personal identification number given to all Danish citizens at birth or at immigration allowed reliable linkage of data between the different registers.

Statistical analysis

A multiple logistic regression model was used to analyze the association between the odds of surgical intervention and gestational age groups (5th to 6th, 7th, 8th, and 9th week), maternal age groups (15–19, 20–24, 25–29, 30–34, 35–39, 40–49 years), and reproductive history including previous medically induced abortions, previous surgically induced abortions, and previous deliveries. Calendar time was included in the model. Reported were adjusted odds ratios (ORs) with 95% confidence intervals (CIs). To illustrate the effect of maternal age on a continuous age scale, a second multiple logistic regression analysis was performed, where instead of maternal age groups, a restricted cubic spline was used to model the effect of maternal age on the odds of surgical intervention. The number and placement of knots was chosen according to suggestions in Harrell.¹⁴ The other variables in the model were unchanged. The result of the restricted cubic spline analysis was reported graphically as the risk of surgical intervention with corresponding pointwise 95% CIs according to maternal age, stratified by gestational age groups for given values of reproductive history and calendar time. A subgroup analysis was made on first-time medical abortions.

Linear trends of time since last induced abortion and of the number of previous induced abortions were analyzed in subgroups of women with previous surgical abortion and previous medical abortion, respectively, by entering the variables as numeric (one degree of freedom) in a multiple logistic regression model also including gestational age group, maternal age group, previous deliveries, and calendar time. Similarly, the effects of time since last delivery as well as number of deliveries were analyzed in subgroups of women with only previous vaginal deliveries with spontaneous delivery of placenta

TABLE 1
Maternal age and reproductive history according to gestational age in women undergoing an early medical abortion

Gestational age, d	28–41	42–48	49–55	56–62	28–62
Gestational age, wk	5–6th	7th	8th	9th	5–9th
Number of abortions	21,969	32,632	22,050	9786	86,437
Percent distribution	25.4	37.8	25.5	11.3	100
Maternal age, y					
15–19	2594 (11.8)	4058 (12.4)	2827 (12.8)	1344 (13.7)	10,823 (12.5)
20–24	5405 (24.6)	7903 (24.2)	5364 (24.3)	2454 (25.1)	21,126 (24.4)
25–29	4546 (20.7)	6589 (20.2)	4411 (20.0)	1973 (20.2)	17,519 (20.3)
30–34	4091 (18.6)	6140 (18.8)	4218 (19.1)	1787 (18.3)	16,236 (18.8)
35–39	3457 (15.7)	5267 (16.1)	3623 (16.4)	1579 (16.1)	13,926 (16.1)
40–49	1876 (8.5)	2675 (8.2)	1607 (7.3)	649 (6.6)	6807 (7.9)
Previous deliveries					
Nulliparous	10,374 (47.2)	16,210 (49.7)	11,170 (50.7)	5041 (51.5)	42,795 (49.5)
Only vaginal deliveries, spontaneous delivery of placenta	8913 (40.6)	12,810 (39.3)	8566 (38.8)	3728 (38.1)	34,017 (39.4)
≥1 cesarean delivery	2365 (10.8)	3167 (9.7)	2031 (9.2)	893 (9.1)	8456 (9.8)
≥1 manual removal of placenta	317 (1.4)	445 (1.4)	283 (1.3)	124 (1.3)	1169 (1.4)
Previous medical abortions					
No	16,988 (77.3)	26,311 (80.6)	17,799 (80.7)	7723 (78.9)	68,821 (79.6)
≥1	4981 (22.7)	6321 (19.4)	4251 (19.3)	2063 (21.1)	17,616 (20.4)
Previous surgical abortions					
No	17,689 (80.5)	26,359 (80.8)	17,858 (81.0)	7831 (80.0)	69,737 (80.7)
≥1, <56 days of gestation	1049 (4.8)	1311 (4.0)	803 (3.6)	299 (3.1)	3462 (4.0)
≥1, ≥56 days of gestation	2611 (11.9)	4173 (12.8)	2884 (13.1)	1424 (14.6)	11,092 (12.8)
Both <56 and ≥56 days of gestation	620 (2.8)	789 (2.4)	505 (2.3)	232 (2.4)	2146 (2.5)

Column percentages are shown in parentheses.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. Am J Obstet Gynecol 2019.

and women with at least 1 previous cesarean delivery, respectively.

To test the predictive value of gestational age, maternal age, previous deliveries, and history of induced abortion, the cohort of early medical abortions was divided into a training data set, including abortions induced in the years 2005–2012, and a validation data set with abortions induced in 2013–2015. The logistic regression model including gestational age groups, maternal age groups, previous deliveries, previous medical abortions, previous surgical abortions, and calendar time was then fitted on the training data set and tested on the validation data set. The same was done for the logistic regression model

including maternal age modeled as a continuous variable with restricted cubic spline. Reported were the areas under the receiver operating characteristic curves (AUCs) with 95% CIs. As a sensitivity analysis, we also randomly split the data into a training and validation data set, both of the same size as for the calendar year-based split. All analyses were performed in R.¹⁵

Ethics approval

The study was approved by the Danish Data Protection Agency and the Danish Health Data Board.


Ethics approval from the Danish National Committee on Health Research

Ethics was not required due to the study being register-based.

Results

We identified 86,437 medical abortions induced before 63 days of gestation during the 11-year-long study period among women aged 15–49 years. Of these, 5320 (6.2%) received a surgical intervention within 8 weeks from mifepristone administration, the majority being uterine vacuum aspirations and only 57 being hysteroscopic excisions of anticipated retained tissue. Characteristics of the women at time of the medical induction are shown in Table 1.

FIGURE 1
Adjusted ORs of surgical intervention in women undergoing an early medical abortion



The adjusted ORs were derived from a multiple logistic regression model including gestational age groups, maternal age groups, previous deliveries, previous medical abortions, previous surgical abortions, and calendar time.

OR, odds ratio.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.


The proportion of surgical intervention increased from 3.5% in abortions induced at a gestational age of 28–41 days to 10.3% in abortions being 56–62 days at induction, OR, 3.2 (95% CI, 2.9–3.6; $P < .001$; Figure 1).

Compared with women aged 15–19 years, the risk of surgical intervention

increased with increasing maternal age until the age group 30–34 years, OR, 1.7 (1.5–1.9; $P < .001$); hereafter, the risk declined (Figure 1). Figure 2 illustrates the absolute risk of surgical intervention for each maternal age according to the gestational age at time of the induction. Previous deliveries increased the risk of

surgical intervention compared with nulliparous women (Figure 1). The OR of surgical intervention for women with at least 1 previous cesarean delivery compared with women with a history of only vaginal deliveries (with spontaneous delivery of placenta) was 1.3 (1.2–1.5; $P < .001$). Women who had

FIGURE 2
Absolute risks of surgical intervention according to maternal age and stratified by gestational age



The assessment of the risks was based on a multiple logistic regression model including maternal age modeled as a continuous variable via a restricted cubic spline, gestational age groups, previous deliveries, previous medical abortions, previous surgical abortions, and calendar time.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.

experienced at least 1 manual removal of placenta had a doubled risk of surgical intervention compared to nulliparous women (Figure 1).

Although previous experience with medically induced abortion reduced the risk of surgical intervention, OR, 0.84 (0.78–0.91; $P < .001$), a history of surgically induced abortion increased the risk compared with women with no experience in surgical abortion. Women with a previous surgical abortion induced at less than 56 days of gestation had a greater risk of surgical intervention compared with women with a history of surgical abortion induced at or after 56 days of gestation (Figure 1).

A subgroup analysis of only first-time medical abortions showed no significant change in adjusted OR of surgical intervention associated to gestational age groups, maternal age groups, previous deliveries, and history of surgical abortions (Supplemental Figure 1).

A subgroup analysis of women having previously experienced a medically induced abortion showed that increasing number of previous medical abortions

reduced the risk of surgical intervention (Supplemental Figure 2, $P < .001$), whereas increasing time since last medical abortion reduced the protective effect of a previous medical abortion on the risk of surgical intervention (Supplemental Figure 2, $P = .010$). The trend was opposite for women with a previous surgical abortion. In a subgroup analysis of women having previously experienced a surgically induced abortion, increasing number of previous surgical abortions increased the risk of surgical intervention (Supplemental Figure 3, $P < .001$), whereas increasing time since last surgical abortion reduced the negative impact of a previous surgical abortion on the risk of surgical intervention (Supplemental Figure 3, $P = .003$).

A subgroup analysis of women with a history of only vaginal deliveries (with spontaneous delivery of placenta) showed no association between number of previous vaginal deliveries ($P = .24$) or time since last delivery ($P = .71$) and risk of surgical intervention (Supplemental Figure 4).

A similar subgroup analysis of women with at least 1 previous cesarean delivery showed no effect of time since last cesarean delivery ($P = .54$). However, the odds of surgical intervention of an early medical abortion increased with increasing number of previous cesarean deliveries ($P = .053$, Supplemental Figure 5).

Figure 3 shows the receiver operating characteristic curve for the prediction test performed on the calendar year-based division of the cohort. Characteristics of the cohort according to each data set are provided in Supplemental Table 2. The AUC was found to be 0.63 (95% CI, 0.62–0.64). The AUC was similar for the model including maternal age as a continuous variable modeled by a restricted cubic spline, AUC 0.63 (95% CI, 0.62–0.64). The AUC did not change significantly when calculated in sensitivity analyses where the division of the cohort in validation and training data set was done randomly.

Discussion

This nationwide cohort study of 86,437 early medical abortions showed gestational age, maternal age, previous deliveries, and history of induced abortion to influence the risk of surgical intervention of early medical abortions.

Complying with other studies on early medical abortions induced by 200 mg mifepristone followed 24–48 hours later by 0.8 mg vaginal misoprostol, we found a prevalence of surgical intervention of 6.2%.^{16–20}

Of the variables studied, gestational age at time of the medical induction showed to be the most significant risk factor for surgical intervention, tripling the odds for medical abortions induced in the 9th gestational week compared with induction in week 5–6. Few studies have shown an increase in risk of surgical intervention with increasing gestational age.^{16,17,21} A Cochrane review by Kulier et al,⁴ however, could not confirm this association.

The restricted cubic spline modeling of maternal age showed a u-shaped association between maternal age and risk of surgical intervention, the risk peaking at its highest for women in their


mid-thirties. Maternal age previously has been proposed as a risk factor for surgical intervention of early medical abortions.^{5,7} To our knowledge, this is the first study allowing a detailed assessment of the association between maternal age and risk of surgical intervention also in very young women and for women aged 40–49 years, thereby revealing a u-shaped curve.

We found previous deliveries to be a risk factor for surgical intervention. While women with only previous vaginal deliveries with spontaneous delivery of placenta had a slight increase in risk of surgical intervention compared to nulliparous women, previous experience with cesarean delivery increased the risk by around 50%, whereas previous necessity of a manual removal of placenta doubled the risk compared with nulliparous. Studies on early medical abortions induced by other medical regimes have suggested parity and cesarean delivery to be risk factors.^{6,22,23} However, we did not find any study investigating the effect of previous placental retention on the risk of surgical intervention.

Multiple studies have not been able to show an association between history of induced abortion and risk of surgical intervention.^{5–7,22} However, none of these studies distinguished between the different types of induction. When stratifying into previous medically induced abortion, previous surgical abortion induced before or at/after 56 days of gestation, respectively, we found previous medical abortions to reduce the risk of surgical intervention, whereas previous surgical abortions increased the risk, with a history of surgical abortions induced before 56 days of gestation increasing the risk the most by approximately 50%.

To our knowledge, this is the first nationwide study on risk factors for surgical intervention of early medical abortions. The obvious strengths of the study are the size of the included population, the lack of selection bias due to the inclusion of all early medical abortions induced in Denmark, as well as the full follow-up of all included abortions. A main limitation is the absence of information on the indication for each

FIGURE 3
The ROC curve of the prediction performance of the multiple logistic regression model including gestational age groups, maternal age groups, previous deliveries, previous medical abortions, previous surgical abortions, and calendar time



The training data set, on which the multiple regression model was fitted, consisted of the abortions induced during the period 2005–2012, whereas the validation data set, on which the model was tested, consisted of abortions induced during 2013–2015.

ROC, receiver operating characteristic.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.

surgical intervention. This is due to the lack of systematic application of diagnosis codes on reason for surgical intervention in the everyday clinical practice, causing the information on indication to be missing or have inconsistent validity.

The decision to surgically intervene an early medical abortion is rarely based on medical necessity.²⁴ Ongoing pregnancy and health-threatening hemorrhage are not common observations in the course of early medical abortions and, therefore, rarely the indication for surgical intervention.^{21,24} Often, the decision to surgically intervene is based on a clinical estimate that depends on the individual woman's symptoms, complains, and acceptability of the procedure as well as the physician's interpretation of the clinical and ultrasound findings. Thus, not knowing the exact indication for each surgical intervention made in the cohort limits the possibility to fully understand the causalities of the associations found. However, existing evidence as well as the subgroup analyses provided

by this study contribute to the understanding of the nature of the associations. Ashok et al⁸ observed an elimination of the association between gestational age and risk of surgical intervention by offering a second dose of misoprostol to women who did not achieve a complete abortion after the first dose, suggesting that increasing gestational age increase the risk of surgical intervention due to an increased risk of retained tissue.

The finding of a u-shaped association between maternal age and risk of surgical intervention may indicate multicausality. Ashok et al⁸ showed the induction-to-abortion interval, defined as the time from administration of prostaglandins to passage of products of conception, to increase with increasing maternal age. In contrast, Suhonen et al²⁵ found a negative correlation between age and pain evoked by medical abortion. It is known that women's acceptability of the early medical abortion procedure influences the clinical

decision to surgically intervene, with low acceptability increasing the risk of surgical intervention.^{24,26,27} Thus, when possible reasons exist for both increasing and decreasing risk of surgical intervention with increasing maternal age, the u-shaped association could be plausibly explained.

In a study of the association of ultrasonographic parameters of cesarean scar defect and outcome of early termination of pregnancy, Au et al²⁸ found that ultrasonographically visible cesarean scar defect was associated to an increased risk of surgical intervention of early medical abortion. In the current study, we observed the trend of increased risk of surgical intervention with increasing number of previous cesarean deliveries, whereas no trend was observed for time since last cesarean delivery. These findings suggest and support that the association between previous cesarean delivery and risk of surgical intervention may be anatomical, eg, related to scar formation.

Increasing number of previous surgical abortions was found to increase the risk of surgical intervention, a risk, however, that decreased over time. Increasing number of previous medical abortions had a protective effect, which also decreased by time. Women with a history of surgical abortions have experienced a different abortion procedure with less bleeding and pain experience. This may cause an expectation of less bleeding and pain during an early medical abortion, thereby less acceptability. If a woman is familiar with the sometimes-extensive bleeding and pain accompanying a medical abortion due to previous experience, she may have greater acceptability. The impact of such previous experiences could mean less with time. We also found that women with a previous surgical abortion induced before a gestation of 56 days had a greater risk of surgical intervention compared with women with a history of surgical abortions induced at a gestation of 56 days or more. Since abortion providers in Denmark do not recommend surgical abortion for the termination of pregnancies with a gestational age of less than 56 days, most women with a history

of such have gone against medical recommendations, possibly indicating a relatively low acceptability of medical abortions.

Although the study identified risk factors for surgical intervention of early medical abortions, the prediction performance of these risk factors was found to be low. We consider this finding to represent the aforementioned complex, diverse, and multicausal nature of the indication for surgical intervention. Despite the low prediction performance, we believe that the knowledge of the existence of risk factors may contribute to a reduction of surgical interventions of early medical abortions. ■

References

1. World Health Organization, Department of Reproductive Health and Research. Safe abortion: technical and policy guidance for health systems. Geneva: World Health Organization; 2012.
2. Singh S, Remez L, Sedgh G, Kwok L, Onda T. Abortion worldwide 2017: uneven progress and unequal access. New York: Guttmacher Institute; 2018.
3. Kahn JG, Becker BJ, Maclsaal L, et al. The efficacy of medical abortion: a meta-analysis. *Contraception* 2000;61:29–40.
4. Kulier R, Kapp N, Gülmezoglu AM, Hofmeyr GJ, Cheng L, Campana A. Medical methods for first trimester abortion. *Cochrane Database Syst Rev* 2011;11:CD002855.
5. Reeves MF, Monmaney JA, Creinin MD. Predictors of uterine evacuation following early medical abortion with mifepristone and misoprostol. *Contraception* 2016;93:119–25.
6. Chien LW, Liu WM, Tzeng CR, Au HK. Effect of previous live birth and prior route of delivery on the outcome of early medical abortion. *Obstet Gynecol* 2009;113:669–74.
7. Haimov-Kochman R, Arbel R, Sciaky-Tamir Y, Brzezinski A, Laufer N, Yagel S. Risk factors for unsuccessful medical abortion with mifepristone and misoprostol. *Acta Obstet Gynecol Scand* 2007;86:462–6.
8. Ashok PW, Templeton A, Wagaarachchi PT, Flett GM. Factors affecting the outcome of early medical abortion: a review of 4132 consecutive cases. *BJOG* 2002;109:12819.
9. Knudsen UB. First trimester abortion with mifepristone and vaginal misoprostol. *Contraception* 2001;63:247–50.
10. Danish National Clinical Guideline on Medical Abortion, 1. Trimester. Updated 2014. Available at: <https://static1.squarespace.com/static/5467abcce4b056d72594db79/t/548e08cee4b06f8a5b8130d7/1418594510349/14-11-29+Ab+med+guide+14.pdf>. Accessed Oct 8, 2018.

11. Tølbøll Blenstrup L, Knudsen LB. Danish registers on aspects of reproduction. *Scand J Public Health* 2011;39:79–82.

12. Lyngø E, Sandegaard JL, Rebolj M. The Danish National Patient Register. *Scand J Public Health* 2011;39:30–3.

13. Knudsen LB, Olsen J. The Danish Medical Birth Registry. *Dan Med Bull* 1998;45:320–3.

14. Harrell FE. Regression modeling strategies, with applications to linear models, survival analysis and logistic regression. London: Springer; 2001.

15. R Core Team. R: A language and environment for statistical computing. R Foundation for statistical Computing. Vienna, Austria, 2017. Available at: <https://www.R-project.org/>. Accessed January 10, 2019.

16. Von Hertzen H, Piaggio G, Wojdyla D, et al. Two mifepristone doses and two intervals of misoprostol administration for termination of early pregnancy: a randomised factorial controlled equivalence trial. *Br J Obstet Gynaecol* 2009;116:381–9.

17. Tang OS, Chan CC, Ng EH, Lee SW, Ho PC. A prospective, randomized, placebo-controlled trial on the use of mifepristone with sublingual or vaginal misoprostol for medical abortions of less than 9 weeks gestation. *Hum Reprod* 2003;18:2315–8.

18. Shannon C, Wiebe E, Jacot F, et al. Regimens of misoprostol with mifepristone for early medical abortion: a randomised trial. *Br J Obstet Gynaecol* 2006;113:621–8.

19. Chawdhary R, Rana A, Pradhan N. Mifepristone plus vaginal misoprostol vs vaginal misoprostol alone for medical abortion in gestation 63 days or less in Nepalese women: a quasi-randomized controlled trial. *J Obstet Gynaecol Res* 2009;35:78–85.

20. Middleton T, Schaff E, Fielding SL, et al. Randomized trial of mifepristone and buccal or vaginal misoprostol for abortion through 56 days of last menstrual period. *Contraception* 2005;72:328–32.

21. Ashok PW, Penney GC, Flett GM, Templeton A. An effective regimen for early medical abortion: a report of 2000 consecutive cases. *Hum Reprod* 1998;13:2962–5.

22. Bartley J, Tong S, Everington D, Baird DT. Parity is a major determinant of success rate in medical abortion: a retrospective analysis of 3161 consecutive cases of early medical abortion treated with reduced doses of mifepristone and vaginal gemeprost. *Contraception* 2000;62:297–303.

23. Dehendorf CE, Fox EE, Ali RF, et al. Medication abortion failure in women with and without previous cesarean delivery. *Contraception* 2015;92:463–8.

24. Allen RH, Westhoff C, De Nonno L, Fielding SL, Schaff EA. Curettage after mifepristone-induced abortion: Frequency, timing, and indications. *Obstet Gynecol* 2001;98:101–6.

25. Suhonen S, Tikka M, Kivinen S, Kauppila T. Pain during medical abortion: predicting factors from gynecologic history and medical staff

evaluation of severity. *Contraception* 2011;83:357–61.

26. Spitz I, Bardin W, Benton L, Robbins A. Early pregnancy termination with mifepristone in the United States. *N Engl J Med* 1998;338:1241–6.

27. Winikoff B, Sivin I, Coyaji K, et al. Safety, efficacy, and acceptability of medical abortion in China, Cuba, and India: a comparative trial of mifepristone-misoprostol versus surgical abortion. *Am J Obstet Gynecol* 1997;176:431–7.

28. Au H, Liu C, Tzeng C, Chien L. Association between ultrasonographic parameters of

Cesarean scar defect and outcome of early termination of pregnancy. *Ultrasound Obstet Gynecol* 2016;47:506–10.

Author and article information

From the Department of Gynaecology, Rigshospitalet, Faculty of Health and Medical Sciences (Drs Meaidi and Lidegaard) and Department of Public Health, Section of Biostatistics (Mr Gerds), University of Copenhagen, Copenhagen, Denmark; and Institute of Statistics, Ulm University, Ulm, Germany (Dr Friedrich).

Received Oct. 29, 2018; revised Jan. 16, 2019; accepted Feb. 9, 2019.

The authors report no conflict of interest.


The study was supported by Department of Gynecology, Copenhagen University Hospital, Rigshospitalet, Denmark.

Presented at the 13th annual conference of the International Federation of Professional Abortion and Contraception Associates (FIAPAC), Nantes, France, Sept. 14–15, 2018.

Corresponding author: Amani Meaidi, MD. amani-meaidi@live.dk

SUPPLEMENTAL FIGURE 1

Adjusted ORs of surgical intervention in women with no previous experience with medically induced abortion




The ORs were estimated by a multiple logistic regression model including maternal age groups, gestational age groups, previous deliveries, previous surgical abortions, and calendar time.

CI, confidence interval; OR, odds ratio.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. Am J Obstet Gynecol 2019.

SUPPLEMENTAL FIGURE 2

Adjusted ORs of surgical intervention in women having experienced at least 1 previous medical abortion




The ORs were estimated by a multiple logistic regression model including number of previous medical abortions, time since last medical abortion, maternal age groups, gestational age groups, previous deliveries, previous surgical abortions, and calendar time.

CI, confidence interval; OR, odds ratio.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.

SUPPLEMENTAL FIGURE 3

Adjusted ORs of surgical intervention in women having experienced at least one previous surgical abortion




The ORs were estimated by a multiple logistic regression model including number of previous surgical abortions, time since last surgical abortion, maternal age groups, gestational age groups, previous deliveries, previous medical abortions, and calendar time.

CI, confidence interval; OR, odds ratio.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.

SUPPLEMENTAL FIGURE 4

Adjusted ORs of surgical intervention in women having experienced only previous vaginal deliveries (with spontaneous delivery of placenta)




The ORs were estimated by a multiple logistic regression model including number of previous vaginal deliveries (with spontaneous delivery of placenta), time since last previous vaginal delivery (with spontaneous delivery of placenta), maternal age groups, gestational age groups, previous medical abortions, previous surgical abortions, and calendar time.

CI, confidence interval; OR, odds ratio.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.

SUPPLEMENTAL FIGURE 5

Adjusted ORs of surgical intervention in women having experienced at least 1 previous cesarean delivery



The ORs were estimated by a multiple logistic regression model including number of previous cesarean deliveries, time since last cesarean delivery, maternal age groups, gestational age groups, previous medical abortions, previous surgical abortions, and calendar time.

CI, confidence interval; OR, odds ratio.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.

SUPPLEMENTAL TABLE 1

Data sources as well as diagnosis and treatment codes used to identify variables of interest

Variable	Data source	Codes from <i>The International Classification of Diseases and Related Health Problems, 10th Revision</i> , The Nordic Medico-Statistical Committee Classification of Surgical Procedures, and The Danish Classification System for Non-Surgical Procedures
Medically induced abortion, study unit	The Register of Legally Induced Abortions ¹¹	D004/D006 + BKHD40 + BKHD41
Previous medically induced abortion	The Register of Legally Induced Abortions	D004-06 + BKHD40/BKHD44 + BKHD41/BKHD45
Previous surgically induced abortion	The Register of Legally Induced Abortions	D004-06 + KLCH03/KLCH00
Previous delivery	The Danish Medical Birth Register ¹³	Each observation in the data source consists of 1 delivery
Cesarean delivery	The Danish National Patient Register ¹²	KMCA00-96
Manual removal of placenta	The Danish National Patient Register	KMBA30
Surgical intervention subsequent to a medically induced abortion	The Danish National Patient Register	KMBA 00, KMBA03, KLCH00, KLCH03, KLCH13, KLCB98, KLCB25, KULC02

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.

SUPPLEMENTAL TABLE 2

Gestational age, maternal age, and reproductive history according to training and validation data set for the primary prediction performance test

Variable	Training set n (%)	Validation set n (%)	Total n (%)
Abortions	57,259 (66.2)	29,178 (33.8)	86,437 (100)
Gestational age, wk			
5th-6th	12,835 (22.4)	9134 (31.3)	21,969 (25.4)
7th	21,818 (38.1)	10,814 (37.1)	32,632 (37.8)
8th	15,591 (27.2)	6459 (22.1)	22,050 (25.5)
9th	7015 (12.3)	2771 (9.5)	9786 (11.3)
Maternal age, y			
15-19	7329 (12.8)	3494 (12.0)	10,823 (12.5)
20-24	13,315 (23.3)	7811 (26.8)	21,126 (24.4)
25-29	11,000 (19.2)	6519 (22.3)	17,519 (20.3)
30-34	11,205 (19.6)	5031 (17.2)	16,236 (18.8)
35-39	9807 (17.1)	4119 (14.1)	13,926 (16.1)
40-49	4603 (8.0)	2204 (7.6)	6807 (7.9)
Previous deliveries			
Nulliparous	28,272 (49.4)	14,523 (49.8)	42,795 (49.5)
Only vaginal deliveries, spontaneous delivery of placenta	22,727 (39.7)	11,290 (38.7)	34,017 (39.4)
Cesarean delivery (≥ 1)	5565 (9.7)	2891 (9.9)	8456 (9.8)
Manual removal of placenta (≥ 1)	695 (1.2)	474 (1.6)	1169 (1.4)
Previous medical abortions			
No	47,110 (82.3)	21,711 (74.4)	68,821 (79.6)
≥ 1	10,149 (17.7)	7467 (25.6)	17,616 (20.4)
Previous surgical abortions			
No	46,216 (80.7)	23,521 (80.6)	69,737 (80.7)
≥ 1 , <56 days of gestation	2359 (4.1)	1103 (3.8)	3462 (4.0)
≥ 1 , ≥ 56 days of gestation	7319 (12.8)	3773 (12.9)	11,092 (12.8)
Both <56 and ≥ 56 days of gestation	1365 (2.4)	781 (2.7)	2146 (2.5)

The training and validation data set are based on a calendar-time division of the original cohort.

Meaidi et al. Risk factors for surgical intervention of early medical abortion. *Am J Obstet Gynecol* 2019.