



## Original Article

## Sleep, energy balance, and meal timing in school-aged children

Andrea M. Spaeth<sup>a,\*</sup>, Nicola L. Hawley<sup>b</sup>, Hollie A. Raynor<sup>c</sup>, Elissa Jelalian<sup>d</sup>, Ashley Greer<sup>e</sup>, Scott E. Crouter<sup>f</sup>, Donna L. Coffman<sup>g</sup>, Mary A. Carskadon<sup>d</sup>, Judith A. Owens<sup>h</sup>, Rena R. Wing<sup>d</sup>, Chantelle N. Hart<sup>e,i</sup>

<sup>a</sup> Department of Kinesiology and Health, Rutgers University, New Brunswick, NJ, USA

<sup>b</sup> Department of Chronic Disease Epidemiology, Yale School of Public Health, New Haven, CT, USA

<sup>c</sup> Department of Nutrition, College of Education, Health and Human Sciences, The University of Tennessee-Knoxville, Knoxville, TN, USA

<sup>d</sup> Department of Psychiatry and Human Behavior, School of Medicine, Brown University, Providence, RI, USA

<sup>e</sup> Center for Obesity Research and Education, College of Public Health, Temple University, Philadelphia, PA, USA

<sup>f</sup> Department of Kinesiology, Recreation and Sport Studies, The University of Tennessee-Knoxville, Knoxville, TN, USA

<sup>g</sup> Department of Epidemiology and Biostatistics, College of Public Health, Temple University, Philadelphia, PA, USA

<sup>h</sup> Department of Neurology, Boston Children's Hospital, Boston, MA, USA

<sup>i</sup> Department of Social and Behavioral Sciences, College of Public Health, Temple University, Philadelphia, PA, USA



## ARTICLE INFO

## Article history:

Received 15 July 2018

Received in revised form

31 January 2019

Accepted 1 February 2019

Available online 16 February 2019

## Keywords:

Pediatric

Obesity

Risk factors

Sleep timing

Sleep duration

## ABSTRACT

**Objective:** To determine associations among objectively-measured nocturnal sleep time, bedtime and obesogenic behaviors, including dietary intake, timing of intake, and physical activity, in a diverse sample of school-aged children who presented for behavioral treatment to enhance sleep duration.

**Methods:** Eighty-seven children (8–11 y, 66.7% female, zBMI:  $0.86 \pm 1.0$ ) who self-reported sleeping <9.5 h/night were studied for one week using wrist actigraphy to estimate sleep; hip-worn accelerometers to measure physical activity; and 24 h dietary recalls to capture dietary intake and meal timing. Pearson and Spearman's rho correlations and linear regressions controlling for age, gender and race were used for statistical analyses.

**Results and conclusion:** Mean bedtime was 10:31 PM ( $\pm 58.2$  min) and mean nocturnal sleep time was 7.7 h ( $\pm 37.5$  min). Although later bedtime was associated with shorter sleep time ( $r = -0.61, p < 0.001$ ), the two variables were differentially related to obesity risk factors. Later bedtime, but not sleep time, correlated with greater daily fat intake, later first meal of the day, and greater after-dinner intake (all  $p < 0.05$ ). Nocturnal sleep time, but not bedtime, correlated with zBMI ( $p = 0.04$ ). Both sleep time and later bedtime were associated with a later last meal of the day (all  $p < 0.05$ ). Findings remained consistent after controlling for demographic factors. In short-sleeping school-aged children, bedtime may be more predictive of dietary obesity risk factors whereas sleep duration may be more predictive of zBMI. Results suggest that health providers should consider both bedtime and sleep duration for reducing obesity risk in children.

**Clinical trial:** Enhancing Sleep Duration: Effects on Children's Eating and Activity Behaviors, NCT03186508, <https://clinicaltrials.gov/ct2/show/NCT03186508>.

© 2019 Published by Elsevier B.V.

Parental reports show that 31% of 6–11 year olds sleep less than the recommended 9–11 h/night [1,2]. The fact that one-third of school-aged children in the United States may be habitually sleep deprived is problematic given the association between insufficient sleep and various adverse outcomes related to cognitive performance, mood and health [3,4]. Increased risk for obesity has been

consistently associated with habitual short sleep in children [5–9]; a recent meta-analysis examining prospective associations between sleep duration and weight status in children and adolescents found that those who sleep for a shorter duration had twice the risk for being overweight/obese compared with those who sleep longer [10]. Given that children who are short sleepers are at increased risk for weight gain, it is important to examine relationships between sleep and obesogenic behaviors in this group as it provides important information on children who may present for intervention.

\* Corresponding author. 72 Lipman Dr, New Brunswick NJ, 08901, USA.

E-mail address: [ams853@kines.rutgers.edu](mailto:ams853@kines.rutgers.edu) (A.M. Spaeth).

Recent observational studies have also focused on how later bedtime, independent from sleep duration, relates to weight gain and increased risk for obesity [11]. Short sleep duration and later bedtime have been associated with dietary (eg, increased daily caloric intake, lower fruit/vegetable consumption, greater soda and added sugar intake) and physical activity (eg, more sedentary/light activity) risk factors for obesity in children [12–20]. A common limitation of these studies, however, is the reliance on self/parent-report to assess sleep, which may not accurately reflect sleep duration and/or bedtime.

Growing evidence indicates that, in addition to the amount of calories consumed, the timing of caloric intake plays an important role in weight regulation [21]. Although a limited number of studies have examined meal timing and weight regulation in children, some evidence shows decreased morning caloric intake (eg, breakfast skipping) and increased evening caloric intake are associated with overweight/obesity [22–28]. Other evidence suggests that sleep may influence meal timing. For example, Thivel and colleagues found that children who consumed breakfast everyday had an earlier bedtime than those who did not, suggesting a relationship between bedtime and the timing of the first meal of the day [17]. In addition, a recent counterbalanced crossover experimental study found that compared to a week of sleep extension (sleep was prolonged by 1.5 h/night), children reported increased caloric intake during a week of sleep restriction (sleep was decreased by 1.5 h/night) which was primarily due to extra calories consumed during the three additional evening hours that children were awake in the restriction condition [29]. Finally, an observational study among adults showed that later sleep timing was associated with later meal timing (time of the last meal/snack of the day) and greater caloric intake after 8:00 PM [30]. Therefore, sleep may impact obesity risk by affecting the timing of food intake. A study examining the relationship between objectively-measured sleep, focusing on both duration and timing, and obesity risk factors, including meal timing, in school-aged children is therefore needed.

To build upon previous studies examining sleep and obesity risk in children, the current study examined relationships among objectively-measured sleep duration and bed time and obesity risk factors, including meal timing, in a racially diverse sample of school-aged children (8–11 years) who presented for behavioral treatment to enhance sleep. This enabled focusing on a developmental period when children are at risk for both short sleep and obesity, yet is prior to adolescence (when circadian shifts may influence sleep patterns). To our knowledge, few of the previous studies have focused on examining the relationship between sleep behaviors and obesity risk factors in short-sleeping children. Given evidence that short-sleeping children are at increased risk for overweight/obesity, it is particularly important to understand how sleep affects eating and activity behaviors in this group. Furthermore, gaining a better understanding of associations between sleep, eating and physical activity behaviors in short-sleeping children can inform refinement of current intervention approaches aimed at improving sleep and health in this age group. We hypothesized that shorter sleep duration and later bedtime would be associated with higher zBMI (body mass index for age z-score), increased daily caloric intake, less time spent in moderate-to-vigorous physical activity, later meal timing, and greater evening caloric intake.

## 1. Materials and methods

### 1.1. Participants

Children 8–11 years old who reported sleeping  $\leq 9.5$  h per night most days of the week were eligible to participate in a randomized

controlled trial to determine the relative efficacy of a brief behavioral intervention to enhance sleep [31]. Data presented in these analyses are from the baseline assessment which occurred prior to randomization. Parents had to be at least 18 years old, the child's primary caretaker, and able to understand and complete the study questionnaires in English. Children were ineligible if parents reported diagnosis of a medical or psychiatric condition (including a sleep disorder) or current medication that could affect sleep, eating behaviors, or weight status. Eighty-seven children [ $9.6 \pm 1.0$  y,  $n = 58$  (66.7%) female] provided complete sleep data at the baseline assessment. The sample was diverse [Race:  $n = 40$  Black (46.0%),  $n = 32$  White (36.8%),  $n = 14$  Other (16.3%),  $n = 1$  Not reported; Ethnicity:  $n = 75$  Not Hispanic (86.2%),  $n = 12$  Hispanic (13.8%)] and included children with normal weight [ $n = 47$  (54.0%)] and overweight/obesity [ $n = 39$  (44.8%),  $n = 1$  Not measured].

### 1.2. Procedure

Families were recruited through direct mailings and posted advertisements. Trained staff screened families by phone and children who appeared eligible were scheduled for an orientation visit that included additional eligibility assessments. All procedures were approved by the Institutional Review Boards at the Miriam Hospital and Temple University. During the orientation visit, study aims and procedures were described, written consent and assent were obtained, and initial eligibility was determined. If eligible, families were provided with an Actiwatch 2 (Mini Mitter, A Respiromics Company, Bend, OR) and an ActiGraph GT3X (ActiGraph, Pensacola, FL) for the child to wear and were instructed when, where, and how to wear the devices. Families were also provided with a 7-day sleep diary and instructed how to complete the diary as well as call-in to the study voicemail box to report sleep and wake times twice daily for one week. Families were contacted on three random days (two weekdays, one weekend day) during this week by phone to complete a 24-h dietary recall. All families were instructed to continue with their child's typical sleep schedule for this week. At the end of the week, families returned to the Center and research staff measured the child's height and weight. Actiwatch 2, sleep diary and call-in data were reviewed with the family to confirm sleep-wake periods [32] and ActiGraph GT3X data were reviewed for wear time.

### 1.3. Measures

#### 1.3.1. Sleep

Actigraphy is a widely used, reliable and valid device for measuring sleep, which monitors motor activity to obtain continuous recordings of sleep-wake states [32,33]. Actiwatch 2, a watch-size monitor worn on the wrist, was used with standard procedures (children wore the device on their non-dominant wrist across the 24-h period, medium sensitivity threshold) [32,34,35]. Actiwatch 2s were configured to store data in 1-min epochs and 7 days of data were gathered during each assessment. Epochs were scored as sleep or wake by Actiware 5.0 software using an algorithm that interprets activity counts during a single epoch by activity produced in the surrounding 2-min time period [36]. In comparison to polysomnography, the sleep-wake algorithm is excellent at detecting sleep (sensitivity = 0.93) [34]. Sleep diaries were completed by the child and parent; variables included bedtime, night wakings, wake up time, daytime naps, caffeine/medication use, and illness (eg, colds). Diary information was used to verify and score Actiwatch 2 data [37]. The primary variables of interest for the current study were weekly average bedtime (actigraph-estimated sleep onset) and weekly average total sleep time (minutes of Actiwatch 2-scored sleep during the Actiwatch 2-estimated sleep period).

### 1.3.2. Physical activity

Physical activity was measured using an ActiGraph GT3x, a valid objective assessment of physical activity in children [38–40]. Families were instructed that the device should be worn on the child's right hip during all waking hours (except during activities where the device would become wet). The ActiGraph GT3x has a dynamic measurement range of  $\pm 6$  g's and the acceleration is digitized using a 12-bit analog to digital converter. Data are sampled at a user selected rate between 30 and 100 Hz (30 Hz was selected for the current study) that are low-pass filtered at a proprietary cutoff and stored and downloaded as raw acceleration. Time spent in moderate to vigorous physical activity (MVPA) was calculated using validated child-specific equations [38,41]. For inclusion in analyses, children had to have worn the accelerometer a minimum of four days for a minimum of eight waking hours/day ( $n = 86$  (99%) children met this criteria). The variable of interest for the current study was weekly average time spent in MVPA (minutes); weekly average wear time (minutes) was entered as a covariate to account for individual differences in wear time.

### 1.3.3. Dietary intake

Three 24 h dietary recalls (two weekdays, one weekend day) during the 1-week baseline assessment period using multiple pass methodology were used to measure dietary intake. Staff at the Cincinnati Center for Nutritional Research and Analysis (CCNRA) completed the recalls over the phone with the child and parent together (to enable capture of food intake when the child was not in the presence of his/her parent) using a laptop computer with the Nutrition Data System Software (NDS) developed by the Nutrition Coordinating Center at the University of Minnesota [42–44]. Variables of interest included 3-day average daily caloric intake and 3-day average daily percentages of calories from protein, carbohydrate and fat.  $N = 84$  (97%) children completed all three dietary recalls, the remaining three participants only completed two dietary recalls so data for those children reflects a 2-day average.

Meal information for each eating episode, including the time and label (provided by participants, ie, breakfast, lunch, snack, dinner, other), was also collected during each dietary recall to provide caloric and macronutrient content for each meal/snack. Meal timing was operationalized as the 3-day average time of the first eating episode of the day and the 3-day average time of the last eating episode of the day, based on previous findings linking both morning [23,25–27] and evening [22,24,28] intake with health outcomes in children. An eating episode was defined as any amount of calories consumed via food or beverage. We also examined 3-day average caloric and macronutrient (kcal from protein, carbohydrate, fat) intake after the “dinner” eating episode, as previous studies in adults have shown that sleep patterns may be particularly important for caloric intake and fat intake during evening/late-night hours [45–48]. Macronutrient intake after dinner was measured in terms of calories consumed (rather than as the percentage of total after-dinner intake) because a number of participants ( $n = 24$ ) did not consume any calories after dinner, thus preventing the percent calculation.

### 1.3.4. Anthropometrics

Weight and height were measured in duplicate by trained research staff. Child weight was obtained on a digital scale and height was measured using a wall-mounted stadiometer. All measurements were obtained with children dressed in light clothes and without shoes. zBMI (BMI-for-age z-score) was calculated using CDC age and gender-specific population norms [49].

### 1.3.5. Statistical analysis

Both sleep variables and zBMI were normally distributed whereas physical activity, daily and after-dinner intake, and meal timing variables were not. Therefore, Pearson and Spearman's rho (for non-normally distributed variables) correlations were used to assess associations between sleep, energy balance, and meal timing. Preliminary analyses revealed differences in sleep, energy balance, and meal timing variables between demographic groups (age, gender, and race). Therefore, general linear regression models, controlling for age, gender and race, were also run to control for potential confounders. Statistical analyses were performed with IBM SPSS Statistics for Windows (version 20.0; IBM).

## 2. Results

Later bedtime was associated with shorter total sleep time ( $r = -0.61$ ,  $p < 0.001$ , see Table 1 for baseline sleep information). However, the two variables were differentially associated with energy balance measures. Later bedtime, but not total sleep time, correlated with less daily carbohydrate (% kcal;  $r_s = -0.25$ ,  $p = 0.022$ ) and greater daily fat (% kcal;  $r_s = 0.30$ ,  $p = 0.005$ ) intake. Findings remained consistent after controlling for age, gender and race (bedtime and carbohydrate intake:  $B = -0.02$ ,  $p = 0.05$ ; bedtime and fat intake:  $B = 0.02$ ,  $p = 0.019$ ). Shorter total sleep time, but not bedtime, correlated with higher zBMI ( $r = -0.22$ ,  $p = 0.04$ ). Again, this finding remained after controlling for demographic variables ( $B = -0.008$ ,  $p = 0.017$ ). Neither sleep variable was associated with time spent in moderate-to-vigorous physical activity, daily caloric intake or protein intake.

When examining relationships between sleep and meal timing, we observed that shorter total sleep time ( $r_s = -0.25$ ,  $p = 0.02$ ) and later bedtime ( $r_s = 0.48$ ,  $p < 0.001$ ), correlated with a later last meal of the day; findings remained consistent after controlling for age, gender and race (total sleep time and last meal:  $B = -0.46$ ,  $p = 0.023$ ; bedtime and last meal:  $B = 0.57$ ,  $p < 0.001$ ). Later bedtime, but not total sleep time, correlated with a later first meal of the day ( $r_s = 0.61$ ,  $p < 0.001$ ). Later bedtime was also marginally associated with more calories consumed after the dinner meal ( $r_s = 0.20$ ,  $p = 0.063$ ); however, this did not reach statistical significance and specifically with more after-dinner calories from protein ( $r_s = 0.21$ ,  $p = 0.048$ ) and fat ( $r_s = 0.22$ ,  $p = 0.041$ ). Again,

**Table 1**  
Sleep, energy balance and meal timing in school-aged children ( $n = 87$ ).

	Mean $\pm$ SD	Range
Sleep Period (min)	520.8 $\pm$ 39.9	402.6–590.4
Total Sleep Time (min)	463.8 $\pm$ 37.5	364.7–532.7
Bedtime	10:31 PM $\pm$ 58.2 min	8:29 PM – 12:58 AM
Wake Time	7:12 AM $\pm$ 41.5 min	5:44 AM – 9:45 AM
Wake After Sleep Onset (min)	57.0 $\pm$ 18.9	25.2–133.8
Sleep Efficiency (Total Sleep Time/Sleep Period)	89.1 $\pm$ 3.4%	76.5–94.8%
zBMI	0.86 $\pm$ 1.0	–1.55–2.44
Daily caloric intake (kcal)	1807.9 $\pm$ 400.6	1057.8–2914.2
Daily protein intake (% kcal)	14.5 $\pm$ 3.0	6.9–24.7
Daily carbohydrate intake (% kcal)	54.4 $\pm$ 5.1	42.6–67.2
Daily fat intake (% kcal)	31.1 $\pm$ 4.5	20.4–44.0
% wake time in MVPA	11.2 $\pm$ 5.2	4.1–26.9
Time of first meal	8:29 AM $\pm$ 67.0 min	6:25 AM–12:45 PM
Time of last meal	7:56 PM $\pm$ 64.5 min	6:05 PM–10:27 PM
Intake after 'Dinner' (kcal)	143.4 $\pm$ 166.9	0.0–700.6
Protein intake after 'Dinner' (kcal)	12.3 $\pm$ 20.6	0.0–99.9
Carbohydrate intake after 'Dinner' (kcal)	78.0 $\pm$ 83.1	0.0–332.4
Fat intake after 'Dinner' (kcal)	56.5 $\pm$ 77.7	0.0–345.1

these findings remained after controlling for age, gender and race (bedtime and first meal:  $B = 0.76, p < 0.001$ ; bedtime and caloric intake after dinner:  $B = 0.81, p = 0.014$ ; bedtime and protein intake after dinner:  $B = 0.08, p = 0.041$ ; bedtime and fat intake after dinner:  $B = 0.37, p = 0.015$ ). Neither sleep variable was associated with carbohydrate intake after the dinner meal.

### 3. Discussion

Our findings demonstrate that, although moderately correlated with each other, bedtime and total sleep time are differentially associated with obesity risk factors in school-aged children who presented for behavioral treatment to enhance sleep duration. Specifically, later bedtime was associated with greater fat intake, later meal timing, and increased caloric intake after the dinner meal whereas shorter total sleep time was associated with higher zBMI and a later last meal of the day. These findings build upon previous research by using objective measures of sleep, physical activity and zBMI, focusing on a diverse sample of children at increased risk for weight gain, and by demonstrating that beyond sleep duration, bedtime may be an important correlate of eating behaviors, including meal timing.

Observed associations between bedtime and both daily fat intake (% daily kcal) and evening fat intake (kcal from fat consumed after the dinner meal) are consistent with previous experimental studies in adults [45–47,50]. Healthy adults undergoing sleep restriction via a delayed bedtime in a laboratory setting exhibited an increase in daily fat intake [46,47] and consumed a greater percentage of calories from fat during late-night hours compared to morning/afternoon and evening hours [45]. Fat intake during evening hours may be particularly problematic for weight management; Baron and colleagues found that the percentage of fat consumed after 22:00 was associated with greater total caloric intake and a higher BMI in adults [51]. Future studies are needed to examine how meal timing and macronutrient intake interact to influence obesity risk and what mechanisms underlie the relationship between sleep timing and fat intake in school-aged children. Alterations in brain areas associated with responses to food stimuli, such as those within the salience or reward networks, may represent one mechanism as these areas have been shown to be disrupted in sleep-restricted adults [52–54].

Bedtime was also associated with delayed meal timing (operationalized as the time of the first and last meal of the day) and was associated with greater evening intake (kcal consumed after the dinner meal). Breakfast skipping [23,25,27], which implies a delay in the timing of the first meal of the day, and evening intake [22,24,28] have been associated with weight gain, increased adiposity, a larger waist circumference and higher BMI in pediatric populations. For example, children who reported a habit of skipping breakfast exhibited a greater increase in BMI during a 2-year longitudinal study, and those who also skipped lunch (suggesting an even greater delay in the time of the first meal of the day) experienced a larger increase in BMI during that time [27]. In a large sample of 7–11 year olds, the proportion of daily intake consumed at dinner or as a night snack was significantly associated with increased adiposity (% fat mass) [28].

There is growing evidence from experimental studies in adults that sleep patterns influence meal timing [28]. Healthy adults undergoing sleep restriction via a delayed bedtime in a laboratory setting exhibit an increase in evening/late-night intake and a subsequent decrease in morning intake [45,48] suggesting that later sleep timing leads to a shift in the timing of caloric intake. Furthermore, in one sleep restriction protocol, bedtime was delayed until 04:00 AM for five consecutive nights (time-in-bed: 04:00 AM–08:00 AM) and then participants were provided with

two consecutive nights of recovery sleep (time-in-bed: 10:00 PM–10:00 AM) [45]. Interestingly, daily caloric intake was only increased during days when bedtime was delayed – it was not increased during the fifth day of sleep restriction, when participants transitioned to the recovery sleep schedule and bedtime was 10:00 PM [45]. Furthermore, school-aged children exhibited increased daily energy intake (134 kcal/day) during one week of sleep restriction (sleep was decreased by 1.5 h/night) compared to one week of sleep extension (sleep was prolonged by 1.5 h/night) and the difference in intake seemed to be primarily due to the consumption of additional calories (103 kcal/day) during the three additional evening hours that children were awake in the restriction condition [29]. This pattern suggests that bedtime, or total hours of wakefulness, may be better predictors of daily caloric intake compared to the amount of sleep obtained the previous night. In the current study, bedtime, but not total sleep time, was associated with caloric intake after the dinner meal in school-aged children. Collectively findings suggest that interventions promoting a consistent early bedtime may reduce obesity risk by promoting earlier meal timing and reduced evening intake in children.

Numerous cross-sectional and prospective association studies have demonstrated a relationship between self-reported short sleep duration and higher BMI in pediatric populations [5–10]. Furthermore, a recent counterbalanced crossover experimental study found that school-aged children exhibited increased daily energy intake and greater weight gain during a week of sleep restriction (sleep was decreased by 1.5 h/night) compared to a week of sleep extension (sleep was prolonged by 1.5 h/night) [29]. Consistent with these findings, in the current study, we observed a negative correlation between total sleep time, assessed using an Actiwatch, and zBMI. In our sample, sleeping 30 additional minutes per night was associated with a 0.24 lower zBMI. These results suggest that modest differences in sleep duration are associated with meaningful differences in zBMI [55]. Research is needed to examine potential mechanisms that underlie the relationship between sleep duration and weight status and the efficacy of sleep interventions for promoting weight management in school-aged children. It is possible that delayed sleep onset and total sleep time impact obesity risk through different mechanisms; although we did not observe a relationship between our measures of caloric intake or moderate-to-vigorous physical activity with total sleep time; other factors, such as the regulation of lipolysis, gastric or intestinal satiation signals (eg, gastric distention, cholecystokinin, glucagon-like peptide-1), or long-term adiposity signals (eg, leptin and insulin) may be affected by sleep duration and not bedtime [56]. There is limited evidence linking sleep duration and the regulation of metabolic hormones in adults [57]; however, whether this is also true for school-aged children remains under-studied.

Findings should be considered within the context of strengths and limitations. Strengths of the study include the diversity of the sample in terms of race/ethnicity and weight status as well as its focus on children with short sleep who presented for behavioral intervention. Thus, findings generalize to a diverse, treatment-seeking population and can inform potential intervention targets for improving sleep and promoting weight management. Despite these strengths, it is important to acknowledge that these results may not generalize to children who obtain sufficient sleep. In addition, it is important to note that caloric intake was based upon self-report via 24-h dietary recalls, which may be subject to reporter bias. Finally, we examined cross-sectional associations between baseline sleep patterns and obesity risk factors. Therefore, we cannot determine if differences in sleep duration or timing lead to changes in eating behaviors and weight regulation or if differences in eating behaviors and weight regulation lead to changes in sleep patterns. Future experimental studies are needed to deduce cause and effect.

#### 4. Conclusions

In a sample of school-aged children (8–11 years) who reported sleeping  $\leq 9.5$  h per night, we found that bedtime and total sleep time were differentially associated with obesity risk factors. Later bedtime was associated with greater fat intake, later meal timing, and increased caloric intake after the dinner meal whereas shorter total sleep time was associated with higher zBMI. These findings, using an objective measure of sleep duration and timing, provide additional evidence for the importance of sleep hygiene for weight regulation in children. In addition to focusing on sufficient sleep duration, health providers should also consider bedtime when working with children to reduce obesity risk.

#### Funding

This work was supported by the National Institutes of Health NHLBI R01 HL092910, 2011–2016 (PI: Hart).

#### Acknowledgments

We would like to thank all of the families who participated in this study and the staff at Cincinnati's Center for Nutritional Research for conducting 24-h dietary recalls. We would also like to thank all staff who worked on this study, including Amanda Samuels, Victoria Mathieu, Zeeley Denmat, Elizabeth Kuhl, Isabella Casell, Risha Kheterpal, Ashley Greer, Heather Polonksy, Andrew Pool, and Courtney DeAngelis.

#### Conflict of interest

The ICMJE Uniform Disclosure Form for Potential Conflicts of Interest associated with this article can be viewed by clicking on the following link: <https://doi.org/10.1016/j.sleep.2019.02.003>.

#### References

- [1] Hirschowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health* 2015;1:40–3.
- [2] National Sleep Foundation. Sleep in American polly sleep in the modern family summary of findings. 2014. Accessed at Monday 8/1/2016, <https://sleepfoundation.org/sites/default/files/2014-NSF-Sleep-in-America-poll-summary-of-findings-FINAL-Updated-3-26-14-.pdf>.
- [3] Vriend JL, Davidson FD, Corkum PV, et al. Manipulating sleep duration alters emotional functioning and cognitive performance in children. *J Pediatr Psychol* 2013;38(10):1058–69.
- [4] Lo JC, Ong JL, Leong RL, et al. Cognitive performance, sleepiness, and mood in partially sleep deprived adolescents: the need for sleep study. *Sleep* 2016;39(3):687–98.
- [5] Cappuccio FP, Taggart FM, Kandala NB, et al. Meta-analysis of short sleep duration and obesity in children and adults. *Sleep* 2008;31(5):619–26.
- [6] Katzmarzyk PT, Barreira TV, Broyles ST, et al. Relationship between lifestyle behaviors and obesity in children ages 9–11: results from a 12-country study. *Obesity (Silver Spring)* 2015;23(8):1696–702.
- [7] Magee L, Hale L. Longitudinal associations between sleep duration and subsequent weight gain: a systematic review. *Sleep Med Rev* 2012;16(3):231–41.
- [8] Ruan H, Xun P, Cai W, et al. Habitual sleep duration and risk of childhood obesity: systematic review and dose-response meta-analysis of prospective cohort studies. *Sci Rep* 2015;5:16160.
- [9] Wu Y, Gong Q, Zou Z, et al. Short sleep duration and obesity among children: a systematic review and meta-analysis of prospective studies. *Obes Res Clin Pract* 2017;11(2):140–50.
- [10] Fatima Y, Doi SA, Mamun AA. Longitudinal impact of sleep on overweight and obesity in children and adolescents: a systematic review and bias-adjusted meta-analysis. *Obes Rev* 2015;16(2):137–49.
- [11] Miller AL, Lumeng JC, LeBourgeois MK. Sleep patterns and obesity in childhood. *Curr Opin Endocrinol* 2015;22(1):41–7.
- [12] Bornhorst C, Wijnhoven TM, Kunesova M, et al. WHO European Childhood Obesity Surveillance Initiative: associations between sleep duration, screen time and food consumption frequencies. *BMC Public Health* 2015;15:442.
- [13] Franckle RL, Falbe J, Gortmaker S, et al. Insufficient sleep among elementary and middle school students is linked with elevated soda consumption and other unhealthy dietary behaviors. *Prev Med* 2015;74:36–41.
- [14] Hjorth MF, Quist JS, Andersen R, et al. Change in sleep duration and proposed dietary risk factors for obesity in Danish school children. *Pediatr Obes* 2014;9(6):e156–9.
- [15] Kjeldsen JS, Hjorth MF, Andersen R, et al. Short sleep duration and large variability in sleep duration are independently associated with dietary risk factors for obesity in Danish school children. *Int J Obes (Lond)* 2014;38(1):32–9.
- [16] Golley RK, Maher CA, Matricciani L, et al. Sleep duration or bedtime? Exploring the association between sleep timing behaviour, diet and BMI in children and adolescents. *Int J Obes* 2013;37(4):546–51.
- [17] Thivel D, Isacco L, Aucouturier J, et al. Bedtime and sleep timing but not sleep duration are associated with eating habits in primary school children. *J Dev Behav Pediatr* 2015;36(3):158–65.
- [18] Williams SM, Farmer VL, Taylor BJ, et al. Do more active children sleep more? A repeated cross-sectional analysis using accelerometry. *PLoS One* 2014;9(4).
- [19] Asarnow LD, McGlinchey E, Harvey AG. Evidence for a possible link between bedtime and change in body mass index. *Sleep* 2015;38(10):1523–7.
- [20] Olds TS, Maher CA, Matricciani L. Sleep duration or bedtime? Exploring the relationship between sleep habits and weight status and activity patterns. *Sleep* 2011;34(10):1299–307.
- [21] Allison KC, Goel N, Ahima RS. Delayed timing of eating: impact on weight and metabolism. *Curr Obes Rep* 2014;3(1):91–100.
- [22] Eng S, Wagstaff DA, Kranz S. Eating late in the evening is associated with childhood obesity in some age groups but not in all children: the relationship between time of consumption and body weight status in US children. *Int J Behav Nutr Phys* 2009;6.
- [23] Lehto R, Ray C, Lahti-Koski M, et al. Health behaviors, waist circumference and waist-to-height ratio in children. *Eur J Clin Nutr* 2011;65(7):841–8.
- [24] Thompson OM, Ballew C, Resnicow K, et al. Dietary pattern as a predictor of change in BMI z-score among girls. *Int J Obes* 2006;30(1):176–82.
- [25] Timlin MT, Pereira MA, Story M, et al. Breakfast eating and weight change in a 5-year prospective analysis of adolescents: project EAT (eating among teens). *Pediatrics* 2008;121(3):E638–45.
- [26] Diederichs T, Rossbach S, Herder C, et al. Relevance of morning and evening energy and macronutrient intake during childhood for body composition in early adolescence. *Nutrients* 2016;8(11).
- [27] Tin SP, Ho SY, Mak KH, et al. Breakfast skipping and change in body mass index in young children. *Int J Obes (Lond)* 2011;35(7):899–906.
- [28] Maffei C, Provera S, Filippi L, et al. Distribution of food intake as a risk factor for childhood obesity. *Int J Obes* 2000;24(1):75–80.
- [29] Hart CN, Carskadon MA, Considine RV, et al. Changes in children's sleep duration on food intake, weight, and leptin. *Pediatrics* 2013;132(6):e1473–80.
- [30] Baron KG, Reid KJ, Kern AS, et al. Role of sleep timing in caloric intake and BMI. *Obesity* 2011;19(7):1374–81.
- [31] Hart CN, Hawley NL, Wing RR. Development of a behavioral sleep intervention as a novel approach for pediatric obesity in school-aged children. *Sleep Med Clin* 2016;11(4):515–+.
- [32] Acebo C, Sadeh A, Seifer R, et al. Estimating sleep patterns with activity monitoring in children and adolescents: how many nights are necessary for reliable measures? *Sleep* 1999;22(1):95–103.
- [33] Sadeh A, Sharkey KM, Carskadon MA. Activity-based sleep-wake identification – an empirical-test of methodological issues. *Sleep* 1994;17(3):201–7.
- [34] Meltzer LJ, Walsh CM, Traylor J, et al. Direct comparison of two new actigraphs and polysomnography in children and adolescents. *Sleep* 2012;35(1):159–66.
- [35] Fallone G, Seifer R, Acebo C, et al. How well do school-aged children comply with imposed sleep schedules at home? *Sleep* 2002;25(7):739–45.
- [36] Actiwatch communication and sleep analysis software: instruction manual [computer software and manual]. In: Company MMRAR. Bend, OR: Mini Mitter; 2006.
- [37] Acebo C. Actigraphy. *Sleep Compr Handb* 2006:1035–8.
- [38] Crouter SE, Horton M, Bassett Jr DR. Validity of ActiGraph child-specific equations during various physical activities. *Med Sci Sports Exerc* 2013;45(7):1403–9.
- [39] Hanggi JM, Phillips LR, Rowlands AV. Validation of the GT3X ActiGraph in children and comparison with the GT1M ActiGraph. *J Sci Med Sport* 2013;16(1):40–4.
- [40] Kelly LA, McMillan DG, Anderson A, et al. Validity of actigraphs uniaxial and triaxial accelerometers for assessment of physical activity in adults in laboratory conditions. *BMC Med Phys* 2013;13(1):5.
- [41] Crouter SE, Horton M, Bassett DR. Use of a two-regression model for estimating energy expenditure in children. *Med Sci Sports Exerc* 2012;44(6):1177–85.
- [42] Stunkard AJ, Waxman M. Accuracy of self-reports of food intake. *J Am Diet Assoc* 1981;79(5):547–51.
- [43] Dennis B, Ernst N, Hjortland M, et al. The NHLBI nutrition data system. *J Am Diet Assoc* 1980;77(6):641–7.
- [44] Feskanich D, Sielaff BH, Chong K, et al. Computerized collection and analysis of dietary intake information. *Comput Methods Programs Biomed* 1989;30(1):47–57.
- [45] Spaeth AM, Dinges DF, Goel N. Effects of experimental sleep restriction on weight gain, caloric intake, and meal timing in healthy adults. *Sleep* 2013;36(7):981–90.

- [46] Spaeth AM, Dinges DF, Goel N. Sex and race differences in caloric intake during sleep restriction in healthy adults. *Am J Clin Nutr* 2014;100(2):559–66.
- [47] St-Onge MP, Roberts AL, Chen J, et al. Short sleep duration increases energy intakes but does not change energy expenditure in normal-weight individuals. *Am J Clin Nutr* 2011;94(2):410–6.
- [48] Markwald RR, Melanson EL, Smith MR, et al. Impact of insufficient sleep on total daily energy expenditure, food intake, and weight gain. *Proc Natl Acad Sci U S A* 2013;110(14):5695–700.
- [49] Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. *Adv Data* 2000;(314):1–27.
- [50] Capers PL, Fobian AD, Kaiser KA, et al. A systematic review and meta-analysis of randomized controlled trials of the impact of sleep duration on adiposity and components of energy balance. *Obes Rev* 2015;16(9):771–82.
- [51] Baron KG, Reid KJ, Horn LV, et al. Contribution of evening macronutrient intake to total caloric intake and body mass index. *Appetite* 2013;60(1):246–51.
- [52] Fang Z, Spaeth AM, Ma N, et al. Altered salience network connectivity predicts macronutrient intake after sleep deprivation. *Sci Rep* 2015;5:8215.
- [53] St-Onge MP, McReynolds A, Trivedi ZB, et al. Sleep restriction leads to increased activation of brain regions sensitive to food stimuli. *Am J Clin Nutr* 2012;95(4):818–24.
- [54] Greer SM, Goldstein AN, Walker MP. The impact of sleep deprivation on food desire in the human brain. *Nat Commun* 2013;4:2259.
- [55] Kolotourou M, Radley D, Chadwick P, et al. Is BMI alone a sufficient outcome to evaluate interventions for child obesity? *Child Obes* 2013;9(4):350–6.
- [56] Cummings DE, Overduin J. Gastrointestinal regulation of food intake. *J Clin Invest* 2007;117:13–23.
- [57] Leproult R, Van Cauter E. Role of sleep and sleep loss in hormonal release and metabolism. *Endocr Dev* 2010;17:11–21.