



CLINICAL REVIEW

Cognitive behavioral therapy for insomnia: A meta-analysis of long-term effects in controlled studies

Tanja van der Zweerde ^{a,*}, Lampros Bisdounis ^b, Simon D. Kyle ^c, Jaap Lancee ^{b,d}, Annemieke van Straten ^a

^a Department of Clinical Psychology, Amsterdam Public Health, VU University, Amsterdam, the Netherlands

^b Department of Clinical Psychology, University of Amsterdam, Amsterdam, the Netherlands

^c Sleep and Circadian Neuroscience Institute, Nuffield Department of Clinical Neurosciences, University of Oxford, United Kingdom

^d PsyQ Amsterdam, Amsterdam, the Netherlands

ARTICLE INFO

Article history:

Received 20 November 2018

Received in revised form

25 April 2019

Accepted 7 August 2019

Available online 12 August 2019

Keywords:

Behavior therapy

Cognitive therapy

Cognitive behavior therapy

Long-term

Randomized clinical trial

Insomnia

Sleep initiation or maintenance disorder

SUMMARY

Cognitive behavioral therapy for insomnia (CBT-I) is a treatment with moderate to large effects. These effects are believed to be sustained long-term, but no systematic meta-analyses of recent evidence exist. In this present meta-analysis, we investigate long-term effects in 30 randomized controlled trials (RCTs) comparing CBT-I to non-active control groups. The primary analyses ($n = 29$ after excluding one study which was an outlier) showed that CBT-I is effective at 3-, 6- and 12-mo compared to non-active controls: Hedges g for Insomnia severity index: 0.64 (3 m), 0.40 (6 m) and 0.25 (12 m); sleep onset latency: 0.38 (3 m), 0.29 (6 m) and 0.40 (12 m); sleep efficiency: 0.51 (3 m), 0.32 (6 m) and 0.35 (12 m). We demonstrate that although effects decline over time, CBT-I produces clinically significant effects that last up to a year after therapy.

© 2019 Elsevier Ltd. All rights reserved.

Introduction

Insomnia constitutes a serious and common mental health problem with a prevalence of around 6% in the general population (i.e., DSM-IV-TR diagnosis [1]). Furthermore, 30% of the general population suffers from symptoms of insomnia without meeting the criteria of a diagnosis [2]. Insomnia disorder is characterized by a persistent difficulty initiating or maintaining sleep, for three months or longer and for at least three nights a week, resulting in impaired daytime functioning and significant distress [3]. The disorder is associated with high (societal) costs [4] and affects daily life in various domains, such as fatigue, mood changes, declined cognitive ability, physical wellbeing, social relationships and daily tasks [5]. Untreated insomnia often persists for many years [6].

Cognitive behavioral therapy (CBT) is an effective treatment for insomnia. Several systematic reviews and meta-analyses have reported moderate to large short-term post-test effects of CBT for insomnia (CBT-I) [7–9]. These results include robust improvement in insomnia severity ($g = 0.98$; [9]) and sleep efficiency ($g = 0.71$ [9]; $g = 0.91$ [8]; and 9.9% increase in [7]).

In the short term, CBT-I is as effective as pharmacotherapy [10]. However, CBT-I is the preferred treatment according to recommendations in European and American guidelines [11,12]. One reason for advocating CBT-I as first line treatment for insomnia is the risk sleep medication poses of patients developing serious side effects (i.e., dizziness, drowsiness, addiction, and relapse when medication is discontinued). Also, there is insufficient evidence for long-term effects of pharmacotherapy, and therefore long-term use of pharmacotherapy is not recommended [11–14]. Based on this lack of evidence for pharmacotherapy's long-term benefits, CBT-I is now regarded the better choice in the long-term [13]. In other words, it is assumed that CBT-I has long-term effects. Several literature reviews indeed report CBT-I's effects may be durable [10,15,16], but they lack a quantitative data synthesis of recent evidence. To our knowledge, no meta-analysis has been published

* Corresponding author. Department of Clinical Psychology, Faculty of Behavior and Movement Sciences, VU University, Van der Boerhorststraat 7, 1081 BT, Amsterdam, the Netherlands.

E-mail address: t.vander.zweerde@vu.nl (T. van der Zweerde).

Glossary of terms			
BT	behavioral therapy	PI	paradoxical intention
CBT-I	cognitive behavioral therapy for insomnia	PR	progressive relaxation
Comm	community	PSQI	Pittsburgh sleep quality index
CMI	comorbid insomnia	RCTs	randomized controlled trials
CT	cognitive therapy	SC	stimulus control
DSM	diagnostic and statistical manual of mental disorders	SD	standard deviation
DSISD	Duke structured interview for sleep disorders	SE	sleep efficiency
ICSD	International classification of sleep disorders	SH	sleep hygiene
ES	effect size	SIS-D	structured interview for sleep disorders for DSM-III-R
F2F	face-to-face	SOL	sleep onset latency
ISI	insomnia severity index	SR	sleep restriction
MA	meta-analysis	SQ	sleep quality
NWAK	number of awakenings	TST	total sleep time
PE	psycho-education	WASO	wake after sleep onset
		WL	wait-list

that included controlled studies reporting on long-term effects of (partial) CBT-I. Thus, it is unclear what the actual long-term effects of CBT-I are.

In the present meta-analysis, we aim to fill this gap in the literature by including all available RCTs reporting on the controlled long-term effects of CBT-I (at three, six and 12 mo) and quantifying these long-term effects of CBT-I. The focus of the present meta-analysis is on subjective sleep outcomes, in terms of both self-reported symptoms and sleep diaries.

Method

Protocol

Details of the protocol for this meta-analysis were registered on PROSPERO and can be accessed at www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42018094459.

Search strategy

Using our previous meta-analysis [9] as a starting point, we first checked whether the included studies ($n = 87$) had reported controlled follow-up measurements in the original papers or had published follow-up data since. This process included screening the original papers for mention of follow-up measurements done or planned, and literature search for publications from the different research groups in the five years following the first publication, using search terms “long”, “long(-)term” and “follow(-)up”. Additionally, we performed searches in Web of Science, Pubmed and PsycINFO for publications by the authors of the original article on short-term effects to see if since then long-term effects had been published.

We then performed a new search, covering the period from the end of the search (December 2015) of the previous meta-analysis [9] until May 2018. An extensive literature search was carried out in PubMed, PsycINFO, EMBASE and the Cochrane central register of controlled trials, using the same search strategy as in the previous meta-analysis [9]. Terms indicative of insomnia (i.e., insomnia, sleep disorders, sleep initiation and maintenance disorders) were paired with terms indicating psychological treatments (i.e., psychotherapy, cognitive therapy, behavior therapy). For example, our PsycINFO database query was “(DE=(“sleep disorders” or “insomnia”)) and(DE=(“psychotherapy” or “behavior therapy” or

“cognitive behavior therapy” or “cognitive therapy”))”, and specifying we looked for results from December 2015 onwards.

Titles and abstracts were screened by two persons individually (TvdZ and LB) and then crosschecked. Records definitely not meeting criteria (e.g., not a randomized trial), not aimed at insomnia, not psychological but a biological or medical treatment, not an original research report (e.g., a meta-analysis) were excluded based on title and/or abstract. We retrieved the full papers of the remaining 70 references. Two researchers assessed the papers independently (TvdZ and LB). When there was disagreement, the paper was discussed (TvdZ, LB, JL, AvS) until consensus was reached.

Inclusion and exclusion criteria

Inclusion criteria were: 1) randomized controlled trial (RCT); 2) investigating CBT-I or at least one component of it (see below); 3) in adults (18 y and older); 4) with self-reported and/or formally diagnosed insomnia complaints (see Table 1); 5) compared to a non-active control group (e.g., waitlist control, care-as-usual, or a minimal intervention (e.g., education about sleep or sleep hygiene information)); 6) including data for sleep diary outcomes; 7) reporting controlled follow-up data for 12 or more weeks after post-test; 8) providing suitable data to calculate effect sizes.

The following components were identified as being part of CBT-I: relaxation (RE), sleep restriction therapy (SRT), stimulus control therapy (SC), paradoxical intention (PI), and identifying and challenging dysfunctional thoughts (about sleep), i.e., cognitive therapy (CT). All monotherapy (only one CBT-I-component. e.g., RE or SRT only) studies were included in line with the previous M-A [9], to ensure not overestimating effects by only taking full (current) CBT-I into account. All other therapies were excluded (e.g., interpersonal therapy, bright light therapy, exercise, tai chi, biofeedback). Studies on treatment in children or adolescents, on tapering medication use or aimed at treating a different mental health disorder and reporting on insomnia secondarily were also excluded.

Data extraction

We coded the following characteristics of the studies: 1) publication year, 2) recruitment setting (community, primary care, other care facilities, university), 3) the insomnia definition used, 4) comorbidity (e.g., insomnia in breast cancer patients), 5) age group

Table 1
Characteristics of the included studies on (elements of) CBT for insomnia.

Study	Recruitment	Definition insomnia	FU (3, 6 and/or 12 mo)	Total sample size at baseline ^a	Completed follow-up ^a	Co-morbidity	Age	Intervention	format	F2F sess	Control group
Alessi 2016 [52]	Comm (veterans)	ICSD-2 criteria	6 m + 12 m	159	FU1: Tx 92/106 (87%); C 52/53 (98%) FU2: Tx 89/106 (84%); C 51/53 (96%)	OSA or severe mental disorder excluded	60+	CBT-I	Indiv Group	2 f2f, 2 phone	Psychoeducation
Arnedt 2013 [45]	Care	wake time > 60 m + SE < 85% Average SOL ≥ 30 min	3 m	33	Tx 15/15 (100%); C 12/15 (80%)	Excluded	18–65	CBT-I	Phone	4–8	Info
Borkovec & Weerts1976 [24]	Univ		12 m	24	n/a (Tx: n = 11; C: n = 5)	Allowed	–	Relaxation	Group	4	No txt
Casault 2015 [51]	Care	ISI ≥ 8	3 m + 6 m	38	FU1 and FU2: Tx: 15/20 exp (75%); C: 16/18 (88.89%)	Cancer patients	18–75	CBT-I	Self-help + phone consult	n/a	No txt
Creti 2005 [35]	Comm	No formal criteria. Poor sleepers were defined as having trouble initiating or maintaining sleep. Participants were reported to have sleep onset insomnia (>30 min of undesired wakefulness >2 times per week, problem duration >6 mo; 10% of sample), sleep maintenance insomnia (duration of awakenings after sleep onset >30, >2 times per week, problem duration > 6 mo; 49% of sample), or both (41% of sample)."	12 m	27	51% of the sample	Excluded	55+	Relaxation	Self-help	n/a	No txt
Currie 2000 [30]	Care	DSM-III insomnia diagnosis with SIS-D and ICSD diagnosis	3m	60	Tx: 28/32 (88%); C 23/28 (82%)	Pain	<60	CBT-I	Group	7	WL
Edinger 2005 [36]	Comm	mean WASO ≥ 60 m	6 m	36	Tx: 6/18 (33%); C-SH: 7/18 (39%); C-TAU 7/11 (64%)	Fibromyalgia	21–65	Behavioral	Indiv	6	SH/TAU
Edinger 2009 [41]	Care	mean SOL + WASO ≥ 60 m	6 m	81	Tx: 33/41 (80%); C: 33/40 (83%)	Excl (PI) Allow (CMI)	–	Behavioral	Indiv	4	Info
Espie 2001 [23]	Care	ICSD difficulty falling/ maintaining sleep, ≥ 4 n/w, ≥ 3 mo + PSQI ≥ 5	12 m	139	78% of the total sample	Excluded	–	CBT-I	Group	6	WL
Espie 2007 [39]	Care	ICSD/DSM-IV criteria of insomnia	6 m	201	Tx: 76/107 (71%); C 67/94 (71%)	Allowed	–	CBT-I	Group	5	No txt
Espie 2008 [40]	Care	SOL or WASO ≥ 30 m, ≥ 3 n/w, ≥ 3 mo + PSQI ≥ 5	6 m	150	Tx: 67/100 (67%); C:39/50 (78%)	Cancer	18+	CBT-I	Group	5	No txt
Friedman 2000 [31]	Comm	SE < 80%, SOL >30m, TST < 6hr, WASO > 30 m, ≥ 5 n/2w	3 m	39	Tx: 88%; C: 100%	Excluded	55+	Behavioral	Indiv	5	Info
Fuller 2016 [53]	Pharmacy care	ISI ≥ 8	3 m	50	Tx: 19/22 (86%); C: 17/28 (61%)	Terminal illness excluded	18+	Behavioral	Indiv	4 weekly session (2 f2f, 2 phone)	TAU

(continued on next page)

Table 1 (continued)

Study	Recruitment	Definition insomnia	FU (3, 6 and/or 12 mo)	Total sample size at baseline ^a	Completed follow-up ^a	Co-morbidity	Age	Intervention	format	F2F sess	Control group
Irwin 2014 [46]	Comm	SOL or WASO \geq 3 n/w, \geq 3 mo + daytime imp	6 m + 12 m	75	Tx: 46/50 (92%); C: 23/25 (92%)	Excluded	55+	PE + SC + CT + Relaxation	Group	16	Info
Jansson 2012 [42]	Care	SOL or WASO > 30 m, \geq 3 n/w, \geq 6 mo + daytime imp	3m	32	Tx: 15/17 (88%); C: 15/15 (100%)	Hearing impairment	18–65	CBT-I	Indiv	7	WL
Jernelov 2012 [43]	Comm	ISI > 10 + poor sleep \geq 4 wk	3 m	133	Tx1: 41/44 (93%); Tx2: 39/45 (87%); C: 39/44 (89%)	Excluded	18+	CBT-I	Self-help	N/A	WL
Jungquist 2010 [22]	Care	SOL or WASO > 30 m, > 3 n/w, > 6 mo	3 m+6 m	28	Both FU's: Tx: 15/19 (79%); C: 5/9 (56%)	Pain	25+	CBT-I	Indiv	8	Plac
Kaldo 2015 [49]	Comm	Difficulty initiating or maintaining sleep + daytime imp + ISI > 10	6 m + 12 m	148	Tx: 54/73 (74%); C: 53/75 (71%)	Excluded	18+	CBT-I	Self-help	n/a	Plac
Lacks 1983 [29]	Comm	WASO \geq 30 m, \geq 1 n/w, \geq 6 mo	3 m	64	Tx: 7/15 (47%); C: 8/16 (50%)	Excluded	17–59	SC	Group	4	Plac
Lichtstein 2000 [32]	Comm	SOL or WASO \geq 30 m, \geq 3 n/w, \geq 6 mo + daytime imp	3 m	44	Tx: 22/23 (96%); C: 17/21 (81%)	Illness	58+	Relaxation + SC	Indiv	4	WL
Lichtstein 2001 [33]	Comm	SOL or WASO \geq 30 m, \geq 3 n/w, \geq 6 mo + daytime imp	12 m	89	83% of the total sample	Excluded	59+	Relaxation + SR	Indiv	6	Plac
Lovato 2014 [47]	Comm	WASO > 30 m, \geq 3 n/w, > 6 mo + daytime imp	3 m	118	Tx: 72/86 (84%); C: 27/32 (84%)	Excluded	Older	CBT-I	Group	4	WL
McCurry 2016 [54]	Comm	ISI > 11	3 m	106	Tx: 44/53 (83%); C: 42/53 (79%)	Menopausal Sleep interfering major illness excluded	40–65	CBT-I	Phone	n/a	Menopause education
Morin 2005 [37]	Comm	Diagnostic criteria for insomnia: (1) complaint of poor sleep quality or dissatisfaction regarding sleep; (2) symptoms of initial, maintenance, or late insomnia at least 3 nights per week; (3) presence of psychological distress or daytime impairment related to sleep difficulties; and (4) presence of the sleep difficulties for at least 1 mo.	6 m	192	Tx: 81/96 (84%); C: 86/96 (90%)	Allowed	18+	CBT-I	Self-help	n/a	No txt
Rybarczyk 2002 [34]	Care	SOL \geq 45 m or WASO \geq 60 m or TST \leq 5 h, \geq 3 n/w	3 m	24	Tx: 10/16 (63%); Tx2: 13/18 (72%); C: 12/17 (71%)	Illnesses	55+	CBT-I	Group/ Self-help	8 n/a	WL
Savard 2016 [48]	Care	ISI \geq 8 or \geq 2 nights of sleep medication in last 2 wk	3m + 6m + 12m	242	Tx1: 61/81 (75%); Tx2: 49/80 (61%); C: 49/81 (60%)	Cancer	18–75	CBT-I	Indiv/video	6 n/a	No txt

Smith 2015 [50]	Care	SOL or WASO > 30 m, ≥ 2 n/w, > 1 mo	3 m + 6 m	100	Tx: 35/50 (70%); C: 38/50 (76%)	Knee osteoarthritis	–	CBT-I	Indiv	8	Plac
Swift 2012 [44]	Comm	Self-referred; complaints of insomnia. No formal criteria. Baseline ISI M(SD) = 16.73 (5.03); 96.7% of participants report > subclinical insomnia symptoms ISI cut-off.	3 m	151	Tx: 49/75 (65%); C: 63/76 (83%)	Allowed	18+	CBT-I workshop	Group	1 d	WL
Wong 2017 [55]	Care	Chronic primary insomnia based on DSM-5 and ICD-10	3 m + 6 m	216	Tx: 93/111 (84%); C: 82/105 (78%)	Excluded	18+	CBT-I + mindfulness	Group	8 weekly 2.5 h sessions	PE + stretching
Wu 2006 [38]	Comm	SOL or WASO ≥ 30 m, ≥ 6 m + daytime imp	3 m + 6 m	36	0% of the total sample	Excluded	–	CBT-I	Group	16	Plac

Abbreviations: Comm = community; CMI = comorbid insomnia; CBT-I = cognitive behavior therapy for insomnia; CT = cognitive therapy; f2f = face-to-face; ICSD = International Classification of Sleep Disorders; ISI = insomnia severity index; NWAK = number of awakenings; PE = psycho-education; PI = primary insomnia; PSQI = Pittsburgh Sleep Quality Index; SC = stimulus control; SE = sleep efficiency; SH = sleep hygiene; SIS-D = Structured Interview for Sleep Disorders for DSM-III-R; SR = sleep restriction; TST = total sleep time; WL = waitlist.

^a For investigated groups in the study, does not include groups not included in meta-analysis.

(e.g., anyone over 18, or older adults only), 6) the treatment format (individual, group or self-help), 7) number of sessions, 8) the control group (e.g., waitlist, no treatment) and 9) type of intervention. We categorized studies in four categories of interventions: 1) full CBT-I, including an educational component as well as a behavioral and cognitive one, 2) behavioral therapy, including both stimulus control and sleep restriction, 3) relaxation therapy only, and 4) “other”, including e.g., stimulus control only or paradoxical intention only. Two independent assessors coded the studies (TvdZ and LB). Differences were discussed amongst the review team until consensus was reached (TvdZ, LB, JL and AvS).

Quality assessment

Using the criteria suggested in the Cochrane handbook [17], we assessed the validity of the studies: 1) adequate sequence generation, 2) concealment of allocation, 3) adequate handling of incomplete outcome data, and 4) selective reporting of data. We did not assess the blinding of patients or therapists since this is not possible in psychotherapy research nor did we assess blinding of outcome assessors since all reported outcomes are based on self-report. Two reviewers conducted the quality assessment independently of each other (TvdZ and LB) and then crosschecked their findings.

Meta-analysis

We focused on the effects on insomnia severity primarily (measured through questionnaires), and secondarily on sleep onset latency (SOL) and sleep efficiency (SE) measured through sleep diaries because these are among the most important sleep outcomes and yielded the largest number of comparisons. We defined an effect as “long-term” if it was measured at least 12 wk after the end of treatment. We created three subgroups, grouped in 1) three months (between 12- and 17.5-wk post-treatment), 2) six months (between 18 and 35 wk) and 3) 12 mo (more than 35 wk) for pragmatic reasons. There were no studies reporting outcomes after one year. Where the exact timing of the follow-up assessment was unclear from the articles (i.e., post-baseline or post-treatment), we contacted the authors. When we did not receive a reply (n = 1), we assumed the reported follow-up period was post-treatment (however, this assumption did not influence the group assignment of this study).

We computed Hedges' g to determine between group effect sizes. Hedges' g is a measure of standardized mean differences (similar to Cohen's d), after adjusting for small sample sizes [18,19]. This effect size (based on the differences between conditions at the different time points) can be interpreted as the difference between the mean scores of the two groups expressed in the number of weighted pooled standard deviations. Effect sizes (ES) are commonly interpreted as either large (>0.56), moderate (0.33–0.55) or small (0–0.32) [20]. The available statistics (means, standard deviations, standard errors, 95% confidence intervals, and interquartile ranges) from the included studies were used to compute the ESs using the metafor package in R [21].

We checked for outliers by visually inspecting forest plots for all analyses. We defined outliers as studies in which we found a 95% confidence interval around the ES that did not show overlap with the 95% confidence interval of the pooled effect size. One study was identified to be an outlier [22] on all variables it included (SOL, WASO, SE, TST and ISI at 3 mo and 6 mo). Based on author consensus we removed this study from the analysis because we judged it to be a significant and pronounced outlier.

In the 12 mo follow-up, one study had to be excluded from the analysis because means were not provided and could not be

calculated from what was reported (WASO and TST) [23]. For SOL, two studies did not report a measure of dispersion [23,24], but effect sizes were calculated using other available statistics.

Expecting the studies to show significant heterogeneity, we used the random effects model in the metafor package [21] to estimate the weighted pooled effect sizes for the different outcome measures. Heterogeneity was inspected under the fixed effects model using I^2 , describing the variance between studies as a proportion of the total variance. A value of 0% means there is no observed heterogeneity, with larger percentages indicating more heterogeneity [25]. The 95% confidence intervals around I^2 were calculated.

To address treatment heterogeneity in the included studies we performed sensitivity analyses, exploring differences in effects of full CBT-I compared to effects of partial CBT-I or monotherapy.

To assess potential publication bias, we visually inspected the funnel plot and conducted Egger's test for all variables, at the three different time points [26]. The Duval and Tweedie [27] trim and fill procedure was used to adjust the effect size for publication bias and to provide an indication of the number of studies that might have been missing from the analysis.

Results are reported in accordance with the PRISMA statement and the PRISMA checklist can be found in the supplement [28]. Results in the expected and desired direction are reported as positive results, results in the undesired direction (i.e., worsening of symptoms) are reported as negative results.

Results

Selection of included studies

We selected studies using two strategies. First, we included articles identified in the previous meta-analysis [9], which had controlled 12 wk or more follow-up data. A total of 25 studies were found using this method [23,24,29–50]. Secondly, we performed a new literature search for the period since the previous meta-analysis [9]. We screened 420 titles and abstracts and excluded 324 as not relevant and/or not meeting criteria based on their titles and abstracts. In total, we then assessed 96 full texts. Of these, we excluded 66 articles because 1) duplicates between the first and second search strategy ($n = 5$), 2) not a controlled RCT ($n = 24$), 3) no CBT component ($n = 3$), 4) not focused on insomnia ($n = 9$), 5) no sleep diary ($n = 14$), 6) no follow-up ($n = 3$), 7) no data we could use for calculations of the effect sizes ($n = 5$) or 8) not a results paper ($n = 3$). The remaining 30 papers were included in this meta-analysis: the 25 already included in our previous meta-analysis and five new papers reporting follow-up results of an RCT on a component of) CBT-I compared to an inactive control group [51–55]. A flowchart of the inclusion and exclusion process is in Fig. 1.

Two studies compared different active interventions in comparison to a control condition. Therefore, the meta-analysis included 32 comparisons in total. Some studies ($n = 9$) reported multiple follow-up measurements and not all studies reported the same sleep variables. The numbers of comparisons vary in the different analyses and are listed in the tables.

Study characteristics

Eighteen studies reported three months follow-up data [22,29–32,34,38,42–45,47,48,50,51,53–55], fourteen studies reported data for six months follow-up [22,36–41,46,48–52,55] and eight studies reported 12 mo follow-up data [23,24,33,35,46,48,49,52]. The oldest study in our sample was published in 1976 [24]. The large majority was published after 2000 ($n = 28$;

93%). Fourteen studies (47%) investigated community samples, 15 studies (50%) recruited patients from care settings and one study recruited within a university (3%). Around 50% of the studies ($n = 16$) excluded patients showing (specific) comorbidities. The remaining 14 studies did not, or did not report it. Out of the 30 studies included in the qualitative assessments, twenty studies (67%) offered full CBT-I, the others ($n = 10$; 33%) offered one or more components of CBT-I but not the full package. Treatment was offered in group format in 13 studies, in individual format in 13 studies and in self-help format in six studies (total $n = 32$ treatment groups across studies). Treatment groups were compared to wait-list ($n = 8$), no treatment ($n = 7$), care-as-usual ($n = 2$), placebo ($n = 6$) or minimal intervention ($n = 8$) control groups (total $n = 31$ control groups across studies). See Table 1 for details.

Quality assessments

Sixteen out of the 30 studies included in the qualitative assessments (53%) generated randomization sequences adequately, whereas 14 (47%) did not, or did not report on it. A total of 13 studies (43%) reported adequate concealing of the random allocation, for the other 17 studies (57%) this was not reported. Twenty studies (67%) reported handling missing data by performing intent-to-treat analysis, six studies (20%) did not conduct intent-to-treat analysis and four (13%) did not report how they handled missing data. Eight studies (27%) seemed to be selective in reporting of data based on comparisons of study protocols to results reported.

Three to 12 mo effects on insomnia severity, sleep efficiency and sleep onset latency

Three months after CBT-I finished the effect compared to the non-treated controls on ISI was statistically significant and large ($g = 0.64$; $N_c = 13$). The effect remained present after three months, with a moderate between group effect size at six months ($g = 0.40$; $N_c = 8$) and a small between group effect size at 12 mo ($g = 0.25$; $N_c = 4$).

Three months after CBT-I was finished the effect on sleep efficiency was moderate ($g = 0.51$; $N_c = 21$) and remained relatively stable over time (six months: $g = 0.32$; $N_c = 16$); 12 mo $g = 0.35$; $N_c = 8$). The effect size of CBT-I on SOL declined from moderate at three months to small at six months but returned to moderate at 12 mo ($g = 0.38, 0.29$ and 0.40 respectively with $N_c = 21, 16$, and 10). Forest plots for ISI, SE and SOL can be found in Figs. S1–S3.

Three to 12 mo effects on secondary outcomes

For the secondary variables, treated groups outperformed the control groups at three months on most variables (PSQI: $g = 0.80$; $N_c = 6$; WASO: $g = 0.42$; $N_c = 20$; SQ $g = 0.49$; $N_c = 5$). Effects on TST ($g = 0.06$; $N_c = 21$) and NWAK ($g = 0.08$; $N_c = 4$) were not significant (see Table 2). At six months, effects were present, but most had become slightly smaller (WASO: $g = 0.27$; $N_c = 13$; PSQI: $g = 0.48$; $N_c = 3$). Effects on SQ ($g = 0.09$; $N_c = 2$) and TST ($g = 0.05$; $N_c = 15$) were non-significant. NWAK data were not available at six months. At 12 mo, results are less reliable due to the small number of studies included in the analysis (WASO: $g = 0.26$; $N_c = 8$; NWAK: $g = 0.52$; $N_c = 2$). Effects on PSQI, SQ and TST were not significant (PSQI: $g = 0.22$; $N_c = 2$; SQ: $g = 0.24$; $N_c = 3$; TST: $g = 0.03$; $N_c = 7$).

Differences between post-test effects of studies with long-term effects vs all studies

We calculated the post-test effect sizes of the 29 studies (excluding one outlier) in the current meta-analysis (ISI, SE, SOL).

We then divided the studies based on follow-up length and compared the post-test effects of studies with 3 mo follow-up, 6 mo follow-up and 12 mo follow-up to the overall post-test effect sizes (results are in [Supplementary Table S1](#)).

Sensitivity analysis

We made a comparison between the sample with all studies included to the sample with only the studies with full CBT-I. Due to variety in follow-up length and reported variables there were different samples only for ISI (3m), SOL (3/6/12m), and SE (3/12m; see [Supplemental Table S2](#) for Nc's and effect sizes). The differences in g ranged from 0.01 to 0.05. We also compared the sample with all studies included with the sample excluding monotherapies. There were different samples only for SE (Nc = 8 versus 6) and SOL (Nc = 10 vs 8), both at 12 m. The difference in g was 0.03 for both (see [Supplemental Table S2](#)).

Publication bias

We found indications for publication bias in only one of the analyses performed: the three-month follow-up effects on total

sleep time. Egger's test for funnel plot asymmetry was significant ($t = 2.38$, $df = 19$, $p = .028$). We performed a trim and fill analysis and found the estimated number of studies missing on the left side of the funnel plot to be $n = 1$. The effect size adjusted for publication bias for three-month TST was 0.039 ($p = .601$), slightly smaller than the 0.061 ($p = .391$) unadjusted effect size, and remained statistically non-significant.

Discussion

We performed a meta-analysis on 29 RCTs (removing one study which was an outlier) to investigate the long-term effects of cognitive and/or behavioral treatments for insomnia. We studied three, six and 12-mo follow-up data in separate analyses. Three months after treatment, the severity of insomnia complaints (primary outcome) was considerably better for patients treated with CBT-I than for patients without an active treatment (ISI: $g = 0.64$). Effects were of moderate size at six months (ISI: $g = 0.40$). At 12 mo the treated group still outperformed the control group, albeit showing a smaller effect size (ISI: $g = 0.25$). The sleep diary variables (sleep efficiency and sleep onset latency; secondary outcomes) showed a similar pattern, although effect sizes were smaller




Fig. 1. PRISMA 2009 flow diagram.

Table 2
Effects of insomnia treatments at 3-, 6- and 12 mo^a follow-up (outlier removed).

	3 mo follow-up			6 mo follow-up			12 mo follow-up					
	Nc	g (95% CI)	I ² (95% CI)	p	Nc	g (95% CI)	I ² (95% CI)	p	Nc	g (95% CI)	I ² (95% CI)	p
ISI	13	0.64 (0.42–0.86)	70.85 (40.34–88.37)	<0.001	8	0.40 (0.23–0.56)	45.26 (0.00–86.42)	<0.001	4	0.25 (0.05–0.46)	39.77 (0.00–95.50)	0.016
SE	21	0.51 (0.34–0.69)	62.97 (35.57–84.46)	<0.001	16	0.32 (0.17–0.47)	51.62 (9.80–82.12)	<0.001	8	0.35 (0.21–0.49)	n/a	<0.001
SOL	21	0.38 (0.20–0.57)	65.34 (38.59–84.90)	<0.001	16	0.29 (0.13–0.46)	62.38 (27.74–86.38)	<0.001	10	0.40 (0.23–0.57)	26.30 (0.00–92.45)	<0.001
PSQI	6	0.78 (0.34–1.25)	69.57 (21.10–94.33)	<0.001	3	0.48 (0.22–0.75)	57.96 (0.00–98.91)	<0.001	2	0.23 (0.05–0.50)	0 (0.00–99.56)	0.107
WASO	20	0.42 (0.25–0.59)	58.66 (28.04–84.70)	<0.001	13	0.27 (0.15–0.39)	23.16 (0.00–83.65)	<0.001	8	0.26 (0.15–0.46)	0 (0.00–70.59)	<0.001
SQ	5	0.49 (0.02–0.96)	71.04 (15.44–96.97)	0.039	2	0.09 (–0.30–0.48)	69.66 (0.00–99.97)	0.657	3	0.24 (–0.08–0.56)	28.83 (0.00–97.59)	0.138
NWAK	4	0.08 (–0.44–0.28)	42.98 (0–96.03)	0.679	0	n/a	n/a	n/a	2	0.52 (0.12–0.93)	0 (0.00–99.73)	0.012
TST	21	0.06 (–0.08–0.20)	37.76 (0–74.34)	0.391	15	0.05 (–0.05–0.15)	<0.01 (0–77.89)	0.311	7	–0.02 (–0.17–0.14)	0 (0.00–76.78)	0.848
Adjusted TST ^b	21	0.04 (–0.11–0.18)		0.600								

Note. g = Hedges g effect size; ISI = Insomnia Severity Index; PSQI = Pittsburgh Sleep Quality Index; SE = sleep efficiency; WASO = wake after sleep onset; SOL = sleep onset latency; SQ = sleep quality; NWAK = number of awakenings; TST = total sleep time; Nc = number of comparisons.

^a The times to follow up were defined as 3 mo when >12 wk and <17.5 wk post-treatment, 6 mo as >18 wk and <35 wk post-treatment, and 12 mo as >40 wk post-treatment.

^b Publication bias detected using Egger's test ($t = 2.38$, $df = 12$, $p = .028$), estimated number of studies missing $n = 1$.

for three months outcomes (SE: $g = 0.51$; SOL: $g = 0.38$). In general, there seems to be a steady decline over time, indicating that the long-term effects of CBT-I are smaller than effects observed in the short term [7–9]. We identify three possible reasons for this decline in effects over time.

Firstly, it could be an artefact specific to this meta-analysis. The analyses at the different time points include different studies and this could lead to biased effect size estimates. For instance, the decline in effect may be related to our decision to include only a subset of studies, those with *controlled* follow-up data. In our view this is a major strength of our analysis since it allowed us to infer whether treatment effects at long-term are actually due to the treatment and not caused by the passage of time. However, the lower effect sizes could reflect this meta-analysis not being a representative sample of studies on CBT-I post-test effects.

To address this, we compared the post-test effects of studies with 3, 6- and 12-mo follow-up to overall post-test effect sizes (See Supplemental Table S1). When compared to the post-test effects in our previous meta-analysis (ISI, $g = 0.98$) [9], the present sample showed relatively similar effect sizes and overlapping confidence intervals at post-test: 3 mo, $g = 0.91$, 6 mo, $g = 0.62$; 12 mo, $g = 0.79$; all studies in the current meta-analysis, $g = 0.83$. For the sleep diary variables, the differences were also relatively small: sleep efficiency effect size was $g = 0.71$ in the previous meta-analysis [9]. Now effect sizes ranged between 0.53 and 0.74 (all with overlapping confidence intervals). The same pattern was observed for SOL with an effect size of $g = 0.57$ in [9], effect sizes now ranging between 0.45 and 0.58 (with overlapping confidence intervals). Overall, the post-test effect sizes of the studies including controlled follow-up studies seem to be in a somewhat lower range. However, an artefact effect in our sample is not likely to be the sole explanation.

A second possible reason is that the patients in the control conditions start sleeping better over time, either because they have sought treatment elsewhere or merely as a result of time passing. When looking at the data from the individual studies included in this meta-analysis, we do observe control participants reporting a small decrease in insomnia symptoms. We do not know whether control patients have undergone treatment. As the time of follow-up increases, so does the window of opportunity to seek treatment elsewhere. Earlier research on the natural course of insomnia reports that insomnia is often persistent when untreated [56], suggesting the symptom decrease in control participants is unlikely to have happened spontaneously. A three-year follow-up study by Blom and colleagues showed that during the follow-up phase, control group participants were indeed likely to use sleep medication and seek additional insomnia treatment elsewhere [57]. Our results may therefore be on the conservative side: if control participants have sought effective help elsewhere, this may have led to a smaller difference between the treatment conditions and the controls.

A third possible reason is an increase in the intervention participants' insomnia symptoms over time. Again, when looking at the data for the individual studies included in this meta-analysis, we see that on average intervention participants report a slight return of symptoms over time. This would mean the effects of the intervention in the long run are not as pronounced as is often thought. This explanation seems plausible: the interventions are relatively short, and focus on behavioral changes (e.g., lifestyle, bedtimes, sleep hygiene). These behavior changes can be hard to stick to, as we know from extensive research in other areas of preventive care where lifestyle changes are necessary (e.g., increasing physical activity, smoking cessation, reducing alcohol consumption [58]).

Our results, therefore, may be explained by an interplay between these three explanations. To answer this question more definitively, we need more randomized controlled trials that include a long and controlled follow-up with multiple measurements (e.g., 3 mo and 12 mo) reporting on what happens in the follow-up phase as well. It would also be good to adopt a more uniform research approach in reporting on (long-term) treatment trials. For this meta-analysis, included studies showed substantial heterogeneity. Furthermore, not all variables investigated were reported in all included studies, limiting the number of comparisons. Pragmatic choices had to be made in choosing outcome variables that present a balanced picture, but SE in particular might not be an ideal outcome to report [59]. Given the small number of studies reporting 12-mo follow-up data ($n = 8$), results at 12 mo should be interpreted with caution. To enhance uniformity, we advise CBT-I researchers to include the Carney consensus diary variables [60] and the ISI [61] in future research. This would increase the opportunities for meaningful pooling of evidence and also allow meaningful subgroup analyses providing insight into the variables influencing magnitude of effect size.

The heterogeneity of included studies is a potential limitation of this meta-analysis. Studies differ in terms of comorbid populations, delivery mode and treatment content. We decided to include these studies for several reasons: 1) Current diagnostic practice no longer makes a distinction between primary and secondary/comorbid insomnia. 2) previous research shows delivery mode is not an important factor determining effects [62,63]. 3) the previous meta-analysis by van Straten and colleagues (2018) showed full CBT-I effects did not differ from those of partial CBT-I [9]. Sensitivity analysis on the current dataset supported our choice (Supplemental Table S2).

Importantly, although the effects decline somewhat over time, we did find sustained long-term CBT-I effects. These established long-term effects strengthen the claim that CBT-I outperforms pharmacotherapy in the long run and is the preferred treatment for insomnia (e.g., [13]). The sustained and (tentatively) clinically relevant effects ($ES > 0.25$ for depression treatment according to Cuijpers and colleagues [64]) are of particular interest. It must be noted however that relatively few meta-analyses on other psychological disorders have established long-term controlled treatment effects [65–67]. Meta-analyses are often limited to short-term effects, due to difficulties interpreting varied follow-up intervals and potential for other treatments or life events during the longer follow-up phase [67]. The few meta-analyses that have investigated long-term effects have generally reported similar findings: sustained (but somewhat declined) long-term treatment effects for CBT for depression [68,69], anxiety [70] and PTSD [71].

To enhance long-term effects, perhaps we need to put more emphasis on relapse prevention within CBT-I. Currently, relapse prevention is a component of most interventions at the end of treatment, making patients aware of the potential of relapse. They are advised to return to the exercises in the intervention when this happens. This could be improved, for example, by adding a booster session after six months or asking patients to continue keeping a sleep diary for a longer period of time.

Taken together, the results of the present meta-analysis show favorable effects of CBT-I at follow-up, still present (albeit smaller) at 12 mo after treatment. Establishing this long-term effect is of major importance: it provides a strong argument for the clinical recommendation of offering CBT-I.

Ultimately, the main goal of CBT-I is to improve daytime functioning, improving quality of life and reducing societal costs due to absenteeism and work productivity losses. Research on long-term

effects of CBT-I on these measures is currently scarce, although daytime impairment is the main reason patients seek treatment [2,72]. This meta-analysis indicates that CBT-I does show the often-claimed long-term effectiveness, but it is not without its limitations. In further insomnia research, we need to aim for more uniformity and more controlled studies with a longer follow-up (preferably one year). This would enable confidently stating that CBT-I has the often-proclaimed long term effects, improves both insomnia severity and daytime functioning and should be the first treatment of choice, in line with recent recommendations for the treatment of insomnia symptoms [12,13].

Author contributions

TZ, AVS, SD and JL conceived of and designed the study. TZ, AVS and JL performed the literature search. TZ and LB screened the search results, completed the in- and exclusion process, and extracted and cross-validated the data, supervised by AVS and JL. TZ and LB drew up the analysis plan under supervision of AVS and JL. LB coded the R-scripts under supervision of TZ. TZ drafted the manuscript. All authors contributed to and approved the manuscript.

Practice points

- Cognitive behavioral therapy for insomnia is effective on insomnia severity, sleep efficiency and sleep onset latency at three, six- and 12-mo follow-up.
- Long-term effects provide support for international guidelines recommending cognitive behavioral therapy as the first treatment option for insomnia.

Research agenda

Future studies on cognitive behavioral treatment for insomnia should aim to:

- Include long-term controlled follow-ups
- Provide information on daytime functioning as a consequence of insomnia and changes in functioning after treatment
- Be uniform in reporting on randomized controlled trials and adequately monitor control participants during the follow-up phase.

Acknowledgements

All authors declare that they have no competing interests and did not receive financial support to perform this meta-analysis. SDK is supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.smr.2019.08.002>.

References

- [1] Ohayon MM. Epidemiology of insomnia: what we know and what we still need to learn. *Sleep Med Rev* 2002;6:97–111.
- *[2] Morin CM, LeBlanc M, Daley M, Grégoire JP, Merette C. Epidemiology of insomnia: prevalence, self-help treatments, consultations, and determinants of help-seeking behaviors. *Sleep Med* 2006;7:123–30.
- [3] American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. 2013. Washington, DC.
- [4] Daley M, Morin CM, Le Blanc M, Gregoir JP, Savard J. The economic burden of insomnia: direct and indirect costs for individuals with insomnia syndrome, insomnia symptoms, and good sleepers. *Sleep* 2009;32:55–64.
- [5] Kyle SD, Espie CA, Morgan K. Quality of life and daytime functioning in insomnia. *Behav Sleep Med* 2010;8:123–40.
- [6] Morin CM, Vallières A, Guay B, Ivers H, Savard J, Mérette C, et al. Cognitive behavioral therapy, singly and combined with medication, for persistent insomnia: a randomized controlled trial. *JAMA* 2009;301:2005–15.
- *[7] Trauer JM, Qian MY, Doyle JS, Rajaratnam SM, Cunnington D. Cognitive behavioral therapy for chronic insomnia: a systematic review and meta-analysis. *Ann Intern Med* 2015;163:191–204.
- *[8] Wu JQ, Appleman ER, Salazar RD, Ong JC. Cognitive behavioral therapy for insomnia comorbid with psychiatric and medical conditions: a meta-analysis. *JAMA Intern Med* 2015;175:1461–72.
- *[9] van Straten A, van der Zweerde T, Kleiboer A, Cuijpers P, Morin CM, Lancee J. Cognitive and behavioral therapies in the treatment of insomnia: a meta-analysis. *Sleep Med Rev* 2018;38:3–16.
- [10] Smith MT, Perlis ML, Park A, Smith MS, Pennington J, Giles DE, et al. Comparative meta-analysis of pharmacotherapy and behavior therapy for persistent insomnia. *Am J Psychiatry* 2002;159:5–11.
- *[11] Qaseem A, Kansagara D, Forcica MA, Cooke M, Denberg TD. Management of chronic insomnia disorder in adults: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2016;165:125–33.
- *[12] Riemann D, Baglioni C, Bassetti C, Bjorvatn B, Dolenc Groselj L, Ellis JG, et al. European guideline for the diagnosis and treatment of insomnia. *J Sleep Res* 2017;26:675–700.
- [13] Buscemi N, Vandermeer B, Friesen C, Bialy L, Tubman M, Ospina M, et al. The efficacy and safety of drug treatments for chronic insomnia in adults: a meta-analysis of RCTs. *J Gen Intern Med* 2007;22:1335.
- [14] Riemann D, Perlis ML. The treatments of chronic insomnia: a review of benzodiazepine receptor agonists and psychological and behavioral therapies. *Sleep Med Rev* 2009;13:205–14.
- [15] Wang MY, Wang SY, Tsai PS. Cognitive behavioural therapy for primary insomnia: a systematic review. *J Adv Nurs* 2005;50:553–64.
- [16] Mitchell MD, Gehrman P, Perlis M, Umscheid CA. Comparative effectiveness of cognitive behavioral therapy for insomnia: a systematic review. *BMC Fam Pract* 2012;13:40.
- [17] Higgins JPT, Altman DG. Assessing risk of bias in included studies. In: Higgins JPT, Green S, editors. *Cochrane Handbook for systematic reviews of interventions*. Chichester, UK: Wiley-Blackwell; 2008 [Chapter 8].
- [18] Cooper H, Hedges LV. *The Handbook of research synthesis*. New York: Russell Sage Foundation Press; 1994.
- [19] Hedges LV, Olkin I. *Statistical methods for meta-analysis*. San Diego, CA: Academic Press; 1985.
- [20] Lipsey MW, Wilson DB. The efficacy of psychological, educational and behavioural treatment. Confirmation from meta-analysis. *Am Psychol* 1993;48:1181–209.
- [21] Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw* 2010;36:1–48.
- [22] Jungquist CR, Tra Y, Smith MT, Pigeon WR, Matteson-Rusby S, Xia Y, et al. The durability of cognitive behavioral therapy for insomnia in patients with chronic pain. *Sleep disorders* 2012. <https://doi.org/10.1155/2012/679648>. Article ID 679648.
- [23] Espie CA, Inglis SJ, Tessier S, Harvey L. The clinical effectiveness of cognitive behaviour therapy for chronic insomnia: implementation and evaluation of a sleep clinic in general medical practice. *Behav Res Ther* 2001;39:45–60.
- [24] Borkovec TD, Weerts TC. Effects of progressive relaxation on sleep disturbance: an electroencephalographic evaluation. *Psychosom Med* 1976;38:173–80.
- [25] Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557.
- [26] Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
- [27] Duval SJ, Tweedie RL. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56:455–63.
- [28] Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–9.
- [29] Lacks P, Bertelson AD, Sugerman J, Kunkel J. The treatment of sleep maintenance insomnia with stimulus-control techniques. *Behav Res Ther* 1983;21:291–5.
- [30] Currie SR, Wilson KG, Pontefract AJ, deLaplante L. Cognitive-behavioral treatment of insomnia secondary to chronic pain. *J Consult Clin Psychol* 2000;68:407–16.
- [31] Friedman L, Benson K, Noda A, Zarcone V, Wicks DA, O'Connell K, et al. An actigraphic comparison of sleep restriction and sleep hygiene treatments for insomnia in older adults. *J Geriatr Psychiatry Neurol* 2000;13:17–27.
- [32] Lichstein KL, Wilson NM, Johnson CT. Psychological treatment of secondary insomnia. *Psychol Aging* 2000;15:232–40.
- [33] Lichstein KL, Riedel BW, Wilson NM, Lester KW, Aguillard RN. Relaxation and sleep compression for late-life insomnia: a placebo-controlled trial. *J Consult Clin Psychol* 2001;69:227–39.
- [34] Rybarczyk B, Lopez M, Benson R, Alsten C, Stepanski E. Efficacy of two behavioral treatment programs for comorbid geriatric insomnia. *Psychol Aging* 2002;17:288–98.
- [35] Creti L, Libman E, Bailes S, Fichten CS. Effectiveness of cognitive behavioral insomnia treatment in a community sample of older individuals: more questions than conclusions. *J Clin Psychol Med Settings* 2005;12:153–64.
- [36] Edinger JD, Wohlgenuth WK, Krystal AD, Rice JR. Behavioral insomnia therapy for fibromyalgia patients. *Arch Intern Med* 2005;165:2527–35.
- [37] Morin CM, LeBlanc M, Savard J. Self-help treatment for insomnia: a randomized controlled trial. *Sleep* 2005;28:1319–27.
- [38] Wu R, Bao J, Zhang C, Deng J, Long C. Comparison of sleep condition and sleep-related psychological activity after cognitive-behavior and pharmacological therapy for chronic insomnia. *Psychother Psychosom* 2006;75:220–8.
- [39] Espie CA, MacMahon KMA, Kelly H, Broomfield NM, Douglas NJ, Engleman HM, et al. Randomized clinical effectiveness trial of nurse administered small-group cognitive behavior therapy for persistent insomnia in general practice. *Sleep* 2007;30:574–84.
- [40] Espie CA, Fleming L, Cassidy J, Samuel L, Taylor LM, White CA, et al. Randomized controlled clinical effectiveness trial of cognitive behavior therapy compared with treatment as usual for persistent insomnia in patients with cancer. *J Clin Oncol* 2008;26:4651–8.
- [41] Edinger JD, Olsen MK, Stechuchak KM, Means MK, Lineberger MD, Kirby A, et al. Cognitive behavioral therapy for patients with primary insomnia or insomnia associated predominantly with mixed psychiatric disorders: a randomized clinical trial. *Sleep* 2009;32:499–510.
- [42] Jansson-Fröjmark M, Linton SJ, Flink IK, Granberg S, Danermark B, Norell-Clarke A. Cognitive-behavioral therapy for insomnia co-morbid with hearing impairment: a randomized controlled trial. *J Clin Psychol Med Settings* 2012;19:224–34.
- [43] Jernelov S, Lekander M, Blom K, Rydh S, Ljotsson B, Axelsson J, et al. Efficacy of a behavioral self-help treatment with or without therapist guidance for co-morbid and primary insomnia: a randomized trial. *BMC Psychiatry* 2012;12:5.
- [44] Swift N, Stewart R, Andiappan M, Smith A, Espie CA, Brown JSL. The effectiveness of community day-long CBT-I workshops for participants with insomnia symptoms: a randomised controlled trial. *J Sleep Res* 2011;21:270–80.
- [45] Arnedt JT, Cudihy L, Swanson LM, Pickett S, Aikens J, Chervin RD. Randomized controlled trial of telephone-delivered cognitive behavioral therapy for chronic insomnia. *Sleep* 2013;36:353–62.
- [46] Irwin MR, Olmstead R, Carrillo C, Sadeghi N, Breen EC, Witarama T, et al. Cognitive behavioral therapy vs. Tai Chi for late life insomnia and inflammatory risk: a randomized controlled comparative efficacy trial. *Sleep* 2014;37:1543–52.
- [47] Lovato N, Lack L, Wright H, Kennaway DJ. Evaluation of a brief treatment program of cognitive behavior therapy for insomnia in older adults. *Sleep* 2014;37:117–26.
- [48] Savard J, Ivers H, Savard MH, Morin CM. Long-term effects of two formats of cognitive behavioral therapy for insomnia comorbid with breast cancer. *Sleep* 2016;39:813–23.
- [49] Kaldov V, Jernelov S, Blom K, Ljotsson B, Brodin M, Jorgensen M. Guided internet cognitive behavioral therapy for insomnia compared to a control treatment: a randomized trial. *Behav Res Ther* 2015;71:90–100.
- [50] Smith MT, Finan PH, Buenaver LF, Robinson M, Haque U, Quain A, et al. Cognitive behavioral therapy for insomnia in knee osteoarthritis: a randomized, double-blind, active placebo-controlled clinical trial. *Arthritis Rheum* 2015;67:1221–33.
- [51] Casault L, Savard J, Ivers H, Savard MH. A randomized-controlled trial of an early minimal cognitive-behavioural therapy for insomnia comorbid with cancer. *Behav Res Ther* 2015;67:45–54.
- [52] Alessi C, Martin JL, Fiorentino L, Fung CH, Dzierzewski JM, Rodriguez Tapia JC, et al. Cognitive behavioral therapy for insomnia in older veterans using nonclinician sleep coaches: randomized controlled trial. *J Am Geriatr Soc* 2016;64:1830–8.
- [53] Fuller JM, Wong KK, Hoyos C, Krass I, Saini B. Dispensing good sleep health behaviours not pills—a cluster-randomized controlled trial to test the feasibility and efficacy of pharmacist-provided brief behavioural treatment for insomnia. *J Sleep Res* 2016;25:104–15.
- [54] McCurry SM, Guthrie KA, Morin CM, Woods NF, Landis CA, Ensrud KE, et al. Telephone-based cognitive behavioral therapy for insomnia in perimenopausal and postmenopausal women with vasomotor symptoms— a MsFLASH randomized clinical trial. *JAMA Intern Med* 2016;176:913–20.
- [55] Wong SYS, Zhang DX, Li CCK, Yip BHK, Chan DCC, Ling YM, et al. Comparing the effects of mindfulness-based cognitive therapy and sleep psycho-

* The most important references are denoted by an asterisk.

- education with exercise on chronic insomnia: a randomised controlled trial. *Psychother Psychosom* 2017;86:241–53.
- [56] Morin CM, Bélanger L, LeBlanc M, Ivers H, Savard J, Espie CA, et al. The natural history of insomnia: a population-based 3-year longitudinal study. *Ann Intern Med* 2009;169:447–53.
- [57] Blom KLP, Jernešević S, Rück C, Lindefors N, Kaldo V. Three-year follow-up of insomnia and hypnotics after controlled internet treatment for insomnia. *Sleep* 2016;39:1267–74.
- [58] Afshin A, Babalola D, Mclean M, Yu Z, Ma W, Chen CY, et al. Information technology and lifestyle: a systematic evaluation of internet and mobile interventions for improving diet, physical activity, obesity, tobacco, and alcohol use. *J Am Heart Assoc* 2016;5:e003058.
- [59] Schwartz DR, Carney CE. Mediators of cognitive-behavioral therapy for insomnia: a review of randomized controlled trials and secondary analysis studies. *Clin Psychol Rev* 2012;32:664–75.
- *[60] Carney CE, Buysse DJ, Ancoli-Israel S, Edinger JD, Krystal AD, Lichstein KL, et al. The consensus sleep diary: standardizing prospective sleep self-monitoring. *Sleep* 2012;35:287–302.
- *[61] Bastien CH, Vallières A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med* 2001;2:297–307.
- [62] Andersson G, Cuijpers P, Carlbring P, Riper H, Hedman E. Guided Internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: a systematic review and meta-analysis. *World Psychiatry* 2014;13:288–95.
- [63] Carlbring P, Cuijpers P, Riper H, Hedman E, Rozental A, Shafraan R, et al. Internet-based vs. Face-to-Face CBT: systematic review and meta-analysis. In: 52nd annual association for behavioral and cognitive therapies convention, Washington, USA; November 15–18, 2018.
- [64] Cuijpers P, Turner EH, Koole SL, Van Dijke A, Smit F. What is the threshold for a clinically relevant effect? The case of major depressive disorders. *Depress Anxiety* 2014;31:374–8.
- [65] van Straten A, Geraedts A, Verdonck-de Leeuw I, Andersson G, Cuijpers P. Psychological treatment of depressive symptoms in patients with medical disorders: a meta-analysis. *J Psychosom Res* 2010;69:23–32.
- *[66] Hofmann SG, Asnaani A, Vonk IJ, Sawyer AT, Fang A. The efficacy of cognitive behavioral therapy: a review of meta-analyses. *Cogn Ther Res* 2012;36:427–40.
- [67] Cuijpers P, Berking M, Andersson G, Quigley L, Kleiboer A, Dobson KS. A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *Can J Psychiatry* 2013;58:376–85.
- *[68] Tolin DF. Is cognitive-behavioral therapy more effective than other therapies? A meta-analytic review. *Clin Psychol Rev* 2010;30:710–20.
- [69] Karyotaki E, Smit Y, Henningsen KH, Huijbers MJH, Robays J, de Beurs D, et al. Combining pharmacotherapy and psychotherapy or monotherapy for major depression? A meta-analysis on the long-term effects. *J Affect Disord* 2016;194:144–52.
- [70] Gil PJM, Carrillo FXM, Meca JS. Effectiveness of cognitive-behavioural treatment in social phobia: a meta-analytic review. *Psychol Spain* 2001;5:17–25.
- [71] Kline AC, Cooper AA, Rytwinski NK, Feeny NC. Long-term efficacy of psychotherapy for posttraumatic stress disorder: a meta-analysis of randomized controlled trials. *Clin Psychol Rev* 2017;59:30–40.
- [72] Kyle SD, Crawford MR, Morgan K, Spiegelhalter K, Clark AA, Espie CA. The Glasgow Sleep Impact Index (GSII): a novel patient-centered measure for assessing sleep-related quality of life impairment in insomnia disorder. *Sleep Med* 2013;14:493–501.