



Small molecule ONG201 inhibits HIV-1 replication in macrophages via FOXO3a and TRAIL

Runze Zhao^a, Yuju Li^{a,c}, Santhi Gorantla^a, Larisa Y. Poluektova^a, Hai Lin^{a,b}, Fengtong Gao^a, Hongyun Wang^a, Jeffrey Zhao^a, Jialin C. Zheng^{a,c,**}, Yunlong Huang^{a,c,*}

^a Department of Pharmacology & Experimental Neuroscience, University of Nebraska Medical Center, Omaha, NE, 68198, United States

^b Guang'anmen Hospital, China Academy of Chinese Medical Sciences, Beijing, China

^c Center for Translational Neurodegeneration and Regenerative Therapy, Shanghai Tenth People's Hospital Affiliated to Tongji University School of Medicine, Shanghai, China

ARTICLE INFO

Keywords:

HIV-1
Reservoir
ONG201
Macrophages
FOXO3a
TRAIL

ABSTRACT

Despite the success of antiretroviral therapy (ART), eradication of HIV-1 from brain reservoirs remains elusive. HIV-1 brain reservoirs include perivascular macrophages that are behind the blood-brain barrier and difficult to access by ART. Macrophages express transcription factor FOXO3a and the TNF superfamily cytokine TRAIL, which are known to target HIV-1-infected macrophages for viral inhibition. ONG201 is a novel and potent FOXO3a activator capable of inducing TRAIL. It can cross the blood-brain barrier, and has shown antitumor effects in clinical trials. We hypothesized that activation of FOXO3a/TRAIL by ONG201 will inhibit HIV-1 replication in macrophages. Using primary human monocyte-derived macrophages, we demonstrated that ONG201 dose-dependently decreased replication levels of both HIV-1 laboratory strain and primary strains as determined by HIV-1 reverse transcriptase activity assay. Consistent with data on HIV-1 replication, ONG201 also reduced intracellular and extracellular p24, viral RNA, and integrated HIV-1 DNA in infected macrophages. Blocking TRAIL or knockdown of FOXO3a with siRNA reversed ONG201-mediated HIV-1 suppression, suggesting that ONG201 inhibits HIV-1 through FOXO3a and TRAIL. The anti-HIV-1 effect of ONG201 was further validated *in vivo* in NOD/scid-IL-2Rgenull mice. After intracranial injection of HIV-1-infected macrophages into the basal ganglia, we treated the mice daily with ONG201 through intraperitoneal injection for six days. ONG201 significantly decreased p24 levels in both the macrophages and the brain tissues, suggesting that ONG201 suppresses HIV-1 *in vivo*. Therefore, ONG201 can be a promising drug candidate to combat persistent HIV-1 infection in the brain.

1. Introduction

HIV continues to be a major global public health issue affecting more than 1.1 million people in the USA and 36.7 million people worldwide (Deeks et al., 2017; Hall et al., 2008). Although application of antiretroviral therapy (ART) dramatically reduces the rate of HIV-1 replication in infected patients, it is not able to eradicate HIV-1 from cellular reservoirs in the lymphatic system, gut, and central nervous system (CNS). Although many types of cells can be infected by HIV-1, cells that form HIV reservoirs are typically long-lived, including memory CD4⁺ T cells (Chomont et al., 2009), brain perivascular macrophages, and microglia (Pierson et al., 2000). In the CNS,

macrophages and microglia are the main cell types that can be productively infected by HIV-1. The persistent infection of these cells often causes complications such as HIV-1 associated neurocognitive disorders (HAND). Because many ART drugs are not able to cross the blood-brain barrier (BBB) and reach therapeutic concentrations within CNS macrophages and microglia, they can produce replication-competent HIV-1, which may spread to other tissues and fuel the emergence of drug-resistant viral strains. Thus, strategies targeting these CNS reservoir cells for HIV-1 eradication are urgently needed.

Vulnerabilities of HIV-1-infected brain macrophages and microglia include the FOXO3a pathway and its transcriptional target TRAIL. FOXO3a is a conserved transcription factor downstream of the PI3K/

Abbreviations: ART, antiretroviral therapy; CNS, central nervous system; HAND, HIV-1 associated neurocognitive disorders; BBB, blood brain barrier; MDM, monocyte-derived macrophages; RTase, reverse transcriptase; ERK1/2, extracellular signal-regulated kinase 1/2

* Corresponding author. Department of Pharmacology & Experimental Neuroscience, University of Nebraska Medical Center, Omaha, NE, 68198-5930, USA.

** Corresponding author. Shanghai Tenth People's Hospital affiliated to Tongji University School of Medicine, Shanghai, China.

E-mail addresses: jialinzheng@tongji.edu.cn (J.C. Zheng), yhuan1@unmc.edu (Y. Huang).

<https://doi.org/10.1016/j.antiviral.2019.05.015>

Received 30 November 2018; Received in revised form 29 May 2019; Accepted 30 May 2019

Available online 31 May 2019

0166-3542/ © 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Akt-1 pathway (Kaestner et al., 2000; Kaufmann and Knochel, 1996). In the absence of environmental signals or growth factors, FOXO3a is in an unphosphorylated state and remains transcriptionally active. Growth factors activate the PI3K/Akt-1 signaling pathway, phosphorylating FOXO3a and abrogating its transcriptional activity (Brunet et al., 2001). FOXO3a is predominantly expressed in peripheral lymphoid organs but the specific immunomodulatory role of FOXO3a remains largely undefined (Lin et al., 2004). In memory CD4⁺ T cells under HIV-1 infection, multiple signals activate I κ B kinase, which phosphorylates FOXO3a, decreasing its ability to induce expression of various pro-apoptotic genes (van Grevenynghe et al., 2008). Our previous publications have demonstrated that FOXO3a targets HIV-1-infected macrophages for apoptosis (Cui et al., 2008). Furthermore, we have determined that FOXO3a is phosphorylated (inactive) during neuroinflammation and expression of FOXO3a inhibits reactive astrogliosis in a neuroinflammatory mouse model (Cui et al., 2011). Intriguingly, the transcriptional targets of FOXO3a include TRAIL, which is a member of the TNF superfamily and is also known as TNF Superfamily Member 10 (Pitti et al., 1996; Wiley et al., 1995). TRAIL interacts with at least five unique receptors. TRAIL receptor 1 (R1) and receptor 2 (R2) have death domains and induce cellular apoptosis following ligand binding (Pan et al., 1997; Seol et al., 2001; Walczak et al., 1997). TRAIL-R3, TRAIL-R4, and the fifth receptor named osteoprotegerin do not possess death domains and instead act as decoy receptors (Degli-Esposti et al., 1997; Emery et al., 1998; Pan et al., 1998). Originally thought to target only tumor cells (Pitti et al., 1996; Wiley et al., 1995), TRAIL has been shown to reduce viral burden in HIV-1-infected macrophage and resting memory CD4⁺ T cells (Huang et al., 2006; Lum et al., 2001b). While its receptors are expressed in macrophage and microglia, TRAIL is expressed in much lower levels in the CNS compared with the lymphoid tissues (Dorr et al., 2002; Huang et al., 2009; Peng et al., 2005; Ryan et al., 2004). The deficiency of FOXO3a signaling during HIV-1 infection and lack of TRAIL expression in the CNS may inadvertently facilitate the forming of HIV-1 brain reservoirs. Therefore, targeting the FOXO3a-TRAIL pathway is a novel strategy to combat HIV reservoirs within the CNS.

Targeting the FOXO3a-TRAIL pathway has been difficult due to a lack of drug candidates. However, recent drug development provides ONC201 as a potent and stable small molecule that can activate FOXO3a and transcriptionally induce TRAIL expression (Allen et al., 2013, 2015; Kline et al., 2016). ONC201 is orally active and can cross the BBB (Allen et al., 2013). It has shown an efficacious antitumor effect and is currently being tested in clinical trials as a promising anticancer agent (Allen et al., 2016). In this study, we have investigated the anti-HIV-1 activity of ONC201. We demonstrate that ONC201 inhibits the HIV-1 infection both *in vitro* in monocyte-derived macrophages and *in vivo* in NOD/scid-IL-2R γ_c ^{null} (NSG) mouse brains engrafted with HIV-1-infected macrophages. This newly found activity of ONC201 suggests that it can serve as a drug candidate to combat persistent HIV-1 infection in the CNS.

2. Materials and methods

2.1. Ethics statement and mice

Primary monocyte-derived macrophages (MDM) and peripheral blood lymphocytes (PBL) were used in full compliance with the University of Nebraska Medical Center and National Institutes of Health ethical guidelines, with the Institutional Review Board (IRB) #: 162-93-FB. For animal experiments, all mice were housed in the Comparative Medicine Animal Facilities at the University of Nebraska Medical Center. All procedures were conducted in accordance with the protocol (15-040-06) approved by the Institutional Animal Care and Use Committee at the University of Nebraska Medical Center. Fourteen four-week old male NOD.Cg-Prkdc^{scid}IL2rg^{tm1Wjl}/SzJ mice were purchased from Jackson Laboratories and HIV-1-infected human MDM

(5×10^5 cells, 3 μ l) was injected intracranially (IC) into the basal ganglion of the mice. The coordinates for inoculation is: 0.5 mm posterior to bregma, 3.5 mm lateral and a depth of 3.6 mm ONC201 (MedKoo Bioscience, Morrisville, NC) was administered daily through intraperitoneal (IP) injection at 50 mg/kg. Animals were sacrificed seven days after the IC injection.

2.2. Isolation and culture of primary MDM and PBL

Human monocytes and PBL were isolated by counter-current centrifugal elutriation from peripheral blood mononuclear cells obtained through leukopheresis from healthy donors (Gendelman et al., 1988). Primary monocytes were cultured as adherent monolayers at a density of 1.1×10^6 cells/well in 24-well plates or 0.25×10^6 cells/well in 96-well plates and differentiated for seven days in differentiation medium consisting of Dulbecco's Modified Eagle's Medium (DMEM) (Sigma Chemical Co., St. Louis, MO) supplemented with 10% human serum, 50 μ g/ml gentamicin (GIBCO Invitrogen Corp.), 10 μ g/ml ciprofloxacin (Thermo Fisher Scientific, Waltham, MA), and 1000 U/ml recombinant human macrophage-colony stimulating factor (M-CSF). After seven days in differentiation, the cells became MDM and M-CSF was removed since the culture can produce M-CSF through autocrine secretion. PBL were stimulated in phorbol 12-myristate 13-acetate (PMA, 20 ng/ml, Sigma #p8139) and ionomycin (1 mM, Sigma #10634) for 5 h in a T75 flask before being reseeded on 96-well-plates and cultured in RPMI-1640 medium (Gibco, Gaithersburg, MD) supplemented with 10% fetal bovine serum, 100 U/ml penicillin, 100 μ g/ml streptomycin, 2 mM L-glutamine, and 100 U/ml IL-2 (R&D Systems, Minneapolis, MN).

2.3. HIV-1 infection of MDM and ONC201 treatment

MDM were infected with HIV-1_{ADA}, a macrophage-tropic strain of HIV-1, at a multiplicity of infection (MOI) of 0.1 virus/target cell (Gendelman et al., 1988). Briefly, viral stocks were diluted in MDM medium (same with the differentiation medium but without M-CSF) for overnight incubation with MDM. On the second day, the medium was removed and substituted with MDM medium and either DMSO (Sigma), ONC201, or ONC201 isomer (MedKoo Bioscience). For selected experiments, a recombinant human TRAIL R2/Fc Chimera Protein (R&D) was added to the cultures along with ONC201. The cultures were re-treated with the same drugs in fresh MDM medium every two days. At four days after infection, the cultures were treated with the same drugs in 0.5 ml/well fresh serum-free Neurobasal™ medium (Thermo Fisher Scientific) for 24 h before all samples were collected.

2.4. Assessment of cell viability

MDM, glioblastoma (ATCC, CRL-1620), and PBL viability was determined by a colorimetric CellTiter 96[®] Aqueous One Solution Assay (Promega, Madison, WI) based on the manufacturer's instruction. Assays were performed by adding 20 μ l of the CellTiter 96[®] Aqueous One Solution Reagent, which contained a tetrazolium compound[3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfo-phenyl)-2H-tetrazolium, inner salt; MTS] and an electron coupling reagent (phenazine ethosulfate; PES), directly to each of 100 μ l MDM medium in 96-well plates. After 2-h incubation at 37 °C in a humidified 5% CO₂ atmosphere, the absorbance of each well was recorded at 490 nm using a 96-well plate reader (BioTek, Winooski, VT).

2.5. HIV-1 reverse transcriptase (RTase) activity assay

HIV-1 replication levels were determined by HIV-1 RTase activity assay as described previously (Gendelman et al., 1988). RT activity was determined in triplicate samples of cell culture fluids. For this assay, 10 μ l of supernatant was incubated in a reaction mixture of 0.05% Nonidet P-40, 10 μ g of poly(A)/ml, 0.25 μ g of oligo(dT)/ml, 5 mM DTT,

150 mM KCl, 15 mM MgCl₂, and [3H]TTP in Tris-HCl buffer (pH 7.9) for 24-h at 37 °C. Radiolabeled nucleotides were precipitated with ice-cold 10% trichloroacetic acid on UniFilter-96 GF/C (FilterMate Harveser, Waltham, MA) in an automatic cell harvester (FilterMate Harveser, Waltham, MA) and washed with 95% ethanol. Radioactivity was determined by liquid scintillation spectroscopy.

2.6. Western blots

MDM or tissues of injection sites were homogenized with a PowerGen Homogenizer (Thermo Fisher Scientific) and then lysed by M-PER™ Mammalian Protein Extraction Reagent (Thermo Fisher Scientific, Waltham, MA). Protein concentrations were determined by Bradford protein assay (Bio-Rad, Hercules, CA). Cell lysates were subjected to SDS-PAGE to separate proteins and electrophoretically transferred to Polyvinylidene fluoride membranes (Bio-Rad, Hercules, CA). Membranes were incubated overnight at 4 °C with polyclonal antibodies for FOXO3a (Cat#, 75D8, Cell Signaling Technology, Danvers, MA), phosphorylated FOXO3a (Cat#, 9466s, Cell Signaling Technology, Danvers, MA), HIV-1 p24 (Ref#, M0857, DAKO Corp, Carpinteria, CA), human CD68 (Ref#, M0876, DAKO Corp, Carpinteria, CA), caspase-3 (Cat#, 9662, Cell Signaling Technology), and β-actin (Sigma-Aldrich, St. Louis, MO), followed by horseradish peroxidase-linked secondary anti-rabbit or anti-mouse secondary antibodies (Cell signaling Technology, Danvers, MA). Antigen-antibody complexes were visualized by Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific). For quantification of the data, films were scanned with a CanonScan 9950 F scanner (Canon Inc., Tokyo, Japan) and images were analyzed using the public domain NIH Image J program (developed at the U.S. National Institutes of Health and available on the internet at <http://rsb.info.nih.gov/ni-image/>).

2.7. RNA and DNA extraction and real time RT-PCR analysis of HIV-1 gag

Total RNA was isolated with TRIzol (Thermo Fisher Scientific) and RNeasy Mini Kit according to the manufacturer's protocol (Qiagen Inc., Valencia, CA). Following RNA extraction, DNA was isolated from the remaining organic phase using the DNeasy Blood and Animal Spin-Column kit according to manufacturer's instructions (Qiagen). Assays-on-demand primers for TRAIL (ID#, Hs00921974_m1), FOXO3a (ID#, Hs00818121_m1), 18S rRNA (Ref#, 4352930#) and GAPDH (ID#, Hs01922876_u1) were purchased from Thermo Fisher Scientific. For quantification of HIV-1 gag, primers and probe were purchased from Thermo Fisher Scientific with the following sequences: forward, 5' ACA TCA AGC CAT GCA AAT-3'; reverse, 5'-ATC TGG CCT GGT GCA ATA GG-3'; probe, 5'-FAM-CAT CAA TGA GGA AGC TGC AGA ATG GGA TAG A-TAMRA-3'. Real-time reverse-transcription polymerase chain reaction (RT-PCR) was performed in a volume of 10 µl using the one-step quantitative TaqMan Real-time RT-PCR system (Applied Biosystems Inc.). For analysis of HIV-1 gag DNA, real-time PCR was carried out directly on the DNA samples using the TaqMan Universal Master Mix and a StepOne Plus Real-Time PCR System (Thermo Fisher Scientific). Relative FOXO3a, TRAIL, CD4, CCR5 and HIV-1 gag levels were determined and standardized with GAPDH or 18S rRNA internal control using comparative ΔΔCT method. All primers used in the study were tested for amplification efficiencies and the results were similar.

2.8. Analysis of HIV-1 proviral DNA integration

Two-step nested PCR assays were used for quantitative HIV-1 DNA integration analysis as previously described (Friedrich et al., 2010; Iordanskiy et al., 2010). The first round was performed in a 25 µl reaction mix to pre-amplify the genomic DNA from infected macrophages. The following primers were used: Alu forward, 5'-GCC TCC CAA AGT GCT GGG ATT ACA G-3'; and HIV-1 gag reverse, 5'-GCT CTC GCA CCC ATC TCT CTC C-3'. The PCR reaction contained 10x EX Taq

Buffer, dNTP mix (2.5 mM), EX Taq (all from TaKaRa Bio Inc., Otsu, Shiga, Japan), ddH₂O, 100 nM Alu forward primer, 600 nM gag reverse primer, and 100 ng of genomic DNA. The DNA Engine Peltier Thermal Cycler (Bio Rad, Hercules, CA) was programmed to perform a 2-min hot start at 94 °C, followed by 30 steps of denaturation at 93 °C for 30 s, annealing at 50 °C for 1 min, and extension at 70 °C for 2 min. The second round real-time quantitative PCR was performed by using 5.6 µl of the reaction mix from the pre-amplification step. The amplified DNA was standardized with an internal GAPDH control. The sequences of the primers and probe were as follows: LTR forward, 5'-GCC TCA ATA AAG CTT GCC TTG A-3'; LTR reverse, 5' -TCC ACA CTG ACT AAA AGG GTC TGA-3'; probe: 5'-FAM-GCG AGT GCC CGT CTG TTG TGT GAC TCT GGT AAC TAG CTC GC-BHQ-3'.

2.9. ELISA

Supernatants were collected and protein levels of HIV-1 p24, TRAIL, RANTES (CCL5), MIP-1α (CCL3), and IP-10 (CXCL10) were analyzed by in-house ELISAs (paired antibodies, R&D Systems, Minneapolis, MN) as previously described (Erichsen et al., 2003) and per manufacturer's instructions.

2.10. siRNA transfection

Predesigned siRNA duplexes targeting FOXO3a (ID: 144672) and a negative control siRNA (catalog no. AM4611) were purchased from Thermo Fisher Scientific. At 72 h after HIV-1 infection, cells were transfected with 100 nM siRNA duplexes for 4 h in DMEM culture medium without serum, in the presence of siIMPORTER reagent (EMD Millipore Corporation, Billerica, MA) according to the manufacturer's instructions. At 72 h post-transfection, cells were harvested and total RNA was extracted for the detection of FOXO3a and TRAIL levels by real-time RT-PCR.

2.11. Free-floating immunohistochemistry and image analyses

Animals were euthanized under deep anesthesia and perfused with phosphate-buffered saline (PBS) and then with 4% paraformaldehyde (PFA) in PBS. The brains were removed and immersed in freshly depolymerized 4% PFA in PBS for 48 h and then cryoprotected by 30% sucrose for 48 h. The fixed, cryoprotected brains were frozen and sectioned in the coronal plane at 30 µm using a Cryostat (Leica Microsystems Inc., Bannockburn, IL), with sections collected serially in PBS as previously described (Zhu et al., 2012). Brain sections were then incubated overnight at 4 °C with primary antibodies, followed by secondary antibodies (Molecular Probes, Eugene, OR, 1:1000) for 1 h at 25 °C. Primary antibodies included P24 (DAKO, 1:1000) and human CD68 (DAKO, 1: 1000). All antibodies were diluted in 5% goat serum in PBS. Cells were counterstained with DAPI (Sigma-Aldrich, 1:2000) to identify the nuclei. Images were taken using a Zeiss Meta 710 confocal microscope (Carl Zeiss MicroImaging, LLC) (20 × object, tile scan 4 × 4 mode). Section images from eight mice were imported into Image-ProPlus, version 7.0 (Media Cybernetics, Silver Spring, MD) for quantification of p24/DAPI and CD68/DAPI double positive staining. The assessors were blinded during image acquisition and quantification.

2.12. Statistical analysis

Data are expressed as means ± SD unless otherwise specified. Statistical analysis was performed by using one-way ANOVA, followed by the Tukey's-post-test for paired observations, or two-way ANOVA when two independent variables are considered. The two-tailed Student's *t*-test was used to compare two groups. Significance was determined by a *p* value < 0.05. All experiments were performed with cells from at least three donors to account for any donor-specific

differences. Assays were performed at least three times in triplicate or quadruplicate within each assay.

3. Results

3.1. ONC201 inhibits HIV-1 infection of macrophages *in vitro*

Because ONC201 was originally identified as a TRAIL-inducing anti-tumor agent (Allen et al., 2013), we first sought to confirm its activity in tumor cells. We treated A-172 human glioblastoma cells with doses of ONC201 or its isomer ranging from 0.03 to 30 μM for 72-h and then determined the cell viability through an MTS assay (Supplemental Fig. 1A). Treatment with ONC201, but not ONC201 isomer, induced dose-dependent cytotoxicity in glioblastoma cells, confirming the bioactivity of ONC201. Next, we performed cytotoxicity evaluation of ONC201 on relevant HIV-1 susceptible cells, including primary MDM and PBL. We treated MDM and PBL with either ONC201 or its isomer for five days at concentrations ranging from 0.03 to 30 μM . Treatment with either ONC201 or its isomer did not significantly change cell viability of MDM or PBL at all concentrations tested, as measured by MTS assays, compared with those in the vehicle control DMSO group (0 μM uninfected MDM, Fig. 1A and Supplemental Fig. 1B). Therefore, ONC201 appears to have specific cytotoxicity to tumor cells but not primary human MDM and PBL. In MDM infected with laboratory strain HIV-1_{ADA}, ONC201 treatment significantly increased the cell viability, compared to isomer treatment of the same dosage (Fig. 1A), which is possibly due to the significant decrease of HIV-1 replication at higher doses (10–30 μM) of ONC201 treatment (Fig. 1B). The HIV-1 RTase activity assay determines the levels of reverse transcriptase activity in cultural supernatants and is a semi-quantitative measurement of HIV-1 replication levels (Gorantla et al., 2005). Therefore, ONC201 is safe to uninfected human MDM, even at high concentrations, and it has specific antiviral activity against HIV-1 in human MDM.

To confirm the antiviral effect of ONC201 on different strains of HIV-1 infection, we infected MDM with two primary HIV-1 Clade B strains (G0048CPX and 2562BG) and one primary HIV Clade C strain (2873MVC), which we characterized previously (Constantino et al., 2011). These HIV strains displayed different replication kinetics and the infection peaked at different times after viral inoculation (Fig. 2). However, for all the strains, ONC201 treatment reduced the viral replication in a dose-dependent manner, which is consistent with the results in HIV-1_{ADA}. Before the peak infection, ONC201 treatment achieved higher degree of viral inhibition compared with those in earlier time points (Fig. 2), suggesting that extending the treatment time has a positive effect on viral inhibition.

To further characterize the antiviral effect of ONC201 in MDM, we determined both intracellular and extracellular levels of HIV-1 capsid protein p24 through Western blots and ELISA, respectively. Treatment with ONC201 significantly decreased p24 levels in the infected MDM (Fig. 3A–C). Consistent with the p24 data, treatment with ONC201 significantly decreased the levels of HIV-1 gag RNA and DNA when compared with those in the isomer or vehicle control DMSO groups (Fig. 3D and E). Furthermore, we determined the HIV-1 integration through a two-step Alu-based nested PCR method and found that treatment with ONC201 significantly decreased the level of integrated HIV-1 DNA in MDM when compared with those in isomer or DMSO groups (Fig. 3F). Together, ONC201 appears to reduce multiple viral intermediates and products in HIV-1 lifecycle, suggesting it is an effective agent to limit HIV-1 replication in human macrophages.

The HIV-1 replication in MDM is productive and cytopathic. To determine whether ONC201 is effective in latently HIV-1 infected cells, we used the pro-monocytic U1 cells (NIH AIDS Reagent Program), which carries latently infected HIV-1 genome that can be stimulated to produce virus (Folks et al., 1987). We treated U1 cells with ONC201 or its isomer in the presence of TNF, a cytokine known to up-regulate viral expression in the cell type, and measured the viral production by RTase




Fig. 1. ONC201 has an antiviral effect on macrophages. A) Human MDM were plated in 96-well plates and infected with HIV-1 for 24 h before incubation with doses of ONC201 or its isomer ranging from 0.03 to 30 μM . Uninfected MDM were treated with the same ONC201 and isomer dosages in parallel for comparison. At 5-day post ONC201 and isomer treatments, cell viability was determined by the MTS assays. Data were analyzed by two-way ANOVA, and results shown are the means \pm SD of 3 experiments. B) At the same experimental end point as the MTS assays, supernatants were removed and the replication levels of HIV-1 were monitored by RTase activity assay. Results shown are from representative experiments performed with three different donors. Data were analyzed by two-way ANOVA: ** denotes $p < 0.01$, *** denotes $p < 0.001$, compared to the HIV-1+ONC201 isomer group of the same treatment dosage.

activity assay and p24 ELISA. In both assays, ONC201 treatment decreased the level of HIV-1 virus in the culture supernatant in a dose-dependent manner (Supplemental Figs. 2A and B). Notably, the inhibitory effect of ONC201 was seen at concentrations as low as 0.3 μM , suggesting a higher potency of ONC201 against HIV-1 induction from latency compared with the viral inhibition in primary macrophages.

3.2. ONC201 has an anti-HIV-1 effect on macrophages *in vivo*

To further determine whether ONC201 has an anti-HIV-1 effect *in vivo*, we used an established HIV-1 infected NOD/SCID/IL2R γ^{null} (NSG) model (Chen et al., 2015; Wu et al., 2009; Ye et al., 2013). HIV-1 infected MDM were intracranially injected into basal ganglion of the NSG immunocompromised mice. At 24-h post MDM engraftment, the mice




Fig. 2. ONC201 has an antiviral effect against different primary HIV strains. A-C) Human MDM were plated in 96-well plates and infected with different primary HIV strains, including 2873MVC (HIV-1 Clade C strain, A), G0048CPX (HIV Clade B strain, B) and 2562BG (HIV Clade B strain, C) for 24 h before incubation with doses of ONC201 or its isomer ranging from 3 to 30 μM. At 2-, 6-, 10-, and 14-day post ONC201 and isomer treatments, supernatants were collected and the replication levels of HIV-1 were monitored by RTase activity assay. Values represent SEM of biological replicates. Data were analyzed by two-way ANOVA: ** denotes $p < 0.01$, *** denotes $p < 0.001$, compared to the HIV-1 + ONC201 isomer group of the same treatment dosage.

were administrated daily with either ONC201 (50 mg/kg) or vehicle control via IP for six days. At 7-day post MDM engraftment, the mice were sacrificed for HIV-1 detection (Fig. 4A). To visualize the HIV-1-infected cells in basal ganglion, we first performed immunohistochemistry in serial para-coronal brain slices encompassing the injection track with DAPI (Fig. 4B and E) and p24 (Fig. 4C and F). Both DAPI and p24 clearly marked the injection tracks and their signals were in close proximity to each other in the injection tracks (Fig. 4D and G). To confirm that the p24-positive cells are CD68-positive macrophages, we performed CD68 staining. Because CD68 and p24 antibodies are from the same species (mouse) and alternative antibodies from other species worked poorly in the co-immunostaining (data not shown), we individually labeled the injection site with either CD68 or p24 along with DAPI. CD68 staining revealed a substantial number of macrophages remained in the injection track in both DMSO and

ONC201 treatment groups (Fig. 4H and I). We quantified the p24-positive and CD68-positive cells in consecutive slides through a previously established stereological method (Tian et al., 2015; Wang et al., 2017). As expected, we observed that ONC201 administration significantly decreased p24-positive cells as a percentage to total CD68-positive cells (Fig. 4J), compared with those in the vehicle control group. These data suggest that ONC201 reduces the number of HIV-1-infected macrophages in the *in vivo* NSG mouse model. In order to better quantitatively determine the levels of HIV-1 inhibition, we homogenized the tissues that contained injection sites and subjected the tissue lysates to Western blots (Fig. 4K). The quantification data revealed that p24 levels significantly decreased in the ONC201 group compared with the control group (Fig. 4L). In contrast, CD68 levels were comparable in all of the brain lysates. As a result, the ratio of p24 to CD68 significantly decreased after ONC201 treatment (Fig. 4M), which is consistent with the p24 and CD68 staining data. Together, these data in HIV-1-infected NSG mice demonstrate that ONC201 is able to inhibit HIV-1 replication in macrophages *in vivo*.

3.3. ONC201 activates FOXO3a and induces TRAIL expression in HIV-1 infected macrophages

After showing the antiviral effect of ONC201 *in vitro* and *in vivo*, we investigated the underlying mechanism of viral inhibition by focusing on the FOXO3a-TRAIL pathway. There are two forms of FOXO3a: the phosphorylated form is inactive and kept in the cytoplasm; once dephosphorylated, it moves to the nucleus and becomes active, inducing pro-apoptotic genes. To assess the status of FOXO3a in HIV-1 infected macrophages after ONC201 treatment, we measured the levels of total FOXO3a, phosphorylated FOXO3a, and β-actin by Western blots. We first confirmed that HIV-1 p24 was indeed reduced by the ONC201 treatment (Fig. 5A). Analysis of FOXO3a phosphorylation revealed that the phosphorylation of FOXO3a, as determined by the ratio of phosphorylated FOXO3a over total FOXO3a, was decreased in a dose-dependent manner after ONC201 treatment compared to the vehicle control DMSO group (Fig. 5A). In contrast, treatment with ONC201 isomer did not significantly change the ratio of phosphorylated FOXO3a to total FOXO3a in HIV-1-infected macrophages. This result indicates that FOXO3a is indeed dephosphorylated and activated by ONC201 in MDM. Next, we determined whether ONC201 treatment increases TRAIL expression by measuring soluble TRAIL secretion in the culture supernatants. We found that ONC201 treatment increased TRAIL levels in a dose-dependent manner (Fig. 5B). In contrast, treatment with ONC201 isomer did not significantly change the TRAIL levels in HIV-1-infected macrophages. Because HIV-1 infection is known to decrease FOXO3a phosphorylation and increase TRAIL production in macrophages (Cui et al., 2008; Huang et al., 2009), the changes of FOXO3a phosphorylation and TRAIL production after ONC201 treatment could be due to a change in HIV-1 infection levels. To determine the direct effect of ONC201 on FOXO3a phosphorylation and TRAIL production, we treated uninfected MDM with ONC201 and found that ONC201 decreased FOXO3a phosphorylation (Supplemental Fig. 3A) and increased soluble TRAIL levels (Supplemental Fig. 3B) similar to the data in HIV-1-infected macrophages (Fig. 5). Together, these data confirm that ONC201 activates FOXO3a and induces the downstream expression of TRAIL in HIV-1 infected macrophages.

3.4. The antiviral effect of ONC201 in HIV-1 infected macrophage is dependent on FOXO3a and TRAIL expressions

Because downstream targets of ONC201 include TRAIL, which belongs to the TNF superfamily of cytokines, one plausible mechanism for HIV-1 inhibition by ONC201 is through ONC201-induction of cytokines that interfere with the HIV-1 lifecycle. RANTES (CCL5) and MIP-1α (CCL3) are both chemokines that bind to CCR5, which function as a primary co-receptor for HIV-1 entry in MDM (Cocchi et al., 1995,




Fig. 3. ONC201 inhibits HIV-1 p24, gag RNA, DNA, and integrated LTR DNA in macrophages. Human MDM were plated in 24-well plates in triplicate and then infected with HIV-1_{ADA} for 24 h before incubation with ONC201 or ONC201 isomer at 30 μ M for 5 days. A, B) Cell lysates were collected and subjected to SDS-PAGE and immunoblotting for HIV-1 p24. Actin was used as the loading control. Densitometric quantifications of p24 in MDM were presented as a ratio to actin and normalized as fold changes to the vehicle control DMSO group. C) Supernatants were collected and subjected to ELISA for HIV-1 p24. Values represent means \pm SEM of three biological replicates. D-F) RNA (D) and DNA (E, F) were isolated from the samples. HIV-1 gag was detected through quantitative real time RT-PCR and real time PCR, respectively. Relative HIV-1 gag and LTR DNA levels were determined and standardized with 18S rRNA or GAPDH internal control. Values represent means \pm SEM of three biological replicates. ANOVA analysis: * denotes $p < 0.05$ compared to the vehicle control DMSO group.

2000). IP-10 (CXCL10) plays a role in chemotaxis of immune cells. We tested the CCL5, CCL3, and CXCL10 levels after ONC201 treatment and found that ONC201 did not significantly increase the levels of these chemokines in the supernatants (Supplemental Fig. 4), suggesting that ONC201 is unlikely to inhibit HIV-1 replication through inducing HIV-1 suppressive chemokines in MDM. To determine if the antiviral effect of ONC201 is mediated by FOXO3a, knockdown of FOXO3a was performed by siRNA targeting FOXO3a in HIV-1-infected MDM. A non-targeting control siRNA was used as the control. We assayed FOXO3a and TRAIL expression at 72-h after siRNA transfection and found that FOXO3a-targeting siRNA reduced the mRNA levels of FOXO3a by 83% (Fig. 6A) and protein levels of FOXO3a by 65% (Supplemental Figs. 5A and B). Similarly, TRAIL expression was downregulated by 85% in HIV-1 infected macrophages (Fig. 6B), suggesting that the siRNA is effective in reducing FOXO3a and its downstream gene transcription. ONC201 significantly decreased the HIV-1 replication levels, as measured by HIV-1 RTase activity assay, compared with those in the isomer treatment group (Fig. 6C). However, FOXO3a knockdown reversed HIV-1 replication to the same level as those in the isomer treated group (Fig. 6C). Consistent with the RTase data, FOXO3a knockdown increased the intracellular p24 level in the ONC201-treated HIV-1 infected MDM (Supplemental Fig. 5C). Together, these data suggest that FOXO3a is required for the antiviral effects of ONC201.

To further determine if TRAIL mediates the antiviral effect of ONC201, we used 50 ng/ml or 100 ng/ml of the soluble TRAIL receptor to act as decoys blocking the activities of both soluble and membrane-associated TRAIL in macrophages. This soluble TRAIL receptor is effective in blocking TRAIL-mediated HIV-1 inhibition in macrophages (Supplemental Fig. 6). In ONC201-treated MDM, we found that

treatment of soluble TRAIL receptors reversed the HIV-1 inhibition to the same level as those in the isomer-treated group (Fig. 7A). Consistent with the RTase activity data, treatment with soluble TRAIL receptors also significantly increased intracellular HIV-1 p24 levels (Fig. 7B), suggesting that the antiviral effect of ONC201 in HIV-1 infected macrophages is dependent on TRAIL expressions.

3.5. ONC201 pretreatment increases its efficacies against HIV-1 infection in macrophages

In cancer cells, activation of the FOXO3a-TRAIL pathway by ONC201 leads to cell death and apoptosis, while in HIV-1 infected MDM, treatment of human recombinant TRAIL induces cell death and apoptosis (Allen et al., 2013; Huang et al., 2006). In our data, ONC201 treatment in HIV-1-infected MDM did not significantly impact MDM viability (Fig. 1A). This is likely due to the unique action of ONC201, which reportedly requires at least 72 h of downstream signaling to induce TRAIL production (Allen et al., 2013). The lagged production of TRAIL by ONC201 during viral propagation in MDM likely undermined the effectiveness of antiviral activities by ONC201. To ensure full TRAIL production, we pretreated MDM with ONC201 for five days before the HIV-1 infection. ONC201 and isomer treatments were continuous both during and after HIV-1 infection. At three-day post infection, we found that ONC201 pretreatment reduced cell viability in a dose-dependent manner (Fig. 8A). Furthermore, cell death of HIV-1 infected macrophages was accompanied by significant increase of caspase cleavage (Fig. 8B and C), indicating that ONC201 pretreatment induces apoptosis in HIV-1 infected macrophages. As expected, analysis of HIV-1 replication by RTase activity assay showed that the ONC201 pretreatment




Fig. 4. ONC201 reduces HIV-1 infection levels in NSG mouse brains engrafted with infected human macrophages. HIV-1-infected human macrophages were intracranially injected into the basal ganglia of NOD/scid-IL-2R γ null (NSG) mice. Mice were administrated daily with ONC201 (50 mg/kg) or solvent control DMSO through intraperitoneal injections. A) Timeline of experimental procedures and sample collection in the HIV-1-infected NSG mouse model. B–J) Immunostaining of p24 and CD68 in the injection sites of mouse brains: co-immunostaining of p24 and DAPI (B–G); co-immunostaining of CD68 and DAPI (H–I); quantification of the p24/CD68 ratio per area (J). K–M) Brain tissues that contained the injection sites were homogenized for detection of HIV-1 p24 and human CD68 in Western blots. Data were evaluated statistically by an unpaired Student's t-test, $n = 6$ per treatment group.

dose-dependently lowered viral levels in the cell culture (Fig. 8D). At 30 μ M, ONC201 pretreatment dramatically reduced HIV-1 replication by 96%, a much higher efficacy compared with those in ONC201 treatment post infection. Taken together, these data demonstrate that ONC201 pretreatment increases its efficacies against HIV-1 infection in macrophages and the pretreatment is associated with significant apoptosis in infected macrophages.

4. Discussion

Targeting long-term HIV persistence with novel approaches is essential to combat AIDS in the post-cART era (Carpenter et al., 2000; Krebs et al., 2000; McArthur et al., 1999). In the present study, a novel small molecule ONC201 was tested for its potential antiviral effect on the HIV-infected macrophages. We demonstrated that ONC201 has





Fig. 5. ONC201 activates FOXO3a and induces TRAIL expression in HIV-1 infected macrophages. Human MDM were plated in 24-well plates and then infected with HIV-1_{ADA} for 24 h before incubation with ONC201 or ONC201 isomer at 3, 10, or 30 μM for five days. A) Cell lysates were collected and subsequently subjected to SDS-PAGE and immunoblotting for the detection of phospho-FOXO3a, total FOXO3a, and p24. Actin was used as the loading control. Densitometric quantifications of phospho-FOXO3a in MDM were presented as a ratio to total FOXO3a and normalized as fold changes to the vehicle control DMSO group. B) Soluble TRAIL concentrations after ONC201 or ONC201 isomer treatment were determined from cell culture supernatants using ELISA. Data were normalized with total cellular protein concentrations. Results shown are from representative experiments performed with three different donors. ANOVA analysis: * denotes p < 0.05, ** denotes p < 0.01, compared to the vehicle control DMSO group.

little toxicity in uninfected human macrophages and exert an anti-HIV-1 effect on infected macrophages *in vitro*. In a murine HIV Encephalitis model, ONC201 treatment significantly reduced HIV-1 levels in engrafted macrophages, indicating the antiviral effect of ONC201 in the brain reservoir cell type *in vivo*. We further identified that the antiviral effect of ONC201 is mediated by the FOXO3a/TRAIL pathway, as ONC201 activated FOXO3a and induced TRAIL expression in HIV-1-infected macrophages, and knockdown of FOXO3a or blocking TRAIL abrogated the antiviral activity of ONC201.

Although variants of recombinant TRAIL and the longer-lived TRAIL receptor agonist antibodies have been developed and applied in clinical trials (Camidge et al., 2010; Greco et al., 2008; Leong et al., 2009; Mom et al., 2009; Plummer et al., 2007; Tolcher et al., 2007; Trarbach et al., 2010), those protein-based therapeutics are costly and therapeutic



(caption on next page)

concentrations remain difficult to maintain. Therefore, increasing endogenous TRAIL protein production via pharmacological means is a more practical approach. ONC201 was originally screened from the National Cancer Institute (NCI) Diversity Set II as one of the small molecules capable of up-regulating endogenous TRAIL gene transcription (Allen et al., 2015). There are other potent TRAIL-inducing compounds such as TIC9 reported from the screening. However, ONC201

Fig. 6. Antiviral effect of ONC201 is dependent upon FOXO3a in HIV-1-infected macrophages. Human MDM were plated in 24-well plates and then infected with HIV-1_{ADA} for 24 h before incubation with ONC201 or ONC201 isomer at 30 μ M for five days. At two days after HIV-1 infection, MDM were transfected with either non-targeting control siRNA or FOXO3a siRNA. A-B) At the experimental end point, FOXO3a and TRAIL mRNA were detected in total cellular RNA through real time RT-PCR. Data were normalized to 18S rRNA and presented as fold change compared to control siRNA group with isomer treatment. C) HIV-1 infection levels were determined by the RTase activity assay. Values represent means \pm SEM of three biological replicates. ANOVA analysis: *** denotes $p < 0.001$, compared to the control siRNA with isomer treatment group; # denotes $p < 0.05$, ### denotes $p < 0.001$, compared to the control siRNA with ONC201 treatment group.




Fig. 7. Antiviral effect of ONC201 in HIV-1 infected macrophages is dependent on TRAIL expression. HIV-1-infected MDM were incubated with ONC201 or ONC201 isomer at 30 μ M, along with 100 ng/ml, 50 ng/ml soluble TRAIL receptor, or 100 ng/ml soluble TNF receptor as a control for 5 days. A) HIV-1 infection levels were determined by RTase activity assay. B) Cell lysates were subjected to SDS-PAGE and immunoblotting for HIV-1 p24. Actin was used as the loading control. Densitometric quantifications of p24 in MDM were presented as a ratio to actin and normalized as fold changes to the control group. ANOVA analysis: *** denotes $p < 0.001$, compared to the control group with isomer treatment; # denotes $p < 0.05$, ## denotes $p < 0.01$, compared to the control group with ONC201 treatment.

seems to uniquely potentiates tumor cell death while sparing normal cells and lacks genotoxicity in normal fibroblasts, which could be the result of a relatively milder induction of death receptor 5 compared with those induced by TIC9 (Allen et al., 2015). This characteristic of ONC201 allows it to specifically target HIV-1 infected macrophages and

be further considered for clinical applications.

ONC201 has been well-known in cancer research for its capacity to induce sustained TRAIL upregulation and apoptosis in tumor cells both *in vitro* and *in vivo* (Allen et al., 2013, 2015). Our study for the first time demonstrates that ONC201 inhibits HIV-1 replication in macrophages both *in vitro* and *in vivo*. The evidence for viral inhibition after ONC201 treatment is multifold. First, the HIV-1 RTase activity assay is a relatively accurate measurement of HIV-1 virion production in the supernatants (Gorantla et al., 2005). Notably, ONC201 decreases HIV-1 RTase activities in MDM through FOXO3a and TRAIL but likely not through a direct inhibition of HIV-1 RTase activities since ONC201 is not able to suppress HIV-1 when directly incubated with HIV-1 virions (Supplemental Fig. 7). Second, two-step Alu-based nested PCR determines the level of HIV-1 integration by measuring HIV-1 LTR DNA as the proviral DNA integrated in the host genome (Brady et al., 2013). ONC201 is able to lower the levels of HIV-1 integration into the genome. Third, ONC201 also decreases intracellular/extracellular HIV-1 p24, GAG RNA, and GAG DNA in HIV-1-infected MDM. Despite its specific effect on HIV-1 replication, at the current working concentration (30 μ M), ONC201 is not able to completely eradicate HIV-1 from the cultures. This could be due to the expression of TRAILshort in MDM that confers TRAIL resistance upon ONC201 treatment (Nie et al., 2018), or lack of optimal trimeric TRAIL conformation that is most potent for the antiviral effect (Mongkolsapaya et al., 1999). Thus, other ART drugs may be required along with ONC201 for optimal HIV-1 inhibition.

As a small molecule, ONC201 shows an outstanding capability of BBB penetration, which makes it particularly attractive for combating CNS viral infection. For brain tumors such as glioblastoma, ONC201 penetrates the intact BBB with five times higher concentrations in brain tissue relative to plasma in rodents (Allen et al., 2013). Therefore, after a successful dose escalation phase I trial (Stein et al., 2015), many ongoing clinical trials (clinicaltrials.gov) focus on brain tumors. In the present study, ONC201 is injected through IP into the NSG mice and its antiviral effect against the HIV-1 infected macrophages in the brains indicate that it is able to cross the BBB and achieve the therapeutic outcome in the case of CNS HIV-1 infection.

Our studies identify that the antiviral effect of ONC201 in HIV-1 infected macrophage is dependent on FOXO3a and TRAIL expressions. However, the exact molecular interactions between ONC201 and FOXO3a remain to be determined. Besides being a FOXO3a activator, ONC201 has also been proposed as a selective dopamine receptor 2 (DRD2) antagonist (Arrillaga-Romany et al., 2017; Kline et al., 2018; Stein et al., 2017). Previous studies have shown ONC201 inactivates both Akt-1 and ERK1/2 to activate FOXO3a and induce the transcription of TRAIL in cancer cell lines (Allen et al., 2013). We observed that the phosphorylation of FOXO3a is decreased after ONC201 treatment in the HIV-1 infected MDM, indicating ONC201 targets the same signaling pathway in cancer cells. However, ONC201 is also known to induce endoplasmic reticulum (ER) stress-related or integrated stress response (ISR)-related genes, such as ATF4, CHOP, GADD34, and TRIB3 (Endo Greer and Lipkowitz, 2016). Whether these genes work in tandem with FOXO3a/TRAIL pathway or not warrants future investigations.

The exact mechanism by which ONC201-activated TRAIL signaling reduces HIV-1 replication remains an active research question. One potential mechanism is that TRAIL-mediated apoptosis clears the HIV-1 infected cells while spares the normal cells. This is supported by previous reports demonstrating that leucine-zipper TRAIL, agonistic anti-TRAIL receptor antibodies, and activated NK cell-derived TRAIL induce apoptosis in MDM and peripheral blood lymphocytes and consequently reduce viral burden in HIV-infected individuals (Lum et al., 2001b, 2004; Shepard et al., 2008). Another potential mechanism of TRAIL is that it may exert its antiviral effect through apoptosis-independent mechanisms. TRAIL signaling may change the status of immune cells by interfering with the activation of NF- κ B, PKB/Akt and MAPKs, which are relevant factors for HIV-1 replication (Falschlehner et al., 2007).




Fig. 8. ONC201 pretreatment increases its efficacies against HIV-1 infection in macrophages. Human MDM were plated in 96-well plates and incubated with doses of ONC201 or its isomer ranging from 3 to 30 μM for 5 days before infection with HIV-1. ONC201 and isomer treatments continued during and after the infection. A) At three-day post infection, cell viability was determined by the MTS assays. B) Cell lysates were collected and subsequently subjected to SDS-PAGE and immunoblotting for the detection of cleaved caspase-3 and pro-caspase-3. Actin was used as the loading control. C) Densitometric quantifications of cleaved caspase-3 in MDM were presented as a ratio to pro-caspase-3 and normalized as fold changes to the vehicle control DMSO group. D) Supernatants were collected and the replication levels of HIV-1 were monitored by RTase activity assay. Results shown are from representative experiments performed with three different donors. Data were analyzed by two-way ANOVA: * denotes $p < 0.05$, ** denotes $p < 0.01$, *** denotes $p < 0.001$, compared to the vehicle control DMSO group.

Our current data support the apoptosis theory as a mechanism of action for ONC201. In HIV-1 infected macrophages with ONC201 post-infection treatment, ONC201-mediated TRAIL production inhibits viral replication. However, viral inhibition could also work to negate the effect of HIV-1-induced cell death as seen in Fig. 1A. In HIV-1 infected macrophages with ONC201 pre-infection treatment, ONC201 dramatically lowered HIV-1 replication levels, which is accompanied by lower cell viabilities and increased cleavage of caspase-3. The ONC201-mediated cytotoxicity in HIV-1 infected MDM raises the question of the safety and potential side effects of ONC201. However, in uninfected MDM and PBL, ONC201 treatment does not cause cytotoxicity, possibly due to the reportedly lower expression levels of death receptors on these cells compared to the infected cells (Lum et al., 2001a). Therefore, these data suggest that while cautions need to be considered, ONC201 is relatively safe to normal cells and will be useful in future antiviral therapies.

Immune-activating agents, such as interleukin 2 and anti-CD3 antibody, or epigenetic modulation, such as Histone deacetylases inhibitors, often failed to purge the virus *in vivo* (Archin et al., 2012; Chun et al., 1999; Elliott et al., 2014; Prins et al., 1999; Rasmussen et al., 2014). Therefore, new therapeutic agents are warranted to combat persistent HIV-1 infection. Our study demonstrates for the first time that ONC201, a FOXO3a activator and TRAIL inducer, inhibits HIV-1 replication in macrophages both *in vitro* and *in vivo*. Notably, ONC201 is the founding member of the imipridone class of compounds that share a unique imidazo pharmacophore. New compounds continue to be developed in this class (Lev et al., 2017; Wagner et al., 2017). These new group of compounds provide us with more drug candidates to modulate immunity and fight viral infections.

Funding

This work was supported in part by research grants by the National Institutes of Health: R01 NS097195 (JCZ), R24 OD018546 (LP, SG), R03 NS094071 (YH), 2P30MH062261-Developmental (YH), the State Key Program of the National Natural Science Foundation of China

(#81830037 to JCZ), the National Basic Research Program of China (973 Program Grant No. 2014CB965001 to JCZ), the Joint Research Fund for Overseas Chinese, Hong Kong and Macao Young Scientists of the National Natural Science Foundation of China (#81329002 to JCZ). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

5. Conflicts of interests/financial disclosures

None.

Author contribution statement

R.Z., J.C.Z., and Y.H., designed the research; R.Z., Y.L., S.G., L.P. H.L., F.G., H.W., J.Z., J.C.Z., and Y.H. performed data acquisition and analysis; R.Z., J.C.Z., and Y.H. drafted the manuscript; all authors approved the final version of the manuscript to be published.

Acknowledgements

The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HIV-1 infected U937 Cells (U1) from Dr. Thomas Folks. We kindly thank Li Wu, Na Ly, and Myhanh Che for providing technical support; Justin Peer and Lenal Bottoms for proofreading the manuscript; Julie Ditter, Johna Belling, and Robin Taylor for providing outstanding administrative and secretarial support.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.antiviral.2019.05.015>.

References

- Allen, J.E., Kline, C.L., Prabhu, V.V., Wagner, J., Ishizawa, J., Madhukar, N., Lev, A., Baumeister, M., Zhou, L., Lulla, A., Stogniew, M., Scholop, L., Benes, C., Kaufman, H.L., Pottorf, R.S., Nallagananchu, B.R., Olson, G.L., Al-Mulla, F., Duvic, M., Wu, G.S., Dicker, D.T., Talekar, M.K., Lim, B., Elemento, O., Oster, W., Bertino, J., Flaherty, K., Wang, M.L., Borthakur, G., Andreeff, M., Stein, M., El-Deiry, W.S., 2016. Discovery and clinical introduction of first-in-class imipridone ONC201. *Oncotarget* 7, 74380–74392.
- Allen, J.E., Krigsfeld, G., Mayes, P.A., Patel, L., Dicker, D.T., Patel, A.S., Dolloff, N.G., Messaris, E., Scata, K.A., Wang, W., Zhou, J.Y., Wu, G.S., El-Deiry, W.S., 2013. Dual inactivation of Akt and ERK by TIC10 signals Foxo3a nuclear translocation, TRAIL gene induction, and potent antitumor effects. *Sci. Transl. Med.* 5, 171ra117.
- Allen, J.E., Krigsfeld, G., Patel, L., Mayes, P.A., Dicker, D.T., Wu, G.S., El-Deiry, W.S., 2015. Identification of TRAIL-inducing compounds highlights small molecule ONC201/TIC10 as a unique anti-cancer agent that activates the TRAIL pathway. *Mol. Cancer* 14, 99.
- Archin, N.M., Liberty, A.L., Kashuba, A.D., Choudhary, S.K., Kuruc, J.D., Crooks, A.M., Parker, D.C., Anderson, E.M., Kearney, M.F., Strain, M.C., Richman, D.D., Hudgens, M.G., Bosch, R.J., Coffin, J.M., Eron, J.J., Hazuda, D.J., Margolis, D.M., 2012. Administration of vorinostat disrupts HIV-1 latency in patients on antiretroviral therapy. *Nature* 487, 482–485.
- Arrillaga-Romany, I., Chi, A.S., Allen, J.E., Oster, W., Wen, P.Y., Batchelor, T.T., 2017. A phase 2 study of the first imipridone ONC201, a selective DRD2 antagonist for oncology, administered every three weeks in recurrent glioblastoma. *Oncotarget* 8, 79298–79304.
- Brady, T., Kelly, B.J., Male, F., Roth, S., Bailey, A., Malani, N., Gijbsbers, R., O'Doherty, U., Bushman, F.D., 2013. Quantitation of HIV DNA integration: effects of differential integration site distributions on Alu-PCR assays. *J. Virol Methods* 189, 53–57.
- Brunet, A., Park, J., Tran, H., Hu, L.S., Hemmings, B.A., Greenberg, M.E., 2001. Protein kinase SGK mediates survival signals by phosphorylating the forkhead transcription factor FKHL1 (FOXO3a). *Mol. Cell Biol.* 21, 952–965.
- Camidge, D.R., Herbst, R.S., Gordon, M.S., Eckhardt, S.G., Kurzrock, R., Durbin, B., Ing, J., Tohny, T.M., Sager, J., Ashkenazi, A., Bray, G., Mendelson, D., 2010. A phase I safety and pharmacokinetic study of the death receptor 5 agonistic antibody PRO95780 in patients with advanced malignancies. *Clin. Cancer Res.* 16, 1256–1263.
- Carpenter, C.C., Cooper, D.A., Fischl, M.A., Gatell, J.M., Gazzard, B.G., Hammer, S.M., Hirsch, M.S., Jacobsen, D.M., Katzenstein, D.A., Montaner, J.S., Richman, D.D., Saag, M.S., Schechter, M., Schooley, R.T., Thompson, M.A., Vella, S., Yeni, P.G., Volberding, P.A., 2000. Antiretroviral therapy in adults: updated recommendations of the International AIDS Society-USA Panel. *JAMA, J. Am. Med. Assoc.* 283, 381–390.
- Chen, Q., Zhang, M., Li, Y., Xu, D., Wang, Y., Song, A., Zhu, B., Huang, Y., Zheng, J.C., 2015. CXCR7 mediates neural progenitor cells migration to CXCL12 independent of CXCR4. *Stem Cell.* 33, 2574–2585.
- Chomont, N., El-Far, M., Ancuta, P., Trautmann, L., Procopio, F.A., Yassine-Diab, B., Boucher, G., Boulassel, M.R., Ghattas, G., Brencnjelj, J.M., Schacker, T.W., Hill, B.J., Douek, D.C., Routy, J.P., Haddad, E.K., Sekaly, R.P., 2009. HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation. *Nat. Med.* 15, 893–900.
- Chun, T.W., Davey Jr., R.T., Engel, D., Lane, H.C., Fauci, A.S., 1999. Re-emergence of HIV after stopping therapy. *Nature* 401, 874–875.
- Cocchi, F., DeVico, A.L., Garzino-Demo, A., Arya, S.K., Gallo, R.C., Lusso, P., 1995. Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8+ T cells. *Science* 270, 1811–1815.
- Cocchi, F., DeVico, A.L., Yarchoan, R., Redfield, R., Cleghorn, F., Blattner, W.A., Garzino-Demo, A., Colombini-Hatch, S., Margolis, D., Gallo, R.C., 2000. Higher macrophage inflammatory protein (MIP)-1alpha and MIP-1beta levels from CD8+ T cells are associated with asymptomatic HIV-1 infection. *Proc. Natl. Acad. Sci. U.S.A.* 97, 13812–13817.
- Constantino, A.A., Huang, Y., Zhang, H., Wood, C., Zheng, J.C., 2011. HIV-1 clade B and C isolates exhibit differential replication: relevance to macrophage-mediated neurotoxicity. *Neurotox. Res.* 20, 277–288.
- Cui, M., Huang, Y., Tian, C., Zhao, Y., Zheng, J., 2011. FOXO3a inhibits TNF- α and IL-1 β -induced astrocyte proliferation: implication for reactive astrogliosis. *Glia* 59, 641–654.
- Cui, M., Huang, Y., Zhao, Y., Zheng, J., 2008. Transcription factor FOXO3a mediates apoptosis in HIV-1-infected macrophages. *J. Immunol.* 180, 898–906.
- Deeks, S.G., Lewin, S.R., Bekker, L.G., 2017. The end of HIV: still a very long way to go, but progress continues. *PLoS Med.* 14, e1002466.
- Degli-Esposti, M.A., Smolak, P.J., Walczak, H., Waugh, J., Huang, C.P., DuBose, R.F., Goodwin, R.G., Smith, C.A., 1997. Cloning and characterization of TRAIL-R3, a novel member of the emerging TRAIL receptor family. *J. Exp. Med.* 186, 1165–1170.
- Dorr, J., Bechmann, I., Waiczies, S., Aktas, O., Walczak, H., Kramer, P.H., Nitsch, R., Zipp, F., 2002. Lack of tumor necrosis factor-related apoptosis-inducing ligand but presence of its receptors in the human brain. *J. Neurosci.* 22, RC209.
- Elliott, J.H., Wightman, F., Solomon, A., Ghneim, K., Ahlers, J., Cameron, M.J., Smith, M.Z., Spelman, T., McMahon, J., Velayudham, P., Brown, G., Roney, J., Watson, J., Prince, M.H., Hoy, J.F., Chomont, N., Fromentin, R., Procopio, F.A., Zeidan, J., Palmer, S., Odeval, L., Johnstone, R.W., Martin, B.P., Sinclair, E., Deeks, S.G., Hazuda, D.J., Cameron, P.U., Sekaly, R.P., Lewin, S.R., 2014. Activation of HIV transcription with short-course vorinostat in HIV-infected patients on suppressive antiretroviral therapy. *PLoS Pathog.* 10, e1004473.
- Emery, J.G., McDonnell, P., Burke, M.B., Deen, K.C., Lyn, S., Silverman, C., Dul, E., Appelbaum, E.R., Eichman, C., DiPrinzio, R., Dodds, R.A., James, I.E., Rosenberg, M., Lee, J.C., Young, P.R., 1998. Osteoprotegerin is a receptor for the cytotoxic ligand TRAIL. *J. Biol. Chem.* 273, 14363–14367.
- Endo Greer, Y., Lipkowitz, S., 2016. ONC201: stressing tumors to death. *Sci. Signal.* 9, fs1.
- Ericksen, D., Lopez, A.L., Peng, H., Niemann, D., Williams, C., Bauer, M., Morgello, S., Cotter, R.L., Ryan, L.A., Ghorpade, A., Gendelman, H.E., Zheng, J., 2003. Neuronal injury regulates fractalkine: relevance for HIV-1 associated dementia. *J. Neuroimmunol.* 138, 144–155.
- Falschlehner, C., Emmerich, C.H., Gerlach, B., Walczak, H., 2007. TRAIL signalling: decisions between life and death. *Int. J. Biochem. Cell Biol.* 39, 1462–1475.
- Folks, T.M., Justement, J., Kinter, A., Dinarello, C.A., Fauci, A.S., 1987. Cytokine-induced expression of HIV-1 in a chronically infected promonocyte cell line. *Science* 238, 800–802.
- Friedrich, B., Li, G., Dziuba, N., Ferguson, M.R., 2010. Quantitative PCR used to assess HIV-1 integration and 2-LTR circle formation in human macrophages, peripheral blood lymphocytes and a CD4+ cell line. *Virol. J.* 7, 354.
- Gendelman, H.E., Orenstein, J.M., Martin, M.A., Ferrua, C., Mitra, R., Phipps, T., Wahl, L.A., Lane, H.C., Fauci, A.S., Burke, D.S., 1988. Efficient isolation and propagation of human immunodeficiency virus on recombinant colony-stimulating factor 1-treated monocytes. *J. Exp. Med.* 167, 1428–1441.
- Gorantla, S., Che, M., Gendelman, H.E., 2005. Isolation, propagation, and HIV-1 infection of monocyte-derived macrophages and recovery of virus from brain and cerebrospinal fluid. *Methods Mol. Biol.* 304, 35–48.
- Greco, F.A., Bonomi, P., Crawford, J., Kelly, K., Oh, Y., Halpern, W., Lo, L., Gallant, G., Klein, J., 2008. Phase 2 study of mapatumumab, a fully human agonistic monoclonal antibody which targets and activates the TRAIL receptor-1, in patients with advanced non-small cell lung cancer. *Lung Cancer* 61, 82–90.
- Hall, H.I., Song, R., Rhodes, P., Prejean, J., An, Q., Lee, L.M., Karon, J., Brookmeyer, R., Kaplan, E.H., McKenna, M.T., Janssen, R.S., Group, H.I.V.I.S., 2008. Estimation of HIV incidence in the United States. *J. Am. Med. Assoc.* 300, 520–529.
- Huang, Y., Erdmann, N., Peng, H., Herek, S., Davis, J.S., Luo, X., Ikezu, T., Zheng, J., 2006. TRAIL-mediated apoptosis in HIV-1-infected macrophages is dependent on the inhibition of Akt-1 phosphorylation. *J. Immunol.* 177, 2304–2313.
- Huang, Y., Walstrom, A., Zhang, L., Zhao, Y., Cui, M., Ye, L., Zheng, J.C., 2009. Type I interferons and interferon regulatory factors regulate TNF-related apoptosis-inducing ligand (TRAIL) in HIV-1-infected macrophages. *PLoS One* 4, e5397.
- Iordanskiy, S., Waltke, M., Feng, Y., Wood, C., 2010. Subtype-associated differences in HIV-1 reverse transcription affect the viral replication. *Retrovirology* 7, 85.
- Kaestner, K.H., Knochel, W., Martinez, D.E., 2000. Unified nomenclature for the winged helix/forkhead transcription factors. *Genes Dev.* 14, 142–146.
- Kaufmann, E., Knochel, W., 1996. Five years on the wings of fork head. *Mech. Dev.* 57, 3–20.
- Kline, C.L., Van den Heuvel, A.P., Allen, J.E., Prabhu, V.V., Dicker, D.T., El-Deiry, W.S., 2016. ONC201 kills solid tumor cells by triggering an integrated stress response dependent on ATF4 activation by specific eIF2alpha kinases. *Sci. Signal.* 9, ra18.
- Kline, C.L.B., Ralff, M.D., Lulla, A.R., Wagner, J.M., Abbosh, P.H., Dicker, D.T., Allen, J.E., El-Deiry, W.S., 2018. Role of dopamine receptors in the anticancer activity of ONC201. *Neoplasia* 20, 80–91.
- Krebs, F.C., Ross, H., McAllister, J., Wigdahl, B., 2000. HIV-1-associated central nervous system dysfunction. *Adv. Pharmacol.* 49, 315–385.
- Leong, S., Cohen, R.B., Gustafson, D.L., Langer, C.J., Camidge, D.R., Padavic, K., Gore, L., Smith, M., Chow, L.Q., von Mehren, M., O'Bryant, C., Hariharan, S., Diab, S., Fox, N.L., Miceli, R., Eckhardt, S.G., 2009. Mapatumumab, an antibody targeting TRAIL-R1, in combination with paclitaxel and carboplatin in patients with advanced solid malignancies: results of a phase I and pharmacokinetic study. *J. Clin. Oncol.* 27, 4413–4421.
- Lev, A., Lulla, A.R., Wagner, J., Ralff, M.D., Kiehl, J.B., Zhou, Y., Benes, C.H., Prabhu, V.V., Oster, W., Astsaturov, I., Dicker, D.T., El-Deiry, W.S., 2017. Anti-pancreatic cancer activity of ONC212 involves the unfolded protein response (UPR) and is reduced by IGF1-R and GRP78/BIP. *Oncotarget* 8, 81776–81793.
- Lin, L., Hron, J.D., Peng, S.L., 2004. Regulation of NF-kappaB, Th activation, and auto-inflammation by the forkhead transcription factor Foxo3a. *Immunology* 21, 203–213.
- Lum, J.J., Pilon, A.A., Sanchez-Dardon, J., Phenix, B.N., Kim, J.E., Mihowich, J., Jamison, K., Hawley-Foss, N., Lynch, D.H., Badley, A.D., 2001a. Induction of cell death in human immunodeficiency virus-infected macrophages and resting memory CD4 T cells by TRAIL/Apo2L. *J. Virol.* 75, 11128–11136.
- Lum, J.J., Pilon, A.A., Sanchez-Dardon, J., Phenix, B.N., Kim, J.E., Mihowich, J., Jamison, K., Hawley-Foss, N., Lynch, D.H., Badley, A.D., 2001b. Induction of cell death in human immunodeficiency virus-infected macrophages and resting memory CD4 T cells by TRAIL/Apo2L. *J. Virol.* 75, 11128–11136.
- Lum, J.J., Schnepfle, D.J., Nie, Z., Sanchez-Dardon, J., Mbisa, G.L., Mihowich, J., Hawley, N., Narayan, S., Kim, J.E., Lynch, D.H., Badley, A.D., 2004. Differential effects of interleukin-7 and interleukin-15 on NK cell anti-human immunodeficiency virus activity. *J. Virol.* 78, 6033–6042.
- McArthur, J.C., Sacktor, N., Selnes, O., 1999. Human immunodeficiency virus-associated dementia. *Semin. Neurol.* 19, 129–150.
- Mom, C.H., Verweij, J., Oldenhuis, C.N., Gietema, J.A., Fox, N.L., Miceli, R., Eskens, F.A., Loos, W.J., de Vries, E.G., Sleijfer, S., 2009. Mapatumumab, a fully human agonistic monoclonal antibody that targets TRAIL-R1, in combination with gemcitabine and cisplatin: a phase I study. *Clin. Cancer Res.* 15, 5584–5590.
- Mongkolsapaya, J., Grimes, J.M., Chen, N., Xu, X.N., Stuart, D.I., Jones, E.Y., Screaton, G.R., 1999. Structure of the TRAIL-DR5 complex reveals mechanisms conferring specificity in apoptotic initiation. *Nat. Struct. Biol.* 6, 1048–1053.
- Nie, Z., Aboulnasr, F., Natesampillai, S., Burke, S.P., Krogman, A., Bren, G.D., Chung, T.D.Y., Anderson, J.R., Smart, M.K., Katzmann, D.J., Rajagopalan, G., Cummins, N.W., Badley, A.D., 2018. Both HIV-infected and uninfected cells express TRAILshort, which confers TRAIL resistance upon bystander cells within the microenvironment. *J.*

- Immunol. 200, 1110–1123.
- Pan, G., Ni, J., Yu, G.-L., Wei, Y.-F., Dixit, V.M., 1998. TRUND D a new member of the TRAIL receptor family that antagonizes TRAIL signalling. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 424, 41–45.
- Pan, G., O'Rourke, K., Chinnaiyan, A.M., Gentz, R., Ebner, R., Ni, J., Dixit, V.M., 1997. The receptor for the cytotoxic ligand TRAIL. *Science* 276, 111–113.
- Peng, H., Huang, Y., Duan, Z., Erdmann, N., Xu, D., Herek, S., Zheng, J., 2005. Cellular IAP1 regulates TRAIL-induced apoptosis in human fetal cortical neural progenitor cells. *J. Neurosci. Res.* 82, 295–305.
- Pierson, T., McArthur, J., Siliciano, R.F., 2000. Reservoirs for HIV-1: mechanisms for viral persistence in the presence of antiviral immune responses and antiretroviral therapy. *Annu. Rev. Immunol.* 18, 665–708.
- Pitti, R.M., Marsters, S.A., Ruppert, S., Donahue, C.J., Moore, A., Ashkenazi, A., 1996. Induction of apoptosis by Apo-2 ligand, a new member of the tumor necrosis factor cytokine family. *J. Biol. Chem.* 271, 12687–12690.
- Plummer, R., Attard, G., Pacey, S., Li, L., Razak, A., Perrett, R., Barrett, M., Judson, I., Kaye, S., Fox, N.L., Halpern, W., Corey, A., Calvert, H., de Bono, J., 2007. Phase I and pharmacokinetic study of lexatumumab in patients with advanced cancers. *Clin. Cancer Res.* 13, 6187–6194.
- Prins, J.M., Jurriaans, S., van Praag, R.M., Blaak, H., van Rij, R., Schellekens, P.T., ten Berge, L.J., Yong, S.L., Fox, C.H., Roos, M.T., de Wolf, F., Goudsmit, J., Schuitemaker, H., Lange, J.M., 1999. Immuno-activation with anti-CD3 and recombinant human IL-2 in HIV-1-infected patients on potent antiretroviral therapy. *AIDS* 13, 2405–2410.
- Rasmussen, T.A., Tolstrup, M., Brinkmann, C.R., Olesen, R., Erikstrup, C., Solomon, A., Winkelmann, A., Palmer, S., Dinarello, C., Buzon, M., Lichterfeld, M., Lewin, S.R., Ostergaard, L., Sogaard, O.S., 2014. Panobinostat, a histone deacetylase inhibitor, for latent-virus reactivation in HIV-infected patients on suppressive antiretroviral therapy: a phase 1/2, single group, clinical trial. *Lancet HIV* 1, e13–21.
- Ryan, L.A., Peng, H., Erichsen, D.A., Huang, Y., Persidsky, Y., Zhou, Y., Gendelman, H.E., Zheng, J., 2004. TNF-related apoptosis-inducing ligand mediates human neuronal apoptosis: links to HIV-1 associated dementia. *J. Neuroimmunol.* 148, 127–139.
- Seol, D.W., Li, J., Seol, M.H., Park, S.Y., Talanian, R.V., Billiar, T.R., 2001. Signaling events triggered by tumor necrosis factor-related apoptosis-inducing ligand (TRAIL): caspase-8 is required for TRAIL-induced apoptosis. *Cancer Res.* 61, 1138–1143.
- Shepard, B.D., De Forni, D., McNamara, D.R., Foli, A., Rizza, S.A., Abraham, R.S., Knutson, K., Wettstein, P.J., Lori, F., Badley, A.D., 2008. Beneficial effect of TRAIL on HIV burden, without detectable immune consequences. *PLoS One* 3, e3096.
- Stein, M.N., Bertino, J.R., Kaufman, H.L., Mayer, T., Moss, R., Silk, A., Chan, N., Malhotra, J., Rodriguez, L., Aisner, J., Aiken, R.D., Haffty, B.G., DiPaola, R.S., Saunders, T., Zloza, A., Damare, S., Beckett, Y., Yu, B., Najmi, S., Gabel, C., Dickerson, S., Zheng, L., El-Deiry, W.S., Allen, J.E., Stogniew, M., Oster, W., Mehnert, J.M., 2017. First-in-Human clinical trial of oral ONC201 in patients with refractory solid tumors. *Clin. Cancer Res.* 23, 4163–4169.
- Stein, M.N., Mayer, T.M., Moss, R.A., Silk, A.W., Chan, N., Haffty, B.G., DiPaola, R.S., Beckett, Y., Bentleywski, E., Zheng, L., Fang, B., Allen, J., Mehnert, J.M., 2015. First-in-human dose escalation study of oral ONC201 in advanced solid tumors. *J. Clin. Oncol.* 33 TPS2623-TPS2623.
- Tian, C., Li, Y., Huang, Y., Wang, Y., Chen, D., Liu, J., Deng, X., Sun, L., Anderson, K., Qi, X., Li, Y., Mosley, R.L., Chen, X., Huang, J., Zheng, J.C., 2015. Selective generation of dopaminergic precursors from mouse fibroblasts by direct lineage conversion. *Sci. Rep.* 5, 12622.
- Tolcher, A.W., Mita, M., Meropol, N.J., von Mehren, M., Patnaik, A., Padavic, K., Hill, M., Mays, T., McCoy, T., Fox, N.L., Halpern, W., Corey, A., Cohen, R.B., 2007. Phase I pharmacokinetic and biologic correlative study of mapatumumab, a fully human monoclonal antibody with agonist activity to tumor necrosis factor-related apoptosis-inducing ligand receptor-1. *J. Clin. Oncol.* 25, 1390–1395.
- Trarbach, T., Moehler, M., Heinemann, V., Kohne, C.H., Przyborek, M., Schulz, C., Sneller, V., Gallant, G., Kanzler, S., 2010. Phase II trial of mapatumumab, a fully human agonistic monoclonal antibody that targets and activates the tumour necrosis factor apoptosis-inducing ligand receptor-1 (TRAIL-R1), in patients with refractory colorectal cancer. *Br. J. Canc.* 102, 506–512.
- van Grevenynghe, J., Procopio, F.A., He, Z., Chomont, N., Riou, C., Zhang, Y., Gimmig, S., Boucher, G., Wilkinson, P., Shi, Y., Yassine-Diab, B., Said, E.A., Trautmann, L., El Far, M., Balderas, R.S., Boulassel, M.R., Routy, J.P., Haddad, E.K., Sekaly, R.P., 2008. Transcription factor FOXO3a controls the persistence of memory CD4(+) T cells during HIV infection. *Nat. Med.* 14, 266–274.
- Wagner, J., Kline, C.L., Ralff, M.D., Lev, A., Lulla, A., Zhou, L., Olson, G.L., Nallaganchu, B.R., Benes, C.H., Allen, J.E., Prabhu, V.V., Stogniew, M., Oster, W., El-Deiry, W.S., 2017. Preclinical evaluation of the imipridone family, analogs of clinical stage anti-cancer small molecule ONC201, reveals potent anti-cancer effects of ONC212. *Cell Cycle* 16, 1790–1799.
- Walczak, H., Degli-Esposti, M.A., Johnson, R.S., Smolak, P.J., Waugh, J.Y., Boiani, N., Timour, M.S., Gerhart, M.J., Schooley, K.A., Smith, C.A., Goodwin, R.G., Rauch, C.T., 1997. TRAIL-R2: a novel apoptosis-mediating receptor for TRAIL. *EMBO J.* 16, 5386–5397.
- Wang, Y., Li, Y., Zhao, R., Wu, B., Lanoha, B., Tong, Z., Peer, J., Liu, J., Xiong, H., Huang, Y., Zheng, J., 2017. Glutaminase C overexpression in the brain induces learning deficits, synaptic dysfunctions, and neuroinflammation in mice. *Brain Behav. Immun.* 66, 135–145.
- Wiley, S.R., Schooley, K., Smolak, P.J., Din, W.S., Huang, C.P., Nicholl, J.K., Sutherland, G.R., Smith, T.D., Rauch, C., Smith, C.A., et al., 1995. Identification and characterization of a new member of the TNF family that induces apoptosis. *Immunity* 3, 673–682.
- Wu, Y., Peng, H., Cui, M., Whitney, N.P., Huang, Y., Zheng, J.C., 2009. CXCL12 increases human neural progenitor cell proliferation through Akt-1/FOXO3a signaling pathway. *J. Neurochem.* 109, 1157–1167.
- Ye, L., Huang, Y., Zhao, L., Li, Y., Sun, L., Zhou, Y., Qian, G., Zheng, J.C., 2013. IL-1beta and TNF-alpha induce neurotoxicity through glutamate production: a potential role for neuronal glutaminase. *J. Neurochem.* 125, 897–908.
- Zhu, B., Xu, D., Deng, X., Chen, Q., Huang, Y., Peng, H., Li, Y., Jia, B., Thoreson, W.B., Ding, W., Ding, J., Zhao, L., Wang, Y., Wavrin, K.L., Duan, S., Zheng, J., 2012. CXCL12 enhances human neural progenitor cell survival through a CXCR7- and CXCR4-mediated endocytotic signaling pathway. *Stem Cell.* 30, 2571–2583.