



Predicting latent classes of drug use among adolescents through parental alcohol use and parental style: a longitudinal study

Juliana Y. Valente¹ · Hugo Cogo-Moreira² · Zila M. Sanchez¹

Received: 4 April 2018 / Accepted: 3 December 2018 / Published online: 12 December 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Purpose The present study examined the roles of parental alcohol use and parental style as predictors of adolescent patterns of drug use.

Methods 6391 students in the 7th and 8th grades at 72 Brazilian public schools participated in a three-wave randomized controlled trial to evaluate a school drug-use prevention program. Patterns of drug use were identified through two latent class analyses using measures of the adolescents' past-year drug use. Multinomial logistic regression analyses examined whether parental alcohol use and parenting style at baseline predicted patterns of drug use in waves 2 and 3 of the study after controlling for sociodemographic covariates.

Results In each of the two waves, three latent classes of drug use were identified among the students, defining three different groups of individuals: (1) abstainers/low users, (2) alcohol users/binge drinkers, and (3) polydrug users. First, parenting style (especially monitoring) was the strongest predictor for the prevention of polydrug use among adolescents. Second, occasional alcohol use by parents can act as a central predictor for adolescent alcohol use and binge drinking. Above all, maternal episodes of drunkenness were involved in the predictive models for both drug use classes in both waves.

Conclusion Parental alcohol use and parenting style seem to be important predictors of adolescent's likelihood of belonging to different latent classes of drug use. This conclusion may point to the importance of considering the inclusion of parenting skills and parental alcohol use within the scope of adolescents' preventive interventions.

Keywords Adolescence · Drug use · Latent class analysis · Parental alcohol use · Parenting style

Introduction

Adolescence is the key period for the initiation of alcohol consumption, with 55.5% of 13–15-year-old students in Brazil reporting having consumed alcohol and 9.0% reporting having taken illicit drugs [1]. Early-onset use of alcohol and other drugs places adolescents at an increased risk of engaging in heavy, frequent drinking and raises their probability of experiencing drug-related problems in later adolescence [2–5]. An early onset of alcohol and drug use is also a key predictor of drug-related problems later in life, such as adult

mental disorders [6], cognitive impairment [7] and substance use problems and dependence in adulthood [8–12]. Considering the international public health impact of adolescent drug use [13], it is important to identify the predictors that underlie this behavior to prevent harm [14].

Three systematic reviews have demonstrated that parental drinking plays a central role in the development of risk for both an early onset of drinking and increased later alcohol use [15, 16], as well as negative alcohol-related consequences in adolescence [17]. Some of the studies have shown that even moderate maternal alcohol consumption is a strong risk factor for the development of alcohol problems in adolescence [18], while other studies associate only heavy episodic drinking [19] or parental alcohol-related problems [20] with adolescent alcohol use. Despite the consistent literature demonstrating the association between parental drinking and adolescent drinking, there is insufficient evidence for causal inference [21]. In addition, some studies did not find such an association between parental

✉ Zila M. Sanchez
zila.sanchez@unifesp.br

¹ Department of Preventive Medicine, Universidade Federal de São Paulo, Rua Botucatu, 740/404, São Paulo 04023-062, Brazil

² Department of Psychiatry, Universidade Federal de São Paulo, São Paulo, Brazil

and adolescent drinking, showing instead that higher levels of parental problem drinking were related to lower levels and later onset of teen drinking [22]. Little is known about the association between parental alcohol use and adolescent drug use [23], and the findings on the subject are still very controversial [24, 25].

Evidence also points to the importance of parenting styles in the risk of drug use by offspring [26]. Studies have indicated that the authoritative parenting style is the most protective against adolescent substance use [27], while the neglectful parenting style is associated with elevated rates of drug use by adolescents [28]. However, research on indulgent and permissive parenting remains inconclusive [29–31]. Nevertheless, most studies investigating these interrelationships do not include parental alcohol use as a possible confounder variable in the analysis [32]. In addition, most studies on the subject do not consider the association between parental risk and protective factors with the multiple latent class patterns of drug use among adolescents [33]; only a few studies have considered capturing the complexity involved in the heterogeneity of drug use [34–37]. Latent class analysis (LCA) is a mixture model [38–40] that makes it possible to analyze simultaneously the consumption of multiple substances [41] and allows an improved understanding of the outcomes of different substance use profiles [42, 43].

To the best of our knowledge, this study is the first one aiming to analyze simultaneously the roles of parental alcohol use and parental style as predictors of adolescent patterns of drug use, using LCA methods to define unobserved underlying groups of adolescent drug use.

Materials and methods

This study presents the results of a three-wave longitudinal (baseline, 9-month follow-up and 21-month follow-up) randomized controlled trial to evaluate a school drug-use prevention program. Thus, the study was originally designed as a cluster randomized controlled parallel-group trial among 6,391 7th- and 8th-grade public school students in six Brazilian cities (São Paulo, São Bernardo do Campo, Federal District, Florianópolis, Tubarão and Fortaleza). In partnership with the Brazilian branch of the United Nations Office on Drugs and Crime (UNODC), the Brazilian Ministry of Health decided to undertake a culturally adapted version of the European drug prevention program Unplugged, renamed #Tamojunto, to be applied in Brazilian public schools [44], and the evaluation was performed by an independent team from two universities [45].

Randomization was performed at the school level via an Excel macro [command RAND]. Seventy-two schools were randomly selected in proportion to the number of schools in each municipality (stratum). A second allocation step

used a random list to determine whether each school would be assigned to the control or intervention group according to a random list, maintaining a 1:1 allocation ratio per municipality.

This article examined the data from the baseline assessment conducted in February 2014 and the data from two follow-ups (9 months and 21 months after baseline assessment). The study was registered in the Brazilian Ministry of Health's Brazilian Registry of Clinical Trials (Registro Brasileiro de Ensaios Clínicos–REBEC) under the number RBR-4mnv5g. The study protocol was approved by the Federal University of São Paulo's research ethics committee (protocol #473.498).

Population and sample size

Based on Lwanga and Lemeshow's [46] calculation of sample sizes for longitudinal studies, the sample size necessary in this study for a power of 80%, a significance level of 5%, and a difference between groups of 1.5% points (5% vs 3.5%), was calculated to be 2835 participants per group. Assuming that 50% would be lost to follow-up, the sample had to include 4253 participants in each group. The parameters used were based on a previous pilot study and the expected results of the randomized controlled trial. Details of the study design and a flowchart of the sampling process have been presented in a prior publication [45].

Instruments and variables

To collect the data, we used an anonymous self-report questionnaire developed and tested by the European Drug Addiction Prevention Trial (EU-DAP) program and used in previous studies on the effectiveness of Unplugged [47]. As our trial was conducted in Brazil, we used a version of the questionnaire that had been translated and adapted to Portuguese, with some questions replaced by items from two questionnaires that have been widely used in various studies of Brazilian students: a World Health Organization questionnaire, used by the Brazilian Center for Information about Psychotropic Drugs (Centro Brasileiro de Informações Sobre Drogas Psicotrópicas–CEBRID) [48], and the questionnaire of the National Survey of Student Health (Pesquisa Nacional de Saúde do Escolar–PENSE), used by the Brazilian Ministry of Health [49].

In the present study, the explanatory variables (predictors) used were three sets of variables from the wave 1 (baseline) data assessment: (1) Control variables: age, gender, socioeconomic class, randomized group; use (yes/no) of alcohol, tobacco, marijuana, and inhalants within the past year; and binge drinking (yes/no) within the past year. Binge drinking was defined as the consumption of 5 or more doses of alcohol during a 2-h period. (2) Parental alcohol use: These

variables have been collected through four dichotomous questions (yes or no), asking the participant whether his/her father or mother drinks occasionally and whether his/her father or mother has episodes of drunkenness. (3) Parental style (neglectful, authoritative, authoritarian, or indulgent) was assessed through demandingness and responsiveness scales.

The students’ socioeconomic class was assessed using the scale of the Brazilian Association of Research Companies (Associação Brasileira de Empresas de Pesquisa–ABEP) [50], which takes into account the education of the head of the household and the goods and services used, with scores ranging from 0 to 46 or in categories from A to E; higher scores indicate better economic standing, and socioeconomic classes are ranked from A (highest) to E (lowest).

The data relating to parenting styles were collected through scales measuring demandingness and responsiveness [51]. The results were used to define four parenting styles (authoritarian, authoritative, neglectful, and indulgent) based on Maccoby and Martin’s theoretical model [52]. The instrument consists of two scales that measure the orthogonal dimensions of demandingness and responsiveness. The scales are structured into six items on the demandingness dimension and ten items on the responsiveness dimension, each assessed by means of a three-point Likert scale such that values closer to three indicate greater perceived demandingness and responsiveness. Parents are classified as high or low in demandingness and high or low in responsiveness. The scale is corrected based on the median scores for each subscale, with the parents who score at or above the median for demandingness or responsiveness being classified as high in demandingness or responsiveness, respectively, whereas parents who score at or below the median were classified as low in demandingness or responsiveness. Parenting styles are classified into four categories, based on the combination of these two dimensions: authoritative (parents scoring high

on demandingness and responsiveness), authoritarian (those scoring high on demandingness and low on responsiveness), indulgent (parents scoring low on demandingness and high on responsiveness), and neglectful (those scoring low on both demandingness and responsiveness) [29].

The assessed outcome variables were from wave 2 (9-month follow-up) and wave 3 (21-month follow-up): (1) Occurrence (yes/no) of alcohol use, tobacco use, marijuana use, inhalant use and binge drinking within the past year. Binge drinking was defined as the consumption of 5 or more doses of alcohol during a 2-h period.

We choose to include alcohol, tobacco, marijuana and inhalants since they are the most prevalent drugs among Brazilian adolescents considering the last Brazilian National Survey Among School Students. Inhalants are the most consumed illicit drugs among Brazilian adolescents [48].

Figure 1 depicts the conceptual model connecting the indicator variables (past-year drug use), the latent classes, and the baseline predictors for the two latent class models in waves 2 and 3.

Statistical analysis

LCA was used to identify groups with similar patterns of drug use. The two LCA models were constructed with the observed variables from past-year drug use (alcohol, tobacco, marijuana, inhalants, and binge drinking) from the two waves (9 and 21 months). The enumeration process extracted 1–5 classes, and due to the study’s multilevel sampling, the standard errors were corrected as described in Asparouhov [53], taking the school (second level) as the cluster indicator. Mplus version 7.4 [54] was used for all analyses. The extraction of latent classes ceased when the addition of a new class yielded little additional information. The model was adjusted based on the “goodness-of-fit” criterion and took into consideration the parsimony and




Fig. 1 Latent class model of past-year drug use, with age, gender, socioeconomic status, group, baseline past year drug use, parental drinking and parental style as baseline predictors

interpretability of the classes. That is, in addition to the statistical indices presented below, the decision regarding the best solution for the number of latent classes took into consideration whether each solution had a logical substantive interpretation. The goodness-of-fit statistics included the following: the Akaike information criterion (AIC), the Bayesian information criterion (BIC), the sample-size-adjusted Bayesian information criterion (SSABIC), and the Vuong–Lo–Mendell–Rubin (VLMR) test. Finally, to assess how well discriminated the latent classes underlying the best solution were, we used entropy, which is based on an a posteriori probability and indicates the accuracy of the classification; values close to 1 indicate clear and very precise classifications. We emphasize that entropy in itself was not used to decide the best solution for the number of latent classes.

Because large amounts of data relating to parental consumption, parental style and adolescent drug use at 9- and 21-month follow-up were lost due to missing data, they were imputed in Mplus through sequential imputation [54]. The following were used as variables in the unrestricted model: group, school, gender, age, past-year drug use at baseline (alcohol, binge drinking, cigarettes, inhalants, marijuana), and ABEP classification. Five imputed datasets were generated. Subsequently, two multinomial logistic regressions [55] were performed in Mplus using the R3STEP option of the AUXILIARY command [56] with the baseline covariant variables affecting the outcome. One regression was performed using the baseline variables impacting the latent classes in wave 2, and the other regression was performed using the baseline variables impacting the latent classes in wave 3.

The descriptive statistics are weighted percentages (wgt%) based on random levels of the sample subjects and records of the expected population taken from official data from the Anisio Teixeira National Institute of Educational Studies and Research (Instituto Nacional de Estudos e Pesquisa Educacionais Anisio Teixeira–INEP). Inferential point estimates are given as adjusted odds ratios (aORs) with their respective 95% CIs and *p* values. The adopted level of significance was 5%.

Results

72 schools participated in the study, totalizing 261 classes. 6391 students answered the baseline questionnaire, 5957 answered the follow-up questionnaire 9 months after baseline, and 4434 answered the follow-up questionnaire 21 months after baseline. The dropout rate was due to refusal and absence. Due to maximum information likelihood, latent classes analysis included in the cross-sectional analysis those participants who had at least one

measurement of past year drug use in each wave ($n = 4231$ in wave 2 and 3635 in wave 3). In the latent class of wave 2, we had 33.80% of losses and in the latent class of wave 3 we had 43.12% of losses, considering the baseline assessment. Due to the intense, but expected missingness, we imputed the drug use missing values achieving with the 6391 participants who were analyzed in the two-waves latent classes.

Table 1 presents the characteristics of the students participating in the study ($N = 6,391$). The obtained data show that, at baseline, the majority of students were girls (51.21%) aged between 11 and 12 years (53.80%) who belonged to the middle socioeconomic class (53.98%), and the drug most used by the students over the past year was alcohol (30.94%). Moreover, alcohol use was more prevalent in the fathers than in the mothers of the adolescents, both for episodic alcohol use (30.52% and 21.24%, respectively) and for drunkenness (9.94% and 2.44%, respectively), and the most prevalent parenting style was the neglectful style (37.84%). In wave 2 and wave 3, alcohol continued to be the most prevalent drug used by the adolescents over the past year (35.5% and 47.75%, respectively).

Latent classes were created based on the drug-use indicator variables reported as occurring during the past year. The three classes distinguished polydrug users, alcohol users/binge drinkers and abstainers/low users. The latent classes were identified based on the variables related to drug consumption over the past year (alcohol, binge drinking, cigarettes, inhalants, and marijuana). Adolescents classified as polydrug users exhibited the highest probabilities of having engaged in all five categories of drug use. Those classified as alcohol users/binge drinkers had high probabilities of self-reported binge drinking and alcohol use in the past year; however, they had lower probabilities of having used cannabis, cigarettes, and inhalants. The third class exhibited very low probabilities of alcohol use, binge drinking, and use of tobacco or cannabis (see Figs. 2, 3). The relative proportions of the classes were relatively stable across the two waves. The polydrug user class was consistently the smallest (from 6.12 to 7.92%), the alcohol users/binge drinkers class was the next smallest (from 29.42 to 21.42%), and the abstainers/low users class was the largest (64.45–70.61%).

In each wave, a total of five classes were defined. Table 2 shows values of the information criteria. In wave 2, the fit indices (lower BIC, SSABIC and AIC values) suggested that the four-class model was slightly superior to the other. However, a careful examination of four-class model solutions led us to select the three-class model because it was the most coherent solution in terms of theoretical interpretation of the drug-use phenomenon and fit index. There would not be a conceptual explanation for the maintenance of this fourth class, which accounts for a small proportion of subjects (0.5% and 0.2% in wave 2 and wave 3, respectively) and is

Table 1 Sociodemographic characteristics of students participating in a study evaluating a school-based program for drug-use prevention (N=6,391)

Variables	N	wgt%	wgt 95% CI
Baseline measures (wave 1)			
Group			
Intervention	3.243	50.26	[34.65; 65.82]
Control	3.148	49.74	[34.18; 65.35]
Gender			
Male	3.130	48.79	[47.03; 50.56]
Female	3.261	51.21	[49.44; 52.96]
Average age		12.61	[12.56; 12.67]
ABEP score		27.67	[26.92; 28.41]
A (35–42)	244	3.78	[2.80; 5.11]
B (23–34)	2.467	36.64	[33.54; 39.85]
C (14–22)	3.343	53.98	[50.41; 57.49]
D/E (0–13)	322	5.60	[4.60; 6.80]
Adolescent past-year drug use^a			
Alcohol	2.015	30.94	[28.67; 33.30]
Binge drinking	1.006	16.50	[15.10; 18.01]
Tobacco	243	4.02	[3.31; 4.85]
Inhalants	525	8.22	[7.45; 9.07]
Marijuana	156	2.56	[2.05; 3.20]
Parenting style^a			
Authoritative	1447	28.69	[26.65; 30.83]
Authoritarian	960	19.66	[18.56; 20.80]
Indulgent	662	13.81	[12.72; 14.98]
Neglectful	1863	37.84	[35.66; 40.07]
Family alcohol use^a			
Paternal alcohol use	1.913	30.52	[28.03; 33.14]
Paternal drunkenness	600	09.94	[09.10; 10.84]
Maternal alcohol use	1313	21.24	[19.50; 23.10]
Maternal drunkenness	151	2.44	[2.00; 2.96]
9-month follow-up measures (wave 2)			
Adolescent past-year drug use^a			
Alcohol	1492	35.50	[34.06; 36.95]
Binge drinking	726	7.36	[16.24; 18.53]
Tobacco	216	5.15	[4.52; 5.86]
Inhalants	422	10.07	[9.19; 11.02]
Marijuana	202	4.81	[4.20; 5.50]
21-month follow-up measures (wave 3)			
Adolescent past-year drug use^a			
Alcohol	1731	47.75	[46.18; 49.38]
Binge drinking	908	25.19	[23.80; 26.64]
Tobacco	252	6.99	[6.20; 7.87]
Inhalants	377	10.45	[9.49; 11.48]
Marijuana	276	7.67	[6.84; 8.58]

^aMissing values of these variables were imputed for the inferential analyses




Fig. 2 Weighted probabilities of occurrence over the past year of alcohol use, binge drinking, tobacco use, inhalant use, and marijuana use according to the model of the three latent classes with imputation data among adolescents who participated in waves 2 and 3 (9- and 21-months follow-up data) in a study evaluating a school-based drug-use prevention program

derived from the class of “polydrug users”. For the three-class solution, the value of entropy was 0.851.

In wave 3, the BIC value suggested that the three-class model was slightly superior to the others in that it showed a lower value, while the SSABIC value more clearly suggested the four-class model. A careful examination of both the three- and four-class model solutions led us to select the three-class model because, taking the BIC value as one of the most reliable measures [57] and considering the same theoretical interpretation criteria used for latent class solutions in wave 2, the model with three latent classes was chosen as the most parsimonious. For the three-class solution, the value of entropy was 0.82. Therefore, for each of the two waves, the best model solution identified three latent classes.

Tables 3 (for the second wave) and 4 (for the third wave) show the results of the integrative models, which were multinomial logistic regression models (univariate and multivariate) using the latent classes as the outcomes. The class of “abstainers/low users” was used as a reference.

Girls were more likely than boys to belong to the “alcohol users/binge drinkers” class rather than the “abstainers/low users” class in waves two and three (aOR = 1.63, 95% CI




Fig. 3 Weighted probabilities of occurrence over the past year of alcohol use, binge drinking, tobacco use, inhalant use, and marijuana use according to the model of the three latent classes without imputation data among adolescents who participated in waves 2 and 3 (9- and 21-months follow-up data) in a study evaluating a school-based drug-use prevention program

[1.29; 2.05], and $aOR = 1.73$, 95% CI [1.35; 2.21], respectively). In wave three, each increase of point on the socioeconomic scale (which varied from 0 to 42) corresponded to an increase of 2% ($aOR = 1.02$, 95% CI [1.00; 1.04]) in the probability that a student would belong to the “alcohol users/binge drinkers” class rather than the “abstainers/low users” class. The adolescents who reported that their mothers or fathers drank were more likely than the children of abstainers to be in the “alcohol users/binge drinkers” class rather than the “abstainers/low users” class in wave two ($aOR = 1.57$, 95% CI [1.20; 2.07], and $aOR = 1.59$, 95% CI [1.30; 1.95], respectively) and wave three ($aOR = 1.62$, 95% CI [1.01; 2.60], and $aOR = 1.36$, 95% CI [1.12; 1.65], respectively). The same result was found for parental drunkenness: adolescents who reported that their mothers or fathers had episodes of drunkenness were more likely than the children of abstinent parents to be in the “alcohol users/binge drinkers” class rather than the “abstainers/low users” class ($aOR = 1.74$, 95% CI [1.02; 2.98], and $aOR = 1.59$, 95% CI [1.19; 2.16], respectively) in wave two. In wave three, only the adolescents who reported at baseline that their

mothers had episodes of drunkenness were more likely than the children of abstainers to be in the “alcohol users/binge drinkers” class rather than the “abstainers/low users” class ($aOR = 2.10$, 95% CI [1.31; 3.35]).

In wave three, girls were also more likely than boys to belong to the “polydrug users” class rather than the “abstainers/low users” class ($aOR = 1.43$, 95% CI [1.14; 1.80]). In wave two, for each additional year of age, the probability of belonging to the “polydrug users” group rather than the “abstainers/low users” group was multiplied by a factor of 1.43 (95% CI [1.16; 1.74]). The students who reported that their mothers had episodes of drunkenness at baseline were more likely than the children of abstainers to belong to the “polydrug users” group rather than to the “abstainers/low users” group in waves two and three ($aOR = 3.73$, 95% CI [1.61; 8.60], and $aOR = 2.79$, 95% CI [1.52; 5.12], respectively). With regard to parenting styles, it was found that students whose parents adopted authoritative or authoritarian styles were less likely than the children of neglectful parents to belong to the “polydrug users” group rather than to the “abstainers/low users” group in wave two ($aOR = 0.47$, 95% CI [0.26; 0.86], and $aOR = 0.30$, 95% CI [0.12; 0.75], respectively) and wave three ($aOR = 0.62$, 95% CI [0.44; 0.89], and $aOR = 0.46$, 95% CI [0.31; 0.69], respectively).

In wave 2, no effect of the program was identified regarding the probability of belonging to the “alcohol users/binge drinkers” or “polydrug users” class rather than to the “abstainers/low users” class ($aOR = 1.29$, $p = 0.058$, and $aOR = 1.03$, $p = 0.903$, respectively). Similarly, in wave 3, no effect of the program was identified regarding to the probability of belonging to the “alcohol users/binge drinkers” or “polydrug users” class rather than to the “abstainers/low users” class ($aOR = 1.20$, $p = 0.300$, and $aOR = 1.51$, $p = 0.334$, respectively).

Discussion

The strength of this large-scale, three-wave prospective study is that it applies established knowledge of LCA to illuminate the roles of parents’ alcohol use and parenting styles in predicting adolescent drug use. A solution with three latent classes (“abstainers/low users”, “alcohol users/binge drinkers,” and “polydrug users”) provided the best explanation for the patterns of drug use among those adolescents surveyed in the two waves. This study identified differences between the “abstainers/low users” class and the two other classes (“alcohol users/binge drinkers” and “polydrug users”) with regard to parenting style and parent’s alcohol use. In this study, we analyze early adolescence (baseline average age 12.61) since it was a period before the average age of onset of drug use in Brazil (13 years-old). Average age of onset for the different substances between

Table 2 Goodness-of-fit statistics for the number of latent classes of drug use over the past year among students participating in the 9-month follow-up phase of a study to evaluate the effect of a school-based program for drug-use prevention

Model	Factor correction	Log likelihood	AIC	BIC	SSABIC	VLMR-LRT	LMR Adjusted LRT	Polydrug users (n, %)	Final count based on most likely membership	
									Alcohol drinkers (n, %)	Abstainers/low users (n, %)
9 months (N=4,231)										
1 Class	2.29	-7693.84	15,397.69	15,429.44	15,413.55					
2 Classes	1.57	-6308.88	12,639.75	12,709.61	12,674.65	<0.001				
3 Classes	1.41	-6209.99	12,453.99	12,561.94	12,507.93	<0.001	259 (6.12%)	1245 (29.42%)	2727 (64.45%)	0.851
4 Classes	1.26	-6177.48	12,400.97	12,547.02	12,473.94	0.0002				
5 Classes	1.23	-6174.09	12,406.18	12,590.33	12,498.18	0.5620				
21 months (N=3,635)										
1 Class	1.69	-7638.96	15,287.92	15,318.91	15,303.02					
2 Classes	1.41	-6254.99	12,531.99	12,600.18	12,565.22	<0.001				
3 Classes	1.25	-6134.69	12,303.31	12,408.75	12,354.74	<0.001	288 (7.92%)	780 (21.45%)	2567 (70.61%)	0.820
4 Classes	1.24	-6116.83	12,279.66	12,422.22	12,349.13	0.071				
5 Classes	1.12	-6115.40	12,288.79	12,468.55	12,376.40	0.335				

AIC Akaike information criterion, BIC Bayesian information criteria, SSABIC sample-size-adjusted Bayesian information criterion, VLMR-LRT Vuong-Lo-Mendell-Rubin, LRT likelihood ratio test, LMR Lo-Mendell-Rubin

Table 3 Characteristics of polysubstance drug use and binge drinking (BD) identified through latent class analysis in a sample of students participating in the 9-month follow-up (wave 2) of a study to evaluate the effect of a school-based program for drug-use prevention ($N=6,391$)

Group	Latent classes, wave 2			<i>p</i>	Univariate analysis ^b			Multivariate analysis ^b								
	Abstainers/low users ^c	Alcohol users/ binge drinkers	Polydrug users		Abstainers/low users vs. alcohol users/binge drinkers	Abstainers/low users vs. polydrug users	Abstainers/low users vs. alcohol users/binge drinkers	Abstainers/low users vs. polydrug users								
	%	%	%		cOR	95% CI	<i>p</i>	cOR	95% CI	<i>p</i>	aOR	95% CI	<i>p</i>			
Intervention	46.83	48.03	44.40	0.531	1.27	[0.96; 1.57]	0.927	1.02	[0.66; 1.67]	0.927	1.29	[0.99; 1.68]	0.058	1.03	[0.67; 1.58]	0.903
Control	53.17	51.97	55.60													
Sex				0.697												
Girl	50.06	50.92	52.51		1.63	[1.29; 2.05]	<0.001	1.27	[0.85; 1.90]	0.234	1.63	[1.29; 2.05]	<0.001	1.33	[0.89; 1.97]	0.164
Boy	49.94	49.08	47.49													
Age				0.537												
12.60±0.76		12.56±0.77	12.49±0.73		1.06	[0.93; 1.20]	0.368	1.49	[1.20; 1.85]	<0.001	1.06	[0.94; 1.21]	0.329	1.43	[1.16; 1.74]	0.001
ABEP				0.766												
27.79±7.66		27.93±7.71	28.01±7.44		1.01	[0.99; 1.02]	0.340	1.00	[0.98; 1.02]	0.957	1.01	[0.99; 1.02]	0.270	1.00	[0.98; 1.03]	0.547
Parenting style ^a				0.556												
Neglectful	31.19	32.10	32.43													
Indulgent	14.66	13.56	15.68		0.81	[0.60; 1.09]	0.167	0.62	[0.29; 1.36]	0.235	0.86	[0.64; 1.17]	0.349	0.67	[0.31; 1.45]	0.313
Authoritarian	20.69	23.07	17.30		0.98	[0.79; 1.23]	0.891	0.47	[0.26; 0.83]	0.010	0.93	[0.74; 1.16]	0.512	0.47	[0.26; 0.86]	0.014
Authoritative	33.47	31.27	34.59		0.86	[0.65; 1.12]	0.256	0.27	[0.11; 0.67]	0.005	0.83	[0.64; 1.08]	0.165	0.30	[0.12; 0.75]	0.010
Parents alcohol use ^a																
Maternal abstinence	75.98	75.31	74.39	0.311												
Maternal alcohol use	21.30	22.93	23.58		1.56	[1.16; 2.06]	0.001	1.74	[1.04; 2.89]	0.034	1.57	[1.20; 2.07]	0.001	1.96	[0.99; 2.89]	0.053
Maternal drunkenness	2.72	1.76	2.03		1.71	[0.99; 2.98]	0.055	3.75	[1.61; 8.74]	0.002	1.74	[1.02; 2.98]	0.043	3.73	[1.61; 8.60]	0.002
Paternal abstinence	58.10	58.90	56.10	0.732												
Paternal alcohol use	31.87	32.33	34.15		1.61	[1.32; 1.98]	<0.001	0.79	[0.53; 1.17]	0.245	1.59	[1.30; 1.95]	<0.001	0.90	[0.59; 1.35]	0.607
Paternal drunkenness	10.03	8.77	9.76		1.56	[1.15; 2.12]	0.004	0.88	[0.47; 1.63]	0.687	1.59	[1.19; 2.16]	0.002	0.91	[0.48; 1.73]	0.777

^aMissing values of these variables were imputed^bAnalysis controlled for drug use at baseline^cAbstainers/low users was considered the reference class

Table 4 Characteristics of polysubstance drug use and binge drinking identified through latent class analysis in a sample of students participating in the 21-month follow-up (wave 3) of a study to evaluate the effect of a school-based program for drug-use prevention (N = 6391)

	Latent classes, wave 3				p	Univariate analysis ^b				Multivariate analysis ^b							
	Abstainers/low users ^c		Polydrug users			Abstainers/low users vs. alcohol users/binge drinkers		Abstainers/low users vs. polydrug users		Abstainers/low users vs. alcohol users/binge drinkers		Abstainers/low users vs. polydrug users					
	%	%	%	%		cOR	95% CI	p	cOR	95% CI	p	aOR	95% CI	p			
Group																	
Intervention	48.54	49.26	52.48	52.48	0.132	1.27	[0.96; 1.57]	0.927	1.65	[0.88; 1.54]	0.289	1.20	[0.85; 1.70]	0.300	1.51	[0.86; 1.53]	0.334
Control	51.46	50.74	47.52	47.52													
Sex					<0.001												
Girl	49.07	58.02	52.27	52.27		1.70	[1.36; 2.15]	<0.001	1.39	[1.11; 1.74]	0.004	1.73	[1.35; 2.21]	<0.001	1.43	[1.14; 1.80]	0.002
Boy	50.93	41.98	47.73	47.73													
Age					0.000												
ABEP	12.56 ± 0.80	12.71 ± 0.82	12.94 ± 0.93	12.94 ± 0.93		0.96	[0.77; 1.19]	0.711	1.09	[0.87; 1.37]	0.445	0.97	[0.73; 1.21]	0.775	1.07	[0.84; 1.35]	0.580
	27.85 ± 8.07	28.36 ± 7.82	28.68 ± 9.75	28.68 ± 9.75	<0.001	1.01	[1.00; 1.04]	0.035	1.01	[0.10; 1.04]	0.098	1.02	[1.00; 1.04]	0.033	1.02	[0.10; 1.04]	0.061
Parenting style ^a					<0.001												
Neglectful	26.19	33.47	46.90	46.90		0.92	[0.69; 1.23]	0.595	0.74	[0.48; 1.14]	0.173	0.93	[0.69; 1.25]	0.626	0.75	[0.48; 1.18]	0.219
Indulgent	10.50	10.56	8.47	8.47		1.04	[0.76; 1.42]	0.809	0.63	[0.45; 0.88]	0.007	0.97	[0.71; 1.33]	0.845	0.62	[0.44; 0.89]	0.009
Authoritarian	14.73	15.14	17.56	17.56		0.84	[0.60; 1.20]	0.356	0.46	[0.32; 0.68]	<0.001	0.80	[0.56; 1.13]	0.206	0.46	[0.31; 0.69]	<0.001
Authoritative	25.04	15.47	17.56	17.56													
Parents alcohol use ^c																	
Maternal abstinence	80.04	55.40	58.26	58.26	<0.001												
Maternal alcohol use	15.71	34.78	31.40	31.40		1.61	[1.04; 2.51]	0.035	1.59	[1.00; 2.53]	0.051	1.62	[1.01; 2.60]	0.046	1.58	[0.98; 2.56]	0.062
Maternal drunkenness	0.66	6.14	9.30	9.30		2.00	[1.29; 3.10]	0.002	2.72	[1.49; 4.96]	0.001	2.10	[1.31; 3.35]	0.002	2.79	[1.52; 5.12]	0.001
Paternal abstinence	61.90	41.57	45.45	45.45	<0.001												
Paternal alcohol use	27.45	38.22	33.06	33.06		1.39	[1.14; 1.69]	0.001	0.90	[0.68; 1.20]	0.479	1.36	[1.12; 1.65]	0.002	0.94	[0.70; 1.27]	0.694
Paternal drunkenness	6.62	15.96	19.63	19.63		1.24	[0.92; 1.68]	0.152	0.73	[0.44; 1.23]	0.240	1.30	[0.96; 1.76]	0.095	0.76	[0.46; 1.25]	0.282

^aMissing values of these variables were imputed

^bAnalysis controlled for drug use in the baseline

^cAbstainers/low users was considered the reference class

Brazilian adolescents are: alcohol: 13.0 y.o. (IC: 12.9–13.1); tobacco 13.3 y.o. (IC: 13.2–13.4); inhalants: 13.2 y.o. (IC: 13.1–13.4); and marijuana: 14.6 y.o. (IC: 14.4–14.7) [48]. Collecting the data before the occurrence of the outcomes we would be more able to understand the casual relationships related to them.

The results related to the best solution for the latent classes were in line with previous studies carried out in other countries, which also found the same three latent classes of drug use in adolescents as the best model to describe their patterns of drug use [58, 59].

With respect to adolescents in the class of “alcohol users/binge drinkers”, our findings indicate that any parental alcohol use (episodic or drunkenness) can act as a predictor of belonging to this class, whether it is the mother or the father who drinks. Our results confirm earlier findings that parental alcohol use is predictive of an early age of first alcohol use [15]. Furthermore, this finding adds to the existing evidence linking parental alcohol use and adolescent-onset drinking in that it suggests the existence of a causal association even after controlling for important covariates, such as parenting style. Our results suggest that adolescents’ alcohol use is partially shaped by their parents’ drinking behavior, as social cognitive/learning theory [60] suggests. Parents directly model drinking behavior for their adolescent children [19], and even positive expectations surrounding alcohol use are transmitted from parents to their children [61], enhancing adolescents’ positive views of drinking and subsequent likelihood of initiating use [62]. Another possible explanation is that parents who drink tend to be more permissive about their children’s alcohol use [22] and may facilitate their teenage children’s access to drinking [18]. In addition, genetic susceptibility plays an important role in the relation between parental drinking problems and the alcohol use behavior of their adolescent offspring [63].

This longitudinal analysis identified that the authoritative and authoritarian parenting styles play an important role as protective factors against membership in the “polydrug users” class (at wave 2 and wave 3). Our results, showing that authoritative parenting style is a protective factor and that the neglectful style is a risk factor, are consistent with previous studies [27, 28]. However, the association between authoritative parenting styles and drug use is still controversial and may vary from culture to culture [34, 64]. Our results highlight the importance of parental monitoring [36] in the prevention of adolescent drug use because both styles (authoritative and authoritarian) that exercise this function are associated with a decreased probability that the adolescent will belong to the “polydrug users” class. Parental monitoring involves a set of behaviors related to providing attention to, remain informed about child’s use of free time and also tracking child’s whereabouts, activities and friends [15, 65].

Findings from this study point to maternal episodes of drunkenness as a strong risk factor, predicting adolescents’ likelihood of belonging to the “polydrug users” and “alcohol users/binge drinkers” classes in both waves (wave 2 and wave 3). Therefore, we can suggest that maternal drunkenness is a key element in the risk of adolescent alcohol consumption and other drug use. Previous studies determined the same predictive relationship with adolescent alcohol use [22, 66] but a few studies found an association between parents’ problem use of alcohol and adolescent drug use [23]. Two main pathways for the effect of mothers’ episodes of drunkenness on adolescent drug use can be hypothesized. The first possible explanation is a biological pathway through a genetic factor, while the second pathway is through the well-known role of maternal influences on child development [63]. Problematic maternal substance use may model adolescent drug use by promoting the perception of low levels of risk associated with drug use [67]. Furthermore, exposure of a child to problematic patterns of maternal alcohol use, such as those resulting in noticeable intoxication, is highly harmful to the child’s emotional development, leading to premature involvement in licit and even illicit drug use [68–70].

Our findings highlight that health approaches should target parenting skills and parenting alcohol use to prevent adolescent drug use. The effects of school-based universal programs may be increased by adding parent-based components [71]. The combined student–parent intervention showed substantial effects on the prevention of alcohol [72–74] and other drugs [75]. According to a systematic review, a central focus in all successful programs was on monitoring the children’s activities, however, less is known about the exclusive effect of targeting parenting alcohol use [76].

Another important consideration is that the drug prevention program #Tamojunto had no effect on adolescents’ past-year drug use at 9-month or 21-month follow-up; however, we retained this variable in the analysis to control for any possible confounding effect, since this study is a randomized controlled trial.

This study has some limitations that should be considered. First, we only collected measures provided by the adolescents; thus, we assessed only the adolescents’ perceptions of parenting style and parental alcohol use. However, it is common practice to use adolescents’ perceptions of parenting behaviors as a categorical observed covariate, as we did [27, 64]. Studies that assess the perceptions of both parents and children simultaneously tend to provide more reliable data on parenting styles and drug use, as children tend to have a more negative perception than parents regarding the parent–child relationship [77]. Second, we did not collect data on parents’ illicit drug use because it could be unethical to ask adolescents to report their parents’ illicit behavior. Third, the attrition

rate due to the losses over the time, leading to a lack of information about non-respondents. However, it is worth noting that the absence of some data is an expected limitation in longitudinal studies, especially those with long follow-up periods [78–80]. However, imputation processes offer excellent solutions to these missing-data problems by estimating the missing values [81]. Another limitation that must be considered is that we analyzed data only from adolescents aged 11–15 years old and how they are able to answering the questions. Finally, we must consider that we analyzed data only from adolescents aged 11–15 years old and how they are able to answering the questions. Finally, this study correlation of parenting dimension with adolescent substance use cannot exclusively be interpreted as effects of parenting on the child outcome, once associations are in most cases bidirectional. Although it is common practice to use adolescents' perceptions of parenting behaviors as an observed covariate, this rating by the children limits the possibility to draw conclusions. There remains a possibility that the perception of the parents is shaped by the drug using style of their children.

In conclusion, this study adds to the existing literature suggesting that parents' behavior is an important predictor of latent classes of drug use by adolescents. First, parenting style, especially monitoring, is a strong predictor for the prevention of polydrug use among adolescents. Second, occasional parental alcohol use can act as a central predictor for adolescent alcohol use and binge drinking. Moreover, maternal drunkenness is involved in predictive models for both drug use classes in both waves. Regarding the prevention of adolescent drug use, our findings highlight the importance of comprehensive public health approaches that target parenting skills, especially monitoring, while simultaneously targeting parental alcohol use. To educate parents to be more aware of their role in the prevention of adolescent drug use, school-based prevention programs should also target parenting skills and parental alcohol use, giving special attention to the influence of problematic patterns of maternal drinking on adolescents' risk of drug use.

Acknowledgements We acknowledge Brazilian Health Ministry and UNODC Brazil (United Nations Office on Drugs and Crime) for initiative to implement this prevention program and support this research. This study was funded by the Brazilian Ministry of Health through the TED 89-2014 (PI: Dr. Sanchez) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) through the grant number 2016/11971-5 (M.S. Juliana Y Valente). Special thanks are due to the Brazilian schools for their continuing collaborative efforts and teachers, students and parents who were involved in the study.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflicts of interest.

Human participants and/or animals All procedures in the present study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. This study was approved by the Ethics in Research Committees at the University of São Paulo (#473.498) and the Federal University of Santa Catarina (#711.377).

References

1. IBGE (2016) Pesquisa Nacional de Saúde do Escolar 2015. IBGE, Coordenação de População e Indicadores Sociais, Rio de Janeiro
2. Kim MJ, Mason WA, Herrenkohl TI et al (2017) Influence of early onset of alcohol use on the development of adolescent alcohol problems: a longitudinal binational study. *Prev Sci* 18:1–11. <https://doi.org/10.1007/s11121-016-0710-z>
3. Heron J, Macleod J, Munafò MR et al (2012) Patterns of alcohol use in early adolescence predict problem use at age 16. *Alcohol Alcohol* 47:169–177. <https://doi.org/10.1093/alcalc/agr156>
4. Bonomo Y, Wolfe R, Lynskey M et al (2001) Adverse outcomes of alcohol use in adolescents. *Addiction* 96:1485–1496
5. Squeglia LM, Gray KM (2016) Alcohol and drug use and the developing brain. *Curr Psychiatry Rep* 18:46. <https://doi.org/10.1007/s11920-016-0689-y>
6. Wittchen H-U, Fröhlich C, Behrendt S et al (2007) Cannabis use and cannabis use disorders and their relationship to mental disorders: a 10-year prospective-longitudinal community study in adolescents. *Drug Alcohol Depend* 88:S60–S70. <https://doi.org/10.1016/j.drugalcdep.2006.12.013>
7. Camchong J, Lim KO, Kumra S (2016) Adverse effects of cannabis on adolescent brain development: a longitudinal study. *Cereb Cortex*. <https://doi.org/10.1093/cercor/bhw015>
8. Bonomo YA, Bowes G, Coffey C et al (2004) Teenage drinking and the onset of alcohol dependence: a cohort study over seven years. *Addiction* 99:1520–1528. <https://doi.org/10.1111/j.1360-0443.2004.00846.x>
9. Liang W, Chikritzhs T (2015) Age at first use of alcohol predicts the risk of heavy alcohol use in early adulthood: a longitudinal study in the United States. *Int J Drug Policy* 26:131–134. <https://doi.org/10.1016/j.drugpo.2014.07.001>
10. Moss HB, Chen CM, Yi H ye (2014) Early adolescent patterns of alcohol, cigarettes, and marijuana polysubstance use and young adult substance use outcomes in a nationally representative sample. *Drug Alcohol Depend* 136:51–62. <https://doi.org/10.1016/j.drugalcdep.2013.12.011>
11. Buchmann AF, Schmid B, Blomeyer D et al (2009) Impact of age at first drink on vulnerability to alcohol-related problems: testing the marker hypothesis in a prospective study of young adults. *J Psychiatr Res* 43:1205–1212. <https://doi.org/10.1016/j.jpsychires.2009.02.006>
12. Grant BF, Stinson FS, Harford TC (2001) Age at onset of alcohol use and DSM-IV alcohol abuse and dependence: a 12-year follow-up. *J Subst Abuse* 13:493–504. [https://doi.org/10.1016/S0899-3289\(01\)00096-7](https://doi.org/10.1016/S0899-3289(01)00096-7)
13. Rehm J, Mathers C, Popova S et al (2009) Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet* 373:2223–2233. [https://doi.org/10.1016/S0140-6736\(09\)60746-7](https://doi.org/10.1016/S0140-6736(09)60746-7)
14. Degenhardt L, Stockings E, Patton G et al (2016) The increasing global health priority of substance use in young people. *Lancet Psychiatry* 3:251–264. [https://doi.org/10.1016/S2215-0366\(15\)00508-8](https://doi.org/10.1016/S2215-0366(15)00508-8)

15. Ryan SM, Jorm AF, Lubman DI (2010) Parenting factors associated with reduced adolescent alcohol use: a systematic review of longitudinal studies. *Aust N Z J Psychiatry* 44:774–783. <https://doi.org/10.1080/00048674.2010.501759>
16. Yap MBH, Cheong TWK, Zaravinos-Tsakos F et al (2017) Modifiable parenting factors associated with adolescent alcohol misuse: a systematic review and meta-analysis of longitudinal studies. *Addiction* 112:1142–1162. <https://doi.org/10.1111/add.13785>
17. Grigsby TJ, Forster M, Unger JB, Sussman S (2016) Predictors of alcohol-related negative consequences in adolescents: a systematic review of the literature and implications for future research. *J Adolesc* 48:18–35. <https://doi.org/10.1016/j.adolescence.2016.01.006>
18. Alati R, Najman JM, Kinner SA et al (2005) Early predictors of adult drinking: a birth cohort study. *Am J Epidemiol* 162:1098–1107. <https://doi.org/10.1093/aje/kwi320>
19. Vermeulen-Smit E, Koning IM, Verdurmen JEE et al (2012) The influence of paternal and maternal drinking patterns within two-partner families on the initiation and development of adolescent drinking. *Addict Behav* 37:1248–1256. <https://doi.org/10.1016/j.addbeh.2012.06.005>
20. Mares SHW, van der Vorst H, Engels RCME, Lichtwarck-Aschoff A (2011) Parental alcohol use, alcohol-related problems, and alcohol-specific attitudes, alcohol-specific communication, and adolescent excessive alcohol use and alcohol-related problems: an indirect path model. *Addict Behav* 36:209–216. <https://doi.org/10.1016/j.addbeh.2010.10.013>
21. Rossow I, Keating P, Felix L, McCambridge J (2016) Does parental drinking influence children's drinking? A systematic review of prospective cohort studies. *Addiction* 111:204–217. <https://doi.org/10.1111/add.13097>
22. van der Zwaluw CS, Scholte RHJ, Vermulst AA et al (2008) Parental problem drinking, parenting, and adolescent alcohol use. *J Behav Med* 31:189–200. <https://doi.org/10.1007/s10865-007-9146-z>
23. Kilpatrick DG, Acierno R, Saunders B et al (2000) Risk factors for adolescent substance abuse and dependence: data from a national sample. *J Consult Clin Psychol* 68:19–30. <https://doi.org/10.1037//0022-006X.68.1.19>
24. Li C, Pentz MA, Chou C-P (2002) Parental substance use as a modifier of adolescent substance use risk. *Addiction* 97:1537–1550. <https://doi.org/10.1046/j.1360-0443.2002.00238.x>
25. Hussong AM, Huang W, Serrano D et al (2012) Testing whether and when parent alcoholism uniquely affects various forms of adolescent substance use. *J Abnorm Child Psychol* 40:1265–1276. <https://doi.org/10.1007/s10802-012-9662-3>
26. Čablová L, Pazderková K, Miovský M (2014) Parenting styles and alcohol use among children and adolescents: a systematic review. *Drugs Educ Prev Policy* 21:1–13. <https://doi.org/10.3109/09687637.2013.817536>
27. Berge J, Sundell K, Ojehagen A, Hakansson A (2016) Role of parenting styles in adolescent substance use: results from a Swedish longitudinal cohort study. *BMJ Open* 6:e008979. <https://doi.org/10.1136/bmjopen-2015-008979>
28. Chassin L, Presson CC, Rose J et al (2005) Parenting style and smoking-specific parenting practices as predictors of adolescent smoking onset. *J Pediatr Psychol* 30:333–344. <https://doi.org/10.1093/jpepsy/jsi028>
29. Calafat A, García F, Juan M et al (2014) Which parenting style is more protective against adolescent substance use? Evidence within the European context. *Drug Alcohol Depend* 138:185–192. <https://doi.org/10.1016/j.drugalcdep.2014.02.705>
30. Cerezo F, Méndez I, Ato M (2013) Moderating role of family and friends' factors between disocial behavior and consumption in adolescents. *Int J Clin Heal Psychol* 13:171–180. [https://doi.org/10.1016/S1697-2600\(13\)70021-8](https://doi.org/10.1016/S1697-2600(13)70021-8)
31. Hoffmann JP, Bahr SJ (2014) Parenting style, religiosity, peer alcohol use, and adolescent heavy drinking. *J Stud Alcohol Drugs* 75:222–227
32. Becoña E, Martínez Ú, Calafat A et al (2012) Parental styles and drug use: a review. *Drugs Educ Prev Policy* 19:1–10. <https://doi.org/10.3109/09687637.2011.631060>
33. Lanza ST, Patrick ME, Maggs JL (2010) Latent transition analysis: benefits of a latent variable approach to modeling transitions in substance use. *J Drug Issues* 40:93–120. <https://doi.org/10.1177/002204261004000106>
34. Valente JY, Cogo-Moreira H, Sanchez ZM (2017) Gradient of association between parenting styles and patterns of drug use in adolescence: a latent class analysis. *Drug Alcohol Depend* 180:272–278. <https://doi.org/10.1016/j.drugalcdep.2017.08.015>
35. Cranford JA, McCabe SE, Boyd CJ (2013) Adolescents' non-medical use and excessive medical use of prescription medications and the identification of substance use subgroups. *Addict Behav* 38:2768–2771. <https://doi.org/10.1016/j.addbeh.2013.06.015>
36. Shin Y, Lee JK, Lu Y, Hecht ML (2015) Exploring parental influence on the progression of alcohol use in Mexican-heritage youth: a latent transition analysis. *Prev Sci* 17:188–198. <https://doi.org/10.1007/s11121-015-0596-1>
37. Snyder SM, Smith RE (2015) Do physical abuse, depression, and parental substance use influence patterns of substance use among child welfare involved youth? *Subst Use Misuse* 50:226–235. <https://doi.org/10.3109/10826084.2014.966845>
38. Collins LM, Lanza ST (2009) Latent class and latent transition analysis: with applications in the social, behavioral, and health sciences. Wiley, New York
39. Lazarsfeld P, Henry N (1968) Latent structure analysis. Houghton Mifflin, Boston
40. Lanza ST, Rhoades BL (2013) Latent class analysis: an alternative perspective on subgroup analysis in prevention and treatment. *Prev Sci* 14:157–168. <https://doi.org/10.1007/s11211-011-0201-1>
41. Cho SB, Llana DC, Adkins AE et al (2015) Patterns of substance use across the first year of college and associated risk factors. *Front Psychiatry*. <https://doi.org/10.3389/fpsy.2015.00152>
42. Anderson KG, Ramo DE, Cummins KM, Brown SA (2010) Alcohol and drug involvement after adolescent treatment and functioning during emerging adulthood. *Drug Alcohol Depend* 107:171–181. <https://doi.org/10.1016/j.drugalcdep.2009.10.005>
43. Scheier LM, Abdallah AB, Inciardi JA et al (2008) Tri-city study of Ecstasy use problems: a latent class analysis. *Drug Alcohol Depend* 98:249–263. <https://doi.org/10.1016/j.drugalcdep.2008.06.008>
44. Pedrosa RT, Abreu S, Kinoshita RT (2015) Aprendizagens da intersectorialidade entre saúde e educação na prevenção do uso de álcool e outras drogas. *Textura* 33:9–24
45. Sanchez ZM, Valente JY, Sanudo A et al (2017) The #Tamojuntó drug prevention program in Brazilian schools: a randomized controlled trial. *Prev Sci* 18:772–782. <https://doi.org/10.1007/s11211-017-0770-8>
46. Lwanga SK, Lemeshow S (1991) Sample size determination in health studies: a practical manual. World Health Organization, Geneva
47. Faggiano F, Vigna-Taglianti FD, Versino E et al (2008) School-based prevention for illicit drugs use: a systematic review. *Prev Med (Baltim)* 46:385–396. <https://doi.org/10.1016/j.ypmed.2007.11.012>
48. Carlini EL, de A, Noto, AR, Sanchez Z van der M, et al (2010) VI Levantamento nacional sobre o consumo de drogas psicotrópicas entre estudantes do ensino fundamental e médio das redes pública e privada de ensino nas 27 capitais brasileiras. SENAD–Secretaria Nacional de Políticas sobre Drogas, Brasília

49. IBGE (2013) Pesquisa Nacional da Saúde do Escolar 2012 (PeNSE). Instituto Brasileiro de Geografia e Estatística–IBGE, Rio de Janeiro
50. ABEP (2012) Critério de Classificação Econômica do Brasil [Criteria for Economic Classification in Brazil]. In: Ibope. <http://www.abep.org/criterio-brasil>. Accessed 15 Feb 2017
51. Lamborn SD, Mounts NS, Steinberg L, Dornbusch SM (1991) Patterns of competence and adjustment among adolescents from authoritative, authoritarian, indulgent, and neglectful families. *Child Dev* 62:1049–1065. <https://doi.org/10.1111/j.1467-8624.1991.tb01588.x>
52. Maccoby E, Martin JA (1983) Socialization in the context of the family: parent-child interaction. In: Hetherington EM (ed) Socialization, personality, and social development. Wiley, New York, pp 1–101
53. Asparouhov T (2006) General multi-level modeling with sampling weights. *Commun Stat Theory Methods* 35:439–460. <https://doi.org/10.1080/03610920500476598>
54. Muthén L, Muthén BO (2010) Mplus statistical analysis with latent variables user's guide (Sixth Edition)
55. Hosmer DW, Lemeshow S, Sturdivant RX (2013) Applied logistic regression. Wiley, New York
56. Asparouhov T, Muthén B (2014) Auxiliary variables in mixture modeling: three-step approaches using M plus. *Struct Equ Model A Multidiscip J*. <https://doi.org/10.1080/10705511.2014.915181>
57. Nylund KL, Asparouhov T, Muthén BO (2007) Deciding on the number of classes in latent class analysis and growth mixture modeling: a Monte Carlo simulation study. *Struct Equ Model A Multidiscip J* 14:535–569. <https://doi.org/10.1080/10705510701575396>
58. Chung T, Kim KH, Hipwell AE, Stepp SD (2013) White and black adolescent females differ in profiles and longitudinal patterns of alcohol, cigarette, and marijuana use. *Psychol Addict Behav* 27:1110–1121. <https://doi.org/10.1037/a0031173>
59. Kelly AB, Evans-Whipp TJ, Smith R et al (2015) A longitudinal study of the association of adolescent polydrug use, alcohol use and high school non-completion. *Addiction* 110:627–635. <https://doi.org/10.1111/add.12829>
60. Bandura A (1977) Social learning theory. Pearson Education, US
61. Jones BT, Corbin W, Fromme K (2001) Conceptualizing addiction: a review of expectancy theory and alcohol consumption. *Addiction* 96:57–72. <https://doi.org/10.1080/09652140020016969>
62. Handley ED, Chassin L (2013) Alcohol-specific parenting as a mechanism of parental drinking and alcohol use disorder risk on adolescent alcohol use onset. *J Stud Alcohol Drugs* 74:684–693. <https://doi.org/10.15288/jsad.2013.74.684>
63. King SM, Keyes M, Malone SM et al (2009) Parental alcohol dependence and the transmission of adolescent behavioral disinhibition: a study of adoptive and non-adoptive families. *Addiction* 104:578–586. <https://doi.org/10.1111/j.1360-0443.2008.02469.x>
64. García F, Gracia E (2009) Is always authoritative the optimum parenting style? Evidence from Spanish families. *Adolescence* 44:101–131
65. Dishion TJ, McMahon RJ (1998) Parental monitoring and the prevention of child and adolescent problem behavior: a conceptual and empirical formulation. *Clin Child Fam Psychol Rev* 1:61–75. <https://doi.org/10.1023/A:1021800432380>
66. Casswell S, Pledger M, Pratap S (2002) Trajectories of drinking from 18 to 26 years: identification and prediction. *Addiction* 97:1427–1437. <https://doi.org/10.1046/j.1360-0443.2002.00220.x>
67. Patrick ME, Maggs JL, Greene KM et al (2014) The link between mother and adolescent substance use: inter-generational findings from the British Cohort Study. *Longit Life Course Stud* 5:56–63. <https://doi.org/10.14301/llics.v5i1.241>
68. Knudsen AK, Ystrom E, Skogen JC, Torgersen L (2015) Maternal heavy alcohol use and toddler behavior problems: a fixed effects regression analysis. *Eur Child Adolesc Psychiatry* 24:1269–1277. <https://doi.org/10.1007/s00787-015-0677-5>
69. Kerr DCR, Capaldi DM, Pears KC, Owen LD (2012) Intergenerational influences on early alcohol use: independence from the problem behavior pathway. *Dev Psychopathol* 24:889–906. <https://doi.org/10.1017/S0954579412000430>
70. Berg L, Bäck K, Vinnerljung B, Hjern A (2016) Parental alcohol-related disorders and school performance in 16-year-olds—a Swedish national cohort study. *Addiction* 111:1795–1803. <https://doi.org/10.1111/add.13454>
71. Newton NC, Champion KE, Slade T et al (2017) A systematic review of combined student- and parent-based programs to prevent alcohol and other drug use among adolescents. *Drug Alcohol Rev* 36:337–351. <https://doi.org/10.1111/dar.12497>
72. Koning IM, Vollebergh WAM, Smit F et al (2009) Preventing heavy alcohol use in adolescents (PAS): cluster randomized trial of a parent and student intervention offered separately and simultaneously. *Addiction* 104:1669–1678. <https://doi.org/10.1111/j.1360-0443.2009.02677.x>
73. Koning IM, Van Den Eijnden RJM, Verdurmen JEE et al (2013) A cluster randomized trial on the effects of a parent and student intervention on alcohol use in adolescents four years after baseline; no evidence of catching-up behavior. *Addict Behav* 38:2032–2039. <https://doi.org/10.1016/j.addbeh.2012.12.013>
74. Koutakis N, Stattin HH, Kerr M (2008) Reducing youth alcohol drinking through a parent-targeted intervention: the Orebro Prevention Program. *Addiction* 103:1629–1637. <https://doi.org/10.1111/j.1360-0443.2008.02326.x>
75. Stormshak EA, Connell AM, Véronneau M-H et al (2011) An ecological approach to promoting early adolescent mental health and social adaptation: family-centered intervention in public middle schools. *Child Dev* 82:209–225. <https://doi.org/10.1111/j.1467-8624.2010.01551.x>
76. Kuntsche S, Kuntsche E (2016) Parent-based interventions for preventing or reducing adolescent substance use—a systematic literature review. *Clin Psychol Rev* 45:89–101. <https://doi.org/10.1016/j.cpr.2016.02.004>
77. Shek DTL (1998) A longitudinal study of Hong Kong adolescents' and parents' perceptions of family functioning and well-being. *J Genet Psychol* 159:389–403. <https://doi.org/10.1080/00221329809596160>
78. Ariza C, Pérez A, Sánchez-Martínez F et al (2013) Evaluation of the effectiveness of a school-based cannabis prevention program. *Drug Alcohol Depend* 132:257–264. <https://doi.org/10.1016/j.drugalcdep.2013.02.012>
79. Newton NC, Teesson M, Vogl LE, Andrews G (2010) Internet-based prevention for alcohol and cannabis use: final results of the climate schools course. *Addiction* 105:749–759. <https://doi.org/10.1111/j.1360-0443.2009.02853.x>
80. Shope JT, Dielman TE, Butchart AT et al (1992) An elementary school-based alcohol misuse prevention program: a follow-up evaluation. *J Stud Alcohol* 53:106–121. <https://doi.org/10.15288/jsa.1992.53.106>
81. Dong Y, Peng C-YJ (2013) Principled missing data methods for researchers. *Springerplus* 2:222. <https://doi.org/10.1186/2193-1801-2-222>