

## Original article

## Sociodemographic and lifestyle factors associated with the neutrophil-to-lymphocyte ratio

Rachel Howard, PhD<sup>\*</sup>, Aaron Scheiner, Peter A. Kanetsky, PhD, Kathleen M. Egan, PhD

Department of Cancer Epidemiology, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL

## ARTICLE INFO

## Article history:

Received 1 April 2019

Accepted 24 July 2019

Available online 1 August 2019

## Keywords:

Inflammation

Biomarkers

Survival

Neutrophil-to-lymphocyte ratio

## ABSTRACT

**Purpose:** The neutrophil-to-lymphocyte ratio (NLR) is a marker of systemic inflammation with established prognostic value in patients with cancer. Although high NLR is associated with poorer clinical outcomes, factors that influence the magnitude of NLR independently of disease are poorly understood. **Methods:** We identified 48,023 adults who participated in the National Health and Nutrition Examination Survey (1999–2016). Demographic, socioeconomic, and lifestyle factors associated with the magnitude of NLR after adjusting for comorbidities including heart disease, cancer, diabetes, and hypertension, and medications including aspirin, were identified. Effect modification by comorbidity status and demographics was explored.

**Results:** Female gender, age less than 60 years, and non-Hispanic black race/ethnicity were associated with lower NLR. Marital statuses of widowed, separated, or never married demonstrated increased NLR as compared with those who were currently married. Never-smoking and moderate alcohol consumption were associated with lower NLR. Participation in physical activity was associated with decreased NLR after adjustment for potential confounders, primarily among non-Hispanic whites.

**Conclusions:** Multiple demographic and lifestyle factors are independently associated with NLR. Sex, age, race, marital status, body mass index, physical activity, smoking history, and alcohol consumption should all be routinely collected and adjusted for to improve the accuracy of assessment of the prognostic power of NLR.

© 2019 Elsevier Inc. All rights reserved.

## Introduction

Acute and chronic inflammation plays an appreciable role in a wide range of medical conditions, including cancer [1–6]. This has led to increased interest in identifying markers of systemic inflammation that can serve as a clinically useful proxy for the overall immune status of the individual. Such markers may have the potential to facilitate risk stratification at diagnosis and guide therapeutic selection.

Alterations in circulating white blood cells accompany systemic inflammation. These alterations include the development of neutrophilia and concurrent relative lymphocytopenia [7], both of which are detectable in the routine complete blood count (CBC).

The neutrophil-to-lymphocyte ratio (NLR) captures the balance between the detrimental effects of neutrophilia and the benefits of an active adaptive immune response and is prognostic of patient outcomes across multiple diseases [8–11].

Most published association studies focus on a baseline NLR measurement obtained when the individual is diagnosed with, or exhibiting symptoms of, disease. Although several additional studies have sought to characterize the range of “normal” NLR in a healthy population [12], little is known about factors that influence the magnitude of NLR independently of disease status. For example, although average NLR differs between subgroups stratified by race [13] and exercise can lower NLR in certain individuals [14], few studies have examined demographic, socioeconomic, and lifestyle correlates of NLR in the general population. Such studies are useful for identifying potential confounders of the strength of association between baseline NLR and disease outcomes. In addition, consideration of lifestyle factors may lead to the identification of targets for behavioral intervention leading to reduced systemic inflammation and potentially impacting the burden of disease in the individual.

Declarations of interest: None.

<sup>\*</sup> Corresponding author. Department of Cancer Epidemiology, H. Lee Moffitt Cancer Center and Research Institute, 12902 USF Magnolia Drive, Tampa, FL, 33612. Tel.: (+1) 813-745-4651; fax: (+1) 813-745-8332.

E-mail address: [rachel.howard@moffitt.org](mailto:rachel.howard@moffitt.org) (R. Howard).

<https://doi.org/10.1016/j.annepidem.2019.07.015>

1047-2797/© 2019 Elsevier Inc. All rights reserved.

The objective of this study was to systematically evaluate sociodemographic and lifestyle influences on the magnitude of NLR after adjustment for potential confounding by comorbidities and medications. The study was based on 48,023 adult participants in the U.S. National Health and Nutrition Examination Survey (NHANES) from 1999 to 2016 for whom survey data and a CBC for NLR determination were available.

## Methods

### Study population

The NHANES aims to assess the health status of adults and children across the United States by obtaining representative samples of the population using multistage sampling [15]. We exported NHANES laboratory and questionnaire data from the 9 surveys between 1999 and 2016 (1999–00, 2001–02, 2003–04, 2005–06, 2007–08, 2009–10, 2011–12, 2013–14, and 2015–16) for all participants aged 18 years or older, resulting in 48,023 unique participants to include in the study (a CONSORT flow chart can be found in [Supplementary Fig. 1](#)).

### Laboratory data

CBC was available for all adults who participated in the NHANES between 1999 and 2016. CBCs for all years were conducted using the Coulter Unicel DxH 800 analyzer, a quantitative, automated hematology analyzer for *in vitro* diagnostic use. The analyzer provided the leukocyte 5-part differential on whole blood, which includes absolute numbers of neutrophils and lymphocytes and their relative percentage of total white blood cells. Samples were run in duplicate, and the average of two measurements was reported. Absolute numbers of lymphocytes and neutrophils were reported in units of 1000 cells per microliter. Full details on the laboratory methods relating to the CBC and components can be found in [16]. Ratios of neutrophils to lymphocytes were calculated for both percentage measures (percent of total white blood cell count) and absolute numbers. Correlation between the two measures was  $R^2 = 0.99$  ([Supplementary Fig. 2](#)), the former was arbitrarily selected for presentation of results. Variation in neutrophil and lymphocyte counts and percentages between survey years were minimal ([Supplementary Fig. 3](#)). The distribution of NLR and its respective components across all included participants can be found in [Supplementary Figure 4](#).

### Questionnaire data

Age, race, and sex were available for all participants. Factors relating to socioeconomic status including education level, marital status, household size, income, poverty income ratio (ratio of family income to poverty threshold), and health insurance were extracted for analysis. Indices relating to intensity and duration of physical activity, as well as ever-smoking status, alcohol use, body mass index (BMI), and diet and nutrition (vegetable consumption, fast food) were also included. We also obtained information on pre-existing health conditions including mental health diagnoses (depression, panic, generalized anxiety disorder), arthritis, heart disease, emphysema, cancer, diabetes, hypertension, and the use of aspirin. The 19 demographic, socioeconomic, and lifestyle factors and additional 13 factors relating to comorbidities and medications are listed in [Table 1](#). Shown are the NHANES variable name, description, and years the variable was included in the survey. For many factors collected as continuous data, responses were clustered rather than distributed uniformly across the available range. As an example, in questions relating to time in minutes, most

participants entered multiples of 15 or 30 minutes with sparse data for other responses. In these cases, we created discrete response categories ([Table 1](#)) to permit more intuitive interpretation of regression results. Where the variable identifier changed between the years of 1999 and 2016, both identifiers are listed. All socioeconomic, lifestyle, and health-related questions included an additional response category for “refused to answer” or “don’t know.” Only BMI was obtained from a physical examination. The remaining variables were collected via questionnaire or interview (self-report).

### Statistical methods

We initially conducted descriptive univariate analyses comparing median NLR across levels of each respective factor, for example, males versus females and age under 60 versus 60 years or older. For factors with two levels, Mann Whitney *U* tests were performed to identify differences in NLR. For variables with more than two levels, the Kruskal-Wallis omnibus test was performed. Where a monotonic increase or decrease in a statistic (median, regression coefficient) was observed across ordinal levels of a factor, *P* values for trend are provided based on the Jonckheere-Terpstra statistic for ordered differences across levels of the variate [17].

Regression analyses were conducted to further evaluate relationships between NLR and sociodemographic and lifestyle factors as well as comorbid conditions factors. All available factors were assessed in univariate regression models. Where several factors addressed the same research question—for example, multiple separate factors relating to aspects of physical activity—the factor with the lowest *P*-value(s) in univariate analysis was included in multivariable modeling. Owing to inherent challenges in imputing missing values on such a large scale [18] and with no overlap in participants between years, only participants with complete data were included in multivariable models.

With the exception of vegetable intake, fast food, depression, generalized anxiety disorder, panic disorder, and aspirin use (collected in only a subset of survey years), all studied factors were included in a primary multivariable model encompassing a total of 20,237 participants with complete data. Where univariate analysis of these six factors omitted from the primary model produced significant results, independent multivariable analyses were conducted including all the factors included in the primary model.

We tested for interactions between demographic risk factors (age, sex, and race) by incorporating interaction terms in the primary multivariable model. We then conducted further subgroup analysis by repeating multivariable analyses in independent subsets of participants stratified by comorbidity status (diagnoses of arthritis, heart disease, cancer, diabetes, hypertension, or emphysema), race, sex, and age.

All statistical tests were two-sided. All statistical analyses were performed using R, version 3.3.2 (R core development team, Vienna, Austria).

## Results

Median NLR across levels of all sociodemographic, lifestyle, and comorbidity factors are presented in [Table 2](#), [Table 3](#), and [Supplementary Table 1](#), respectively. [Table 4](#) presents univariate regression results for all factors included in 5 or more survey years. All factors listed in [Table 4](#) are included in the primary multivariable model; 20,237 participants had complete data for these factors and thus were included in this primary multivariable model. [Supplementary Table 2](#) presents univariate regression results for all factors included in only a small subset of survey years. These factors

**Table 1**

Summary of 32 included sociodemographic, lifestyle, and health factors extracted from NHANES, including NHANES identifier, factor description, discrete response categories, and years included in the study

Subject	Factor	NHANES identifier	Description	Discrete response categories	Survey years
Demographic	Sex	RIAGENDR	Gender	Male, female	1999–2016
	Age	RIDAGEYR	Age at screening	>60, ≥60*	1999–2016
	Race	RIDRETH1	Race/ethnicity	Non-Hispanic white, non-Hispanic black, Mexican American, other Hispanic, other race	1999–2016
Socioeconomic	Education level	DMDEDUC2	What is the highest grade or level of school you have completed or the highest degree you have received?	Less than 12th grade, high school/GED or equivalent, college graduate or above	1999–2016
	Marital status	DMDMARTL	Marital status	Married, widowed, divorced, separated, never married, living with partner	1999–2016
	Household size	DMDHHSIZ	Total number of people in the household	1, 2, 3, 4, 5, 6, 7+	1999–2016
	Income	INDHHINC/INDHHIN2†	Annual household income	< \$20,000, > \$20,000 and ≤ \$44,999, > \$45,000 and ≤ \$74,999, ≥ \$75,000	1999–2016
Lifestyle	Poverty income ratio	INDFMPIR	Poverty income ratio (PIR)	<1, ≥1	1999–2016
	Health insurance	HID010/HIQ011†	Are you covered by health insurance or some other kind of health care plan?	Yes, no	1999–2016
	Ever smoker	SMQ020	Have you smoked at least 100 cigarettes in your entire life?	Yes, no	1999–2016
	Alcohol use I	ALQ120Q/ALQ120U†	In the past 12 mo, how often did you drink any type of alcoholic beverage? (d)	0, >0 and ≤10, >10 and ≤50, >50 and ≤100, >100*	1999–2016
	Alcohol use II	ALQ130	In the past 12 mo, on those days that you drank alcoholic beverages, on average, how many drinks did you have?	1, 2 or 3, 4 or 5, >5*	1999–2016
	BMI	BMXBMI	Body mass index (kg/m <sup>2</sup> )	≤18.5, >18.5 and ≤25, >25 and ≤30, >30*	1999–2016
	Vigorous activity I	PAQ650	Do you do any vigorous-intensity sports, fitness, or recreational activities that cause large increases in breathing or heart rate like running or basketball for at least 10 min continuously?	Yes, no	2007–2016
	Vigorous activity II	PAD660	How much time do you spend doing vigorous-intensity sports, fitness, or recreational activities on a typical day? (h)	≤1, >1 and ≤2, >2 and ≤4, >4*	2007–2016
	Moderate activity I	PAD665	Do you do any moderate-intensity sports, fitness, or recreational activities that cause a small increase in breathing or heart rate such as brisk walking, bicycling, swimming, or golf for at least 10 min continuously?	Yes, no	2007–2016
	Moderate activity II	PAD675	How much time do you spend doing moderate-intensity sports, fitness or recreational activities on a typical day? (h)	≤1, >1 and ≤2, >2 and ≤4, >4*	2007–2016
Comorbidities and medications	Time sitting/reclining	PAD680	How much time do you usually spend sitting or reclining on a typical day? (h)	≤1, >1 and ≤2, >2 and ≤4, >4*	2007–2016
	Vegetable intake	DBD270D	On an average day, how many helpings of the following kinds of foods do you eat? Vegetables, including vegetable salads.	0, 1 or 2, 3 or 4, >5*	1999–2000
	Fast food	DBD900	How many (meals you did not eat at home) did you get from a fast-food or pizza place? (meals per week)	0, >0 and ≤5, >5 and ≤10*	2007–2014
	Depression	CIDDSCOR	Depression score (diagnosis)	Yes, no	1999–2004
	GAD	CIDGSCOR	Generalized anxiety disorder score (diagnosis)	Yes, no	1999–2004
	Panic	CIDPSCOR	Panic disorder score (diagnosis)	Yes, no	1999–2004
	Arthritis	MCQ160A	Has a doctor or other health professional ever told you that you had arthritis?	Yes, no	1999–2016
	Heart disease	MCQ160B-F‡	Has a doctor or other health professional ever told you that you had congestive heart failure OR coronary heart disease OR angina OR myocardial infarction OR a stroke?	Yes, no	1999–2016
	Emphysema	MCQ160G	Has a doctor or other health professional ever told you that you had emphysema?	Yes, no	1999–2016
	Cancer	MCQ220	Have you ever been told by a doctor or other health professional that you had cancer or a malignancy of any kind?	Yes, no	1999–2016

(continued on next page)

Table 1 (continued)

Subject	Factor	NHANES identifier	Description	Discrete response categories	Survey years
	Diabetes	DIQ010	Other than during pregnancy, have you ever been told by a doctor or health professional that you have diabetes or sugar diabetes?	Yes, no, borderline	1999–2016
	Hypertension	BPQ020	Have you ever been told by a doctor or other health professional that you had hypertension, also called high blood pressure?	Yes, no	1999–2016
	Aspirin use (recommended)	RXQ510/RXQ515 <sup>†</sup>	Doctors and other health care providers sometimes recommend that you take a low-dose aspirin each day to prevent heart attacks, strokes, or cancer. Have you ever been told to do this? Are you now following this advice?	Yes, no	2011–2014
	Aspirin use (independent)	RXQ520	On your own, are you now taking a low-dose aspirin each day to prevent heart attacks, strokes, or cancer?	Yes, no	2011–2014
	Aspirin frequency	RXQ525G	How often do you take an aspirin?	Every day, alternate days, another schedule	2011–2014

NHANES = National Health and Nutrition Examination Survey.

<sup>\*</sup> Also available as continuous.

<sup>†</sup> Identifier changed during the 1999–2016 period.

<sup>‡</sup> Heart condition variables combined for purposes of present study.

were omitted from the primary multivariable model as their inclusion reduced the number of available participants with complete data to below 5000. Where these factors were found to have a significant association with NLR in the univariate model, additional multivariable analyses were conducted including the same core set of covariates as the primary model (Table 4). Multivariable model results for these factors are also presented in Supplementary Table 2. Note that in Table 4 and Supplementary Table 2, beta coefficients for levels of each categorical factor represent the change in mean NLR when transitioning from the referent to the exposure level of interest. Where factors are also treated as continuous, this coefficient represents the change in NLR per unit change in the factor.

Tables 5 and 6 presents results from multivariable models for participants stratified by comorbidity status (has comorbid condition vs. does not have comorbid condition), race, age, and sex. Supplementary Table 3 presents results from multivariable models for participants stratified by specific comorbidity (arthritis, heart conditions, cancer, diabetes, and hypertension).

### Comorbidity

Participants with any comorbidity (diagnoses of arthritis, heart disease, cancer, diabetes, hypertension, or emphysema) had higher NLRs than healthy participants (median NLR 2.03 and 1.91, respectively;  $P < 2E-16$ ). This pattern was consistent regardless of gender, age, race, education, and marital status (not shown).

### Demographic factors

#### Sex

Women exhibited slightly lower NLR on average than men (1.96 vs. 1.99,  $P = .03$ ), although both groups remained close to the overall cohort median of 1.98 (Table 2). An association between female sex and lower NLR was observed in univariate regression analysis ( $\beta = -0.03$ ,  $P = .008$ ) and became stronger after multivariate adjustment ( $\beta = -0.13$ ,  $P = 4.13E-12$ ) (Table 4).

NLR associations for sex across demographic strata and lifestyle factors are shown in Tables 5 and 6. Among persons with any comorbidity (Table 5), a lower NLR was observed in women than men, mainly among persons aged 60 or older. This pattern was consistent regardless of race. Among persons without comorbidity (Table 6), a similar pattern was observed for non-Hispanic whites with lower NLR observed in females aged 60 or older and no association by sex in persons under age 60. Weaker associations were observed in African Americans. By contrast, among Hispanics, women under age 60 had higher NLRs than men. Associations between sex and NLR were maintained in subgroups of participants with arthritis, heart disease, cancer, diabetes, and hypertension (Supplementary Table 3).

#### Age

Participants aged 60 years or older demonstrated a higher NLR than participants younger than 60 (2.11 vs. 1.92,  $P < 2.2E-16$ , Table 2). Older age remained a risk factor for higher NLR after multivariate adjustment ( $\beta = 0.12$ ,  $P = 7.05E-07$ ) (Table 4). An association between age and NLR was observed among non-Hispanic white males ( $\beta = 0.39$ ,  $P = 6.43E-07$  with comorbidity,  $\beta = 0.34$ ,  $P = 1.56E-05$  without comorbidity) and Hispanic males ( $\beta = 0.26$ ,  $P = .02$  with comorbidity,  $\beta = 0.18$ ,  $P = .02$  without comorbidity), but not non-Hispanic black males or females (Tables 5 and 6). Associations between age and NLR were also maintained in subgroups of participants with arthritis, heart disease, diabetes, and hypertension, but not in participants with cancer (Supplementary Table 3).

**Table 2**  
Median NLR across levels of each sociodemographic factor examined

Factor <sup>*</sup>	Level	N (level)	N (factor)	NLR median (IQR)	P-value <sup>†</sup>
Sex	Male	23,206	48,023	1.99 (1.18)	.03
	Female	24,817		1.96 (1.17)	
Age	<60	33,039	48,023	1.92 (1.10)	<2.2E-16
	≥60	14,984		2.11 (1.33)	
Race	Non-Hispanic white	21,254	48,023	2.14 (1.23)	<2.2E-16
	Non-Hispanic black	9928		1.60 (1.10)	
	Mexican American	9138		2.00 (1.07)	
	Other Hispanic	3847		1.93 (1.06)	
	Other race	3856		1.86 (1.09)	
Education level	Less than 12th grade	12,430	44,597	1.98 (1.18)	.5 P-trend: .23
	High school/GED or equivalent	22,657		1.99 (1.19)	
	College graduate or above	9452		1.99 (1.15)	
	Refused/don't know	58		1.76 (1.46)	
Marital status	Married	23,641	46,207	2.00 (1.15)	<2.2E-16
	Widowed	3856		2.17 (1.40)	
	Divorced	4465		1.99 (1.18)	
	Separated	1468		1.94 (1.16)	
	Never married	9432		1.92 (1.18)	
	Living with partner	3323		1.88 (1.12)	
	Refused/don't know	22		2.01 (1.73)	
People in household	1	6391	48,023	2.07 (1.34)	<2.2E-16 P-trend: <2.2E-16
	2	14,156		2.02 (1.22)	
	3	8683		1.96 (1.16)	
	4	7857		1.94 (1.12)	
	5	5240		1.93 (1.07)	
	6	2647		1.91 (1.09)	
	7+	3049		1.85 (1.10)	
Income	< \$20,000	10,662	46,175	2.01 (1.26)	6.95E-07 P-trend: 6.58E-09
	\$20,000 to \$44,999	13,860		1.99 (1.20)	
	\$45,000 to \$74,999	8762		1.96 (1.18)	
	≥ \$75,000	10,327		1.95 (1.07)	
	Refused/don't know	2564		1.93 (1.18)	
Poverty income ratio	<1	9686	44,016	1.96 (1.20)	.002
	≥1	34,330		1.99 (1.17)	
Health insurance	Yes	37,185	47,874	2.00 (1.21)	<2.2E-16
	No	10,601		1.91 (1.07)	
	Refused/don't know	88		1.88 (1.06)	

A detailed description of each variable can be found in [Table 1](#).

NLR = neutrophil-to-lymphocyte ratio.

<sup>\*</sup> Detailed descriptions of all factors can be found in [Table 1](#).

<sup>†</sup> P values for difference in median NLR between factor levels: calculated using the Mann Whitney test for 2 levels and the Kruskal-Wallis omnibus test for >2 levels. P-trend obtained from the Jonckheere-Terpstra test for ordered differences among levels.

### Race

Black participants demonstrated lower median NLR than white participants, with intermediate values observed among other races (non-Hispanic white: 2.14; Mexican American: 2.00; other Hispanic: 1.93; other race: 1.86; non-Hispanic black: 1.60;  $P < 2E-16$ ) ([Table 2](#)). In multivariable analysis, non-Hispanic black race remained strongly and independently associated with lower NLR ( $\beta = -0.53$ ,  $P < 2E-16$ ; non-Hispanic white referent) ([Table 4](#)). In all age and sex subgroups (under 60 years, 60 years or older, male, female), non-Hispanic black and Hispanic participants demonstrated lower NLR than non-Hispanic white participants, regardless of presence or absence of comorbidities (not shown). Associations between race and NLR were also maintained in subgroups of participants with arthritis, heart disease, cancer, diabetes, and hypertension ([Supplementary Table 3](#)).

### Socioeconomic status

Higher educational attainment was associated with an increase in NLR in multivariate analysis (high school/general education diploma or equivalent:  $\beta = 0.05$ ,  $P = .02$ ; college graduate or above:  $\beta = 0.08$ ,  $P = .01$ , less than 12th grade referent). Marital statuses of widowed ( $\beta = 0.19$ ,  $P = 9.99E-07$ ), separated ( $\beta = 0.11$ ,  $P = .02$ ), or never married ( $\beta = 0.05$ ,  $P = .04$ ) were all associated with increased NLR after multivariable adjustment, as compared with a status of currently married. Widowed participants had a higher mean NLR

than persons in all other marital groups ( $P < 2E-16$ ) with a median of 2.17, one of the highest values observed in any subgroup. Household size, income, and health insurance status were not independently associated with NLR after multivariate adjustment.

### Lifestyle factors

#### Ever smoker status

Never smokers (participants who had smoked less than 100 cigarettes in their lifetime) demonstrated lower NLR (1.93) than those who had ever smoked (2.04;  $P < 2.2E-16$ ) ([Table 3](#)). In univariate regression analysis, never smoker status was associated with lower NLR ( $\beta = -0.14$ ,  $P < 2E-16$ ); results were attenuated after multivariate adjustment ( $\beta = -0.06$ ,  $P = .0007$ ) ([Table 4](#)). In subgroup analysis, the association between never-smoker status and decreased NLR was strongest among older non-Hispanic white participants with any comorbidity ( $\beta = -0.15$ ,  $P = .02$ ) ([Table 5](#)).

#### Alcohol use

Participants who did not drink at all (zero drinking days per year) and those who drank frequently (>100 drinking days per year) both exhibited a higher NLR (2.06 and 2.01, respectively) than less-frequent drinkers (NLR = 1.95–1.96) ([Table 3](#)). In univariate regression analysis, any alcohol consumption was associated with lower NLR, though without evidence of dose response ( $\beta$

**Table 3**  
Median NLR across levels of each lifestyle factor examined

Factor *	Level	N (level)	N (factor)	NLR median (IQR)	P-value †
Ever smoker	Yes	20,596	45,131	2.04 (1.24)	<2.2E-16
	No	24,494		1.93 (1.12)	
	Refused/don't know	41		1.79 (1.12)	
Alcohol use I (drinking days per year)	None	7988	35,335	2.06 (1.30)	<2.2E-16
	≤10	7995		1.96 (1.16)	
	>10 and ≤50	7209		1.95 (1.13)	
	>50 and ≤100	3838		1.95 (1.11)	
	>100	8263		2.01 (1.18)	
	Refused/don't know	42		2.43 (1.26)	
Alcohol use II (drinks per drinking day)	1	9586	27,346	2.00 (1.19)	2.91E-06
	2 or 3	11,295		1.94 (1.12)	
	4 or 5	3205		1.96 (1.13)	
	>5	3220		1.99 (1.14)	
	Refused/don't know	40		2.04 (1.30)	
BMI	≤18.5	914	47,175	1.91 (1.30)	.008 P-trend: 5.39E-05
	>18.5 and ≤25	14,285		1.95 (1.21)	
	>25 and ≤30	15,676		1.96 (1.14)	
	>30	16,300		2.00 (1.16)	
	Refused/don't know	40		2.04 (1.30)	
Physical activity	None	14,627	28,029	1.99 (1.20)	<2.2E-16
	Moderate only	7136		1.95 (1.11)	
	Vigorous only	2257		1.79 (0.99)	
	Moderate and vigorous	4001		1.84 (0.99)	
	Refused/don't know	8		1.69 (0.43)	
Vigorous activity duration	≤1 h	4025	6248	1.84 (0.97)	.19
	>1 h and ≤2 h	1575		1.78 (1.01)	
	>2 h and ≤4 h	575		1.80 (1.02)	
	>4 h	72		1.94 (1.01)	
	Refused/don't know	1		3.62 (0.00)	
Moderate activity duration	≤1 h	8564	11,115	1.91 (1.05)	.13
	>1 h and ≤2 h	1624		1.89 (1.04)	
	>2 h and ≤4 h	764		1.99 (1.19)	
	>4 h	159		1.98 (1.01)	
	Refused/don't know	4		1.63 (0.42)	
Time sitting/reclining	≤1 h	4371	27,992	1.88 (1.04)	1.67E-07 P-trend: 2.99E-09
	>1 h and ≤2 h	12,820		1.93 (1.14)	
	>2 h and ≤4 h	9789		1.96 (1.16)	
	>4 h	903		2.00 (1.11)	
	Refused/don't know	109		2.00 (1.17)	
Vegetable intake (servings per day)	0	152	1509	2.11 (1.12)	.43
	1 or 2	1163		2.10 (1.31)	
	3 or 4	177		2.22 (1.17)	
	>4	12		2.74 (1.39)	
	Refused/don't know	5		1.93 (1.44)	
Fast food (meals per week)	0	5559	17,309	1.96 (1.16)	.007 P-trend: .02
	>0 and ≤5	10,283		1.95 (1.13)	
	>5 and ≤10	1107		1.94 (1.11)	
	>10	357		1.84 (1.12)	
	Refused/don't know	3		0.99 (0.50)	

A detailed description of each variable can be found in Table 1.

NLR = neutrophil-to-lymphocyte ratio.

\* Detailed descriptions of all factors can be found in Table 1.

† P values for difference in median NLR between factor levels: calculated using the Mann Whitney test for 2 levels and the Kruskal-Wallis omnibus test for >2 levels. P-trend obtained from the Jonckheere-Terpstra test for ordered differences among levels.

coefficients of  $-0.16$ ,  $-0.18$ ,  $-0.19$ , and  $-0.11$  for increasing drink-day categories per year) (Table 4). Associations were attenuated after multivariate adjustment (between 1 and 10 drinking days per year:  $\beta = -0.09$ ,  $P = .001$ ; between 50 and 100 drinking days per year:  $\beta = -0.08$ ,  $P = .009$ ,  $P$  trend = .31; no drinking days per year as referent) (Table 4).

Among participants with comorbidities, an association was demonstrated between drinking <10 drinks per year and decreased NLR ( $\beta = -0.16$ ,  $P = .02$ ) in non-Hispanic white females. In the absence of comorbidities, a similar relationship for light drinking was observed among Hispanic participants (under 60s:  $\beta = -0.19$ ,  $P = .002$ ; males:  $\beta = -0.18$ ,  $P = .03$ ; females:  $\beta = -0.17$ ,  $P = .04$ ; Table 6).

#### Body mass index

A higher BMI was positively associated with NLR in crude analyses ( $P$ -trend =  $5.39E-05$ ) (Table 3). However, after multivariable

adjustment, a higher body weight was associated with reduced NLR. A similar inverse pattern was observed for overweight (BMI greater than 25 and less than or equal to 30) and obese (BMI above 30) BMI categories ( $\beta = -0.17$ ,  $P = .02$ , and  $\beta = -0.16$ ,  $P = .02$ , respectively) (Table 4). In non-Hispanic whites, an association of increased body weight (BMI greater than 25) with decreased NLR was restricted to persons aged 60 or older, with ( $\beta = -0.27$ ,  $P = .0001$ ) or without ( $\beta = -0.29$ ,  $P = .007$ ) comorbidity. A similar association was observed among non-Hispanic blacks with comorbidities ( $\beta = -0.36$ ,  $P = .0001$ ) (Table 5).

#### Physical activity

Two questions each were included relating to moderate and vigorous physical activity: "Do you do any moderate/vigorous activity for at least 10 minutes at a time" and "How much time do you spend doing moderate/vigorous activity per day?" Participants that

**Table 4**  
Univariate and multivariable regression results for association between NLR and all factors included in 5 or more survey years

Factor	Level	Univariate			Multivariable		
		N (level) *	Coefficient	P-value	N (level) *	Coefficient	P-value
Sex	Male	23,206		Ref	10,704		Ref
	Female	24,817	−0.03	.008	9533	−0.13	<b>4.13E-12</b>
Age	<60	33,039		Ref	13,685		Ref
	≥60	14,984	0.28	<2E-16	6552	0.12	<b>7.05E-07</b>
Race	Continuous	48,023	0.0066	<2E-16	20,237		Ref
	Non-Hispanic white	21,254		Ref	9251		Ref
	Non-Hispanic black	9928	−0.59	<2E-16	4066	−0.53	<b>&lt;2E-16</b>
	Mexican American	9138	−0.20	<2E-16	3019	−0.10	<b>3.42E-04</b>
	Other Hispanic	3847	−0.28	<2E-16	2089	−0.16	<b>2.34E-07</b>
Education level	Other race	3856	−0.31	<2E-16	1812	−0.21	<b>2.68E-11</b>
	Less than 12th grade	12,430		Ref	4674		Ref
	High school/GED or equivalent	22,657	0.009	.51	10,707	0.05	<b>.02</b>
Marital status	College graduate or above	9452	−0.003	.84	4843	0.08	<b>.01</b>
	Married	23,641		Ref	10,419		Ref
	Widowed	3856	0.27	<2E-16	1391	0.19	<b>9.99E-07</b>
	Divorced	4465	−0.005	.8	2355	0.03	.28
	Separated	1468	−0.04	.23	683	0.12	<b>.02</b>
People in household	Never married	9432	−0.10	3.64E-11	3691	0.05	<b>.04</b>
	Living with partner	3323	−0.12	1.06E-07	1692	−0.05	.17
	1	6391		Ref	2900		Ref
	2	14,156	−0.06	2.00E-03	6261	0.01	.67
	3	8683	−0.17	<2E-16	3621	0.01	.68
	4	7857	−0.21	<2E-16	3284	0.04	.33
	5	5240	−0.23	<2E-16	2065	0.02	.60
Income	6	2647	−0.21	5.59E-13	1051	0.06	.24
	7+	3049	−0.27	<2E-16	1055	−0.07	.17
	Continuous	48,023	−0.05	<2E-16	20,237		Ref
	<\$20,000	10,662		Ref	4015		Ref
	\$20,000 to \$44,999	13,860	−0.05	.001	5701	0.004	.89
Health Insurance	\$45,000 to \$74,999	8762	−0.08	5.67E-06	3855	−0.04	.17
	≥\$75,000	10,327	−0.13	9.27E-15	5295	−0.06	<b>.05</b>
	Yes	37,185		Ref	15,746		Ref
Ever Smoker	No	10,601	−0.15	<2E-16	4474	−0.03	.18
	Yes	20,596		Ref	10,218		Ref
Alcohol use I (drinking days per year)	No	24,494	−0.14	<2E-16	10,002	−0.06	<b>.0007</b>
	None	7988		Ref	4306		Ref
	≤10	7995	−0.16	1.57E-15	4522	−0.09	<b>.001</b>
	>10 and ≤50	7209	−0.18	<2E-16	4226	−0.05	.09
	>50 and ≤100	3838	−0.19	3.41E-15	2243	−0.08	<b>.009</b>
	>100	8263	−0.11	2.65E-08	4915	−0.02	.44
	Continuous	35,293	−0.02	1.72E-06	20,212		Ref
BMI	≤18.5	914		Ref	307		Ref
	>18.5 and ≤25	14,285	−0.01	.77	5558	−0.10	.14
	>25 and ≤30	15,676	−0.02	.66	6740	−0.17	<b>.02</b>
	>30	16,300	−0.02	.57	7632	−0.16	<b>.02</b>
Physical activity	Continuous	47,175	−0.09	.4	20,237		Ref
	None	14,627		Ref	10,280		Ref
	Moderate only	7136	−0.08	1.67E-05	5360	−0.08	<b>.0002</b>
	Vigorous only	2257	−0.26	<2E-16	1560	−0.13	<b>.0002</b>
Time sitting/reclining	Moderate and vigorous	4001	−0.23	<2E-16	3030	−0.13	<b>8.11E-07</b>
	≤1 h	4371		Ref	2977		Ref
	>1 h and ≤2 h	12,820	0.08	9.47E-05	9412	0.04	.12
	>2 h and ≤4 h	9789	0.15	5.51E-12	7138	0.11	<b>3.76E-05</b>
	>4 h	903	0.16	.0003	648	0.05	.4
Arthritis	Continuous	27,883	0.07	5.88E-13	20,175		Ref
	Yes	11,739		Ref	5470		Ref
Heart condition	No	32,775	−0.19	<2E-16	14,728	−0.02	.43
	Yes	4925		Ref	2165		Ref
Emphysema	No	39,611	−0.43	<2E-16	18,053	−0.22	<b>5.82E-13</b>
	Yes	910		Ref	473		Ref
Cancer	No	43,630	−0.72	<2E-16	19,744	−0.33	<b>1.59E-08</b>
	Yes	3997		Ref	1970		Ref
Diabetes	No	40,553	−0.37	<2E-16	18,250	−0.15	<b>8.95E-07</b>
	Yes	5252		Ref	2491		Ref
Hypertension	No	41,876	−0.20	<2E-16	17,268	−0.16	<b>2.33E-08</b>
	Borderline	862	−0.13	.003	465	−0.17	<b>.00473</b>
	Yes	15,457		Ref	7268		Ref
	No	32,330	−0.17	<2E-16	12,946	−0.07	<b>.00129</b>

Bold font indicates items which remain significant after multivariable adjustment.

NLR = neutrophil-to-lymphocyte ratio.

\* Univariate models include all available participants. Lower N values in multivariable model reflect restriction to only participants with complete data for all included variables. Note that participants reporting “refused/don’t know” are excluded from this table for brevity, and thus slight variations from these participant totals may be observed.

**Table 5**  
Associations between demographic and lifestyle factors (sex, age, ever-smoker status, alcohol use, BMI, and physical activity) and NLR within the independent subgroup of participants with comorbidities classified by race/ethnicity, age, and sex

Subjects with comorbidity													
Factor	Level	Non-Hispanic white				Non-Hispanic black				Hispanic			
		Age		Sex		Age		Sex		Age		Sex	
		<60	>60	M	F	<60	>60	M	F	<60	>60	M	F
		Ref											
Sex	M	Ref				Ref				Ref			
	F	-0.08	<b>-0.50*</b>			<b>-0.18†</b>	<b>-0.30‡</b>			-0.13	<b>-0.38‡</b>		
Age	<60	Ref				Ref				Ref			
	≥60			<b>0.39*</b>	0.01			0.005	0.05			<b>0.26‡</b>	-0.11
Ever smoker	Yes	Ref				Ref				Ref			
	No	-0.05	<b>-0.15‡</b>	-0.12	-0.09	0.04	0.01	0.007	0.08	-0.04	0.01	-0.09	0.05
Alcohol use 1 (drinking days per year)	None	Ref				Ref				Ref			
	≤10	-0.06	0.007	0.14	<b>-0.16‡</b>	-0.02	-0.10	0.02	-0.16	-0.07	-0.17	-0.17	-0.09
	>10 and ≤50	-0.03	0.04	0.14	-0.12	-0.003	-0.07	-0.04	-0.09	0.05	-0.21	-0.22	0.01
	>50 and ≤100	-0.08	-0.008	0.05	-0.10	-0.04	-0.24	-0.13	-0.19	-0.02	0.04	0.04	-0.05
	>100	-0.09	-0.10	-0.01	-0.15	0.09	-0.05	0.06	-0.07	0.06	-0.02	-0.04	0.20
BMI	≤25	Ref				Ref				Ref			
	>25	0.03	<b>-0.27*</b>	<b>-0.13‡</b>	0.01	-0.01	<b>-0.36*</b>	-0.11	-0.13	-0.02	-0.003	-0.01	-0.03
Physical activity	None	Ref				Ref				Ref			
	Moderate only	-0.10	<b>-0.31*</b>	<b>-0.25*</b>	<b>-0.18‡</b>	0.03	-0.14	-0.10	-0.02	-0.02	0.04	-0.04	0.05
	Vigorous only	<b>-0.31‡</b>	-0.15	-0.25	-0.28	-0.14	-0.07	-0.10	-0.10	<b>-0.26‡</b>	-0.13	-0.23	-0.29
	Moderate and vigorous	<b>-0.24‡</b>	-0.24	<b>-0.21‡</b>	<b>-0.28‡</b>	-0.10	-0.12	-0.15	0.02	<b>-0.20‡</b>	<b>-0.53‡</b>	-0.30	<b>-0.29‡</b>

Results represent β coefficient and P value from a multivariable regression model that included all factors included in the multivariable model of Table 4 and also presence of comorbidities (arthritis, heart disease, diabetes, cancer, emphysema, and hypertension).

Bold font indicates items which remain significant after multivariable adjustment.

BMI = body mass index; NLR = neutrophil-to-lymphocyte ratio.

\* Represents P < .001.

† Represents P < .01.

‡ Represents P < .05.

reported no physical activity for at least 10 minutes at a time demonstrated higher NLR (1.99) than those who engaged in at least 10 minutes of moderate activity (1.95), vigorous activity (1.79) or

both (1.84) (Table 3). NLR did not change significantly with increasing time spent engaging in moderate or physical activity (P = .19, P = .13, respectively) (Table 3). In univariate regression

**Table 6**  
Associations between demographic and lifestyle factors (sex, age, ever-smoker status, alcohol use, BMI, and physical activity) and NLR within the independent subgroup of participants without comorbidities classified by race/ethnicity, age, and sex

Subjects without comorbidity													
Factor	Level	Non-Hispanic white				Non-Hispanic black				Hispanic			
		Age		Sex		Age		Sex		Age		Sex	
		<60	>60	M	F	<60	>60	M	F	<60	>60	M	F
		Ref											
Sex	M	Ref				Ref				Ref			
	F	0.04	<b>-0.46*</b>			0.08	-0.23			<b>0.11†</b>	-0.19		
Age	<60	Ref				Ref				Ref			
	≥60			<b>0.34*</b>	-0.15			0.14	-0.18			<b>0.18‡</b>	-0.22
Ever Smoker	Yes	Ref				Ref				Ref			
	No	-0.03	-0.10	-0.007	-0.07	-0.08	-0.09	-0.14	0.007	0.006	-0.27	-0.04	-0.71
Alcohol use 1 (drinking days per year)	None	Ref				Ref				Ref			
	≤10	-0.01	0.004	-0.06	0.06	-0.05	-0.10	-0.07	-0.14	<b>-0.19‡</b>	-0.09	<b>-0.18‡</b>	<b>-0.17‡</b>
	>10 and ≤50	-0.03	-0.08	-0.08	0.05	0.15	-0.08	0.19	-0.02	-0.11	-0.02	-0.08	-0.11
	>50 and ≤100	-0.06	-0.05	-0.12	0.03	-0.02	-0.24	-0.01	-0.07	-0.08	-0.10	-0.13	0.06
	>100	0.05	-0.006	-0.01	0.10	0.13	-0.10	0.12	-0.006	-0.0009	0.23	0.08	-0.08
BMI	≤25	Ref				Ref				Ref			
	>25	0.04	<b>-0.29‡</b>	-0.03	0.04	-0.02	0.11	-0.03	0.02	0.02	-0.20	-0.02	0.02
Physical Activity	None	Ref				Ref				Ref			
	Moderate only	-0.03	-0.03	-0.03	-0.02	-0.03	-0.08	-0.06	-0.05	-0.01	0.14	-0.05	0.02
	Vigorous only	<b>-0.16‡</b>	-0.13	<b>-0.17‡</b>	-0.14	0.16	-0.65	0.15	0.19	-0.009	-0.11	-0.005	-0.02
	Moderate and vigorous	<b>-0.12‡</b>	-0.06	0.009	<b>-0.24*</b>	0.04	-0.14	0.03	0.02	-0.07	0.10	-0.08	-0.02

Results represent β coefficient and P value from a multivariable regression model that included all factors included in the multivariable model of Table 4 and also presence of comorbidities (arthritis, heart disease, diabetes, cancer, emphysema, and hypertension).

Bold font indicates items which remain significant after multivariable adjustment.

BMI = body mass index; NLR = neutrophil-to-lymphocyte ratio.

\* Represents P < .001.

† Represents P < .01.

‡ Represents P < .05.

analysis, physical activity for at least 10 minutes was associated with reduced NLR for moderate activity NLR ( $\beta = -0.08$ ,  $P = 1.67E-05$ ) with further reduction in NLR for vigorous activity alone ( $\beta = -0.26$ ,  $P < 2E-16$ ), or vigorous combined with moderate activity ( $\beta = -0.23$ ,  $P < 2E-16$ ). Associations were attenuated after multivariate adjustment (Table 4). Increasing hours of moderate or vigorous activity was not associated with further reduction in NLR (Supplementary Table 2).

Associations between physical activity and NLR varied by demographic subgroup and comorbidity status (Tables 5 and 6). In non-Hispanic whites with any co-morbidity (Table 5), moderate activity was associated with lower NLR in older persons ( $\beta = -0.31$   $P = 3.64E-06$ ), and in both genders (male:  $\beta = -0.25$   $P = .0007$ , female:  $\beta = -0.18$   $P = .003$ ) whereas only vigorous activity ( $\beta = -0.31$   $P = .006$ ) or a combination of moderate and vigorous ( $\beta = -0.24$   $P = .002$ ) activity was associated with lower NLR in younger participants. Similar, though attenuated, results were observed among non-Hispanic whites without comorbidities (Table 6). Among Hispanics, reductions in NLR with physical activity were observed only in participants with comorbidities: a combination of moderate and vigorous physical activity was associated with decreased NLR for those aged under 60 ( $\beta = -0.20$   $P = .05$ ), aged 60 or older ( $\beta = -0.53$   $P = .03$ ), and in females ( $\beta = -0.29$   $P = .03$ ) only. Vigorous activity alone was also associated with decreased NLR in persons aged under 60 ( $\beta = -0.26$   $P = .04$ ). Physical activity demonstrated no association with NLR in non-Hispanic black participants regardless of age, sex, and comorbidity status. Interestingly, when stratifying by specific comorbidity (Supplementary Table 3), physical activity is the only lifestyle factor for which the association with NLR remains significant.

Greater sedentary behavior—more time spent sitting or reclining in an average day—was associated with higher NLR ( $P$ -trend =  $2.99E-09$ ) (Table 3). As compared with a referent of  $\leq 1$  hour per day, spending 2 to 4 hours per day sitting or reclining was associated with higher NLR ( $\beta = 0.11$   $P = 3.76E-05$ ) after multivariate adjustment. Few subjects reported greater levels of sedentary behavior ( $>4$  hours per day).

#### Diet and aspirin use

Increasing servings per day of vegetables was not found to be associated with increasing NLR; however, data were sparse (only 12 participants reported more than four servings per day) and results were thus imprecise (Supplementary Table 2). An increasing number of meals eaten at fast food restaurants was associated with a lower NLR ( $P$ -trend =  $.02$ ). This result was not maintained after multivariate adjustment.

Participants who were taking regular aspirin on the advice of a medical professional had a significantly higher NLR than those who were not taking regular aspirin ( $P = 1.25E-13$ ) (Supplementary Table 2). For those who were taking a regular aspirin but had not been advised to do so by a medical professional, NLR was not significantly increased as compared with those not taking an aspirin ( $P = .65$ ). Of participants who reported taking aspirin, those who did so at less frequent intervals than once per day tended to have a lower NLR ( $P = .02$ – $.04$ ). None of these results maintained significance after multivariable adjustment (Supplementary Table 2).

#### Discussion

Based on data from over 48,023 participants in the National Health and Nutrition Survey from 1999 to 2016, we evaluated associations between 19 sociodemographic and lifestyle factors and the NLR. Overall, participants with existing comorbidities

(arthritis, heart disease, cancer, diabetes, emphysema, and hypertension) had higher NLR than those without. After adjustment for pre-existing conditions, we found that sex, age, race, and marital status were independently associated with NLR. Furthermore, we observed relationships between NLR and smoking, alcohol use, BMI, and physical activity that differed by age and race, suggesting opportunities for targeted reductions in NLR through changes in lifestyle.

Postmenopausal women have demonstrated lower NLR than men, and this trend has been found to reverse for women aged 50 and younger [19]. In line with these earlier observations, females included in the present study also demonstrated lower NLR than males within the older age group, with a trend toward higher NLR in younger females. This reversal of the association between sex and NLR between age groups was only identified among participants without any comorbidity. If not due to chance, the female advantage with respect to NLR among older subjects may reflect differential effects of aging on the immune system. Females tend to remain immune-privileged later in life, in contrast to males who experience more rapid decreases in lymphocytes with advancing age [20,21].

We observed a lower NLR in non-Hispanic black subjects across all demographic subgroups. When compared with other racial groups, lifestyle appeared to play only a minor role in NLR among black participants: as shown in Table 3, beta coefficients were similar before ( $-0.59$ ) and after ( $-0.53$ ) multivariate adjustment for socioeconomic and lifestyle factors. A lower NLR in non-Hispanic blacks has been reported previously and may be attributed to a higher prevalence of benign ethnic neutropenia in persons of African descent [22]. Mechanisms underlying this phenomenon are unclear [22]. Owing to the limited sample size and nonspecific survey questions, it was not possible to rule-out differences in lifestyle or specific medical conditions as explanations for disparate NLR values by race. Further investigation of this association is warranted.

Participants who were widowed, separated, or never married exhibited higher NLR than married persons. Previous studies suggest that married individuals experience less psychological stress, leading to lower levels of cortisol and improved inflammatory regulation when compared with unmarried individuals [23]; such a potential biologic mechanism is in line with observations in the present study which found reduced NLR in married persons even after controlling for lifestyle and other demographic factors.

The association between smoking and increased levels of inflammatory markers has been previously established [24,25]; here, never smoker status was linked to a lower NLR, only among older non-Hispanic whites with comorbidities. Moderate drinking ( $<10$  drinking days per year) was associated with decreased NLR as compared with complete abstinence from alcohol in non-Hispanic white females with comorbidities and Hispanic individuals without comorbidities. Although potential benefits of moderate alcohol consumption remain under debate [26], the present study offers some limited support for the hypothesis that modest alcohol consumption may favorably impact health through reductions in systemic inflammation.

Physical activity can contribute to reducing markers of systemic inflammation [27]; participating in at least 10 minutes of physical activity at a time was also identified as the modifiable lifestyle factor most strongly associated with NLR in the NHANES cohort. Although no dose-response effect was observed (in that no association was found between increasing duration of moderate or vigorous activity and NLR), participating in at least 10 minutes of physical activity on a regular basis (moderate activity being optimal for those aged 60 or older, and vigorous being optimal for under 60s) led to lower NLR. A combination of both moderate and

vigorous physical activity demonstrated the most universally beneficial effect on NLR. However, the strong association between physical activity and NLR was only observed among non-Hispanic white and Hispanic participants, but not non-Hispanic blacks. This racial difference in systemic response to physical activity also warrants further study. Interestingly, associations between alcohol use and NLR and smoking history and NLR were no longer apparent in subgroups of participants with specific comorbidities (arthritis, heart conditions, cancer, diabetes, and hypertension). Physical activity, however, maintains significant associations with NLR in participants with arthritis, cancer, diabetes, and hypertension, suggesting there may be potential for further study of the impact of physical activity on NLR in the setting of these specific diseases.

Although increased time sitting or reclining per day (2–4 hours as compared with less than or equal to 1 hour) demonstrated an association with increased NLR, the reliability of self-report of this variable is brought into question by the small number of persons reporting 4 or more hours sitting or reclining per day (<5% of participants).

BMI demonstrated a negative association with NLR, notably in participants aged 60 or older. Although a link between adiposity and inflammation has been established [28, 29], lower NLR has also been observed in obese individuals (BMI > 30) as compared with those of normal weight, supporting the findings of the present work [30]. Higher BMI is associated with increased lymphocyte count [31]; if obesity induces a more substantial increase in lymphocytes than neutrophils, this could potentially be driving the observed inverse relationship between BMI and NLR. Greater consumption of vegetables has demonstrated both positive and negative associations with inflammatory biomarkers in earlier studies [32–34]. No conclusions could be drawn from the findings of the present work, particularly with only a limited number of participants reporting 3 or more servings per day. The relationship between fast food and NLR was also not upheld after multivariable adjustment. Further study of the association between NLR and diet and nutrition is thus warranted. Finally, although the association between aspirin use and NLR was not found to be significant in our multivariable models, it is worth noting that participants who were taking aspirin based on the recommendations of a medical professional and those taking aspirin at an increased frequency (once per day as compared with every other day or another schedule) tended to have higher NLR. We may speculate that these participants were taking aspirin in response to an existing inflammatory condition, contributing to the elevated NLR observed in these groups.

Several limitations in the present analysis should be acknowledged. The cross-sectional nature of the NHANES warrants cautious interpretation, as it is not possible to determine temporal relationships between studied factors and NLR. Inherent measurement error in a single blood biomarker may also have biased all results toward the null. Retrospective self-report data present additional challenges, particularly with regard to drinking and smoking habits and other potentially sensitive topic areas. Finally, in spite of the large overall sample size, data were sparse and results imprecise in a number of the demographically homogeneous subgroups examined (Tables 5 and 6). Further studies are needed to confirm the present results and to examine the utility of NLR for evaluating the efficacy of lifestyle interventions. It should also be noted that in the present study, the NLR was only weakly correlated ( $r = 0.25$ ) with C-reactive protein, another established marker of systemic inflammation. This also suggests a need for a clearer understanding of the underlying mechanisms driving alterations in each of these respective biomarkers.

NLR is an established marker of systemic inflammation and is strongly associated with survival outcomes across multiple diseases. Present findings suggest that age, race, sex, and marital

status, as well as a range of comorbidities, are all associated with NLR.

Modifiable exposures including alcohol use, physical activity, and smoking status were also found to impact NLR. Inclusion and appropriate control of these factors in multivariate analyses may improve the accuracy of, and consistency among, studies of the association between NLR and clinical outcomes. Furthermore, differences observed in the magnitude of association according to health status and demographics also offer the potential for targeting recommendations for lifestyle modifications to at-risk populations. Finally, given the already established link between NLR and multiple medical conditions, studies aiming to influence the magnitude of NLR are likely to become more commonplace; the present findings may also provide guidance for future interventional study design by suggesting optimal characteristics for participant matching.

### Financial support

Dr. Rachel Howard was supported by a Ruth L. Kirschstein Institutional National Research Service award from the National Institutes of Health, United States (T32 CA147832). The National Institutes of Health had no direct role in the design of this study, nor the execution, analyses, interpretation of the data or decision to submit results. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### References

- [1] Hunter P. The inflammation theory of disease. The growing realization that chronic inflammation is crucial in many diseases opens new avenues for treatment. *EMBO Rep* 2012;13(11):968–70.
- [2] Krishnamoorthy S, Honn KV. Inflammation and disease progression. *Cancer Metastasis Rev* 2006;25(3):481–91.
- [3] McMillan DC. Systemic inflammation, nutritional status and survival in patients with cancer. *Curr Opin Clin Nutr Metab Care* 2009;12(3):223–6.
- [4] Holmes C, Cunningham C, Zotova E, Woolford J, Dean C, Kerr S, et al. Systemic inflammation and disease progression in Alzheimer disease. *Neurology* 2009;73(10):768–74.
- [5] Gan WQ, Man SFP, Senthilselvan A, Sin DD. Association between chronic obstructive pulmonary disease and systemic inflammation: a systematic review and a meta-analysis. *Thorax* 2004;59:574–80.
- [6] Duncan BB, Schmidt MI, Pankow JS, Ballantyne CM, Couper D, Vigo A, et al. Low-Grade Systemic Inflammation and the Development of Type 2 Diabetes. *Diabetes* 2003;52(7):1799–805.
- [7] Zahorec R. Ratio of neutrophil to lymphocyte counts—rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy* 2001;102(1):5–14.
- [8] Templeton AJ, McNamara MG, Seruga B, Vera-Badillo FE, Aneja P, Ocana A, et al. Prognostic role of neutrophil-to-lymphocyte ratio in solid tumors: a systematic review and meta-analysis. *J Natl Cancer Inst* 2014;106(6):dju124.
- [9] Durmus E, Kivrak T, Gerin F, Sunbul M, Sari I, Erdogan O. Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio are predictors of heart failure. *Arq Bras Cardiol* 2015;105(6):606–13.
- [10] Tamhane UU, Aneja S, Montgomery D, Rogers EK, Eagle KA, Gurm HS. Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary syndrome. *Am J Cardiol* 2008;102(6):653–7.
- [11] Azab B, Daoud J, Naem FB, Nasr R, Ross J, Ghimire P, et al. Neutrophil-to-lymphocyte ratio as a predictor of worsening renal function in diabetic patients (3-year follow-up study). *Ren Fail* 2012;34(5):571–6.
- [12] Forget P, Khalifa C, Defour JP, Latine D, Van Pel MC, De Kock M. What is the normal value of the neutrophil-to-lymphocyte ratio? *BMC Res Notes* 2017;10(1):12.
- [13] Azab B, Camacho-Rivera M, Taioli E. Average values and racial differences of neutrophil lymphocyte ratio among a nationally representative sample of United States subjects. *PLoS One* 2014;9(11):e112361.
- [14] Wang R, Chen PJ, Chen WH. Diet and exercise improve neutrophil to lymphocyte ratio in overweight adolescents. *Int J Sports Med* 2011;32(12):982–6.
- [15] Centers for Disease Control, Prevention (CDC). National Centers for Health Statistics (NCHS). NHANES – National Health and Nutrition Examination Survey. <http://www.cdc.gov/nchs/nhanes.htm>. [Accessed 8 October 2018].
- [16] Centers for Disease Control, Prevention (CDC), National Centers for Health Statistics (NCHS). Complete Blood Count. [http://www.cdc.gov/nchs/data/nhanes/2015-2016/labmethods/CBC\\_I\\_MET\\_Complete\\_Blood\\_Count.pdf](http://www.cdc.gov/nchs/data/nhanes/2015-2016/labmethods/CBC_I_MET_Complete_Blood_Count.pdf). [Accessed 8 October 2018].
- [17] Jonckheere AR. A distribution-free k-sample test against ordered alternatives. *Biometrika* 1954;41:133–45.

- [18] Sterne JAC, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ* 2009;338:b2393.
- [19] Chen Y, Zhang Y, Zhao G, Chen C, Yang P, Ye S, et al. Difference in leukocyte composition between women before and after menopausal age, and distinct sexual dimorphism. *PLoS One* 2016;11(9):e0162953.
- [20] Hirokawa K, Utsuyama M, Hayashi Y, Kitagawam M, Makinodan T, Fulop T. Slower immune system aging in women versus men in the Japanese population. *Immun Ageing* 2013;10(1):19.
- [21] Giefing-Kroll C, Berger P, Lepperdinger G, Grubeck-Loebenstien B. How sex and age affect immune responses, susceptibility to infections, and response to vaccination. *Aging Cell* 2015;14(3):309–21.
- [22] Hsieh M, Chin K, Link B, Stroncek D, Wang E, Everhart J, et al. Benign ethnic neutropenia in individuals of African descent: Incidence, granulocyte mobilization, and gene expression profiling. *Blood* 2005;106(11):858a–9a.
- [23] Chin B, Murphy MLM, Janicki-Deverts D, Cohen S. Marital status as a predictor of diurnal salivary cortisol levels and slopes in a community sample of healthy adults. *Psychoneuroendocrinology* 2017;78:68–75.
- [24] Lee J, Taneja V, Vassallo R. Cigarette smoking and inflammation. Cellular and molecular mechanisms. *J Dent Res* 2012;91(2):142–9.
- [25] Bakhru A, Erlinger TP. Smoking cessation and cardiovascular disease risk factors: results from the third national health and nutrition examination survey. *PLoS Med* 2005;2(6):e160.
- [26] Wilson M. Does moderate alcohol consumption really have health benefits? *BMJ* 2018;362:k3888.
- [27] Tir AMD, Labor M, Plavec D. The effects of physical activity on chronic sub-clinical systemic inflammation. *Arh Hig Rada Toksikol* 2017;68:276–86.
- [28] Forsythe LK, Wallace JMW, Livingstone BE. Obesity and inflammation: the effects of weight loss. *Nutr Res Rev* 2008;21:117–33.
- [29] Visser M, Bouter LM, McQuillan GM, Wener MH, Harris TB. Elevated C-reactive protein levels in overweight and obese adults. *JAMA* 1999;282:2131–5.
- [30] Koca TT. Does obesity cause chronic inflammation? The association between complete blood parameters with body mass index and fasting glucose. *Pak J Med Sci* 2017;33(1):65–9.
- [31] Womack J, Tien PC, Feldman J, Shin JH, Fennie K, Anastos K, et al. Obesity and Immune Cell Counts in Women. *Metabolism* 2007;56(7):998–1004.
- [32] Asgard R, Rytter E, Basu S, Abramsson-Zetterberg L, Moller L, Vessby B. High intake of fruit and vegetables is related to low oxidative stress and inflammation in a group of patients with type 2 diabetes. *Scand J Food Nutr* 2007;51(4):149–58.
- [33] Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB, Willett WC. Fruit and vegetable intakes, C-reactive protein, and the metabolic syndrome. *Am J Clin Nutr* 2006;84:1489–97.
- [34] Lee Y, Kang D, Lee SA. Effect of dietary patterns on serum C-reactive protein level. *Nutr Metab Cardiovasc Dis* 2014;24:1004–11.

## Appendix

**Table S1**Median NLR across levels of each comorbidity and medication factor examined. A detailed description of each variable can be found in [Table 1](#)

Factor *	Level	N (level)	N (factor)	NLR median (IQR)	P-value †
Depression	Yes	141	2107	2.02 (1.38)	.67
	No	1966		2.06 (1.31)	
GAD	Yes	50	2111	2.09 (1.66)	.63
	No	2061		2.06 (1.31)	
Panic	Yes	47	2112	1.95 (1.45)	.73
	No	2065		2.06 (1.31)	
Arthritis	Yes	11,739	44,596	2.08 (1.28)	<2.2E-16
	No	32,775		1.95 (1.14)	
	Refused/don't know	82		2.16 (1.01)	
Heart condition	Yes	4925	44,597	2.27 (1.46)	<2.2E-16
	No	39,611		1.95 (1.15)	
	Refused/don't know	61		2.02 (0.95)	
Emphysema	Yes	910	44,597	2.46 (1.71)	<2.2E-16
	No	43,630		1.98 (1.17)	
	Refused/don't know	57		2.69 (1.38)	
Cancer	Yes	3997	44,597	2.24 (1.42)	<2.2E-16
	No	40,553		1.96 (1.16)	
	Refused/don't know	47		1.99 (0.98)	
Diabetes	Yes	5252	48,020	2.10 (1.29)	<2.2E-16
	No	41,876		1.96 (1.16)	
	Borderline	862		1.96 (1.17)	
Hypertension	Refused/don't know	30	47,862	2.22 (1.29)	<2.2E-16
	Yes	15,457		2.06 (1.27)	
	No	32,330		1.94 (1.13)	
Aspirin use (recommended)	Refused/don't know	75	6198	2.07 (1.21)	1.71E-09
	Yes	1789		2.13 (1.41)	
	No	4401		1.98 (1.14)	
Aspirin use (independent)	Refused/don't know	8	4409	2.36 (1.54)	.83
	Yes	235		2.03 (1.20)	
	No	4168		1.98 (1.14)	
Aspirin frequency	Refused/don't know	6	2132	2.01 (0.69)	.0009
	One every day	1807		2.13 (1.40)	
	One every other day	134		1.89 (1.19)	
	Another schedule	187		1.96 (1.09)	
	Refused/don't know	4		2.86 (1.40)	

NLR = neutrophil-to-lymphocyte ratio.

\* Detailed descriptions of all factors can be found in [Table 1](#).

† P values for difference in median NLR between factor levels: calculated using the Mann Whitney test for 2 levels and Kruskal Wallis omnibus test for &gt;2 levels. P-trend obtained from Jonckheere-Terpstra test for ordered differences among levels.

**Table S2**

Univariate and multivariable regression results for associations between NLR and all factors included in only a small subset of survey years. These factors were omitted from the primary multivariable model (Table 4) as their inclusion reduced the number of available participants with complete data to below 5000. Where these factors were found to have a significant association with NLR in the univariate model, additional multivariable analyses were conducted including the same set of covariates as the primary model

Factor	Level	Univariate			Multivariable*		
		N (level)†	Coefficient	P-value	N (level)†	Coefficient	P-value
Vegetable intake (servings per day)	0	152		Ref			
	1 or 2	1163	0.05	.71			
	3 or 4	177	−0.04	.81			
	>4	12	0.46	.27			
	Continuous	1504	−0.04	.95			
Fast food (meals per week)	0	5559		Ref	1257		Ref
	>0 and ≤5	10,283	−0.04	<b>.03</b>	1970	0.05	.30
	>5 and ≤10	1107	−0.09	<b>.03</b>	145	−0.19	.09
	>10	357	−0.18	<b>.005</b>	44	0.08	.68
	Continuous	17,306	−0.49	<b>.0003</b>	3416		
Depression	Yes	141		Ref			
	No	1966	−0.12	.31			
GAD	Yes	50		Ref			
	No	2061	−0.03	.86			
Panic	Yes	47		Ref			
	No	2065	0.02	.92			
Aspirin use (recommended)	Yes	1789		Ref	1376		Ref
	No	4401	−0.29	<b>1.25E-13</b>	3263	−0.005	.93
Aspirin use (independent)	Yes	235		Ref			
	No	4168	0.04	.65			
Aspirin frequency	One every day	1807		Ref	1401		Ref
	One every other day	134	−0.30	<b>.04</b>	109	−0.31	.06
	Another schedule	187	−0.30	<b>.02</b>	146	−0.04	.77

Bold font indicates items which remain significant after multivariable adjustment.

\* Results presented are from multivariable models including all factors included in the primary multivariable model of Table 4.

† Univariate models include all available participants. Lower N values in the multivariable model reflect restriction to only participants with complete data for all included variables. Note that participants reporting “refused/don’t know” are excluded from this table for brevity, and thus slight variations from these participant totals may be observed.

**Table S3**

Multivariable analysis for associations between demographic, lifestyle and health-related factors and NLR within independent subgroups of the participant population classified by diagnosis with a specific comorbidity. As can be seen, all multivariable models include all factors included in the primary multivariable model of Table 4

Factor	Level	Arthritis (n = 5470)			Heart condition (n = 2165)			Cancer (n = 1970)			Diabetes (n = 2491)			Hypertension (n = 7268)		
		N (level) *	Coefficient	P-value	N (level) *	Coefficient	P-value	N (level) *	Coefficient	P-value	N (level) *	Coefficient	P-value	N (level) *	Coefficient	P-value
Sex	Male	2460		Ref	1344		Ref	999		Ref	1425		Ref	3829		Ref
	Female	3010	-0.34	<2E-16	821	-0.33	<b>1.80E-04</b>	971	-0.47	<b>3.05E-09</b>	1066	-0.23	<b>7.00E-04</b>	3439	-0.26	<b>1.83E-12</b>
Age	<60	2213		Ref	597		Ref	571		Ref	979		Ref	3268		Ref
	≥60	3257	0.11	<b>.02</b>	1568	0.25	<b>.02</b>	1399	0.10	.28	1512	0.19	<b>.01</b>	4000	0.14	<b>.001</b>
Race	Non-Hispanic white	3056		Ref	1209		Ref	1403		Ref	932		Ref	3413		Ref
	Non-Hispanic black	1148	-0.63	<2E-16	466	-0.63	<b>1.89E-09</b>	262	-0.58	<b>1.42E-07</b>	672	-0.69	<2E-16	833	-0.60	<2E-16
	Mexican American	543	-0.11	.11	191	-0.08	.58	104	-0.26	.12	429	-0.27	<b>.005</b>	672	-0.13	.04
	Other Hispanic	471	-0.12	.09	185	-0.18	.21	123	-0.26	.09	273	-0.21	.05	1873	-0.19	.003
	Other race	252	-0.03	.77	114	-0.05	.77	78	-0.22	.22	185	-0.38	.002	477	-0.21	.004
Education level	Less than 12th grade	1426		Ref	681		Ref	360		Ref	822		Ref	1870		Ref
	High school/GED or equivalent	2984	0.10	<b>.04</b>	1150	0.15	.10	1053	0.10	.32	1264	0.13	.07	3982	0.12	<b>.004</b>
	College graduate or above	1056	0.15	<b>.03</b>	332	0.44	<b>.001</b>	557	0.30	<b>.02</b>	401	0.31	<b>.005</b>	1408	0.18	<b>.002</b>
Marital status	Married	2914		Ref	1122		Ref	1156		Ref	1392		Ref	3885		Ref
	Widowed	773	2.40E-01	<b>5.00E-04</b>	381	5.30E-01	<b>5.27E-05</b>	307	0.18	.19	314	0.05	.68	906	0.19	<b>.003</b>
	Divorced	869	-0.07	.27	324	0.05	.69	266	0.06	.64	364	-0.07	.47	1058	-0.01	.81
	Separated	189	0.1	.36	76	0.29	.18	57	0.35	.12	92	0.05	.77	274	0.20	<b>.04</b>
	Never married	467	0.05	.52	169	0.02	.92	124	0.05	.76	225	-0.14	.23	736	0.05	.40
	Living with partner	255	0.03	.73	93	-0.06	.76	60	0.09	.65	102	-0.13	.42	406	-0.01	.89
People in household	1	1168		Ref	503		Ref	464		Ref	473	0.45		1444		Ref
	2	2232	-0.07	.25	926	0.08	.53	956	-0.0006	.99	917	0.01	.92	2797	-0.01	.83
	3	784	-0.05	.47	289	0.09	.54	240	-0.06	.68	408	-0.12	.33	1125	-0.06	.34
	4	603	-0.07	.36	208	-0.14	.39	143	-0.10	.56	313	0.12	.33	849	0.03	.73
	5	356	-0.12	.18	115	-0.10	.62	87	-0.11	.60	176	-0.09	.55	526	-0.01	.89
	6	171	0.01	.92	64	0.04	.89	48	0.09	.73	101	0.12	.50	269	-0.12	.26
	7+	156	-0.17	.19	60	-0.06	.80	32	0.36	.24	103	-0.2	.28	258	-0.11	.29
	Income	< \$20,000	1455		Ref	694		Ref	394		Ref	666		Ref	1781	
\$20,000 to \$44,999	1571	0.007	.90	679	0.03	.77	585	-0.05	.64	775	0.11	.20	2138	0.05	.26	
\$45,000 to \$74,999	959	0.007	.91	358	0.02	.86	369	-0.12	.32	450	-0.09	.36	1305	-0.07	.22	
≥ \$75,000	1137	-0.04	.60	299	-0.13	.36	504	-0.17	.18	422	-0.23	.04	1570	-0.07	.25	
Health insurance	Yes	4870		Ref	1950		Ref	1841		Ref	2146		Ref	6259		Ref
	No	593	-0.04	.49	210	-0.05	.73	127	-0.02	.92	344	-0.004	.97	1002	-0.05	.37
Ever smoker	Yes	3324		Ref	1469		Ref	1184		Ref	1431		Ref	4129		Ref
	No	2141	-0.07	.09	696	-0.16	.06	784	-0.05	.50	1059	-0.11	.10	3132	-0.05	.16
Alcohol use I (drinking days per year)	None	1689		Ref	867		Ref	565		Ref	971		Ref	2136		Ref
	≤10	1295	-0.1	.06	459	-0.19	.07	407	-0.05	.64	635	-0.05	.55	1628	-0.10	<b>.04</b>
	>10 and ≤50	854	-0.01	.85	280	0.04	.77	316	-0.18	.10	373	0.05	.63	1221	-0.07	.21
	>50 and ≤100	431	-0.05	.48	150	-0.12	.45	164	0.10	.48	186	0.13	.31	609	-0.11	.11
	>100	1191	-0.06	.26	404	-0.21	.05	516	-0.13	.21	321	0.02	.83	1658	-0.06	.20
BMI	≤18.5	56		Ref	32		Ref	29		Ref	10		Ref	58		Ref
	>18.5 and ≤25	1095	1E-05	.99	430	-0.01	.97	517	-0.05	.64	300	-0.64	.19	1234	-0.19	.33
	>25 and ≤30	1675	-0.16	.99	692	-0.34	.30	680	-0.20	.10	698	-0.74	.12	2305	-0.33	.08
	>30	2635	-0.14	.35	1011	-0.34	.30	744	-0.28	.48	1483	-0.78	.10	3671	-0.35	.07
Physical activity	None	3338		Ref	1448		Ref	1072		Ref	1613		Ref	4288		Ref
	Moderate only	1518	-0.16	<b>.0004</b>	565	-0.08	.39	658	-0.17	<b>.03</b>	686	-0.18	<b>.01</b>	2043	-0.14	<b>.0006</b>
	Vigorous only	165	-0.15	.18	35	-0.23	.45	75	-0.34	.07	64	-0.04	.83	293	-0.20	<b>.03</b>

	Moderate and vigorous	446	-0.16	<b>.03</b>	115	-0.22	.22	165	-0.31	<b>.02</b>	125	-0.11	.43	642	-0.20	<b>.003</b>
Time sitting/ reclining	≤1 h	662		Ref	218		Ref	183		Ref	318		Ref	884		Ref
	>1 h and ≤2 h	2615	0.05	.45	997	0.11	.42	957	0.11	.39	1130	-0.02	.87	3340	0.07	.22
	>2 h and ≤4 h	1980	0.16	<b>.01</b>	852	0.21	.13	761	0.19	.15	931	0.26	<b>.01</b>	2768	0.16	<b>.005</b>
	>4 h	191	-0.001	.99	89	0.09	.71	62	-0.0004	.99	101	0.09	.61	240	0.13	.23
Arthritis	Yes				1188		Ref	1016		Ref	1155		Ref	3237		Ref
	No				970	-0.08	.32	950	-0.06	.40	1332	0.04	.58	4012	-0.01	.69
Heart condition	Yes	1188		Ref				472		Ref	681		Ref	1599		
	No	4275	-0.22	<b>5.24E-06</b>				1497	-0.05	.55	1804	-0.16	<b>.02</b>	5658	-0.23	<b>1.54E-07</b>
Emphysema	Yes	294		Ref	194		Ref	118		Ref	119		Ref	290		Ref
	No	5166	-0.27	<b>.002</b>	1967	-0.25	.08	1850	-0.56	<b>.00027</b>	2365	-0.12	.40	6960	-0.36	<b>5.71E-05</b>
Cancer	Yes	1016		Ref	472		Ref				367		Ref	1135		Ref
	No	4447	-0.14	<b>.004</b>	1690	0.01	.90				2120	-0.21	<b>.02</b>	6124	-0.17	<b>.0006</b>
Diabetes	Yes	1155		Ref	681		Ref	367		Ref				1739		Ref
	No	4103	-0.12	<b>.03</b>	1393	-0.14	.12	1530	-0.25	<b>.01</b>				5231	-0.15	<b>.0002</b>
	Borderline	208	-0.23	<b>.03</b>	89	0.04	.85	72	-0.53	<b>.009</b>				291	-0.10	.29
Hypertension	Yes	3237		Ref	1599		Ref	1135		Ref	1739		Ref			
	No	2228	-0.08	<b>.05</b>	564	-0.23	<b>.01</b>	832	-0.19	<b>.02</b>	749	-0.07	.29			

Bold font indicates items which remain significant after multivariable adjustment.

BMI = body mass index; NLR = neutrophil-to-lymphocyte ratio.

\* Participant counts in the multivariable model reflect only participants with complete data for all included variables. Note that participants reporting "refused/don't know" are excluded from this table for brevity, and thus slight variations from these participant totals may be observed.

**Table S4**  
Multivariable analysis as presented in Table 4 repeated with age treated as a continuous variable

Factor	Level	Univariate			Multivariable		
		N (level) *	Coefficient	P-value	N (level) *	Coefficient	P-value
Sex	Male	23,206		Ref	10,704		Ref
	Female	24,817	−0.03	.008	9533	−0.13	<b>7.55E-12</b>
Age	<60	33,039		Ref	13,685		
	≥60	14,984	0.28	<2E-16	6552		
Race	Continuous	48,023	0.0066	<2E-16	20,237	0.004	<b>5.59E-07</b>
	Non-Hispanic white	21,254		Ref	9251		Ref
	Non-Hispanic black	9928	−0.59	<2E-16	4066	−0.53	<b>&lt;2E-16</b>
	Mexican American	9138	−0.20	<2E-16	3019	−0.10	<b>5.43E-04</b>
	Other Hispanic	3847	−0.28	<2E-16	2089	−0.15	<b>3.30E-07</b>
Education level	Other race	3856	−0.31	<2E-16	1812	−0.21	<b>3.36E-11</b>
	Less than 12th grade	12,430		Ref	4674		Ref
	High school/GED or equivalent	22,657	0.009	.51	10,707	0.058	<b>.01</b>
Marital status	College graduate or above	9452	−0.003	.84	4843	0.076	<b>.01</b>
	Married	23,641		Ref	10,419		Ref
	Widowed	3856	0.27	<2E-16	1391	0.19	<b>1.25E-06</b>
	Divorced	4465	−0.005	.8	2355	0.03	.31
	Separated	1468	−0.04	.23	683	0.12	<b>.02</b>
People in household	Never married	9432	−0.10	3.64E-11	3691	0.09	<b>.002</b>
	Living with partner	3323	−0.12	1.06E-07	1692	−0.02	.47
	1	6391		Ref	2900		Ref
	2	14,156	−0.06	2.00E-03	6261	0.02	.48
	3	8683	−0.17	<2E-16	3621	0.02	.49
	4	7857	−0.21	<2E-16	3284	0.05	.22
	5	5240	−0.23	<2E-16	2065	0.03	.41
	6	2647	−0.21	5.59E-13	1051	0.07	.14
Income	7+	3049	−0.27	<2E-16	1055	−0.05	.26
	Continuous	48,023	−0.05	<2E-16	20,237		
	<\$20,000	10,662		Ref	4015		Ref
	\$20,000 to \$44,999	13,860	−0.05	.001	5701	0.003	.92
	\$45,000 to \$74,999	8762	−0.08	5.67E-06	3855	−0.04	.13
Health insurance	≥\$75,000	10,327	−0.13	9.27E-15	5295	−0.07	<b>.03</b>
	Yes	37,185		Ref	15,746		Ref
Ever smoker	No	10,601	−0.15	<2E-16	4474	−0.03	.18
	Yes	20,596		Ref	10,218		Ref
Alcohol use 1 (drinking days per year)	No	24,494	−0.14	<2E-16	10,002	−0.06	<b>.001</b>
	None	7988		Ref	4306		Ref
	≤10	7995	−0.16	1.57E-15	4522	−0.08	<b>.002</b>
	>10 and ≤50	7209	−0.18	<2E-16	4226	−0.04	.16
	>50 and ≤100	3838	−0.19	3.41E-15	2243	−0.08	<b>.02</b>
	>100	8263	−0.11	2.65E-08	4915	−0.02	.47
	Continuous	35,293	−0.02	1.72E-06	20,212		
BMI	≤18.5	914		Ref	307		Ref
	>18.5 and ≤25	14,285	−0.01	.77	5558	−0.11	.13
	>25 and ≤30	15,676	−0.02	.66	6740	−0.17	<b>.01</b>
	>30	16,300	−0.02	.57	7632	−0.16	<b>.02</b>
	Continuous	47,175	−0.09	.4	20,237		
Physical activity	None	14,627		Ref	10,280		Ref
	Moderate only	7136	−0.08	1.67E-05	5360	−0.08	<b>.0003</b>
	Vigorous only	2257	−0.26	<2E-16	1560	−0.12	<b>.0006</b>
	Moderate and vigorous	4001	−0.23	<2E-16	3030	−0.12	<b>6.17E-06</b>
Time sitting/reclining	≤1 h	4371		Ref	2977		Ref
	>1 h and ≤2 h	12,820	0.08	9.47E-05	9412	0.04	.11
	>2 h and ≤4 h	9789	0.15	5.51E-12	7138	0.11	<b>2.89E-05</b>
	>4 h	903	0.16	.0003	648	0.05	.37
	Continuous	27,883	0.07	5.88E-13	20,175		
Arthritis	Yes	11,739		Ref	5470		Ref
	No	32,775	−0.19	<2E-16	14,728	−0.01	.6
Heart condition	Yes	4925		Ref	2165		Ref
	No	39,611	−0.43	<2E-16	18,053	−0.21	<b>1.03E-12</b>
Emphysema	Yes	910		Ref	473		Ref
	No	43,630	−0.72	<2E-16	19,744	−0.33	<b>1.51E-08</b>
Cancer	Yes	3997		Ref	1970		Ref
	No	40,553	−0.37	<2E-16	18,250	−0.15	<b>1.60E-06</b>
Diabetes	Yes	5252		Ref	2491		Ref
	No	41,876	−0.20	<2E-16	17,268	−0.15	<b>3.18E-08</b>
	Borderline	862	−0.13	.003	465	−0.17	<b>.004</b>
Hypertension	Yes	15,457		Ref	7268		Ref
	No	32,330	−0.17	<2E-16	12,946	−0.06	<b>.004</b>

Bold font indicates items which remain significant after multivariable adjustment.

BMI = body mass index; NLR = neutrophil-to-lymphocyte ratio.

\* Univariate models include all available participants. Lower N values in the multivariable model reflect restriction to only participants with complete data for all included variables. Note that participants reporting “refused/don’t know” are excluded from this table for brevity, and thus slight variations from these participant totals may be observed.




Fig. S1. CONSORT flow diagram of study cohort identification.




Fig. S3. Median and range of neutrophils and lymphocytes as a percentage of total WBC for the 9 years of NHANES included in the present study.




Fig. S2. Correlation between NLR defined by the ratio of absolute count of neutrophils to absolute count of lymphocytes, and NLR defined by the ratio of percentage neutrophils to percentage lymphocytes (of total WBCs), demonstrating near-perfect correlation.




Fig. S4. Distribution (density) of neutrophils, lymphocytes and NLR across the 48,023 NHANES participants included in the present study, demonstrating skewed (non-normal) distributions for all three variables.