




ELSEVIER

Contents lists available at ScienceDirect

Journal of Neuroimmunology

journal homepage: www.elsevier.com/locate/jneuroim

Review article

Sodium chloride triggers Th17 mediated autoimmunity

Stefanie Haase^a, Nicola Wilck^{b,c,d,e,f}, Markus Kleinewietfeld^g, Dominik N. Müller^{b,c,d,e,f}, Ralf A. Linker^{a,*}^a Department of Neurology, Friedrich-Alexander University Erlangen, Germany^b Experimental and Clinical Research Center, A Joint Cooperation of Max-Delbrück Center for Molecular Medicine and Charité-Universitätsmedizin Berlin, Germany^c Charité-Universitätsmedizin Berlin, Germany^d Max-Delbrück Center for Molecular Medicine in the Helmholtz Association, Germany^e DZHK (German Centre for Cardiovascular Research), Partner Site Berlin, Germany^f Berlin Institute of Health (BIH), Berlin, Germany^g VIB Laboratory of Translational Immunomodulation, Center for Inflammation Research (IRC), Hasselt University, Diepenbeek, Belgium

A B S T R A C T

The detrimental effects of a high-salt diet on human health have received much attention in the past few years. While it has been well established that high dietary salt intake is related to cardiovascular diseases, there is growing evidence that excess salt also affects the immune system and might be considered as a risk factor in autoimmune diseases such as multiple sclerosis (MS). Several studies have implicated T helper 17 cells (Th17) in the pathogenesis of MS. We and others recently demonstrated that excessive salt enhances the differentiation of Th17 cells, inducing a highly pathogenic phenotype that aggravates experimental neuroinflammation. Moreover, a diet rich in sodium affects intestinal microbiota alongside increased intestinal Th17 cells, thus linking the detrimental effects of high salt consumption to the gut-immune axis. First human studies revealed an association of increased MS disease activity with elevated sodium chloride consumption, while more recent epidemiology studies in larger cohorts suggest no correlation between salt intake and MS. However, it is known that ordinary urinary sodium analyses and nutritional questionnaires do not necessarily correspond to the actual sodium load and more sophisticated analyses are needed. Moreover, studies revealed that sodium can temporarily be stored in the body. This review summarizes recent findings on the impact of salt on the immune system and discusses potential challenges investigating dietary salt intake as a risk factor in MS.

1. Introduction

A vast number of animal and human studies revealed a role for CD4+ T helper 17 (Th17) cells and their downstream pathways as important drivers of autoimmune diseases. Indeed, Th17 cells were already implicated in the development of inflammatory bowel disease (Gálvez, 2014), rheumatoid arthritis (Annunziato et al., 2009; Azizi et al., 2013), psoriasis (Annunziato et al., 2009) and also multiple sclerosis (MS). First studies using the murine experimental autoimmune encephalomyelitis (EAE) model revealed that interleukin (IL) 23 deficient mice are protected from the disease (Cua et al., 2003). Thus, IL-23 directly acts on T cells, promoting the induction of Th17 cells (Aggarwal et al., 2003). In accordance, the transfer of myelin antigen-specific Th17 cells was very potent in inducing EAE in naive recipient mice (Jäger et al., 2009; Langrish et al., 2005). Strikingly, Th17 cells most frequently occur in the intestine, where they help to protect against extracellular pathogens and maintain the barrier function of the

gut (Conti et al., 2009). Yet, depending on the modulating factors they are exposed to, Th17 cells can also have pro-inflammatory properties, characterized by the up-regulation of IL-23 receptor (IL-23R) and an increased secretion of IL-17A, granulocyte-macrophage colony-stimulating factor (GM-CSF), tumor necrosis factor (TNF) α and IL-2 (Ghoreschi et al., 2010). Dietary components might represent such modulating factors, implicating the diet as a potential risk factor in autoimmune diseases. The excess consumption of salt (sodium chloride) has repeatedly been associated with an increased risk of developing high blood pressure (Mozaffarian et al., 2014), disturbances in normal kidney function (Fellner et al., 2014), cancer (D'Elia et al., 2014) and chronic inflammation (Sundström et al., 2015). Moreover, there is growing evidence that excess salt affects the immune system, and first studies assume that high salt enhances MS disease activity (Farez et al., 2015). It is thus not surprising that salt restriction is widely viewed as an essential part of a healthy life style. The latest World Health Organization guidelines recommend adults to consume less than 5 g salt or

* Corresponding author at: Department of Neurology, University Hospital Erlangen, Schwabachanlage 6, D-91054 Erlangen, Germany.
E-mail address: ralf.linker@uk-erlangen.de (R.A. Linker).

2 g sodium per day (Guideline: Sodium Intake for Adults and Children, 2012). However, the estimated average salt intake is about two to three times higher as the recommended maximum level (Brown et al., 2009), rendering it a potential risk factor in various diseases, including MS.

2. High-salt effects on immune cells

Several immune cell subtypes are potentially important in mediating the detrimental effects of excess sodium chloride. First studies investigated a possible effect of high salt concentrations on the innate immune system. While dendritic cells do not respond to excess sodium chloride *in vitro* and *in vivo* (Jörg et al., 2016), monocytes acquire a highly pro-inflammatory state (Shapiro and Dinarello, 1995). RNA analyses revealed the induction of a set of pro-inflammatory genes, including chemokines, cytokines, and chemokine receptors, while genes associated with anti-inflammatory functions were down-regulated. More precisely, high-salt activated pro-inflammatory M1 macrophages (Müller et al., 2013; Zhang et al., 2015) while it decreased the ability of regulatory M2 macrophages to suppress effector T cell proliferation (Binger et al., 2015). Moreover, high salt had direct stimulatory effects on the migration of macrophages (Müller et al., 2013). Mechanistically, high salt-induced pro-inflammatory macrophage activation required p38/MAPK and downstream signaling (Jantsch et al., 2015), whereas the blockade of regulatory macrophage activation depended on the impairment of the protein kinase AKT and the mechanistic target of rapamycin (mTOR) (Binger et al., 2015). Further studies revealed that high salt-induced pathways also play an important role in T cell immunology. Already two decades ago, it was shown that increasing sodium chloride conditions by approximately 40 mM boosted IL-2 expression and T cell proliferation (Coimbra et al., 1995; Junger et al., 1994). Moreover, excess salt does not only favor T cell proliferation in general, but also affects the polarization of T cells: High sodium concentrations increased the differentiation of murine and human Th17 cells and induced a highly pathogenic phenotype, characterized by an increased expression of the surface receptors IL-23R and chemokine receptor CCR6 and the upregulation of GM-CSF, IL-2 and TNF α (Kleinewietfeld et al., 2013; Wu et al., 2013). The osmoprotective transcription factor tonicity enhancer binding protein (TonEBP/NFAT5) mediated the pro-inflammatory response by signaling through the serum/glucocorticoid-regulated kinase 1 (SGK1). Other studies also implicated SGK1 in sodium homeostasis and revealed its expression on a wide array of immune cells, including Th17 cells (Wu et al., 2013). Indeed, most studies discussing the mechanism for the detrimental effect of high salt demonstrated a pivotal involvement of pathogenic Th17 cells. In a murine model of colitis, high dietary sodium increased the level of Th17 cells in the gut and exacerbated colitis (Tubbs et al., 2017; Wei et al., 2017). Moreover, high sodium chloride intake enhanced disease severity in mouse models of lupus nephritis by the induction of Th17 and Th1 cells (Yang et al., 2015). In the same study, high salt concentrations up-regulated Th17 cells in CD4+ T cells isolated from systemic lupus erythematosus patients in an SGK1 dependent manner, indicating that high salt may also affect lupus erythematosus in humans (Yang et al., 2015). Immunologic effects of high salt consumption have also been linked to transplantation (Safa et al., 2015): Feeding a high-salt diet in a mouse model of solid organ transplantation accelerated cardiac allograft rejection without affecting blood pressure or serum sodium concentrations. The accelerated rejection was associated with a reduction of regulatory T cells (Tregs) and a significant decrease of Treg proliferation, resulting in an increase of CD4+ T cells in mice fed a high-salt diet (Safa et al., 2015). Accordingly, increased sodium chloride concentrations also impaired murine and human Treg cell function *in vitro* and *in vivo* (Hernandez et al., 2015). High-salt induced a pro-inflammatory Treg phenotype associated with a heightened interferon (IFN)- γ secretion and a diminished suppressive capacity (Hernandez et al., 2015). These studies provide evidence that, in addition to the increased induction of pro-

inflammatory Th17 cells, excess dietary sodium intake can impact autoimmunity by inhibiting the function of Tregs. Since it is well accepted that the autoimmune basis of MS stems from an imbalance between pro-inflammatory Th1 and Th17 cells and anti-inflammatory Tregs (Viglietta et al., 2004), salt may be added to the list of risk factors affecting the disease. Indeed, studies in EAE as an animal model of MS, showed that a high-salt diet augmented disease onset and severity, which was accompanied by an increased infiltration of pathogenic Th17 cells in the spinal cord (Hammer et al., 2017; Jörg et al., 2016; Kleinewietfeld et al., 2013; Wu et al., 2013).

3. High-salt affects the gut microbiome

In vitro studies investigating the effect of high-salt on various immune cells used comparable sodium chloride concentrations to those measured in the intestinal lumen after ingesting a high-salt diet. Moreover, before being absorbed and distributed to tissues, sodium chloride passes the gastrointestinal tract (Jose et al., 2016). Indeed, studies in models of inflammatory bowel disease imply the gut as relevant organ mediating effects of salt on the immune system (Monteleone et al., 2017; Wei et al., 2017). It was shown that a high-salt diet increased the frequency of IL-17A producing cells in the intestinal lamina propria compared to mice on a normal salt diet (Wei et al., 2017). Moreover, ingestion of a high-salt diet significantly inhibited the secretion of IL-10 and the suppressive function of Tregs in the small intestine, while not affecting the percentage of Treg cells (Wei et al., 2017). These results were also corroborated in human studies where exposure of human intestinal mononuclear cells to high sodium chloride concentrations enhanced the production of IL-17A, IL-23R, TNF α and ROR γ t (Monteleone et al., 2017).

While not expecting a direct correlation between the gut and brain autoimmunity, the potential importance of the gut microbiome for the development of MS and EAE has recently been recognized (Berer et al., 2011; Jangi et al., 2016; Ochoa-Repáraz et al., 2009; Yokote et al., 2008). MS patients display an altered microbiome composition compared to healthy controls. Interestingly, gut microbes rapidly respond to fluctuations in dietary composition (David et al., 2014). It is thus likely that dietary factors could lead to alterations of the gut microbiome. Furthermore, diet-induced shifts in microbiome composition can have profound effects on the host immune system (Haghikia et al., 2015; Turnbaugh et al., 2006), including T cells (Arpaia et al., 2013; Monteleone et al., 2017; Wei et al., 2017). Th17 cells are particularly affected by the abundance of specific commensal bacteria (Ivanov et al., 2009). Thus, the gut and the gut microbiota represent potential targets linking high salt intake, Th17 cell induction and MS. Indeed, we recently aimed to unravel a link between high-salt intake, gut microbiome composition and Th17 induced central nervous system autoimmunity (Wilck et al., 2017). High-salt diet significantly increased fecal sodium concentrations, what was paralleled by alterations in the fecal metabolome and gut microbiome composition. Several intestinal bacteria were affected by the high-salt concentrations; particularly a *Lactobacillus* strain was suppressed. We further hypothesized that the depletion of *Lactobacillus* by high salt might be linked to the observed induction of intestinal and systemic Th17 cells. Indeed, treatment with *Lactobacillus* prevented the salt-induced increase of Th17 cells, thereby preventing the exacerbation of EAE. Mice on a high-salt diet concomitantly receiving *Lactobacillus* displayed less severe EAE symptoms, with a reduced infiltration of Th17 cells in the spinal cord and a decreased production of IL-17A in the intestine and spleen. Interestingly, high salt induced increase of Th17 cells was paralleled by reduced concentrations of the fecal tryptophan metabolite indole-3-lactic acid (ILA), a bacterial fermentation product produced by *Lactobacilli* (Zelante et al., 2013) that has been implicated in the suppression of central nervous system autoimmunity (Rothhammer et al., 2016). Finally, we could show that ILA inhibited Th17 cell polarization *in vitro*. ILA may thus link the high-salt diet induced suppression of *Lactobacillus* to the

induction of Th17 cells. Indicating a potential relevance of these findings in humans, we showed that a salt challenge in healthy humans affects the abundance of intestinal *Lactobacillus*, alongside increased frequencies of pro-inflammatory Th17 cells in the blood. Yet, it is unclear whether *Lactobacillus* mediated high-salt effects can be transferred from the EAE model to MS. These limitations may also extend to the recent controversy on more general effects of a high-salt diet in neuroinflammation.

4. Linking excess salt to human disease

Recently published studies investigating high dietary salt intake as a potential risk factor in MS yielded controversial results. In an observational study of two independent cohorts of 70 MS patients, high dietary salt intake resulted in higher relapse rates as well as increased numbers of new MRI lesions compared to patients with moderate dietary salt intake (Farez et al., 2015). Accordingly, high salt has been implicated to influence human immune responses in general. Long-time salt intake positively correlated with the amount of monocytes and was paralleled by a significant increase of IL-6 and IL-23 production while the secretion of IL-10 was decreased (Yi et al., 2015). Intriguingly, IL-6 and IL-23 are major inducers of Th17 cells, also in humans. Although T cell numbers in the blood did not correlate with the amount of dietary salt consumed, plasma levels of IL-17 decreased with a reduction in dietary salt consumption (Yi et al., 2015). The results from a short-duration study on changes in dietary salt intake in humans were largely consistent with these findings reporting salt-related monocyte variations (Zhou et al., 2013). In contrast to this positive correlation between excess salt consumption and MS, two studies of pediatric MS revealed no association between high dietary sodium intake and the disease (McDonald et al., 2016; Nourbakhsh et al., 2016). The investigators correlated an estimated sodium intake with time to relapse (Nourbakhsh et al., 2016) or an increased risk of developing MS (McDonald et al., 2016) and found no association. However, in both studies, sodium intake was estimated using a basal food frequency questionnaire (FFQ), which may be unreliable and underestimate the daily sodium intake (Shim et al., 2014). Furthermore, the questionnaire was administered only at baseline and thus could not account for changes in diet during the period of investigation. General limitations of FFQs might be the long period of investigation of more than 12 months and the missing consideration of salt intake from „dining out“ and takeaway food. It would thus be necessary to establish a good quality FFQ with shorter duration of less than 12 months and the inclusion of regional dietary habits.

Further controversial findings regarding the risk of high salt intake and MS come from novel epidemiological studies investigating the effect of high salt intake on adult MS disease course or activity (Cortese et al., 2017; Fitzgerald et al., 2017). Large-scale epidemiologic studies based on the Nurses' Health Study and Nurses' Health Study II did not show any correlation between salt intake and the risk of MS (Cortese et al., 2017). Sodium intake was assessed from food frequency questionnaires at baseline and every 4 years during follow-up. The study thus relies on a large dataset. However, as already mentioned above, food frequency questionnaires represent a retrospective method that relies upon the respondent's memory. Participants were asked for their average portion size and the frequency of intake of foods covering the main dietary sodium sources. Total sodium intake was therefore only calculated by summing the sodium content of ingested foods. It is thus likely that the estimated sodium intake by food questionnaires is not consistent with the actual total sodium intake, especially in the face of “hidden sodium” in processed food. Furthermore, another study also found no association between excess salt and MS. Based on multiple assessments of sodium in spot urine samples and standardized clinical and multi-resonance imaging (MRI) follow-up, Fitzgerald and colleagues suggested that salt intake does not influence the conversion from clinically isolated syndrome to MS (Fitzgerald et al., 2017). 465

participants in the BENEFIT (Betaferon/Betaseron in Newly Emerging Multiple Sclerosis for Initial Treatment) study provided 14 spot-urine samples over 5 years that were used to estimate the 24-h urinary sodium excretion with the mathematical ‘Tanaka equation’. However, spot urine collections may not represent a valuable tool to assess total salt intake (Dougher et al., 2016; Zhou et al., 2017). Comparing the known 24-h sodium excretion and the estimated from morning spot urine collections by ‘Tanaka equation’ revealed that sodium excretion was about 3 g different from those measured using 24-h sodium urine collections (Zhou et al., 2017). Moreover, every second accurately collected 24-h sodium excretion sample fails to detect a 3-g difference in sodium intake. In line with data from earlier dietary intervention studies (Luft et al., 1982) and more recently published clinical trials (Dougher et al., 2016; Lerchl et al., 2015; Zhou et al., 2017), these findings suggest that multiple 24-h collections are necessary to assess salt intake and single spot urine collections may thus not be a valuable tool. Strikingly, there is a considerable day-to-day variation in 24-h sodium excretion, even when dietary salt intake is known and fixed over several weeks or month (Rakova et al., 2013). Rather, it was shown that an equilibration between sodium intake and sodium excretion follows weekly or even monthly rhythms of accumulated and secreted sodium independent of dietary salt intake (Rakova et al., 2013). Moreover, the prevailing notion that excess dietary sodium leads to direct urinary excretion has been challenged by evidence of periodic sodium tissue storage in both rodents and human. Already 4 decades ago, Ivanova et al. reported that huge amounts of sodium can be stored in the body (Ivanova et al., 1978). This finding has recently been confirmed by others. Long-term studies showed that large amounts of sodium accumulate without commensurate changes in body weight, indicating a sodium storage without water retention (Heer et al., 2009; Rakova et al., 2013; Titze et al., 2002). Novel strategies to determine the body's sodium content revealed a sodium accumulation in the skin interstitium and the muscle, where sodium binds to negatively charged glycosaminoglycans (Titze et al., 2004; Titze et al., 2003). Initial studies in humans, measuring the sodium content by ²³Na-multi resonance imaging (²³Na-MRI), confirmed this specific accumulation in the skin (Kopp et al., 2013; Linz et al., 2015). While the mechanisms that drive dietary sodium accumulation in the skin remain elusive, it is known that the clearance is regulated by the immune system. Sodium accumulation in the skin interstitium activates NFAT5 in macrophages, leading to the secretion of mediators acting on the local lymphatic system (Machnik et al., 2009). Upon chemotactic stimuli, macrophages infiltrate the skin, where they secrete a soluble growth factor for lymphatic vessels, vascular endothelial growth factor-c (VEGF-C), leading to VEGFR-3 induced cutaneous lymph capillary hyperplasia (Joukov et al., 1996; Karkkainen et al., 2004). Disturbances in this mechanism have already been linked to hypertension and elevated blood pressure (Machnik et al., 2010). Whether increased salt concentrations in the skin are directly linked to central nervous system autoimmunity remains to be elucidated. Considering the salt-effects on macrophage immunology during MS and the persisting challenge to accurately determine the salt intake via salt excretion, the skin may represent an important organ for further investigations on salt and MS risk. In summary, current standard methods may be inappropriate to estimate sodium intake precisely. It is thus important to develop novel measures or more defined FFQs or a combination of both to better account for the above mentioned limitations.

5. Conclusion

A major challenge for all studies addressing the effect of dietary habits in MS is that it is not possible to account for all potential confounders: sodium intake may be linked to healthy or unhealthy food that may exaggerate the effect of sodium intake. In fact, processed foods or fast foods are one of the main sources of salt, containing up to 100-times more sodium than homemade meals (Brown et al., 2009). Besides

salt, fast food contains high amounts of fat, and it was recently shown that both ingredients together might represent a risk factor in neuroinflammation (Hammer et al., 2017). While no definite association between dietary restrictions and the modulation of autoimmune diseases have been established yet, a large proportion of patients already consider special diets as alternative therapeutic options. In general, the goal of a nutritional intervention in MS must be the control of inflammation. In part, this may be achieved by modulating the gut microbiome composition and the systemic immune responses.

Observations in animal models of MS, colitis, lupus, transplantation and others argue for a potential role of salt as a risk factor in diseases associated with a Treg an Th17 cell imbalance. Yet, human studies are still sparse or have a limited technical approach to accurately define the association between excess salt intake and MS severity. It is thus necessary to establish new methods to accurately measure the sodium intake and storage in the body. Alternative techniques such as direct non-invasive measurements of tissue sodium content in humans by ²³Na-MRI may provide more concise information on the relation between salt and disease.

Disclosures and acknowledgments

MK was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (640116), by a SALK-grant from the government of Flanders, Belgium and by an Odysseus-grant of the Research Foundation Flanders (FWO) (GOG1216FWO), Belgium.

References

- Aggarwal, S., Ghilardi, N., Xie, M.-H., de Sauvage, F.J., Gurney, A.L., 2003. Interleukin-23 promotes a distinct CD4 T cell activation state characterized by the production of interleukin-17. *J. Biol. Chem.* 278, 1910–1914. <http://dx.doi.org/10.1074/jbc.M207577200>.
- Annunzio, F., Cosmi, L., Liotta, F., Maggi, E., Romagnani, S., 2009. Type 17 T helper cells—origins, features and possible roles in rheumatic disease. *Nat. Rev. Rheumatol.* 5, 325–331. <http://dx.doi.org/10.1038/nrrheum.2009.80>.
- Arpaia, N., Campbell, C., Fan, X., Dikiy, S., van der Veeken, J., Deroos, P., Liu, H., Cross, J.R., Pfeffer, K., Coffey, P.J., Rudenski, A.Y., 2013. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature* 504, 451–455. <http://dx.doi.org/10.1038/nature12726>.
- Azizi, G., Jadidi-Niaragh, F., Mirshafiey, A., 2013. Th17 Cells in Immunopathogenesis and treatment of rheumatoid arthritis. *Int. J. Rheum. Dis.* 16, 243–253. <http://dx.doi.org/10.1111/1756-185X.12132>.
- Berer, K., Mues, M., Koutrolos, M., Rasbi, Z.A., Boziki, M., Johner, C., Wekerle, H., Krishnamoorthy, G., 2011. Commensal microbiota and myelin autoantigen cooperate to trigger autoimmune demyelination. *Nature* 479, 538–541. <http://dx.doi.org/10.1038/nature10554>.
- Binger, K.J., Gebhardt, M., Heinig, M., Rintisch, C., Schroeder, A., Neuhofer, W., Hilgers, K., Manzel, A., Schwartz, C., Kleinewietfeld, M., Voelkl, J., Schatz, V., Linker, R.A., Lang, F., Voehringer, D., Wright, M.D., Hubner, N., Dechend, R., Jantsch, J., Titze, J., Müller, D.N., 2015. High salt reduces the activation of IL-4- and IL-13-stimulated macrophages. *J. Clin. Invest.* 125, 4223–4238. <http://dx.doi.org/10.1172/JCI80919>.
- Brown, I.J., Tzoulaki, I., Candeias, V., Elliott, P., 2009. Salt intakes around the world: implications for public health. *Int. J. Epidemiol.* 38, 791–813. <http://dx.doi.org/10.1093/ije/dyp139>.
- Coimbra, R., Junger, W.G., Liu, F.C., Loomis, W.H., Hoyt, D.B., 1995. Hypertonic/hyperosmotic fluids reverse prostaglandin E2 (PGE2)-induced T-cell suppression. *Shock Augusta Ga* 4, 45–49.
- Conti, H.R., Shen, F., Nayyar, N., Stocum, E., Sun, J.N., Lindemann, M.J., Ho, A.W., Hai, J.H., Yu, J.J., Jung, J.W., Filler, S.G., Masso-Welch, P., Edgerton, M., Gaffen, S.L., 2009. Th17 cells and IL-17 receptor signaling are essential for mucosal host defense against oral candidiasis. *J. Exp. Med.* 206, 299–311. <http://dx.doi.org/10.1084/jem.20081463>.
- Cortese, M., Yuan, C., Chitnis, T., Ascherio, A., Munger, K.L., 2017. No association between dietary sodium intake and the risk of multiple sclerosis. *Neurology* 10 1212/WNL.0000000000004417. <https://doi.org/10.1212/WNL.0000000000004417>.
- Cua, D.J., Sherlock, J., Chen, Y., Murphy, C.A., Joyce, B., Seymour, B., Lucian, L., To, W., Kwan, S., Churakova, T., Zurawski, S., Wiekowski, M., Lira, S.A., Gorman, D., Kastelein, R.A., Sedgwick, J.D., 2003. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* 421, 744–748. <http://dx.doi.org/10.1038/nature01355>.
- David, L.A., Maurice, C.F., Carmody, R.N., Gootenberg, D.B., Button, J.E., Wolfe, B.E., Ling, A.V., Devlin, A.S., Varna, Y., Fischbach, M.A., Biddinger, S.B., Dutton, R.J., Turnbaugh, P.J., 2014. Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 505, 559–563. <http://dx.doi.org/10.1038/nature12820>.
- D'Elia, L., Galletti, F., Strazzullo, P., 2014. Dietary salt intake and risk of gastric cancer. *Cancer Treat. Res.* 159, 83–95. http://dx.doi.org/10.1007/978-3-642-38007-5_6.
- Dougher, C.E., Rifkin, D.E., Anderson, C.A., Smits, G., Persky, M.S., Block, G.A., Ix, J.H., 2016. Spot urine sodium measurements do not accurately estimate dietary sodium intake in chronic kidney disease. *Am. J. Clin. Nutr.* 104, 298–305. <http://dx.doi.org/10.3945/ajcn.115.127423>.
- Farez, M.F., Fiol, M.P., Gaitán, M.I., Quintana, F.J., Correale, J., 2015. Sodium intake is associated with increased disease activity in multiple sclerosis. *J. Neurol. Neurosurg. Psychiatry* 86, 26–31. <http://dx.doi.org/10.1136/jnnp-2014-307928>.
- Fellner, R.C., Cook, A.K., O'Connor, P.M., Zhang, S., Pollock, D.M., Incho, E.W., 2014. High-salt diet blunts renal autoregulation by a reactive oxygen species-dependent mechanism. *Am. J. Physiol. Ren. Physiol.* 307, F33–F40. <http://dx.doi.org/10.1152/ajprenal.00040.2014>.
- Fitzgerald, K.C., Munger, K.L., Hartung, H.-P., Freedman, M.S., Montalbán, X., Edan, G., Wicklein, E.-M., Radue, E.-W., Kappos, L., Pohl, C., Ascherio, A., BENEFIT Study Group, 2017. Sodium intake and multiple sclerosis activity and progression in BENEFIT. *Ann. Neurol.* 82, 20–29. <http://dx.doi.org/10.1002/ana.24965>.
- Gálvez, J., 2014. Role of Th17 cells in the pathogenesis of human IBD. *ISRN Inflamm.* 2014 928461. <https://doi.org/10.1155/2014/928461>.
- Ghoreschi, K., Laurence, A., Yang, X.-P., Tato, C.M., McGeachy, M.J., Konkel, J.E., Ramos, H.L., Wei, L., Davidson, T.S., Bouladoux, N., Grainger, J.R., Chen, Q., Kanno, Y., Watford, W.T., Sun, H.-W., Eberl, G., Shevach, E.M., Belkaid, Y., Cua, D.J., Chen, W., O'Shea, J.J., 2010. Generation of pathogenic T(H)17 cells in the absence of TGF- β signalling. *Nature* 467, 967–971. <http://dx.doi.org/10.1038/nature09447>.
- Guideline: Sodium Intake for Adults and Children, 2012. WHO Guidelines Approved by the Guidelines Review Committee. World Health Organization, Geneva.
- Haghikia, A., Jörg, S., Duscha, A., Berg, J., Manzel, A., Waschbisch, A., Hammer, A., Lee, D.-H., May, C., Wilck, N., Balogh, A., Ostermann, A.I., Schebb, N.H., Akkad, D.A., Grohme, D.A., Kleinewietfeld, M., Kempa, S., Thöne, J., Demir, S., Müller, D.N., Gold, R., Linker, R.A., 2015. Dietary fatty acids directly impact central nervous system autoimmunity via the small intestine. *Immunity* 43, 817–829. <http://dx.doi.org/10.1016/j.immuni.2015.09.007>.
- Hammer, A., Schliep, A., Jörg, S., Haghikia, A., Gold, R., Kleinewietfeld, M., Müller, D.N., Linker, R.A., 2017. Impact of combined sodium chloride and saturated long-chain fatty acid challenge on the differentiation of T helper cells in neuroinflammation. *J. Neuroinflammation* 14, 184. <http://dx.doi.org/10.1186/s12974-017-0954-y>.
- Heer, M., Frings-Meuthen, P., Titze, J., Boschmann, M., Frisch, S., Baecker, N., Beck, L., 2009. Increasing sodium intake from a previous low or high intake affects water, electrolyte and acid-base balance differently. *Br. J. Nutr.* 101, 1286–1294. <http://dx.doi.org/10.1017/S0007114508088041>.
- Hernandez, A.L., Kitz, A., Wu, C., Lowther, D.E., Rodriguez, D.M., Vudattu, N., Deng, S., Herold, K.C., Kuchroo, V.K., Kleinewietfeld, M., Hafler, D.A., 2015. Sodium chloride inhibits the suppressive function of FOXP3+ regulatory T cells. *J. Clin. Invest.* 125, 4212–4222. <http://dx.doi.org/10.1172/JCI81151>.
- Ivanov, I.I., Atarashi, K., Manel, N., Brodie, E.L., Shima, T., Karaoz, U., Wei, D., Goldfarb, K.C., Santee, C.A., Lynch, S.V., Tanoue, T., Imaoka, A., Itoh, K., Takeda, K., Umesaki, Y., Honda, K., Littman, D.R., 2009. Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* 139, 485–498. <http://dx.doi.org/10.1016/j.cell.2009.09.033>.
- Ivanova, L.N., Archibasova, V.K., Shterental, I.S., 1978. Sodium-depositing function of the skin in white rats. *Fiziol. Zh. SSSR Im. I. M. Sechenova* 64, 358–363. <http://dx.doi.org/10.1016/j.cell.2009.09.033>.
- Jäger, A., Dardalhon, V., Sobel, R.A., Bettelli, E., Kuchroo, V.K., 2009. Th1, Th17 and Th9 effector cells induce experimental autoimmune encephalomyelitis with different pathological phenotypes. *J. Immunol. Baltim. Md 1950* 183, 7169–7177. <http://dx.doi.org/10.4049/jimmunol.0901906>.
- Jangi, S., Gandhi, R., Cox, L.M., Li, N., von Glehn, F., Yan, R., Patel, B., Mazzola, M.A., Liu, S., Glanz, B.L., Cook, S., Tankou, S., Stuart, F., Melo, K., Nejad, P., Smith, K., Topçuoğlu, B.D., Holden, J., Kivisäkk, P., Chitnis, T., De Jager, P.L., Quintana, F.J., Gerber, G.K., Bry, L., Weiner, H.L., 2016. Alterations of the human gut microbiome in multiple sclerosis. *Nat. Commun.* 7, 12015. <http://dx.doi.org/10.1038/ncomms12015>.
- Jantsch, J., Schatz, V., Friedrich, D., Schröder, A., Kopp, C., Siegert, I., Maronna, A., Wendelborn, D., Linz, P., Binger, K.J., Gebhardt, M., Heinig, M., Neubert, P., Fischer, F., Teufel, S., David, J.-P., Neufert, C., Cavallaro, A., Rakova, N., Küper, C., Beck, F.-X., Neuhofer, W., Müller, D.N., Schuler, G., Uder, M., Bogdan, C., Luft, F.C., Titze, J., 2015. Cutaneous Na+ storage strengthens the antimicrobial barrier function of the skin and boosts macrophage-driven host defense. *Cell Metab.* 21, 493–501. <http://dx.doi.org/10.1016/j.cmet.2015.02.003>.
- Jörg, S., Kissel, J., Manzel, A., Kleinewietfeld, M., Haghikia, A., Gold, R., Müller, D.N., Linker, R.A., 2016. High salt drives Th17 responses in experimental autoimmune encephalomyelitis without impacting myeloid dendritic cells. *Exp. Neurol.* 279, 212–222. <http://dx.doi.org/10.1016/j.expneurol.2016.03.010>.
- Jose, P.A., Yang, Z., Zeng, C., Felder, R.A., 2016. The importance of the gastrorenal axis in the control of body sodium homeostasis. *Exp. Physiol.* 101, 465–470. <http://dx.doi.org/10.1113/EP085286>.
- Joukov, V., Pajusola, K., Kaipainen, A., Chilov, D., Lahtinen, I., Kukk, E., Saksela, O., Kalkkinen, N., Alitalo, K., 1996. A novel vascular endothelial growth factor, VEGF-C, is a ligand for the Flt4 (VEGFR-3) and KDR (VEGFR-2) receptor tyrosine kinases. *EMBO J.* 15, 290–298.
- Junger, W.G., Liu, F.C., Loomis, W.H., Hoyt, D.B., 1994. Hypertonic saline enhances cellular immune function. *Circ. Shock* 42, 190–196.
- Karkkainen, M.J., Haiko, P., Sainio, K., Partanen, J., Taipale, J., Petrova, T.V., Jeltsch, M., Jackson, D.G., Taliikka, M., Rauvala, H., Betsholtz, C., Alitalo, K., 2004. Vascular endothelial growth factor C is required for sprouting of the first lymphatic vessels from embryonic veins. *Nat. Immunol.* 5, 74–80. <http://dx.doi.org/10.1038/ni1013>.

- Kleinewietfeld, M., Manzel, A., Titze, J., Kvakan, H., Yosef, N., Linker, R.A., Müller, D.N., Hafler, D.A., 2013. Sodium chloride drives autoimmune disease by the induction of pathogenic TH17 cells. *Nature* 496, 518–522. <http://dx.doi.org/10.1038/nature11868>.
- Kopp, C., Linz, P., Dahlmann, A., Hammon, M., Jantsch, J., Müller, D.N., Schmieder, R.E., Cavallaro, A., Eckardt, K.-U., Uder, M., Luft, F.C., Titze, J., 2013. ²³Na magnetic resonance imaging-determined tissue sodium in healthy subjects and hypertensive patients. *Hypertension* 61, 635–640. <http://dx.doi.org/10.1161/HYPERTENSIONAHA.111.00566>.
- Langrish, C.L., Chen, Y., Blumenschein, W.M., Mattson, J., Basham, B., Sedgwick, J.D., McClanahan, T., Kastelein, R.A., Cua, D.J., 2005. IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *J. Exp. Med.* 201, 233–240. <http://dx.doi.org/10.1084/jem.20041257>.
- Lerchl, K., Rakova, N., Dahlmann, A., Rauh, M., Goller, U., Basner, M., Dinges, D.F., Beck, L., Agureev, A., Larina, I., Baranov, V., Morukov, B., Eckardt, K.-U., Vassilieva, G., Wabel, P., Vienken, J., Kirsch, K., Johannes, B., Krannich, A., Luft, F.C., Titze, J., 2015. Agreement between 24-hour salt ingestion and sodium excretion in a controlled environment novelty and significance. *Hypertension* 66, 850–857. <http://dx.doi.org/10.1161/HYPERTENSIONAHA.115.05851>.
- Linz, P., Santoro, D., Renz, W., Rieger, J., Ruehle, A., Ruff, J., Deimling, M., Rakova, N., Müller, D.N., Luft, F.C., Titze, J., Niendorf, T., 2015. Skin Sodium Measured with ²³Na MRI at 7.0 T. *NMR Biomed.* vol. 28. pp. 54–62. <http://dx.doi.org/10.1002/nbm.3224>.
- Luft, F.C., Fineberg, N.S., Sloan, R.S., 1982. Estimating dietary sodium intake in individuals receiving a randomly fluctuating intake. *Hypertension* 4, 805–808. <http://dx.doi.org/10.1161/01.HYP.4.6.805>.
- Machnik, A., Neuhofer, W., Jantsch, J., Dahlmann, A., Tammela, T., Machura, K., Park, J.-K., Beck, F.-X., Müller, D.N., Derer, W., Goss, J., Ziomber, A., Dietsch, P., Wagner, H., van Rooijen, N., Kurtz, A., Hilgers, K.F., Alitalo, K., Eckardt, K.-U., Luft, F.C., Kerjaschki, D., Titze, J., 2009. Macrophages regulate salt-dependent volume and blood pressure by a vascular endothelial growth factor-C-dependent buffering mechanism. *Nat. Med.* 15, 545–552. <http://dx.doi.org/10.1038/nm.1960>.
- Machnik, A., Dahlmann, A., Kopp, C., Goss, J., Wagner, H., van Rooijen, N., Eckardt, K.-U., Müller, D.N., Park, J.-K., Luft, F.C., Kerjaschki, D., Titze, J., 2010. Mononuclear phagocyte system depletion blocks interstitial tonicity-responsive enhancer binding protein/vascular endothelial growth factor C expression and induces salt-sensitive hypertension in rats. *Hypertension* 55, 755–761. <http://dx.doi.org/10.1161/HYPERTENSIONAHA.109.143339>.
- McDonald, J., Graves, J., Waldman, A., Lotze, T., Schreiner, T., Belman, A., Greenberg, B., Weinstock-Guttman, B., Aaen, G., Tillema, J.-M., Hart, J., Lulu, S., Ness, J., Harris, Y., Rubin, J., Candee, M., Krupp, L.B., Gorman, M., Benson, L., Rodriguez, M., Chitnis, T., Mar, S., Barcellos, L.F., Larai, B., Rose, J., Roalstad, S., Simmons, T., Casper, T.C., Waubant, E., 2016. A case-control study of dietary salt intake in pediatric-onset multiple sclerosis. *Mult. Scler. Relat. Disord.* 6, 87–92. <http://dx.doi.org/10.1016/j.msard.2016.02.011>.
- Monteleone, I., Marafini, I., Dinallo, V., Di Fusco, D., Troncone, E., Zorzi, F., Laudis, F., Monteleone, G., 2017. Sodium chloride-enriched diet enhanced inflammatory cytokine production and exacerbated experimental colitis in mice. *J. Crohns. Colitis* 11, 237–245. <http://dx.doi.org/10.1093/ecco-jcc/jjw139>.
- Mozaffarian, D., Fahimi, S., Singh, G.M., Micha, R., Khatibzadeh, S., Engell, R.E., Lim, S., Danaei, G., Ezzati, M., Powles, J., 2014. Global burden of diseases nutrition and chronic diseases expert group, Global sodium consumption and death from cardiovascular causes. *N. Engl. J. Med.* 371, 624–634. <http://dx.doi.org/10.1056/NEJMoa1304127>.
- Müller, S., Quast, T., Schröder, A., Hucke, S., Klotz, L., Jantsch, J., Gerzer, R., Hemmersbach, R., Kolanus, W., 2013. Salt-dependent chemotaxis of macrophages. *PLoS One* 8, e73439. <http://dx.doi.org/10.1371/journal.pone.0073439>.
- Nourbakhsh, B., Graves, J., Casper, T.C., Lulu, S., Waldman, A., Belman, A., Greenberg, B., Weinstock-Guttman, B., Aaen, G., Tillema, J.-M., Hart, J., Ness, J., Rubin, J., Krupp, L., Gorman, M., Benson, L., Rodriguez, M., Chitnis, T., Rose, J., Barcellos, L., Waubant, E., 2016. Network of pediatric multiple sclerosis centers, Dietary salt intake and time to relapse in paediatric multiple sclerosis. *J. Neurol. Neurosurg. Psychiatry* 87, 1350–1353. <http://dx.doi.org/10.1136/jnnp-2016-313410>.
- Ochoa-Repáraz, J., Mielcarz, D.W., Ditrio, L.E., Burroughs, A.R., Foureau, D.M., Haque-Begum, S., Kasper, L.H., 2009. Role of gut microbial microflora in the development of experimental autoimmune encephalomyelitis. *J. Immunol. Baltim. Md.* 1950 (183), 6041–6050. <http://dx.doi.org/10.4049/jimmunol.0900747>.
- Rakova, N., Jüttner, K., Dahlmann, A., Schröder, A., Linz, P., Kopp, C., Rauh, M., Goller, U., Beck, L., Agureev, A., Vassilieva, G., Lenkova, L., Johannes, B., Wabel, P., Moissl, U., Vienken, J., Gerzer, R., Eckardt, K.-U., Müller, D.N., Kirsch, K., Morukov, B., Luft, F.C., Titze, J., 2013. Long-term light simulation reveals infradian rhythmicity in human Na⁺ balance. *Cell Metab.* 17, 125–131. <http://dx.doi.org/10.1016/j.cmet.2012.11.013>.
- Rothhammer, V., Mascanfroni, I.D., Bunse, L., Takenaka, M.C., Kenison, J.E., Mayo, L., Chao, C.-C., Patel, B., Yan, R., Blain, M., Alvarez, J.I., Kébir, H., Anandasabapathy, N., Izquierdo, G., Jung, S., Obholzer, N., Pochet, N., Clish, C.B., Prinz, M., Prat, A., Antel, J., Quintana, F.J., 2016. Type I interferons and microbial metabolites of tryptophan modulate astrocyte activity and central nervous system inflammation via the aryl hydrocarbon receptor. *Nat. Med.* 22, 586–597. <http://dx.doi.org/10.1038/nm.4106>.
- Safa, K., Ohori, S., Borges, T.J., Uehara, M., Batal, I., Shimizu, T., Magee, C.N., Belizaire, R., Abdi, R., Wu, C., Chandraker, A., Riella, L.V., 2015. Salt accelerates allograft rejection through serum- and glucocorticoid-regulated Kinase-1-dependent inhibition of regulatory T cells. *J. Am. Soc. Nephrol. JASN* 26, 2341–2347. <http://dx.doi.org/10.1681/ASN.2014090914>.
- Shapiro, L., Dinarello, C.A., 1995. Osmotic regulation of cytokine synthesis in vitro. *Proc. Natl. Acad. Sci. U. S. A.* 92, 12230–12234.
- Shim, J.-S., Oh, K., Kim, H.C., 2014. Dietary assessment methods in epidemiologic studies. *Epidemiol. Health* 36. <http://dx.doi.org/10.4178/epih/e2014009>.
- Sundström, B., Johansson, I., Rantapää-Dahlqvist, S., 2015. Interaction between dietary sodium and smoking increases the risk for rheumatoid arthritis: results from a nested case-control study. *Rheumatol. Oxf. Engl.* 54, 487–493. <http://dx.doi.org/10.1093/rheumatology/keu330>.
- Titze, J., Mailet, A., Lang, R., Gunga, H.C., Johannes, B., Gauquelin-Koch, G., Kihm, E., Larina, I., Gharib, C., Kirsch, K.A., 2002. Long-term sodium balance in humans in a terrestrial space station simulation study. *Am. J. Kidney Dis. Off. J. Natl. Kidney Found.* 40, 508–516. <http://dx.doi.org/10.1053/ajkd.2002.34908>.
- Titze, J., Lang, R., Ilies, C., Schwind, K.H., Kirsch, K.A., Dietsch, P., Luft, F.C., Hilgers, K.F., 2003. Osmotically inactive skin Na⁺ storage in rats. *Am. J. Physiol. Ren. Physiol.* 285, F1108–F1117. <http://dx.doi.org/10.1152/ajprenal.00200.2003>.
- Titze, J., Shakibaei, M., Schaffhuber, M., Schulze-Tanzil, G., Porst, M., Schwind, K.H., Dietsch, P., Hilgers, K.F., 2004. Glycosaminoglycan polymerization may enable osmotically inactive Na⁺ storage in the skin. *Am. J. Physiol. Heart Circ. Physiol.* 287, H203–H208. <http://dx.doi.org/10.1152/ajpheart.01237.2003>.
- Tubbs, A.L., Liu, B., Rogers, T.D., Sartor, R.B., Miao, E.A., 2017. Dietary salt exacerbates experimental colitis. *J. Immunol. Baltim. Md.* 1950 (199), 1051–1059. <http://dx.doi.org/10.10049/jimmunol.1700356>.
- Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., Gordon, J.I., 2006. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 444, 1027–1131. <http://dx.doi.org/10.1038/nature05414>.
- Viglietta, V., Baecher-Allan, C., Weiner, H.L., Hafler, D.A., 2004. Loss of functional suppression by CD4⁺CD25⁺ regulatory T cells in patients with multiple sclerosis. *J. Exp. Med.* 199, 971–979. <http://dx.doi.org/10.1084/jem.20031579>.
- Wei, Y., Lu, C., Chen, J., Cui, G., Wang, L., Yu, T., Yang, Y., Wu, W., Ding, Y., Li, L., Uede, T., Chen, Z., Diao, H., 2017. High salt diet stimulates gut Th17 response and exacerbates TNBS-induced colitis in mice. *Oncotarget* 8, 70–82. <http://dx.doi.org/10.18632/oncotarget.13783>.
- Wilck, N., Matus, M.G., Kearney, S.M., Olesen, S.W., Forslund, K., Bartolomeaus, H., Haase, S., Mähler, A., Balogh, A., Markó, L., Vvedenskaya, O., Kleiner, F.H., Tsvetkov, D., Klug, L., Costea, P.I., Sunagawa, S., Maier, L., Rakova, N., Schatz, V., Neubert, P., Frätzer, C., Krannich, A., Gollasch, M., Grohme, D.A., Côte-Real, B.F., Gerlach, R.G., Basic, M., Typas, A., Wu, C., Titze, J.M., Jantsch, J., Boschmann, M., Dechend, R., Kleinewietfeld, M., Kempa, S., Bork, P., Linker, R.A., Alm, E.J., Müller, D.N., 2017. Salt-responsive gut commensal modulates TH17 axis and disease. *Nature* 551, 585–589. <http://dx.doi.org/10.1038/nature24628>.
- Wu, C., Yosef, N., Thalhammer, T., Zhu, C., Xiao, S., Kishi, Y., Regev, A., Kuchroo, V.K., 2013. Induction of pathogenic TH17 cells by inducible salt-sensing kinase SGK1. *Nature* 496, 513–517. <http://dx.doi.org/10.1038/nature11984>.
- Yang, X., Yao, G., Chen, W., Tang, X., Feng, X., Sun, L., 2015. Exacerbation of lupus nephritis by high sodium chloride related to activation of SGK1 pathway. *Int. Immunopharmacol.* 29, 568–573. <http://dx.doi.org/10.1016/j.intimp.2015.09.027>.
- Yi, B., Titze, J., Rykova, M., Feurecker, M., Vassilieva, G., Nichiporuk, I., Schelling, G., Morukov, B., Choukèr, A., 2015. Effects of dietary salt levels on monocytic cells and immune responses in healthy human subjects: a longitudinal study. *Transl. Res. J. Lab. Clin. Med.* 166, 103–110. <http://dx.doi.org/10.1016/j.trsl.2014.11.007>.
- Yokote, H., Miyake, S., Croxford, J.L., Oki, S., Mizusawa, H., Yamamura, T., 2008. NKT cell-dependent amelioration of a mouse model of multiple sclerosis by altering gut flora. *Am. J. Pathol.* 173, 1714–1723. <http://dx.doi.org/10.2353/ajpath.2008.080622>.
- Zelante, T., Iannitti, R.G., Cunha, C., De Luca, A., Giovannini, G., Pieraccini, G., Zecchi, R., D'Angelo, C., Massi-Benedetti, C., Fallarino, F., Carvalho, A., Puccetti, P., Romani, L., 2013. Tryptophan catabolites from microbiota engage aryl hydrocarbon receptor and balance mucosal reactivity via interleukin-22. *Immunity* 39, 372–385. <http://dx.doi.org/10.1016/j.immuni.2013.08.003>.
- Zhang, W.-C., Zheng, X.-J., Du, L.-J., Sun, J.-Y., Shen, Z.-X., Shi, C., Sun, S., Zhang, Z., Chen, X.-Q., Qin, M., Liu, X., Tao, J., Jia, L., Fan, H.-Y., Zhou, B., Yu, Y., Ying, H., Hui, L., Liu, X., Yi, X., Liu, X., Zhang, L., Duan, S.-Z., 2015. High salt primes a specific activation state of macrophages, M(Na). *Cell Res.* 25, 893–910. <http://dx.doi.org/10.1038/cr.2015.87>.
- Zhou, X., Zhang, L., Ji, W.-J., Yuan, F., Guo, Z.-Z., Pang, B., Luo, T., Liu, X., Zhang, W.-C., Jiang, T.-M., Zhang, Z., Li, Y.-M., 2013. Variation in dietary salt intake induces coordinated dynamics of monocyte subsets and monocyte-platelet aggregates in humans: implications in end organ inflammation. *PLoS One* 8, e60332. <http://dx.doi.org/10.1371/journal.pone.0060332>.
- Zhou, L., Tian, Y., Fu, J.-J., Jiang, Y.-Y., Bai, Y.-M., Zhang, Z.-H., Hu, X.-H., Lian, H.-W., Guo, M., Yang, Z.-X., Zhao, L.-C., 2017. Validation of spot urine in predicting 24-h sodium excretion at the individual level. *Am. J. Clin. Nutr.* 105, 1291–1296. <http://dx.doi.org/10.3945/ajcn.116.147553>.