

Journal of Voice

Volume 33, 2019

Subject Index

- Abductor spasmodic dysphonia, bilateral vocal fold medialization treatment for, 45–48
- Acoustic analysis
- of classical and popular female singers in sentimental melody, 255.e27–255.e37
 - of crying signal in infants with disabling hearing impairment, 946.e7–946.e13
 - of normal voice patterns in Pakistani adults, 124.e49–124.e58
 - of opera-chant singers, evaluation of supraglottic and subglottic activities including, 255.e1–255.e7
 - and perceptual, of adductor spasmodic dysphonia in Mandarin-speaking Chinese, 333–339
 - signal, dependencies and ill-designed parameters within high-speed videoendoscopy and, 811.e1–811.e12
 - of soccer fans in acute phonotrauma after the match, 129–134
 - voice, of pediatric vocal nodule patients using ratios calculated with biomedical image segmentation, 195–203
- Acoustic parameters
- and aerodynamic in canine larynges, effects of angle of epiglottis on, 627–633
 - and aerodynamic in untrained singers, comparing the exposure-response relationships of physiological and traditional vocal warm-ups on, 420–428
 - aerodynamics and electroglottographic parameters, lingering effects of straw phonation exercises on, 810.e5–810.e11
- Acoustic Voice Quality Index (AVCI)
- and DSI, influence of gender and age on, 340–345
 - in Korean language, validation of, 948.e1–948.e9
 - version 03.01 for Japanese-speaking population, 125.e1–125.e12
- Acoustics
- perceptual correlates of voice in Purohits, 804.e1–804.e4
 - aerodynamic, and self-rating measurements, effects of vocal loading and vocal rest using, 809.e11–809.e18
 - and aerodynamic characteristic of connected speech in patients with and without voice disorders, effects of variably occluded face mask on, 809.e1–809.10
 - and aerodynamics of choral singers, 803.e1–803.e5
 - effect of chant training on lateral thyrohyoid ligament morphology and, 802.e17–802.23
 - in epiglottic cyst, 497–500
 - features and glottal changes define two modes of singing in Peking opera, 583.e9–583.e14
 - perceptual and subjective voice outcomes in therapy success in patients with functional dysphonia, 317–324
 - and perceptual measures of an SATB chorus, effects of straw phonation protocol on, 80–86
 - and perceptual measures of choral and solo sound, effects of musician earplugs on, 87–95
 - and perceptual outcome measures following medialization laryngoplasty, comparison of pitch strength with, 795–800
 - vocal, and auditory-perceptual characteristics during fluctuations in estradiol levels during menstrual cycle, 536–544
 - voice assessments, of responder and nonresponder patients, impact of laryngopharyngeal reflux on, 929–939
- Adductor spasmodic dysphonia
- in Mandarin-speaking Chinese, acoustic and perceptual analyses of, 333–339
 - quality of life and voice changes after single injection in patients with, 721–727
 - reliability and validity of speech evaluation in, (letter), 49–51
 - reliability of high-speed videoendoscopic ratings of essential voice tremor and, 16–26
- ADSD. See Adductor spasmodic dysphonia
- Aerodynamics
- acoustic and self-rating measurements, effects of vocal loading and vocal rest using, 809.e11–809.e18
 - and acoustic characteristic of connected speech in patients with and without voice disorders, effects of variably occluded face mask on, 809.e1–809.10
 - and acoustic parameters in canine larynges, effects of angle of epiglottis on, 627–633
 - and acoustic parameters in untrained singers, comparing the exposure-response relationships of physiological and traditional vocal warm-ups on, 420–428
 - and acoustics of choral singers, 803.e1–803.e5
 - computational models of laryngeal, 385–400
 - electroglottographic and acoustic parameters, lingering effects of straw phonation exercises on, 810.e5–810.e11
 - of growl voice and reinforced falsetto in metal singing, 803.e7–803.e13
 - profile of muscle tension dysphonia, 812.e15–812.e18
 - of syllable and sentence productions in normal speakers, 297–301
 - voice assessments, of responder and nonresponder patients, impact of laryngopharyngeal reflux on, 929–939
- Affective Voice test, Persian, effect of Parkinson disease on emotion perception using, 580.e1–580.e9
- Age, and gender, influence on AVQI and DSI, 340–345
- Aging, with voice disorders, reliability and validity of Aging Voice Index for, 807.e7–807.e12
- Aging Voice Index, for aging with voice disorders, reliability and validity of, 807.e7–807.e12
- Airflow
- patterns of running speech in patients with voice disorders, 277–283
 - transglottal, in vowels and speech, effect of Parkinson disease tremor phenotype on cepstral peak prominence and, 580.e11–580.e19
- Airflow vibrato, dependence on pitch and loudness, 815–830
- Alexander disease, Botox injection for laryngeal dysfunction in, 728–729
- Allergy, in phonation, role of, 811.e19–811.e27
- Ambulatory care facility, private practice community voice, epidemiologic characteristics of treatment-seeking population in, 429–434
- Anatomy, comparative, of pig and human arytenoid cartilage, 620–626
- Anchors, natural voice, on vocal quality evaluation, effect of auditory perceptual training with, 220–225

- Anesthesia, topical, for endoscopic office-based procedures of upper aerodigestive tract, 732–746
- Anterior vocal fold plication, external approach to, 792–794
- Anxiety, preoperative, assessment and management of, 691–696
- Arytenoid cartilage, comparative anatomy of pig and human, 620–626
- Arytenoid dislocation, using pixel-valued cuneiform movement, detection of, 370–374
- Arytenoidectomy, endoscopic partial, for bilateral vocal fold paralysis, 751–758
- Atrophy, vocal folds, efficacy of two-dimensional scanning digital kymography in evaluation of, 554–560
- Attention deficit disorder with hyperactivity, vocal characteristics of school-aged children with and without, 945.e37–945.e45
- Audio recordings, of German standard text readings, detection of diplophonation in, 949.e1–949.e10
- Auditory feedback, and vocal rehabilitation, influence on prelingual hearing-impaired individuals post cochlear implant, 947.e1–947.e9
- Auditory perception
 - characteristics, of voice of nondysphonic school children from 8:0 to 10:0 years old, 946.e1–946.e5
 - evaluation, and vocal acoustic during fluctuations in estradiol levels during menstrual cycle, 536–544
 - training, with natural voice anchors on vocal quality evaluation, effect of, 220–225
- Autoimmune diseases, and microphonotrauma, bamboo nodes as sign of, 357–362
- AVQI. See Acoustic Voice Quality Index
- Behavior, of children with vocal fold nodules, a case-control study, 584.e1–584.e4
- Behavioral tendency, in children with vocal nodules, 945.e1–945.e18
- Biopsy, cost analysis of channeled, distal chip laryngoscope for in-office laryngopharyngeal, 575–579
- BMI. See Body mass index
- Body mass index (BMI), Tanner stage and singing activity in children and adolescents, normative data and associations with, 580.e21–580.e30
- Body position, and emission number, in results of maximum phonation times of adults without vocal complaints, influence of, 831–837
- Botulinum toxins
 - single injection, in patients with ADSD over time, quality of life and voice changes after, 721–727
 - type A, for laryngeal dysfunction in Alexander disease, 728–729
 - type A, multidose, for laryngeal dystonia compared to single-use, cost savings of, 159–161
 - type A, spasmodic dysphonia in multiple sclerosis treatment with, 550–553
- Burnout, psychological, and voice disorder in teachers, 581.e7–581.e16
- CAPE-V. See Consensus Auditory-Perceptual Evaluation of Voice
- Child
 - from 8:0 to 10:0 years old, auditory-perceptual characteristics of voice of nondysphonic school, 946.e1–946.e5
 - 4-year-old, in STEPS study, respiratory tract infections and voice quality in, 801.e21–801.e25
 - and adolescents, normative data and associations with BMI, Tanner stage and singing activity in, 580.e21–580.e30
 - with allergic rhinitis, evaluation of pVHI in, 801.e17–801.e20
 - behavioral with vocal fold nodules, a case-control study, 584.e1–584.e4
 - comprehension of spoken language, impact of dysphonic voices on, 801.e7–801.e16
 - with history of laryngotracheal reconstruction surgery and relationship between clinician perceptual rating of voice quality and parent proxy/child self-report of VRQOL, perceptual evaluation of voice disorder in, 945.e27–945.e35
 - school-aged, with and without attention deficit hyperactivity disorder, vocal characteristics of, 945.e37–945.e45
 - with vocal fold nodules, personality traits in, 801.e1–801.e6
 - with vocal nodules, personality, psychological factors, and behavioral tendencies in, 945.e1–945.e18
- Choir training, and semi-occluded vocal tract on voice with congenital, isolated untreated growth hormone deficiency, effects of therapy with, 808.e1–808.e5
- Choral sound, and solo, effects of musician earplugs on acoustic and perceptual measures of, 87–95
- Cigarette smoking, exposure to pig larynx in an inhalation chamber, 846–850
- Cleft palate, with and without glottal stop in Mandarin, comparative study of vocal attack time in patients with, 803.e15–803.e21
- Clergy, prevalence of and risk factors for self-report voice problems among Hindu temple, 805.e1–805.e12
- Clinicians, and singer perspective on vitamin B12 treatment for vocal benefits, 352–356
- Cochlear implants
 - influence of auditory feedback and vocal rehabilitation on prelingual hearing-impaired individuals post, 947.e1–947.e9
 - postlingual, long-term average speech spectra with, 255.e19–255.e25
- Communication disorders, speech, female, graduate students, perceived desirability of vocal fry among, 805.e21–805.e35
- Computerized tomography
 - morphometric analysis of professional opera singers vocal folds, 583.e1–583.e8
 - study of vocal tract setting in hyperfunctional dysphonia and in belting, 412–419
- Consensus Auditory-Perceptual Evaluation of Voice (CAPE-V)
 - psychometric characteristics of II European Portuguese version of, 582.e5–582.e13
 - reliability and validity of Turkish version of, 382.e1–382.e10
- Contemporary commercial music, and classical styles, emotion recognition from singing voices using, 501–509
- Cost savings, of multidose botulinum toxin A for laryngeal dystonia compared to single-use, 159–161
- Costs, and cost analysis, of channeled, distal chip laryngoscope for in-office laryngopharyngeal biopsies, 575–579
- Cricoarytenoid joint ankylosis, classification and transoral laser microsurgical treatment for, 375–380
- Cricoarytenoid joint dislocation, type, three-dimensional structure of vocal folds in identifying, 611–619
- Cricothyroid approximation, and glottoplasty, long term effects and comparison outcome between, 781–791
- Crying signal, in infants with disabling hearing impairment, acoustic analysis of, 946.e7–946.e13
- Cysts, acoustic characteristics in epiglottic, 497–500
- Deep brain stimulation, subthalamic nucleus and zona incerta, in patients with Parkinson disease, voice tremor outcomes of, 545–549
- Dexamethasone, intracordal and estradiol injection on wound healing on vocal fold injuries, effects of, 759–766

- Diagnosis
 comparing rigid telescopic stroboscopy with direct microlaryngoscopy in epithelial and lamina propria glottic lesions, 143–149
 laryngopharyngeal reflux, dual pH probes without proximal esophageal and pharyngeal impedance in, 697–703
- Digital kymographic system, clinical practicability of, 346–351
- Digital kymography
 scanning, two-dimensional, in evaluation of atrophic vocal folds, efficacy of, 554–560
 two-dimensional, in patients with vocal fold scarring, usefulness of, 906–914
- Diphonation, in audio recordings of German standard text readings, detection of, 949.e1–949.e10
- Diphonia, studied by high speed video and vibrogram analysis, vibratory characteristics of, 7–15
- DoctorVox voice therapy, technique for mutational falsetto, efficacy of, 950.e1–950.e8
- Dr. Speech device, validation of Dysphonia Severity Index in, 948.e23–948.e29
- DSI. See Dysphonia Severity Index
- Dysphagia, in patients with non-neoplastic vocal fold pathology, prevalence of, 708–711
- Dysphonia
 across vowel categories, perceptual and quantitative assessment with, 473–481
 associations between posture, voice and, 124.e1–124.e12
 behavioral, effect of voice therapy program based on taxonomy of vocal therapy in women with, 256.e1–256.e16
 behavioral, effectiveness of voice therapy associated with electromyographic biofeedback in women with, 381.e11–381.e22
 effects of amplification on vocal dose in teachers with, 73–79
 functional, standard instrumental acoustic, perceptual and subjective voice outcomes in therapy success in patients with, 317–324
 hyperfunctional, and in belting, computerized tomography study of vocal tract setting in, 412–419
 muscle tension, aerodynamic profile of, 812.e15–812.e18
 muscle tension, positive effects of manual circumlaryngeal therapy in treatment of, 866–871
 perceived control and psychological distress, relationship between, 682–690
 perceptual method to rate voice, 453–464
 during phonation and speech, control of fundamental frequency in patients, 851–859
 primary muscle tension, using functional magnetic resonance imaging, exploring the neural bases of, 183–194
 spasmodic, in multiple sclerosis treatment with Botulin toxin A, 550–553
 spasmodic, outcome measurement in treatment of, systematic review, 810.e13–810.e39
 utility of pulmonary function testing in patients presenting with, 567–574
 voices and normal voices, comparison of post-therapy, 581.e1–581.e6
 voices on children comprehension of spoken language, impact of, 801.e7–801.e16
- Dysphonia Severity Index (DSI)
 and AVQI, influence of gender and age on, 340–345
 in Dr. Speech device, validation of, 948.e23–948.e29
 for Persian-speaking populations, design and assessment of multiparametric model for, 226–231
- EASE. See Evaluation of the Ability to Sing Easily
- Editorials, 127–128
- Electroglottography
 aerodynamics and acoustic parameters, lingering effects of straw phonation exercises on, 810.e5–810.e11
 estimation of source-filter interaction regions based on, 269–276
 fundamental frequency estimation of, 401–411
- Electromyography
 biofeedback in women with behavioral dysphonia, effectiveness of voice therapy associated with, 381.e11–381.e22
 laryngeal, and VHI findings, vocal fold paresis incidence, and relationship between, 940–944
- Electronic nicotine delivery systems, on voice quality, effect of, 811.e13–811.e17
- EMG. See Electromyography
- Emotion
 perception, using Persian Affective Voice test, effect of Parkinson disease on, 580.e1–580.e9
 from singing voices using contemporary commercial music and classical styles, 501–509
 transient states, modulation of relative fundamental frequency during, 894–899
- Endoscopy
 office-based procedures of upper aerodigestive tract, topical anesthesia for, 732–746
 and phoniatric evaluation in singing students, 135–142
- Epiglottis
 angle on aerodynamic and acoustic parameters in canine larynges, effects of, 627–633
 cyst, acoustic characteristics in, 497–500
- Esophagus, and pharyngeal impedance, in detecting and diagnosing laryngopharyngeal reflux, dual pH probes without, 697–703
- Estradiol
 intracardial, and dexamethasone injection on wound healing on vocal fold injuries, effects of, 759–766
 levels during menstrual cycle, vocal acoustic and auditory-perceptual characteristics during fluctuations in, 536–544
- Evaluation of the Ability to Sing Easily (EASE), scale in development of vocal warm-up program for amateur choir, effectiveness of, 310–316
- Exercises
 induced laryngeal obstruction, perceptual clinical features in, 880–893
 effects of vocal function, 124.e13–124.e34
 semi-occluded vocal tract, analytical model for flow resistance and geometric properties of tubes used in, 585–590
 straw phonation on aerodynamics, electroglottographic and acoustic parameters, lingering effects of, 810.e5–810.e11
- Facial expression, special, classical singers use, 510–515
- Fake Phone Call app, in public practice of voice technique for generalization training, feasibility of, 659–668
- Fundamental frequency
 in dysphonic patients during phonation and speech, control of, 851–859
 relative, during transient emotional states, modulation of, 894–899
 relative, loudness, stress type, vowel identity, and baseline f₀ with healthy voices, effects of, 603–610
 of speaking voice, effect of smoking on, 802.e11–802.e16

- Gender, and age, influence on AVQI and DSI, 340–345
- Glass tubes, in singers, effects of phonation in, 381.e1–381.e9
- Glottal gap, effects of voice therapy using lip trill technique in patients with, 949.e11–949.e19
- Glottal insufficiency, compensation strategies in voice production with, 96–102
- Glottal stop, in Mandarin, comparative study of vocal attack time in patients with cleft palate with and without, 803.e15–803.e21
- Glottic cancer, early, voice outcome following radiation versus laser microsurgery, 176–182
- Glottic insufficiency
under 50 years old, office-based autologous fat injection laryngoplasty for, 747–750
injectable silk protein microparticle-based fillers for, 773–780
- Glottic lesions
epithelial and lamina propria, comparing rigid telescopic stroboscoped laryngoscopy with direct microlaryngoscopy in diagnosing, 143–149
voice profile recovery and quality of life changes after microdirect laryngoscopy in, 382.e11–382.e20
- Glottis, changes and distinct acoustic features define two modes of singing in Peking opera, 583.e9–583.e14
- Glottoplasty, long term effects and comparison outcome between cricothyroid approximation and, 781–791
- Growth hormone deficiency, congenital, isolated untreated, effects of therapy with semi-occluded vocal tract and choir training on voice with, 808.e1–808.e5
- Hearing impairment. See Hearing loss
- Hearing loss
acoustic analysis of crying signal in infants with disabling hearing impairment, 946.e7–946.e13
prelingual, individuals post cochlear implant, influence of auditory feedback and vocal rehabilitation on, 947.e1–947.e9
- High speed video, and vibrogram analysis, vibratory characteristics of diplophonia studied by, 7–15
- High-speed videoendoscopy
and acoustic signal analysis, dependencies and ill-designed parameters within, 811.e1–811.e12
of essential voice tremor and adductor spasmodic dysphonia, reliability of, 16–26
- Hoarseness, lipoid proteinosis a rare cause of, 155–158
- Human papillomavirus type 6 virus-like particle vaccine, nonadjuvant, in recurrent respiratory papillomatosis, safety and immunogenicity of, 363–369
- Hydration, on voice quality in adults, effect of, 125.e13–125.e28
- Immunogenicity, and safety of nonadjuvant human papillomavirus type 6 virus-like particle vaccine in recurrent respiratory papillomatosis, 363–369
- Infants
with disabling hearing impairment, acoustic analysis of crying signal in, 946.e7–946.e13
phonation, registers in, 382.e21–382.e32
- Inflammatory bowel disease, laryngeal manifestations of, 1–6
- Inhalation chamber, cigarette smoke exposure to pig larynx in, 846–850
- Injections
Botox, for laryngeal dysfunction in Alexander disease, 728–729
Botox, single, in patients with ADSD over time, quality of life and voice changes after, 721–727
intra-lesional steroid, in small benign vocal fold lesions, efficacy of, 767–772
intracordal estradiol and dexamethasone, on wound healing on vocal fold injuries, effects of, 759–766
- Knowledge, and perceptions, comparison of laboratory and virtual laryngeal dissection experiences on preservice music educators, 872–879
- Kunqu Opera, voice characteristics of young girl role in, 945.e19–945.e25
- Language
spoken, impact of dysphonic voices on children comprehension of, 801.e7–801.e16
Tamil, newsreaders on varied emotional texts in, 255.e9–255.e17
- Laryngeal dissection, laboratory and virtual, experiences on preservice music educators knowledge and perceptions, comparison of, 872–879
- Laryngeal dysfunction, in Alexander disease, Botox injection for, 728–729
- Laryngeal dystonia, cost savings of botulinum toxin A for, 159–161
- Laryngeal nerve avulsion, unilateral recurrent, polysialylated neural cell adhesion molecule supports regeneration of neurons in nucleus ambiguus after, 52–57
- Laryngeal nodules, role of pepsin in etiology of, 704–707
- Laryngeal obstruction, exercise-induced, perceptual clinical features in, 880–893
- Laryngeal pathology, and vocal, among professional singers, the impact of, 58–65
- Laryngopharyngeal reflux
association between pepsin in saliva and subjective symptoms in patients with, 150–154
dual pH probes without proximal esophageal and pharyngeal impedance in detecting and diagnosing, 697–703
on subjective, aerodynamic, and acoustic voice assessments of responder and nonresponder patients, impact of, 929–939
symptomatic, diagnostic usefulness of salivary pepsin test in, 923–928
- Laryngoplasty
bilateral medialization, for prebylaryngis, treatment outcomes of, 40–44
injection, in patients with puberphonia, 564–566
medialization, comparison of pitch strength with perceptual and other acoustic metric outcome measures following, 795–800
office-based autologous fat injection, for glottic insufficiency under 50 years old, 747–750
- Laryngoscope, channeled, distal chip, for in-office laryngopharyngeal biopsies, cost analysis of, 575–579
- Laryngoscopy
microdirect, in glottis lesions, voice profile recovery and quality of life changes after, 382.e11–382.e20
transnasal flexible, grading system for, 712–715
- Laryngotracheal reconstruction surgery, and relationship between clinician perceptual rating of voice quality and parent proxy/child self-report of VRQOL, perceptual evaluation of voice disorder in children with history of, 945.e27–945.e35
- Larynx
aerodynamics, computational models of, 385–400
canine, effects of angle of epiglottis on aerodynamic and acoustic parameters in, 627–633

- excised canine, evaluation of surgical strategies for bilateral vocal cord paralysis using, 33–39
- manifestations of inflammatory bowel disease, 1–6
- pig, in inhalation chamber, cigarette smoke exposure to, 846–850
- and respiratory function in teachers, 302–309
- web as possible cause for nonabsorption of vocal nodules in boys after puberty, 561–563
- Larynx tilt, on same-pitch phonation of vocal mode metal and density parameters, 806.e9–806.e21
- Laser microsurgery
- versus radiation in early glottic cancer, voice outcome following, 176–182
 - transoral, and classification for cricoarytenoid joint ankylosis, 375–380
- LaxVox exercises, and voiced high-frequency oscillation, in vocally healthy subjects, immediate effects of, 808.e7–808.e14
- LaxVox tubes, in singers, effects of phonation in, 381.e1–381.e9
- Levodopa, effectiveness in patients with early idiopathic Parkinson disease, voice quality and orofacial strength as outcome of, 716–720
- Lip trill technique, in patients with glottal gap, effects of voice therapy using, 949.e11–949.e19
- Lipoid proteinosis of Urban and Wiethe, a rare cause of hoarseness, 155–158
- Long-term average speech spectra, of postlingual cochlear implant users, 255.e19–255.e25
- Loudness, and pitch, airflow vibrato dependence on, 815–830
- Machine learning, approaches for automatic detection of voice disorders, survey on, 947.e11–947.e33
- Magnetic resonance imaging, exploring the neural bases of primary muscle tension dysphonia using, 183–194
- Manual circumlaryngeal therapy, in treatment of muscle tension dysphonia, positive effects of, 866–871
- Menstrual cycle, vocal acoustic and auditory-perceptual characteristics during fluctuations in estradiol levels during, 536–544
- Microlaryngoscopy, direct, in diagnosing epithelial and lamina propria glottic lesions, comparing rigid telescopic stroboscoped laryngoscopy with, 143–149
- Microphonotrauma, bamboo nodes as sign of autoimmune disease and, 357–362
- Mokken scaling, restructuring Vocal Fatigue Index using, 110–114
- Mucosal wave, improved, in unilateral autologous temporal fascia graft in sulcus vocalis type 2 and vocal scars, 915–922
- Multiple sclerosis, treatment with Botulin toxin A, spasmodic dysphonia in, 550–553
- Muscle tonus
- dysphonia, aerodynamic profile of, 812.e15–812.e18
 - dysphonia, positive effects of manual circumlaryngeal therapy in treatment of, 866–871
 - dysphonia, primary, using functional magnetic resonance imaging, exploring the neural bases of, 183–194
- Musculoskeletal pain, and voice self-assessment, in tele-operators, relation between, 948.e11–948.e21
- Music educator, preservice, knowledge and perceptions, comparison of laboratory and virtual laryngeal dissection experiences on, 872–879
- Musical theater, singers in Brazil, immediate impact of vocal demand on, 804.e13–804.e22
- Musician, earplugs on acoustic and perceptual measures of choral and solo sound, effects of, 87–95
- Mutational falsetto, efficacy of DoctorVox voice therapy technique for, 950.e1–950.e8
- Neurons, adhesion molecule supports regeneration of, in nucleus ambiguus after unilateral recurrent laryngeal nerve avulsion, 52–57
- Noise
- resulting from location and conditions of classrooms and schools in upper Egypt on teachers voices, influence of, 802.e1–802.e9
 - on vocal dose in women, influence of, 214–219
- Nucleus ambiguus, after unilateral recurrent laryngeal nerve avulsion, polysialylated neural cell adhesion molecule supports regeneration of neurons in, 52–57
- Outcomes
- between cricothyroid approximation and glottoplasty, long term effects and comparison, 781–791
 - of levodopa effectiveness in patients with early idiopathic Parkinson disease, voice quality and orofacial strength as, 716–720
 - measurement in treatment of spasmodic dysphonia, systematic review, 810.e13–810.e39
 - with perceptual and other acoustic metric measures, following medialization laryngoplasty, comparison of pitch strength with, 795–800
 - standard instrumental acoustic, perceptual and subjective voice, in therapy success in patients with functional dysphonia, 317–324
 - treatment, of bilateral medialization thyroplasty for prebylaryngis, 40–44
 - voice, following radiation versus laser microsurgery in early glottic cancer, 176–182
 - voice tremor, of subthalamic nucleus and zona incerta deep brain stimulation in patients with Parkinson disease, 545–549
- Parkinson disease
- early idiopathic, voice quality and orofacial strength as outcome of levodopa effectiveness in patients with, 716–720
 - on emotion perception using Persian Affective Voice test, effect of, 580.e1–580.e9
 - tremor phenotype on cepstral peak prominence and transglottal airflow in vowels and speech, effect of, 580.e11–580.e19
 - voice tremor in, 526–535
 - voice tremor in (letter), 813.e7
 - voice tremor outcomes of subthalamic nucleus and zona incerta deep brain stimulation in patients with, 545–549
- Pediatric Voice Handicap (pVHI), evaluation in children with allergic rhinitis, 801.e17–801.e20
- Pepsin
- in etiology of laryngeal nodules, role of, 704–707
 - in saliva and subjective symptoms in patients with laryngopharyngeal reflux, association between, 150–154
- Perception
- and acoustic measures of an SATB chorus, effects of straw phonation protocol on, 80–86
 - and acoustic measures of choral and solo sound, effects of musician earplugs on, 87–95
 - of breathiness in voices of pediatric speakers, 204–213
 - and knowledge, comparison of laboratory and virtual laryngeal dissection experiences on preservice music educators, 872–879
 - and other acoustic metric outcome measures following medialization laryngoplasty, comparison of pitch strength with, 795–800

- of patients, and attitudes toward voice therapy, 860–865
 voice quality, effect of telephone transmission on, 649–658
- Perceptual analysis
 and acoustic, of adductor spasmodic dysphonia in
 Mandarin-speaking Chinese, 333–339
 acoustic correlates of voice in Purohits, 804.e1–804.e4
 standard instrumental acoustic, and subjective voice outcomes in
 therapy success in patients with functional dysphonia, 317–324
- Personality traits
 in children with vocal fold nodules, 801.e1–801.e6
 in children with vocal nodules, 945.e1–945.e18
- Pharynx, and esophageal impedance, in detecting and diagnosing
 laryngopharyngeal reflux, dual pH probes without, 697–703
- Phonation
 into glass, plastic and LaxVox tubes in singers, effects of,
 381.e1–381.e9
 maximum times, of adults without vocal complaints, influence of
 body position and emission number in results of, 831–837
 registers in infant, 382.e21–382.e32
 role of allergy in, 811.e19–811.e27
 same-pitch, of vocal mode metal and density parameters. laryngeal
 tilt on, 806.e9–806.e21
 and speech, control of fundamental frequency in dysphonic
 patients during, 851–859
 straw, exercises on aerodynamics, electroglottographic and
 acoustic parameters, lingering effects of, 810.e5–810.e11
 straw, protocol on acoustic and perceptual measures of an
 SATB chorus, effects of, 80–86
- Phoniatric evaluation, and endoscopic, in singing students, 135–142
- Phonotrauma, acute, after the match, acoustic analysis of soccer fans
 in, 129–134
- Pitch, and loudness, airflow vibrato dependence on, 815–830
- Plastic tubes, in singers, effects of phonation in, 381.e1–381.e9
- Polysialic acid, neural cell adhesion molecule supports regeneration of
 neurons in nucleus ambiguus after unilateral recurrent laryngeal
 nerve avulsion, 52–57
- Posture, voice and dysphonia, associations between,
 124.e1–124.e12
- Prebylaryngis, treatment outcomes of bilateral medialization
 thyroplasty for, 40–44
- Psychogenic voice disorder, and efficiency of voice therapy and
 psychological evaluation, clinical features of, 250–254
- Psychological distress, relationship between dysphonia, perceived
 control and, 682–690
- Psychological evaluation, clinical features of psychogenic voice
 disorder and efficiency of voice therapy and, 250–254
- Psychological factors, in children with vocal nodules,
 945.e1–945.e18
- Psychometrics
 of II European Portuguese version of CAPE-V, 582.e5–582.e13
 of Spanish version of Voice Activity and Participation Profile,
 582.e15–582.e22
- Puberphonia, injection laryngoplasty in patients with, 564–566
- Puberty, laryngeal web as possible cause for nonabsorption of vocal
 nodules in boys after, 561–563
- Pulmonary function tests, in patients presenting with dysphonia, utility
 of, 567–574
- PVHI. See Pediatric Voice Handicap Index
- Quality of life
 changes and voice profile recovery after microdirect laryngoscopy
 in glottis lesions, 382.e11–382.e20
- and voice changes, after single injection in patients with ADSD
 over time, 721–727
- Radiation, versus laser microsurgery in early glottic cancer, voice
 outcome following, 176–182
- Reflux Symptom Index, Spanish, translation and validation of,
 807.e1–807.e5
- Regeneration, of neurons in nucleus ambiguus after unilateral recurrent
 laryngeal nerve avulsion, polysialylated neural cell adhesion
 molecule supports, 52–57
- Reproducibility of results
 of Aging Voice Index for aging with voice disorders,
 807.e7–807.e12
 of Bahasa Malaysia version of Voice Handicap Index-10,
 581.e17–581.23
 of Mandarin Chinese version of Voice Activity and Participation
 Profile, 103–109
 of speech evaluation in adductor spasmodic dysphonia (letter),
 49–51
 of Turkish version of CAPE-V, 382.e1–382.e10
- Respiratory papillomatosis, recurrent, safety and immunogenicity of
 nonadjuvant human papillomavirus type 6 virus-like particle
 vaccine in, 363–369
- Respiratory physiological phenomena, and laryngeal function in
 teachers, 302–309
- Respiratory tract infections, and voice quality in 4-year-old children in
 STEPS study, 801.e21–801.e25
- Rhinitis, allergic, evaluation of pVHI in children with,
 801.e17–801.e20
- Rigid telescopic stroboscopy, with direct
 microlaryngoscopy in diagnosing epithelial and lamina propria
 glottic lesions, comparing, 143–149
- Risk factors
 and prevalence of self-reported voice problems among Yakshagana
 artists, 124.e35–124.e47
 for self-report voice problems among Hindu temple priests,
 prevalence of and, 805.e1–805.e12
- Salivary pepsin test, in symptomatic laryngopharyngeal reflux,
 diagnostic usefulness of, 923–928
- Semi-occluded vocal tract
 and choir training on voice with congenital, isolated untreated
 growth hormone deficiency, effects of therapy with,
 808.e1–808.e5
 exercises, analytical model for flow resistance and geometric
 properties of tubes used in, 585–590
- Silk protein microparticle-based fillers, for glottic insufficiency,
 773–780
- Singing
 age and frequency impact the thyroid cartilage in, 284–289
 BMI, and Tanner stage in children and adolescents, normative data
 and associations with, 580.e21–580.e30
 choral, acoustics and aerodynamics in, 803.e1–803.e5
 classical, use special facial expression, 510–515
 classical and popular female, sentimental melody, acoustic
 characteristics of, 255.e27–255.e37
 comparing the exposure-response relationships of physiological
 and traditional vocal warm-ups on aerodynamic and acoustic
 parameters in, 420–428
 distinct acoustic features and glottal changes define two modes of
 opera, 583.e9–583.e14

- effects of phonation into glass, plastic and LaxVox tubes in, 381.e1–381.e9
- female, voice in choral, development, 516–525
- the impact of vocal and laryngeal pathologies among professional, 58–65
- metal, aerodynamic characteristics of growl voice and reinforced falsetto in, 803.e7–803.e13
- musical theater, immediate impact of vocal demand in, 804.e13–804.e22
- opera-chant, evaluation of supraglottic and subglottic activities including acoustic assessment of, 255.e1–255.e7
- professional, unusual etiology of vocal tremor in, 730–731
- professional opera vocal folds, CT-based morphometric analysis of, 583.e1–583.e8
- students, endoscopic and phoniatric evaluation in, 135–142
- vitamin B12 treatment for vocal benefits in, 352–356
- voice, trace and research patterns in, 263–268
- voices using contemporary commercial music and classical styles, emotion recognition from, 501–509
- Singing Voice Handicap Index-10 (SVHI-10), Kannada, adaptation and validation of, 582.e1–582.e4
- Smoking, on fundamental frequency of speaking voice, effect of, 802.e11–802.e16
- Solo sound, and choral, effects of musician earplugs on acoustic and perceptual measures of, 87–95
- Source-filter interaction, regions based on electroglottography, estimation of, 269–276
- Speakers
 - normal, aerodynamic characteristics of syllable and sentence productions in, 297–301
 - pediatric, perception of breathiness in voices of, 204–213
- Speech
 - connected, in patients with and without voice disorders, effects of variably occluded face mask on aerodynamic and acoustic characteristics of, 809.e1–809.10
 - evaluation in adductor spasmodic dysphonia, reliability and validity of (letter), 49–51
 - and phonation, control of fundamental frequency in dysphonic patients during, 851–859
 - running, in patients with voice disorders, airflow patterns of, 277–283
 - and vowels, effect of Parkinson disease tremor phenotype on cepstral peak prominence and transglottal airflow in, 580.e11–580.e19
- Steroid, intra-lesional injection in small benign vocal fold lesions, efficacy of, 767–772
- SToPS. See Sunderland Tracheoesophageal Perceptual Scale
- Stress
 - impact, in water resistance voice therapy, 490–496
 - type, vowel identity, baseline f0 and loudness on relative fundamental frequency with healthy voices, effects of, 603–610
- Students
 - endoscopic and phoniatric evaluation in singing, 135–142
 - perceived desirability of vocal fry among female speech communication disorders graduate, 805.e21–805.e35
- Subthalamic nucleus, and zona incerta deep brain stimulation in patients with Parkinson disease, voice tremor outcomes of, 545–549
- Sulcus vocalis, type 2, and vocal scars, improved mucosal wave in unilateral autologous temporal fascia graft in, 915–922
- Sunderland Tracheoesophageal Perceptual Scale (SToPS), perceptual assessment of tracheoesophageal voice quality with, 465–472
- Tanner stage, BMI and singing activity in children and adolescents, normative data and associations with, 580.e21–580.e30
- Teachers
 - with dysphonia, effects of amplification on vocal dose in, 73–79
 - respiratory and laryngeal function in, 302–309
 - student, voice disorders and impact of voice handicap in, 445–452
 - vocal health practices among school, 812.e1–812.e7
 - voice disorder and burnout syndrome in, 581.e7–581.e16
 - voices, influence of noise resulting from location and conditions of classrooms and schools in upper Egypt on, 802.e1–802.e9
- Tele-operators, relation between musculoskeletal pain and voice self-assessment in, 948.e11–948.e21
- Temporal fascia graft, unilateral autologous, in sulcus vocalis type 2 and vocal scars, improved mucosal wave in, 915–922
- Thyrohyoid ligament, lateral, morphology and acoustic assessment of the voice, effect of chant training on, 802.e17–802.23
- Thyroid cartilage, of professional singers, age and frequency impact the, 284–289
- Thyroid surgery, vocal changes following, 27–32
- Thyroplasty. See Laryngoplasty
- Transsexual Voice Questionnaire, in French, translation, adaptation and preliminary validation of, 807.e13–807.e24
- Tremor, voice, in Parkinson disease (letter), 813.e7
- Trial vocal fold injection, augmentation, evaluation of shorter follow-up time to capture benefit of, 169–175
- Tuberculosis, laryngeal, primary, at tertiary care teaching hospital in Eastern India, 812.e9–812.e14
- Upper aerodigestive tract, topical anesthesia for endoscopic office-based procedures of, 732–746
- Validity and reliability. See Reproducibility of results
- VAPP. See Voice Activity and Participation Profile
- Variably occluded face mask, on aerodynamic and acoustic characteristics of connected speech in patients with and without voice disorders, effects of, 809.e1–809.10
- VHI. See Voice Handicap Index
- Vibrogram, vibratory characteristics of diplophonia studied by high speed video and, 7–15
- Vitamin B12, for vocal benefits, singers and clinician perspectives, 352–356
- Vocal attack time
 - extended analysis of, 258–262
 - in patients with cleft palate with and without glottal stop in Mandarin, comparative study of, 803.e15–803.e21
- Vocal complaints, influence of body position and emission number in results of maximum phonation times of adults without, 831–837
- Vocal cord injuries, effects of intracordal estradiol and dexamethasone injection on wound healing on, 759–766
- Vocal cord lesions, small benign, efficacy of intra-lesional steroid injection in, 767–772
- Vocal cord nodules, personality traits in children with, 801.e1–801.e6
- Vocal cord paralysis
 - bilateral, endoscopic partial arytenoidectomy for, 751–758
 - bilateral, using excised canine larynges, evaluation of surgical strategies for, 33–39
- Vocal cords
 - atrophic, efficacy of two-dimensional scanning digital kymography in evaluation of, 554–560

- CT-based morphometric analysis of professional opera singers, 583.e1–583.e8
- in identifying cricoarytenoid joint dislocation type, three-dimensional structure of, 611–619
- immobile, feasibility study of new approach in medialization and lateralization of, 162–168
- pathology, non-neoplastic, dysphagia in patients with, 708–711
- Vocal demand, on musical theater singers in Brazil, immediate impact of, 804.e13–804.e22
- Vocal dose, in women, influence of noise on, 214–219
- Vocal Fatigue Index
- into Persian, cross-cultural adaptation and validation of, 947.e35–947.e41
- using Mokken scaling, restructuring, 110–114
- Vocal fold lateralization, and medialization, of immobile vocal folds, feasibility study of new approach in, 162–168
- Vocal fold medialization
- bilateral, for abductor spasmodic dysphonia, 45–48
- and lateralization, of immobile vocal folds, feasibility study of new approach in, 162–168
- Vocal fold nodules
- assessment of resonance voice therapy in treatment of, 810.e1–810.e4
- behavioral profile of children with, a case-control study, 584.e1–584.e4
- Vocal fold paralysis, unilateral, imaging modalities in etiological evaluation of, 813.e1–813.e5
- Vocal fold paresis, incidence, and relationship between VHI and laryngeal EMG findings, 940–944
- Vocal fold scarring
- and sulcus vocalis type 2, improved mucosal wave in unilateral autologous temporal fascia graft in, 915–922
- usefulness of two-dimensional digital kymography in patients with, 906–914
- Vocal fry, among female speech communication disorders graduate students, perceived desirability of, 805.e21–805.e35
- Vocal function, exercises, effects of, 124.e13–124.e34
- Vocal health, practices among school teachers, 812.e1–812.e7
- Vocal loading, and vocal rest using aerodynamic, acoustic and self-rating measurements, effects of, 809.e11–809.e18
- Vocal nodules
- in boys after puberty, laryngeal web as possible cause for nonabsorption of, 561–563
- pediatric, using ratios calculated with biomedical image segmentation, voice acoustic analysis of, 195–203
- personality, psychological factors, and behavioral tendencies in children with, 945.e1–945.e18
- Vocal pathology, and laryngeal, among professional singers, the impact of, 58–65
- Vocal rehabilitation, and auditory feedback, influence on prelingual hearing-impaired individuals post cochlear implant, 947.e1–947.e9
- Vocal rest, and vocal loading using aerodynamic, acoustic and self-rating measurements, effects of, 809.e11–809.e18
- Vocal score profile, in Verdi's characteristics, 805.e13–805.e20
- Vocal strain, objective indices of perceived, 838–845
- Vocal tract
- in hyperfunctional dysphonia and in belting, computerized tomography study of, 412–419
- shape via biofeedback, adjustment of the, 482–489
- Vocal Tract Discomfort Scale (VTDS)
- cross-cultural adaptation and validation of Italian version of, 115–123
- and VoiSS in evaluation of patients with voice disorders, 381.e23–381.e32
- Vocal warm-up
- physiological and traditional, on aerodynamic and acoustic parameters in untrained singers, comparing the exposure-response relationships of, 420–428
- program for amateur choir, effectiveness of EASE scale in development of, 310–316
- Voice
- acoustic analysis of pediatric vocal nodule patients using ratios calculated with biomedical image segmentation, 195–203
- acoustic and auditory-perceptual characteristics during fluctuations in estradiol levels during menstrual cycle, 536–544
- benefits, singers and clinicians perspective on vitamin B12 treatment for, 352–356
- changes, and quality of life, after single injection in patients with ADSD over time, 721–727
- characteristics of school-aged children with and without attention deficit hyperactivity disorder, 945.e37–945.e45
- characteristics of young girl role in Kunqu Opera, 945.e19–945.e25
- complex, task without external feedback, video modeling and instruction in acquisition and retention of, 239–249
- with congenital, isolated untreated growth hormone deficiency, effects of therapy with semi-occluded vocal tract and choir training on, 808.e1–808.e5
- dysphonic, on children comprehension of spoken language, impact of, 801.e7–801.e16
- effect of chant training on lateral thyrohyoid ligament morphology and acoustic assessment of, 802.e17–802.e23
- essential, tremor and adductor spasmodic dysphonia, reliability of high-speed videoendoscopic ratings of, 16–26
- growl, reinforced falsetto in metal singing, aerodynamic characteristics of, 803.e7–803.e13
- healthy, effects of stress type, vowel identity, baseline f0 and loudness on relative fundamental frequency with, 603–610
- natural, anchors on vocal quality evaluation, effect of auditory perceptual training with, 220–225
- of nondysphonic school children from 8:0 to 10:0 years old, auditory-perceptual characteristics of, 946.e1–946.e5
- normal, patterns in Pakistani adults, acoustic analysis of, 124.e49–124.e58
- ornamentation in Byzantine chant, 256.e17–256.e34
- outcome following radiation versus laser microsurgery in early glottic cancer, 176–182
- of pediatric speakers, perception of breathiness in, 204–213
- perceptual method to rate dysphonic, 453–464
- post-therapy dysphonic, and normal, comparison of, 581.e1–581.e6
- posture and dysphonia, associations between, 124.e1–124.e12
- production with glottal insufficiency, compensation strategies in, 96–102
- of professional users with no vocal cord lesions similar to non-professional users, 66–72
- profile recovery and quality of life changes after microdirect laryngoscopy in glottis lesions, 382.e11–382.e20
- in Purohits, acoustic-perceptual correlates of, 804.e1–804.e4
- signals based on variation in energy distribution, an objective parameter to classify, 591–602
- singing, female, in choral singing development, 516–525

- singing, using contemporary commercial music and classical styles, emotion recognition from, 501–509
- speaking, effect of smoking on fundamental frequency of, 802.e11–802.e16
- subjective, standard instrumental acoustic and perceptual outcomes in therapy success in patients with functional dysphonia, 317–324
- teachers, influence of noise resulting from location and conditions of classrooms and schools in upper Egypt on, 802.e1–802.e9
- technique for generalization training, feasibility of Fake Phone Call app in public practice of, 659–668
- trace and research patterns in singing, 263–268
- tremor, in Parkinson disease, 526–535
- tremor, outcomes of subthalamic nucleus and zona incerta deep brain stimulation in patients with Parkinson disease, 545–549
- tremor in Parkinson disease (letter), 813.e7
- Voice Activity and Participation Profile (VAPP)
among etiological groups, comparison of, 804.e5–804.e12
psychometric properties of Spanish version of, 582.e15–582.e22
reliability and validity of Mandarin Chinese version, 103–109
- Voice amplification, on vocal dose in teachers with dysphonia, effects of, 73–79
- Voice analyzer, programs for voice quality analysis, improve version of, 642–648
- Voice assessments, of responder and nonresponder patients, impact of laryngopharyngeal reflux on subjective, aerodynamic, and acoustic, 929–939
- Voice Catastrophization Index, development and validation of, 232–238
- Voice Disability Coping Questionnaire (VDCQ), Persian version, cross-cultural adaptation and validation of, 435–440
- Voice disorders
airflow patterns of running speech in patients with, 277–283
in Brazilian community-dwelling older adults, prevalence and associated factors with, 806.e1–806.e7
and burnout syndrome in teachers, 581.e7–581.e16
in children with history of laryngotracheal reconstruction surgery and relationship between clinician perceptual rating of voice quality and parent proxy/child self-report of VRQOL, perceptual evaluation of, 945.e27–945.e35
effects of variably occluded face mask on aerodynamic and acoustic characteristics of connected speech in patients with and without, 809.e1–809.e10
in general population, prevalence of, 900–905
and impact of voice handicap in Norwegian student teachers, 445–452
quality and readability of internet information for, 290–296
reliability and validity of Aging Voice Index for aging with, 807.e7–807.e12
self-report, among Hindu temple priests, prevalence of and risk factors for, 805.e1–805.e12
self-reported, among Yakshagana artists, prevalence and risk factors of, 124.e35–124.e47
survey on machine learning approaches for automatic detection of, 947.e11–947.e33
- VTDS and VoiSS in evaluation of patients with, 381.e23–381.e32
- Voice Handicap Index-10 (VHI-10), validity and reliability study of Bahasa Malaysia version of, 581.e17–581.e23
- Voice Handicap Index (VHI)
cross-cultural adaptation and validation of Danish, 441–444
and laryngeal EMG findings, vocal fold paresis incidence, and relationship between, 940–944
in Norwegian student teachers, voice disorders and impact of, 445–452
- Voice high-frequency oscillation, and LaxVox exercises, in vocally healthy subjects, immediate effects of, 808.e7–808.e14
- Voice pathology, using cepstrum vectors, detection of, 634–641
- Voice quality
in adults, effect of hydration on, 125.e13–125.e28
analysis, improve version of voice analyzer programs for, 642–648
effect of auditory perceptual training with natural voice anchors on, 220–225
effect of electronic cigarettes on, 811.e13–811.e17
and orofacial strength as outcome of levodopa effectiveness in patients with early idiopathic Parkinson disease, 716–720
and parent proxy/child self-report of VRQOL, perceptual evaluation of voice disorder in children with history of laryngotracheal reconstruction surgery and relationship between clinician perceptual rating of, 945.e27–945.e35
of pediatric speakers, perception of, 204–213
perception, effect of telephone transmission on, 649–658
and respiratory tract infections in 4-year-old children in STEPS study, 801.e21–801.e25
tracheoesophageal, with SToPS, perceptual assessment of, 465–472
- Voice rehabilitation, lessons in, 669–681
- Voice-Related Quality of Life (VRQOL)
pediatric, validation of Mandarin Chinese version of, 325–332
perceptual evaluation of voice disorder in children with history of laryngotracheal reconstruction surgery and relationship between clinician perceptual rating of voice quality and parent proxy/child self-report of, 945.e27–945.e35
- Voice self-assessment, and musculoskeletal pain, in tele-operators, relation between, 948.e11–948.e21
- Voice Symptom Scale (VoiSS) and VTDS in evaluation of patients with voice disorders, 381.e23–381.e32
- Voice therapy
associated with electromyographic biofeedback in women with behavioral dysphonia, effectiveness of, 381.e11–381.e22
for elderly when applying conventional and intensive approaches, effectiveness of, 809.e19–809.e26
patients perceptions of and attitudes toward, 860–865
program based on taxonomy of, in women with behavioral dysphonia, effect of, 256.e1–256.e16
and psychological evaluation, clinical features of psychogenic voice disorder and efficiency of, 250–254
resonant, in treatment of vocal fold nodules, assessment of, 810.e1–810.e4
success in patients with functional dysphonia, standard instrumental acoustic, perceptual and subjective voice outcomes in, 317–324
using lip trill technique in patients with glottal gap, effects of, 949.e11–949.e19
water resistance, impact stress in, 490–496
- VoiSS. See Voice Symptom Scale
- Vowels
identity, stress type, baseline f0 and loudness on relative fundamental frequency with healthy voices, effects of, 603–610

- perceptual and quantitative assessment with dysphonia across,
473–481
- and speech, effect of Parkinson disease tremor phenotype on
cepstral peak prominence and transglottal airflow in,
580.e11–580.e19
- VRQOL. See Voice-Related Quality of Life
- VTDS. See Vocal Tract Discomfort Scale
- Wound healing, on vocal fold injuries, effects of intracordal estradiol
and dexamethasone injection on, 759–766
- Zona incerta, and subthalamic nucleus deep brain stimulation in
patients with Parkinson disease, voice tremor outcomes of,
545–549