



# Emotional talk of patients with breast cancer during review appointments with therapeutic radiographers: effects on fears of cancer recurrence

G. Humphris<sup>1,2</sup> · Y. Yang<sup>3</sup> · L. Barracliffe<sup>2</sup> · J. Cameron<sup>1</sup> · C. Bedi<sup>1</sup>

Received: 6 June 2018 / Accepted: 20 September 2018 / Published online: 1 October 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

## Abstract

Fears of cancer recurrence (FCR) in patients with breast cancer may develop during radiotherapy. Radiographer communication with their patients may influence early survivorship FCR level.

**Aim** To investigate the management of emotional talk in patients with breast cancer attending their initial review appointments during radiotherapy and predict FCR at 6–8 weeks follow-up.

**Methods** A mixed-methods observational study was conducted. Patients (consecutive sample,  $n = 60$ ) with breast cancer, attending a major Scottish cancer centre, had their first two review appointments with their therapeutic radiographer (TR) audio-recorded. In addition, FCR was assessed (FCR7) at baseline and at 6–8 weeks following their final radiotherapy visit. Two TRs participated. Audio files were coded by the VR-CoDES system to identify emotional cues and therapeutic radiographer (TR) responses. Linear regression models were tested for fit and to identify factors associated with follow-up FCR, i.e. patient cues, responses by TR.

**Results** Follow-up FCR was predicted negatively (robust estimator,  $p = .01$ ) by level of patient emotional talk at the second review session. The provision of space by the TR, at the second session, to enable patients to expand their emotional utterances was also associated, but negatively ( $p = .01$ ), with follow-up FCR. These effects were maintained after inclusion of covariates: age, treatment received and living conditions.

**Conclusions** Patient's emotional expression and TR responses at the second review meeting predicted follow-up FCR. The study shows the effect of communication processes on this specific distress component of the patient's survivorship experience.

**Trial registration** NCT02599506

**Keywords** Breast cancer patients · Fears of recurrence · Communication · Emotional talk · Survivorship · Distress · Therapeutic radiotherapy · Behavioural coding

## Background

Patients with breast cancer may experience distress in the form of fears of cancer recurrence (FCR) during their recovery from primary treatment [6]. The proportion who report moderate to severe FCR varies from 30 to 70% [23]. These levels are unpleasant for patients and have negative consequences including propensity to depression, sleeplessness, reduced quality of life, requests for more checks and increased health service utilisation [18]. One report has shown that FCR may be present as early as the diagnostic stage of cancer [12]. Systematic review methodology has shown that treatments, such as chemotherapy [31] and radiotherapy [30], on average, have been associated significantly with FCR increases. Admittedly, correlational investigations are of limited value

---

✉ G. Humphris  
gmh4@st-andrews.ac.uk

<sup>1</sup> Edinburgh Cancer Centre, Western General Hospital, Edinburgh, UK

<sup>2</sup> Division of Population and Behavioural Science, Medical School, University of St Andrews, North Haugh, St Andrews, Fife KY16 9TF, UK

<sup>3</sup> Department of Psychiatry and Psychology, Nanfang Hospital, Southern Medical University, Guangzhou 510515, Guangdong, China

especially with high heterogeneity in time locations of FCR assessments. However, the psycho-oncology literature has recognised that the way clinicians communicate to their patients is a vital aid to patient recovery [7]. A detailed model has been developed to assist researchers investigate these processes [26]. Our research focus is radiotherapy treatment [28]. This complex and technically demanding intervention requires substantial resources not only from the health care providers but also including the patient, e.g. multiple regular visits requiring frequent travel. It is also, very often, the final recommended treatment prior to discharge and patients may remember clearly how staff managed them.

The field of health care communication is advancing in oncology with many recommendations for good practice and training programmes [13]. In summary, the clinician is expected to deliver comprehensible descriptions of the disease and treatment, to listen to patients' concerns and acknowledge their concerns in an empathic and considerate manner. The exact combination of these behaviours that may be favourable will vary between clinicians, patients and the treatment context. A qualitative study has reported patients' and oncologists' preferences on how to approach emotional talk [29]. Direct observations, however, may further assist our understanding. Research tools are available to test best evidence for clinicians to ease treatment acceptability and manage patient distress. One tool is the VR-CoDES system for coding emotional concerns and cues [4, 32]. We have reported on the sequence of emotional cues that patients express and the type of response from therapeutic radiographers [16]. In addition, in a mixed-methods study, we found that the expression of certain emotional content was linked to the FCR trajectory over radiotherapy treatment using a daily diary record [1]. Our interest was to explore the specific effects of patients raising emotional issues in appointments with their TR during their treatment. Furthermore, how TRs responded to this emotional content raised by patients might have an influence on FCR development. For example, the patient who expresses some form of distress about their experience of a symptom indicator (e.g. skin inflammation) may react negatively to the TR who distracts them, or moves immediately onto another issue, such as medication. The TR who enables the patient to expand or 'provide space' to talk through their concern might confer a lasting psychological benefit and prevent the patient dwelling or ruminating about the topic [5]. The manner of how these processes may influence outcomes have been outlined [9]. In summary, the patient who is enabled to express distressing content in their review consultation by the clinician may assist the patient in their regulation of emotional processing in the brain. The adoption of such a self-regulation model has been presented in the FCR field [19]. Our choice of outcome was FCR some weeks following treatment when patients had a period of time to recover and adapt to a life without intensive treatment, sometimes referred to as 'early

survivorship' [2]. Previous research is lacking on predicting distress following diagnosis [3] and has tended to concentrate on baseline variables such as clinical and socio-economic factors. Very few studies have focused on how patients are managed and their anxieties recognised or acknowledged during treatment. Hence, the focus of our research is to investigate an aspect of this patient management and establish if there is a link between the communication of TR and patient, during the radiotherapy treatment phase, and FCR approximately 2 months after discharge.

## Aim

To investigate the management of emotional talk in patients with breast cancer attending their initial review appointments during radiotherapy and assess the influence on fears of recurrence at 6–8 weeks follow-up from end of treatment. Our objectives were to test the effects on follow-up FCR of (1) patients disclosing emotional cues in routine review appointments and (2) clinicians responding to emotional cues with reducing space.

## Methods

### Sample

Consecutive patients attending the specialist cancer unit in a major hospital in Scotland were recruited over January to May 2016. Psychiatric disorder, non-English speaking, metastatic disease or consultant withdrawal were exclusion variables. Detailed sampling has been reported elsewhere [1] (see Appendix 1 Fig. 1 for flow chart). Two TRs were recruited from the cancer unit. They had no prior training in the management of FCR.

### Design

A consecutive mixed-methods approach was employed [17]. That is, self-report ratings of FCR were collected at baseline and at 6–8 weeks follow-up from radiotherapy treatment discharge. Codes of patient emotional talk and TR immediate responses were applied to the digital audio files of the first two review appointments. Hence, this was an observational study without randomisation.

### Measures

A study-specific demographic self-report questionnaire was administered to participants to assess the following: age in years, treatment regimen (regular vs  $\pm$  boost, i.e. regular treatment is 15 sessions, whereas boost is given for positive margins and under 50 years of age), received chemotherapy or

not, living alone vs living with others. Fears of cancer recurrence were assessed using the FCR7 which has been previously used with patients with breast, colorectal or head/neck cancer [15]. Reliability is good (Cronbach's alpha in reference breast cancer group 0.92), and there is evidence of validity.

The emotional content of the patient's interaction with the TR was assessed using the Verona CoDES system that recognises emotional 'hints' or cues including frankly expressed concerns of the patient. The system has been endorsed by over 60 scientific articles since publication of manuals in 2011 [5]. Specifically, the coding of the audio files was performed by one of the authors (LB) under the supervision of the study PI who is one of the VR-CoDES originators. The system comprises of two major components. First, the identification of all emotional concerns or more subtle hints, referred to as cues in the consultation. Second, the immediate response in the next utterance by the health provider. The cues are classified into seven defined meaningful categories (cue A to cue G) and the responses by the TR are split into two major types, namely 'providing space' or 'reducing space'. In other words, the TR could deliver a response to the patient to provide more detail through remaining silent, or inviting and encouraging the patient to expand their comments, or alternatively reduce space for the patient to elaborate by use of distraction, changing topic or blocking [20]. The distinction is important as the relatively rare occasions where reducing space has been identified following cancer treatment is associated with subsequent mental distress [8, 22]. Coder training consisted of detailed reading of the coding manual and completion of the on-line exercise material available. In addition, the first five consultations were double coded and discrepancies discussed with GH and precision enhanced. When certain utterances were highlighted by the coder as being difficult to identify, the following procedure was followed. Consensus was achieved after detailed discussion concentrating, first on the list of extensive examples supplied within the manual, and second, an inspection of the current context of the consultation topic. A reliability check of eight randomly chosen audio files was completed by two researchers and achieved a kappa of 0.85 (95% CI = 0.84, 0.86), Spearman's Rho = 0.96 [11].

## Procedure

The research assistant (YY) approached all potential participants and invited written consent following reading of the patient information sheet a minimum of 24 h prior to cancer centre attendance. The baseline questionnaire was administered prior to first radiotherapy treatment and review appointment with their TR. The review appointments were digitally recorded within the clinical setting with an unobtrusive digital studio-quality recorder with SD card unit storage that was transferred on the same day of recording to the 'safe haven' constructed at the Medical School for confidential storage on a

non-networked PC. The audio files were transformed by a propriety software package to be imported into the event logging software Observer XT™v12 for VR-CoDES coding. The eventual files were exported into an Excel spreadsheet with identifiers removed. The TRs were instructed to conduct their review sessions as usual. If the TR became aware of a highly distressed patient, then this individual was referred to the centre's psychological services as per usual protocol of the cancer centre.

The study was approved by East of Scotland Research Ethics Service (NRES reference number: 13/ES/0015) and registered with [ClinicalTrials.gov](https://www.clinicaltrials.gov), ID number: NCT02599506.

## Statistical analysis

Linear regression with a robust maximum likelihood estimator (STATA™v15) was applied [25]. After controlling for baseline FCR ( $R^2$  of 0.38 obtained from previous study [14]) with four test and four further control covariates, the sample size required to detect a 0.1  $R^2$  increase was 56 (*power squared routine* [25]). Two nested models were fitted to explain the dependent variable: FCR at follow-up, controlling for baseline FCR level. The first tested simultaneously the effects of number of cues in the two consecutive sessions and the frequency that patients were not provided space to expand on these emotional cues. The second model added control covariates, namely age in years, treatment regimen (regular vs boost), received chemotherapy or not, living alone vs living with others such as partner or spouse, to determine if the model one effects were maintained. A likelihood ratio test assessed model improvement. Alpha level of 0.05 was used for all tests (2-sided).

## Results

The demographic breakdown, clinical intervention and mode of radiotherapy treatment are presented for the sample (Table 1). The mean total number of cues and concerns identified per patient was about 20. The mean number of cues and concerns for each consultation (first and second) and the frequency of reducing space when this emotional talk was coded was calculated (Table 1). The TR reduced space following an instance of emotional talk on about 9% of occasions. Appendix 2 Table 3 shows examples of patient emotional talk and provider responses. Drop-out analysis of those patients without complete data on the demographic and treatment variables compared with those entered into the regression revealed no differences ( $p > 0.05$ ). The nearest discrepancy was the proportion who received chemotherapy. Patients who were not followed tended to be less likely to have received this additional treatment ( $\chi^2(1) = 2.45, p = 0.12$ ).

**Table 1** Descriptive aggregate data for variables in statistical models (1 and 2)

Variables	Mean	Std. deviation	Range: min–max
FCR7 follow-up	14.88	8.61	6–40
FCR7 baseline	17.21	7.52	6–39
Number of pt. cues: session 1	5.33	7.89	0–43
Number of pt. cues: session 2	5.03	6.42	0–23
Reducing space TR responses: session 1	0.63	0.88	0–3
Reducing space TR responses: session 2	1.08	1.47	0–7
Patient age (in years)	57.92	11.26	35–85
Radiotherapy			
Regular <sup>†</sup>	44 (73%)		
Regular + Boost	16 (27%)		
Treatment received			
No chemotherapy <sup>†</sup>	29 (48%)		
Chemotherapy	31 (52%)		
Social living condition			
With another <sup>†</sup>	48 (80%)		
Alone	12 (20%)		

<sup>†</sup> Reference category (0) in dummy variable assignment

Follow-up FCR7 was regressed on baseline FCR7 and explained 47% of the variance. Appendix Table 4 shows the raw correlation matrix. The a priori linear model demonstrated reasonable fit (model 1) with the raw data ( $R^2 = 0.54$ ;  $F = 35.45$ ;  $df_5, 54$ ;  $p < 0.001$ ). The number of cues and the frequency of reducing space by the health provider in the second consultation were associated with the dependent variable (both  $p$  levels  $< 0.05$ , Table 2). Model 2 included additional

covariates to adjust for potential bias and explained approximately 2% extra variance of follow-up FCR. The likelihood ratio ( $LR$ ) test showed that there was no improvement in model 1 variance explained by the additional four covariates, i.e. model 1 nested in model 2;  $\chi^2(4) = 2.15$ ,  $p > 0.7$ .

A supplementary set of models was run that included not only the number of emotional hints (i.e. cues) but also the less frequent concerns. These regressions varied very little and failed to improve dependent variable explanation. Furthermore, separate models of first and second consultation variables were fitted and gave similar  $p$  levels. The addition of the TR identity into all regression models revealed trivial explanation of FRC follow-up values and communication behaviour.

## Discussion

The overall aim of the study was achieved by collecting a substantial set of 120 consultation recordings for detailed coding with accompanying patient completed FCR ratings and background variables. The first objective was to describe the level of emotional talk in both of the review appointments with the patient's TR. The level of cues and concerns identified was relatively high compared with other samples of patients with cancer, especially when the length of the appointments was no greater than 15 min [10]. For example, in a study of follow-up appointments with the oncologist and adolescent patients, the average number of cues and concerns expressed was 3.6 per consultation that averaged 24 min in duration [21]. The association of cues, in the current study, with the follow-up FCR rating was positive and significant for the second session only. The strength of the effect was within the a priori significance level.

**Table 2** Multiple linear regression (robust estimator) to predict follow-up FCR7

		Model 1			Model 2		
		Beta	95% CI	$p$	Beta	95% CI	$p$
Session 1	FCR7 baseline	0.716	0.48, 0.94	0.0001	0.685	0.47, 0.89	0.0001
	Cues	0.106	-0.11, 0.12	0.32	0.047	-0.19, 0.28	0.69
	Reduce space	0.749	-1.25, 2.74	0.46	1.053	-1.06, 3.16	0.32
Session 2	Cues	-0.514	-0.90, -0.12	0.011	-0.468	-0.91, -0.03	0.037
	Reduce space	1.934	0.45, 3.41	0.011	1.994	0.48, 3.51	0.011
	Age				0.046	-0.15, 0.24	0.64
	Chemotherapy <sup>†</sup>				-2.447	-6.26, 1.37	0.20
	Treatment <sup>†</sup>				-0.256	-5.75, 5.24	0.93
	Live alone <sup>†</sup>				0.912	-3.33, 5.16	0.63
	Constant	1.814	-1.81, 5.52	0.33	3.06	-9.77, 15.9	0.63
	$R^2$	0.54			0.56		
	$LR$ test				$\chi^2(4) = 2.15$		0.71

FCR7 fears of cancer recurrence 7-item scale

<sup>†</sup> Specified as dummy variable 0.1 (see Table 1)

The second related objective was to examine the aggregate responses by the TR to patient emotional talk exhibited in both sessions and inspect the association with follow-up FCR. Again, this association was significant only for the second session but negative. Hence, the first session expression of emotions and their management by the TRs was not important in understanding the patients' FCR level at follow-up.

The introduction of the number of concerns, as part of the emotional talk identified, into the analysis did not improve explanation; hence, we excluded the addition of concerns into the emotional talk quotients and simply used the number of cues (for both sessions). This selection, admittedly post hoc, could be of theoretical significance. Paradoxically, it may be the somewhat more hidden emotional issues that are embedded in the interactions with patients undergoing radiotherapy that may be the most important for the TR to recognise. Therefore, it is not simply the most obviously signalled concerns ('I am worried the treatment isn't working') that are relatively easy to recognise by the health provider that need appropriate responses, but rather the more subtle indicators of emotion, e.g. 'I am wondering what these "electric" twitches I feel now really mean'. Likewise the frequency of closing down of patients was relatively low (9% of all TR responses) but potentially important. This phenomena has been identified as a critical moment in many clinical interviews [5]. Patients' presentation of emotional speech following the first session review meeting, that is during the second and subsequent sessions, may be especially sensitive to being closed down by the TR. As raised previously in the literature, once baseline distress levels are controlled for, additional predictors of distress outcomes are often 'masked' out [3]. Our data would partially support this, except we have identified some processes occurring during treatment to relate to follow-up FCR. The importance of the patient cues presented and TR closing down on occasion at the second review appointment may be interpreted as follows. The initial review may be simply functioning as an information exchange, and emotional expression may assume little significance. The second appointment, however, after the patient has received some active radiotherapy treatment, experienced some side effects and have the desire to share their anxieties. These anxieties, if not expanded upon or shared with the current health care provider, may suppress these potential concerns and generate ruminative processes responsible for FCR development [19]. A recent article that is consistent with this formulation has demonstrated the process of patients experiencing 'symptoms' potentially triggering FCR development and consequent behavioural responses [24]. An accessible model that links these events to FCR has been explicated by Lee-Jones et al. [19]. It states that certain antecedents in a cognitive-behavioural framework operate to raise FCR which can be reinforced through positive or negative consequences. The attendance by the patient to their radiotherapy treatment session may sensitise these patients and raise associated anxiety about cancer risk, success of treatment and

possible recurrence. The experience of the patient who expresses these cues or concerns and then not have them attended to by the clinician may, as a consequence, reinforce psychological avoidance by the patient to subsume the emotional experience. Alternatively, the experience may simply provide an indicator to the patient that these issues are not regarded as important and are better discarded if possible. The longer term psychological consequence of these processes, according to the illness representation model of Lee-Jones et al., is an exacerbation of FCR. The clinical implication of this study would be for therapeutic radiographers to enable their patients to express their concerns whether they are specific to experienced symptoms and side effects, or considerations of cancer recurring.

## Limitations

A modest sample size restricted our ability to test the expression of specific types of emotional cue on FCR development. It would have been interesting to be guided as to whether, for example, the expressed hints to an emotion were focused on issues about external stresses outside the clinic environment (i.e. cue D in the VR-CoDES system). Two members of the therapeutic radiographer staff participated; hence, it is recognised that this cannot be generalised across radiography services. The baseline FCR assessment was conducted prior to the initial consultation by the TR and the first radiotherapy treatment session in an attempt to standardise the measurement process. All other ratings were assessed after the review appointments lending a possible complexity to interpretation. We conducted some checks for representativeness of the patient sample and showed that the participants retained are not substantially different from the total collected; however, the baseline variables were not comprehensive and there is still potential for systematic bias. Furthermore, we expected patients to have no further treatment from the cancer centre following their final radiotherapy session. They may have been in contact with the centre for advice but this was not recorded. Replication is advisable and these results should be treated with caution. Future work should include a more comprehensive inclusion of both patients and staff and include additional potential confounds including severity of illness and staff variables such as seniority and level of training. In addition, we concentrated on the first two review consultations and there is scope to include all four review appointments in the recordings to assess a more extensive longitudinal panel of data.

## Conclusions

The management of emotional talk in patients with breast cancer during their early weekly review appointments with their TR predicts eventual FCR level some 6–8 weeks post discharge from primary treatment. Additional training to increase awareness of these features on the consultations of

these patients may be of benefit. Studies need to be conducted to examine the generalisability of these findings and pinpoint whether there are particular emotional issues that may warrant close attention to prevent the development of disabling FCR levels. A new study protocol to develop a training programme for TRs has been reported [27].

**Acknowledgements** All patients and staff that participated are gratefully acknowledged.

**Funding information** The FORECAST study was supported generously by Breast Cancer Now (Reg. No. SP529).


## Compliance with ethical standards

The study was approved by East of Scotland Research Ethics Service (NRES reference number: 13/ES/0015) and registered with [ClinicalTrials.gov](http://ClinicalTrials.gov), ID number: NCT02599506.

**Conflict of interest** The authors declare that they have no conflict of interest.

## Appendix 1

**Fig. 1** Flow chart for the detailed sampling



## Appendix 2

**Table 3** Examples of patient utterances and provider responses (using VR-CoDES system)

Patient utterances			
Code	Example	Pat. no.	Sess. no.
Concern	‘I’m terrified; what kind of scan this is’	5	2
	‘But it’s him I’m more concerned about, I’m not interested in myself’	6	1
	‘I hate being ill! I’m not a person that likes being ill’	6	2
	‘I feel very stressed out and depressed and worried’	23	1
Cue A	‘It’s just been a bit full on’	97	1
	‘It’s getting to the stage where don-, I honestly don’t think I can do it much longer’	9	2
	‘...if I feel a bit uuohh’	72	1
Cue B	‘It feels like a needle is pinning me’	58	2
	‘What’s the point of taking an antidepressant?!’	38	2
Cue C	‘The chemo was for me the killer’	26	1
	‘Last night I had very little sleep..., so today I’m tired’	32	2
	‘Most of my pain, in fact probably all of my pain is coming from the fissure’	17	2
Cue D	‘Man, I woke up with a massive hot flush’	1	1
	‘We haven’t told the boys because we thought we were going to have to cancel everything when I got diagnosed’	94	2
	‘My husband’s not well, and he won’t take responsibility but, he’s never come with me to an appointment’	9	2
Cue E	‘I try not to bother my younger sister, she’s seven years younger than me, because she’s recovering from cancer and she doesn’t keep very well’	23	1
	‘It’s only when I’ve had the radio that it does it’; ‘And it only happens when I’ve had the radio’	49	2
	‘What should I look out for?’; ‘It’s just nice to sort of know what to look out for’	24	1
	‘Apart from the... time for the bus again!’; ‘Just...come in...get off, get on, go!’	2	1
Cue F	‘Feel slightly more tired on a Monday’; ‘I was probably a wee bit more tired Monday’	83	2
	<i>Inapplicable to audiotapes</i>	–	–
Cue G	‘I was more angry...’	47	1
	‘I was really, really frightened’	19	1
	‘Panicky, like a panic attack...anxious, anxiety, erm, lack of confidence’	32	1
	‘I was a bit concerned I would react’	8	2
Provider responses (CD, closing down; PS, providing space; Pat, patient; Prov, provider)			
Code	Example	Pat. no.	Sess. no.
Ignore (CD)	Pat: ‘Yes yes, struggling a bit this week’; Prov: ‘How are you getting on with your tamoxifen, are you having any more problems with bruising?’	97	1
	Pat: ‘That was the worst bit’; Prov: ‘Yep, so going back to the Letrozole’	5	1
Shutting down (CD)	Pat: ‘Sorry for...’; Prov: ‘No, do not apologise!’	34	2
	Pat: ‘...no, I’m getting worried’; Prov: ‘Noo, not at all!’	89	2
Acknowledgement (PS)	Pat: ‘I’m trying to figure out what normal is!’; Prov: ‘I know, uhuh, mhm’	1	1
	Pat: ‘I was really, really frightened’; Prov: ‘Were you, oh dear’	19	1
Implicit empathy (PS)	Pat: ‘I didn’t expect this on top of it, because it’s not connected seemingly... it’s not connected’; Prov: ‘I know, it’s a lot. A lot for you to get your head round’	72	1
	Pat: ‘I just feel so tired’; Prov: ‘Sounds like you have been fighting a long time, to stay afloat’	9	2
Content exploration (PS)	Pat: ‘I’m still sore inside eh, it’s quite jagged and sore’; Prov: ‘Are you getting sharp shooting pains?’	14	1
	Pat: ‘I, I go to bed and I go to sleep, but I do waken up’; Prov: ‘Right, so a bit more disturbed?’	19	2
	Pat: ‘And it just suddenly hits you, this kind of overwhelming tiredness’; Prov: ‘Yeah, and are you sleeping alright?’	15	2

**Table 3** (continued)

Patient utterances			
	Pat: 'Under my arm's been a bit sore...a bit kinda niggly... inside...inside kinda niggly feeling, I don't know...'; Prov: 'Sort of sharp shooting pains? Or just more a kind of pulling sensation?'	90	2
	Pat: 'Maybe the co-codamol has worn off, I've got to put my other foot forward, because to put my full weight on it, it's painful'; Prov: 'Uhuh, so that's your left hip isn't it?'	92	1
Affective acknowledgement (PS)	Pat: 'my biggest concern is getting here and parking and, and, getting through the traffic and and so on'; Prov: 'It can be stressful'	21	1
Affective exploration (PS)	Pat: 'I've never felt as emotionally raw'; Prov: 'How did you feel after you were told that you'd had the cancer diagnosed, did you feel upset then or..., do you think this is a bit delayed?'	43	1
Affective empathy (PS)	Pat: 'You think for God's sake how did this happen'; Prov: 'I think that's hard to deal with when you are doing all you can to keep yourself healthy'	97	2
	Pat: 'I said I felt very lonely... I feel like everything's my responsibility, I'm the only one that's working'; Prov: 'And that's very, very hard when you are the one that's needing some support and attention yourself'	9	1

## Appendix 3

**Table 4** Pearson correlation matrix

Variable	1	2	3	4	5	6	7	8	9
1 FCR7 follow-up	1.000								
2 FCR7 baseline	0.686	1.000							
3 No. of pt. cues: session 1	0.303	0.351	1.000						
4 No. of pt. cues: session 2	0.005	0.071	0.415	1.000					
5 Reducing space TR responses: session 1	0.271	0.237	0.385	0.318	1.000				
6 Reducing space TR responses: session 2	0.205	0.105	0.318	0.727	0.362	1.000			
7 Age (in years)	-0.134	-0.190	0.114	-0.047	0.065	-0.175	1.000		
8 Treatment received <sup>†</sup>	-0.291	-0.298	-0.122	0.105	0.166	0.078	0.390	1.000	
9 Social living conditions <sup>†</sup>	0.163	0.120	0.160	-0.117	0.019	0.000	0.037	-0.017	1.000
10 Radiotherapy <sup>†</sup>	0.114	0.190	-0.022	0.177	-0.006	0.249	-0.515	-0.096	0.075

<sup>†</sup> See dummy variable key in Table 1

## References

- Barraciff L, Yang Y, Cameron J, Bedi C, Humphris G (2018) Does emotional talk vary with fears of cancer recurrence trajectory? A content analysis of interactions between women with breast cancer and their therapeutic radiographers. *J Psychosom Res* 106:41–48
- Breaden K (1997) Cancer and beyond: the question of survivorship. *J Adv Nurs* 26:978–984
- Cook SA, Salmon P, Hayes G, Byrne A, Fisher PL (2018) Predictors of emotional distress a year or more after diagnosis of cancer: a systematic review of the literature. *Psychooncology* 27:791–801
- Del Piccolo L, de Haes H, Heaven C, Jansen J, Verheul W, Bensing J, Bergvik S, Deveugele M, Eide H, Fletcher I, Goss C, Humphris G, Kim YM, Langewitz W, Mazzi MA, Mjaaland T, Moretti F, Nubling M, Rimondini M, Salmon P, Sibbern T, Skre I, van Dulmen S, Wissow L, Young B, Zandbelt L, Zimmermann C, Finset A (2011) Development of the Verona coding definitions of emotional sequences to code health providers' responses (VR-CoDES-P) to patient cues and concerns. *Patient Educ Couns* 82:149–155
- Del Piccolo L, Finset A, Mellblom A, Figueiredo-Braga M, Korsvold L, Zhou Y, Zimmermann C, Humphris G (2017) Verona coding definitions of emotional sequences (VR-CoDES): conceptual framework and future directions. *Patient Educ Couns* 100(12):2303–2311
- Dunn LB, Langford DJ, Paul SM, Berman MB, Shumay DM, Kober K, Merriman JD, West C, Neuhaus JM, Miaskowski C (2015) Trajectories of fear of recurrence in women with breast cancer. *Support Care Cancer* 23:2033–2043
- Epstein R, Street R (2007) Patient-centered communication in cancer care: promoting healing and reducing suffering. National Cancer Institute, NIH Publication No. 07-6225, Bethesda
- Farrell C, Heaven C, Beaver K, Maguire P (2005) Identifying the concerns of women undergoing chemotherapy. *Patient Educ Couns* 56:72–77
- Finset A (2012) "I am worried, Doctor!" Emotions in the doctor-patient relationship. *Patient Educ Couns* 88:359–363
- Finset A, Heyn L, Ruland C (2013) Patterns in clinicians' responses to patient emotion in cancer care. *Patient Educ Couns* 93:80–85
- Fletcher I, Mazzi M, Nuebling M (2011) When coders are reliable: the application of three measures to assess inter-rater reliability/agreement with doctor-patient communication data coded with the VR-CoDES. *Patient Educ Couns* 82:341–345
- Gross SE, Nitzsche A, Gloede TD, Ansmann L, Street R, Pfaff H, Neumann M, Wirtz M, Baumann W, Schmitz S, Ernstmann N (2015) The initial clinical interview—can it reduce cancer patients' fear? *Support Care Cancer* 23:977–984
- Halkett GKB, Kristjanson LJ, Lobb E, O'Driscoll C, Taylor M, Spry N (2010) Meeting breast cancer patients' information needs during radiotherapy: what can we do to improve the information and support that is currently provided? *Eur J Cancer Care* 19:538–547
- Humphris G, Rogers S (2012) AFTER and beyond: cancer recurrence fears and a test of an intervention in oral and oropharyngeal cancer. *Soc Sci Dentistry* ISSN2040-4263 2:29–38
- Humphris G, Watson E, Sharpe M, Ozakinci G (2018) Unidimensional scales for fears of cancer recurrence and their psychometric properties: the FCR4 and FCR7. *Health Qual Life Outcomes* 16:30
- Humphris G, Yuan Y, Barracliffe L, Cameron J, Bedi C (2017) Therapeutic Radiographer responses to emotional talk in breast cancer patients undergoing radiotherapy: a longitudinal multi-level sequential study. International Conference on Communication in Healthcare, 8–11<sup>th</sup> October. Baltimore, USA
- Johnson R, Onwuegbuzie A (2004) Mixed methods research: a research paradigm whose time has come. *Educ Res* 33:14–26
- Lebel S, Tomei C, Feldstain A, Beattie S, McCallum M (2013) Does fear of cancer recurrence predict cancer survivors' health care use? *Support Care Cancer* 21:901–906
- Lee-Jones C, Humphris G, Dixon R, Bebbington-Hatcher M (1997) Fear of cancer recurrence: a literature review and formulation to explain exacerbation of recurrence fears. *Psycho-Oncology* 6:95–105
- Maguire P, Faulkner A, Booth K, Elliot C, Hillier V (1996) Helping cancer patients disclose their concerns. *Eur J Cancer* 32A:78–81
- Mellblom AV, Finset A, Korsvold L, Loge JH, Ruud E, Lie HC (2014) Emotional concerns in follow-up consultations between paediatric oncologists and adolescent survivors: a video-based observational study. *Psychooncology* 23:1365–1372
- Parle M, Jones B, Maguire P (1996) Maladaptive coping and affective disorders among cancer patients. *Psychol Med* 26:735–744
- Simard S, Thewes B, Humphris G, Dixon M, Hayden C, Mireskandari S, Ozakinci G (2013) Fear of cancer recurrence in adult cancer survivors: a systematic review of quantitative studies. *J Cancer Surviv* 7:300–322
- Soriano EC, Valera R, Pasipanodya EC, Otto AK, Siegel SD, Laurenceau JP (2018) Checking behavior, fear of recurrence, and daily triggers in breast cancer survivors. *Ann Behav Med*. <https://doi.org/10.1093/abm/kay033>
- StataCorp (2017) Stata Statistical Software: Release 15. StataCorp LLC. College Station, Texas
- Street RL Jr (2013) How clinician-patient communication contributes to health improvement: modeling pathways from talk to outcome. *Patient Educ Couns* 92:286–291
- van Beusekom M, Cameron J, Bedi C, Banks B, Kelse T, Humphris G (2018) Development, acceptability and feasibility of a communication skills training package for therapeutic radiographers to reduce fear of recurrence development in breast cancer patients (FORECAST2). *Pilot and Feasibility Studies* 4:148
- van Beusekom M CJ, Bedi C, Banks E, Humphris G (submitted) Communication skills training for the radiotherapy team to manage cancer patients' emotional concerns: a systematic review
- Visser LNC, Schepers S, Tollenaar MS, de Haes H, Smets EMA (2018) Patients' and oncologists' views on how oncologists may best address patients' emotions during consultations: an interview study. *Patient Educ Couns* 101(7):1223–1231
- Yang Y, Cameron J, Humphris G (2017) The relationship between cancer patient's fear of recurrence and radiotherapy: a systematic review and meta-analysis. *Psychooncology* 26:738–746
- Yang Y, Wen Y, Bedi C, Humphris G (2017) The relationship between cancer patient's fear of recurrence and chemotherapy: a systematic review and meta-analysis. *J Psychosom Res* 98:55–63
- Zimmermann C, Del Piccolo L, Bensing J, Bergvik S, De Haes H, Eide H, Fletcher I, Goss C, Heaven C, Humphris G, Kim YM, Langewitz W, Meeuwesen L, Nuebling M, Rimondini M, Salmon P, van Dulmen S, Wissow L, Zandbelt L, Finset A (2011) Coding patient emotional cues and concerns in medical consultations: the Verona coding definitions of emotional sequences (VR-CoDES). *Patient Educ Couns* 82:141–148