



## Research Paper

## Surgical site infection in a tertiary care centre-an overview - A cross sectional study

R. Saravanakumar <sup>b</sup>, B. M. Pabitha Devi <sup>a,\*</sup><sup>a</sup> Dept. of General Surgery, Tirunelveli Medical College, Tirunelveli, Tamilnadu, India<sup>b</sup> Dept. of General Surgery, Pudukkottai Medical College, Pudukkottai, Tamilnadu, India

## ARTICLE INFO

## Article history:

Received 24 March 2019

Accepted 25 September 2019

Available online 4 October 2019

## Keywords:

Surgical site infection

Elective surgery

Emergency surgery

Clean contaminated wound

Antibiotic sensitivity

## ABSTRACT

**Background:** Surgical Site Infections (SSIs) are the third most frequently reported nosocomial infection, accounting for 14 to 16 percent of all nosocomial infections among hospitalized patients according to National Nosocomial Infections Surveillance. SSIs are responsible for the increased morbidity and mortality related to surgery. Surgical wounds are classified as clean, clean contaminated, contaminated and dirty wounds as per CDC criteria. Better understanding of the causal factors of SSIs and prevention will reduce the cost of the surgery.

**Materials and methods:** 1570 patients who underwent surgery in the dept. of General surgery in a tertiary care hospital were taken for the study. Out of which 990 patients underwent elective surgical procedures and 580 underwent emergency procedures. An elaborate study of these cases with regard to date of admission, history, clinical features, type of surgery, emergency or elective, preoperative preparation, type of incision, contamination, procedure done, peroperative findings, drain used and its type and duration of operation and postoperative management and signs of wound infection were done and the findings are tabulated.

**Results:** 115 cases out of 1570 surgeries had surgical site infections. The overall post-operative SSI rate in elective clean and clean contaminated cases is 4.34% and emergency cases is 12.41%. The study showed that the superficial surgical site infections are the commonest type and accounted for about 72.09% in elective and 61.11% in emergency of all the SSI's and deep surgical site infection accounted for about 23% in elective and 30.55% in emergency cases. *E. Coli* is the commonest organism isolated from the surgical wounds that is 30.23% in elective cases. In emergency cases *Proteus mirabilis* is the most common organism involved in SSI followed by *E. Coli*.

**Conclusion:** *E. coli* is the commonest organism isolated from elective surgical wounds and *Proteus mirabilis* is the commonest organism isolated from emergency surgical wounds. A change in the pre operative antibiotics may reduce the incidence even lower. A preexisting medical illness like diabetes mellitus and duration of operating time, class of the wound, emergency surgeries and wound contamination had strong effect on the development of surgical site infection.

© 2019 The Authors. Published by Elsevier Ltd on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Background

Hundreds of Millions of People around the world undergo surgery each year. Infection of the surgical site (formerly referred to as “wound infection”, terminology that is no longer used owing to confusion between infections of surgical incisions and those of traumatic wounds) is a consequence of surgery, but it is not

inevitable. Surgical site infections are the commonest nosocomial infections and are responsible for the morbidity, mortality and significantly prolong the duration of hospitalization, thus adding the economic burden.

Surgical infections are those that occur as a result of a surgical procedure or those that require surgical intervention as part of their treatment. They are characterized by a breach of mechanical/anatomic defense mechanisms (barriers) and are associated with greater morbidity, significant mortality, and increased cost of care.

Despite the advances in surgical sciences post-operative wound infection remains one of the common complication which surgeons

\* Corresponding author.

E-mail addresses: [sarav3085@gmail.com](mailto:sarav3085@gmail.com) (R. Saravanakumar), [pabitha73@gmail.com](mailto:pabitha73@gmail.com) (B.M.P. Devi).

encounter. This problem if not evaluated and treated in a timely manner can have significant sequel. Based on NNIS (National Nosocomial Infections Surveillance) system reports, Surgical Site Infections (SSIs) are the third most frequently reported nosocomial infection, accounting for 14 to 16 percent of all nosocomial infections among hospitalized patients. Among surgical patients, surgical site infections were the most common nosocomial infection, accounting for 38 percent of all such infections. Of these surgical site infections 2/3rd were confined to the incision, and one third involved organs or spaces accessed during surgery.

## 2. Materials and Methods

This study was conducted in DEPARTMENT OF GENERAL SURGERY, in a tertiary care hospital. Elective and emergency general surgical cases were taken up for study. Surgeries on severely immunocompromised patients, patients with incomplete primary closure of the wounds and Re-look surgeries were excluded from the study.

A total number of 1570 patients admitted in general surgical wards for elective and emergency surgery in the study period, out of which 990 were elective cases and 580 were emergency cases, fulfilling our study criteria. Surgeries were classified as clean, clean – contaminated, contaminated and dirty wounds. The elective surgeries fall under first two categories while emergency procedures came under contaminated and dirty wounds. Clean Wound was defined as an uninfected operative wound in which no inflammation is encountered at the respiratory, alimentary, genital or uninfected urinary tracts are not entered. In addition, clean wounds are primarily closed and if necessary, drained with closed drainage. Operative incision wound that follows non-penetrating (blunt) trauma should be included in this category if they meet the criteria. Clean contaminated wounds were defined as operative wounds in which respiratory, alimentary, genital or urinary tracts are entered under controlled conditions and without unusual contamination. Specifically, operations involving the biliary tract, appendix, vagina and oropharynx are included in this category provided no evidence of infection or major breaks in technique is encountered.

All patients received pre-operative antibiotic injection of an aminoglycoside along with metronidazole one hour before surgery. Totally 115 cases had surgical site infections which had been taken up for this study. An elaborate study of these cases with regard to date of admission, history, clinical features, type of surgery, emergency or elective, preoperative preparation, type of incision, contamination, procedure done, peroperative findings, drain used and its type and duration of operation and postoperative management and post-operative findings which included day of 1st dressing and frequency of change of dressing, day of wound infection, signs of wound infection were noted which included fever, erythema, discharge, type and color of the exudates till patient is discharged from hospital, and then followed up the patient for 6 weeks on OPD basis for any signs of wound infection.

All patients with discharge of the wound, culture and sensitivity of the discharge was done and treated accordingly. Symptomatic treatment was given depending upon the combination and severity of various symptoms due to SSI. SSI are defined as incisional (either superficial or deep) infection or organ-space infection according to CDC criteria [1]. Superficial incisional SSI involve skin and subcutaneous tissues only, common stitch mini-abscesses; deep incisional infections involve fascia and muscle; and organ-space infections involve any organ or space other than the incised layer of body wall that was opened or manipulated during surgery [2].

## 3. Results

In our study, the overall post-operative SSI rate in elective clean and clean contaminated cases is 4.34% and emergency cases is 12.41%. It showed that the superficial surgical site infections are the commonest type and accounted for about 72.09% in elective and 61.11% in emergency of all the SSI's and deep surgical site infection accounted for about 23% in elective and 30.55% in emergency cases.

### 3.1. Age distribution in SSI

Age in Years	No of Elective Cases with SSI	Percentage	No of Emergency Cases With SSI	Percentage
<20	2	4.6	6	8.3
21–40	14	32.5	25	34.72
41–60	16	37.2	31	43.05
>60	11	25.5	10	13.88

In our study, most of the cases were in the middle age group, almost 69.7% and 77.7% of the cases were in the age group between 20–60 years in elective and emergency cases respectively. 11% and 10% of the cases were in the elderly age group in elective and emergency cases respectively. Extremes of age is one of the risk factors in SSI as shown in few studies.

Our study did not have significant statistical difference with respect to the relation between sex and SSI. Although the incidence was slightly more in case of men accounting for 53.4% in elective and 58.3% in emergency cases.

### 3.2. Elective cases

Type	No of Cases	Percentage
Superficial SSI	31	72.1
Deep SSI	10	23.25
Intra-Abdominal Abscess	2	4.65

### 3.3. Emergency cases

Types	No of Cases	Percentage
Superficial SSI	44	61.11
Deep SSI	22	30.55
Intra-Abdominal Abscess	6	8.33

From the above observation, it shows that the superficial surgical site infections in the commonest type and accounted for about 72.09% in elective and 61.11% in emergency of all the SSI's and deep surgical site infection accounted for about 23% in elective and 30.55% in emergency cases.

### 3.4. Drain and infection rate

Wound	No of Elective Cases	Percentage	No of Emergency Cases	Percentage
With Drain	29	67.44	48	66.66
Without Drain	14	32.55	24	33.33

Surgical site infection is more common in surgeries when a drain is used.

### 3.5. Co-morbidity and infection rate

Co-Morbidity	No of Elective Cases	Percentage	No of Emergency Cases	Percentage
Diabetes Mellitus	15	34.9	19	26.4
Obesity	7	16.3	15	20.8
Anemia	5	11.6	12	16.6
Hypertension	3	7	4	5.5
Others	13	30.2	22	30.5

Diabetes is the foremost comorbid condition associated with SSI. In our study, 13.04% of elective clean cases, 24.34% of elective clean contaminated cases and 62.6 % of emergency cases developed SSI.

### 3.6. Organisms isolated from SSI wounds

Types of Organism	No of Elective Cases	Percentage	No of Emergency Cases	Percentage
<i>E. coli</i>	13	30.23	14	19.44
<i>Proteus Mirabilis</i>	9	20.93	16	22.22
<i>Klebsiella Pneumoniae</i>	7	16.27	9	12.5
<i>Klebsiella Oxytoca</i>	3	6.97	6	8.33
<i>Proteus Vulgaris</i>	4	9.30	7	9.27
No Growth	5	11.62	15	20.83
Others	2	4.65	5	6.94

*E. coli* is the commonest organism isolated from the surgical wounds that is 30.23% in elective cases. In emergency cases *Proteus mirabilis* is the most common and *E. coli* is second most common organism isolated from the wound.

### 3.7. Relation of organisms with the type of surgery

Type	<i>E. coli</i>	<i>P.Mirabilis</i>	<i>K.Pneumoniae</i>	<i>P.Vulgaris</i>	<i>K.Oxytoca</i>
Elective Clean	33.33%	26.66%	20%	13.33%	6.66%
Elective Clean Contaminated	38.09%	23.80%	19.04%	9.52%	9.52%
Emergency	26.92%	30.76%	17.30%	13.46%	11.53%

Secondary suturing was done in 34.8% of cases and the remaining 65.2% of cases healed by secondary intension. Method used for wound healing was preferred based on the site, size and intensity of the infection. Smaller wounds with less and superficial infections were allowed to heal by secondary intension.

### 3.8. Antibiotic sensitivity

The less than 50% cultured aerobes demonstrated sensitivity to the cephalosporin's tested (Ceftazidime, Cefuroxime and Ceftriaxone) in infected patients. The proportion of bacteria resistant to all antibiotics for which tested was as high as 33.93%. *Proteus* was

mostly sensitive to Amikacin, Gentamycin, Vancomycin, clindamycin and Imipenam. And *E. Coli* was commonly sensitive to piperacillin, clavulanic acid, Imipenam and ceftazidime.

## 4. Discussion

In the present study, the post-operative SSI rate in elective clean and clean contaminated cases is 4.34% and emergency cases is 12.41%, accounting for an overall SSI rate of 7.32%. Reports of SSI from different workers gave different infection rates. Number of studies carried out in India indicate overall infection rate of 4.04–30% for clean surgical cases. In a prospective study conducted in Ethiopia, out of 105 patients who undergone surgical procedure, 20 patients developed SSIs which give overall incidence rate of 19.1% [15]. The finding was similar to studies done in India 20.09% [3], Nigeria 20.3% [4], India 21.66% [5] and Egypt 22.6% [6]. But, lower rate compared to studies done in Tanzania 26% [7], India 33.5% [8] and Mekele, Ethiopia 75% [9]. The finding was higher compared with studies done in USA 7.2% [10], France 2.5% [8], Egypt 9.2% [11] and Sudan 9% [12]. Studies done in Egypt and Sudan included only elective surgery but in our study both elective and emergency surgeries were included. Our results were comparable with US study.

### 4.1. Age

In our study, almost 69.7% and 77.7% of the cases with SSI were in the age group between 20 and 60 years in elective and emergency cases respectively. The study conducted at Ethiopia revealed that patients with age greater than 40 years were 7.72 times more likely to develop SSIs compared with patients in the age range of 19–40 years with AOR = 7.72 (95% CI [1.461–40.810, p = 0.016]). The result was in agreement with other studies [13,14] confirming that as age increases the risk occurrence of SSIs increases [15].

### 4.2. Co-morbidity and infection rate

Sixteen total studies were available to assess the association between hyperglycemia and SSI with ten papers (n = 27,844 procedures) including pre- or intra-operative assessments of blood glucose levels and eleven papers (n = 32,625 procedures) including post-operative assessments of blood glucose levels. The overall estimate for the association between elevated blood glucose in the post-operative period and SSI was 1.45 (95% Predictive Interval 0.77, 3.04). History of diabetes remained a significant risk factor in meta-analyses of studies that controlled for hyperglycemia [16]. In our study also, 34.9% of elective clean and clean contaminated cases and 26.4% of emergency cases with SSI had diabetes mellitus, stating Diabetes as a strong risk factor for the development of SSI irrespective of the type of surgery.

### 4.3. Drain and prosthesis

Infections are more in drained wounds and in the procedure involving implanting prosthesis like mesh. This increased incidence may be due to the effect of the drain itself by acting as a microbial pathway. Implants carry higher risk of infections acting as a foreign body if in case there is breach in the strict aseptic protocols. Umesh S. Kamat et al., in 2007 studied that patients with post-operative drain were 5.8 (2.33–14.66) times more likely to develop SSI compared to those without the drain. In our study, while the proportion of those with postoperative drain acquiring SSI was 2 times higher than those without drain. Further, the infection rate increases with the increasing duration of the drain. Hence, we can avoid keeping the drain, unless otherwise very much indicated. In

indicated cases also, use of a closed drain for a minimum period of time as short as 48 h will reduce the incidence of postoperative SSI.

#### 4.4. Organisms

Giacometti et al. studied 676 surgery patients with signs and symptoms indicative of wound infections, who presented over the course of 6 years. Bacterial pathogens were isolated from 614 individuals. A high preponderance of aerobic bacteria was observed. Among the common pathogens were *Staphylococcus aureus* (28.2%), *Pseudomonas aeruginosa* (25.2%), *Escherichia coli* (7.8%), *Staphylococcus epidermidis* (7.1%), and *Enterococcus faecalis* (5.6%) [17]. *Pseudomonas* was most common isolate in other studies like Mofikoya Bo et al. But in our study, *E. Coli* is the commonest organism isolated from the surgical wounds that is 30.23% in elective cases. The second most common was *Proteus mirabilis* (19.64) followed by *K. Pneumoniae* (16.27%). In emergency cases *Proteus mirabilis* is the most common organism involved in SSI followed by *E. Coli* which is a second most common organism isolated from the wound.

#### 4.5. Antibiotic sensitivity

The introduction of routine surgical antimicrobial prophylaxis (SAP) was a breakthrough in the prevention of SSI. Current guidelines, however, suggest that single-shot administration of a first-or second-generation cephalosporin is sufficient for optimal prevention of SSI in the absence of high rates of resistant bacteria. Due to a limited anaerobic activity of most cephalosporin, treatment is supplemented with metronidazole where indicated. Importantly, two large prospective studies observed the lowest risk of SSI when SAP was given within 30 min prior to incision, and the National Surgical Infection Prevention Project simply recommends administering SAP as close to the incision time as possible. Similar statements are made in European guidelines. The translocation of skin microorganisms into the wound during incision is the first vulnerable phase of surgery, and administering SAP only a few minutes before incision might not be optimal to achieve the tissue levels required to prevent SSI [18]. In our study, the proportion of bacteria resistant to all antibiotics for which tested was as high as 33.93%. *Proteus* demonstrated sensitivity to the aminoglycosides and cephalosporin. *E. Coli* was mostly sensitive to –Piperacillin, clavulanic acid and cephalosporin. Hence, we have to change the prophylactic antibiotics used prior to the surgical procedure accordingly.

### 5. Conclusion

*E.coli* is the commonest organism isolated from elective surgical wounds and *Proteus mirabilis* is the commonest organism isolated from emergency surgical wounds. Overall Amikacin, Gentamycin, Ciprofloxacin, Imipenam and Ceftazidime with Clavulinic acid are the sensitive drugs for SSI's developed in our hospital. Hence, change in the pre operative antibiotics may reduce the incidence even lower. A preexisting medical illness like diabetes mellitus, prolonged operating time, class IV wounds (dirty wounds), emergency surgeries and wound contamination strongly predispose to surgical site infection which has to be taken care of.

#### Limitations of the study

High risk factors for SSI like smoking and increased BMI were not taken into account. Regarding the bacteria, anaerobic bacteria was not cultured. Hence, SSI with negative cultures (17.1%) may be positive for anaerobic bacteria.

### Ethical approval

Ethical approval was given by Institutional Ethical Committee of Tirunelveli Medical College. Ref No:800/GS/2015.

### Funding

No funding.

### Author contribution

Dr. PabithaDevi contributed in Study design, data analysis and writing Dr. Saravanakumar contributed in Data collection and data analysis.

### Conflict of interest statement

No conflict of Interest.

### Guarantor

Dr. Pabitha Devi and Dr. Saravana Kumar.

### Research registration number

4718.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijso.2019.09.008>.

### References

- [1] Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. hospital infection control practices advisory committee. *Infect Control Hosp Epidemiol* 1999;20:250–78. <https://doi.org/10.1086/501620>. quiz 279–280.
- [2] Mujagic Edin, Zwimpfer Tibor, Marti Walter R, Zwahlen Marcel, Hoffmann Henry, Kindler Christoph, et al. Evaluating the optimal timing of surgical antimicrobial prophylaxis: study protocol for a randomized controlled trial. *Trials* 2014;15:188. <https://doi.org/10.1186/1745-6215-15-188>. Published online 2014 May 24.
- [3] Mahesh CB, Shivakumar S, Suresh BS, Chidanand SP. A Prospective study of surgical site infections in a teaching hospital. *J Clin Diagn Res* 2010;4:3114–9.
- [4] Nwankwo EO, Ibeh IN, Enabulele O. Incidence and risk factors of surgical site infection in a tertiary health institution in Kano, Northwestern Nigeria. *Int J Infect Control* 2012;8(4):8–13. <https://doi.org/10.3396/ijic.v8i4.035.12>.
- [5] Kikkeri N, Setty H, Shimoga M, Halumatthigatta D, Giriyaiah S, Ramanagaram N, et al. A study on Surgical Site Infections (SSI) and associated factors in a government tertiary care teaching hospital in Mysore, Karnataka. *Int J Med Public Health* 2014;4(2):171–5. <https://doi.org/10.4103/2230-8598.133126>.
- [6] Afifi IK, Labah EA, Ayad KM. Surgical site infections after elective general surgery in Tanta University Hospital: rate, risk factors and microbiological profile. *Egyptian J Med Microbiol* 2009;18(2):61–72.
- [7] Mawalla B, Mshana SE, Chalya PL, Mirzalioglu C, Mahalu W. Predictors of surgical site infections among patients undergoing major surgery at Bugando Medical Centre in Northwestern Tanzania. *BMC Surg* 2011;11(1):21. <https://doi.org/10.1186/1471-2482-11-21>.
- [8] Amutha B, Viswanathan T, Science I, Association C. A retrospective study on the pattern of pathogens isolated from surgical site wound infection in tertiary care hospital in coimbatore, India. *Int Res J Med Sci* 2014;2(10):1–6. <https://doi.org/10.5455/2320-6012.ijrms20140511>.
- [9] Mengesha RE, Kasa BG-S, Saravanan M, Berhe DF, Wasihun AG. Aerobic bacteria in post surgical wound infections and pattern of their antimicrobial susceptibility in Ayder Teaching and Referral Hospital, Mekelle, Ethiopia. *BMC Res Notes* 2014;7(575):4–9.
- [10] Garey KW, Dao T, Chen H, Amrutkar P, Kumar N, Reiter M, et al. Timing of vancomycin prophylaxis for cardiac surgery patients and the risk of surgical site infections. *J Antimicrobial Chemother* 2006;58:645–50. <https://doi.org/10.1093/jac/dkl279>.

- [11] Wassef MA, Hussein A, Abdul Rahman EM, El-Sherif RH. A prospective surveillance of surgical site infections: study for efficacy of preoperative antibiotic prophylaxis. *Afr J Microbiol Res* 2008;6(12):3072–8.
- [12] Elbur AI, Sayed A, Elsayed A, Abdel-rahman ME. Prevalence and predictors of wound infection in elective clean and clean/contaminated surgery in Khartoum Teaching Hospital, Sudan. *Int J Infect Control* 2012;8(4):1–10. <https://doi.org/10.3396/ijic.v8i4.036.12>.
- [13] Lakshmidivi N, Suchitra Joyce B. Surgical site infections: assessing risk factors, outcomes and antimicrobial sensitivity patterns. *Afr J Microbiol Res* 2009;3(4):175–9.
- [14] Astagneau P, Miliani K. Non-compliance with recommendations for the practice of antibiotic prophylaxis and risk of surgical site infection: results of a multilevel analysis from the INCISO Surveillance Network. *J Antimicrobial Chemother* 2009;64:1307–15. <https://doi.org/10.1093/jac/dkp367>.
- [15] Legesse Laloto Tamrat, Hiko Gameda Desta, Hussen Abdella Sadikalmahdi. Incidence and predictors of surgical site infection in Ethiopia: prospective cohort. *BMC Infect Dis* 2017;17:119.
- [16] Martin Emily T, Kaye Keith S, Knott Caitlin, Nguyen Huong, Santarossa Maressa, Evans Richard, et al. Diabetes and risk of surgical site infection: a systematic review and meta-analysis. *Infect Control Hosp Epidemiol* 2016 Jan;37(1):88–99.
- [17] Spagnolo AM, Ottria G, Amicizia D, Perdelli F, Cristina ML. Operating theatre quality and prevention of surgical site infections. *J Prev Med Hyg* 2013 Sep;54(3):131–7.
- [18] Mujagic E, Zwimpfer T, Marti WR, Zwahlen M, Hoffmann H, Kindler C, et al. Evaluating the optimal timing of surgical antimicrobial prophylaxis: study protocol for a randomized controlled trial. *Trials* 2014;15:188. <https://doi.org/10.1186/1745-6215-15-188>. Published online 2014 May 24.