



Gastroscope guidewire volvulus tube decompression

S. Panteleimonitis^{1,2} · G. Nash¹

Received: 3 December 2018 / Accepted: 21 January 2019 / Published online: 4 February 2019
© Springer Nature Switzerland AG 2019

Trick of the trade

Acute sigmoid volvulus is a surgical emergency [1]. Any delay in treatment increases the risk of bowel ischaemia, perforation, faecal peritonitis, and death. The majority of patients can be successfully treated with non-operative decompression, but elective definitive surgery has been recommended as most volvuli recur [2]. We describe a new technique for treating a sigmoid volvulus.

Traditionally with the patient in the left lateral position, decompression may be possible by passing a flatus tube alongside a flexible sigmoidoscope [3]. Usually there is a rush of liquid faeces and flatus with relief of the obstruction. The tube may be left in place for at least 24 h to maintain decompression, prevent recurrence and give time for the bowel wall to recover. However, it is uncomfortable to have both tubes in the anus and the flatus tube often coils in the rectum. Alternatively, by withdrawing the sigmoidoscope (once it has achieved deflation of the sigmoid) and placing the flatus tube afterwards risks it not reaching the sigmoid, which may falsely reassure us whilst the tube is in the rectum. Converting an acute to a semi-elective operation is the aim of deflation. The flatus tube may be left until a definitive plan, which may include surgery [4] or the reduced mortality of fixation of the colon endoscopically (PEC) [5] is agreed.

We describe a novel alternative that has been used successfully to place a decompressing flatus tube with less pain and greater accuracy. In a patient not needing sedation it is possible to place a paediatric gastroscope (Fig. 1; from Olympus, diameter 5.9 mm) through the lumen of a polyurethane size 24 flatus tube and passing this single combined tube (Fig. 2) anally allows the accurate identification and placement of the flatus tube ensuring decompression as

soon as the gastroscope is withdrawn to allow gas to pass. This technique has been used at our centre with success. The



Fig. 1 Paediatric gastroscope passed through the lumen of a flatus tube



Fig. 2 The end of the gastroscope is fitted just outside the end of the flatus tube during insertion

✉ S. Panteleimonitis
sofoklis_p@hotmail.com

¹ Poole Hospital NHS Trust, Poole, UK

² School of Health Sciences and Social Work, University of Portsmouth, James Watson West, Portsmouth, UK

flatus tube may be kept in for some days by sticking it to the patient's buttock with tape to avoid dislodgement, whilst definitive treatment is considered.

Funding This study received no funding.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. This type of study does not require formal ethical approval.

Informed consent All patients having this procedure sign an informed consent form before having the procedure. No patient data or characteristics were used for this study.

References

1. Tan K-K, Chong C-S, Sim R (2010) Management of acute sigmoid volvulus: an institution's experience over 9 years. *World J Surg* 34:1943–1948. <https://doi.org/10.1007/s00268-010-0563-8>
2. Raveenthiran V, Madiba TE, Atamanalp SS, De U (2010) Volvulus of the sigmoid colon. *Colorect Dis* 12:e1–e17. <https://doi.org/10.1111/j.1463-1318.2010.02262.x>
3. Safioleas M, Chatziconstantinou C, Felekouras E et al (2007) Clinical considerations and therapeutic strategy for sigmoid volvulus in the elderly: a study of 33 cases. *World J Gastroenterol* 13:921–924
4. Suleyman O, Kessaf AA, Ayhan KM (2012) Sigmoid volvulus: long-term surgical outcomes and review of the literature. *S Afr J Surg* 50:9–15
5. Frank L, Moran A, Beaton C (2016) Use of percutaneous endoscopic colostomy (PEC) to treat sigmoid volvulus: a systematic review. *Endosc Int Open* 4:E737–E741. <https://doi.org/10.1055/s-0042-106957>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.