

References

- [1] Manouchehrifar M, Lakestani M, Kashani P, Safari S. Sonographic diameter of optic nerve sheath in differentiation of ischemic and hemorrhagic strokes; a diagnostic accuracy study. *Am J Emerg Med* 2018;36(11):2064–7.
- [2] Rosa N, Lanza M, Borrelli M, De Bernardo M, Palladino A, Di Gregorio MG, et al. Low intraocular pressure resulting from ciliary body detachment in patients with myotonic dystrophy. *Ophthalmology* 2011;118(2):260–4.
- [3] De Bernardo M, Iaccarino G, Russo V, Rosa N. Echographic evaluation of a subconjunctival cystic lesion. *Case Rep Ophthalmol Med* 2017;2017(5401850).
- [4] Rosa N, De Bernardo M. Measurement of the optic nerve in a resource-limited setting. *J Neur Rural Pract* 2017;8(2):310–1.
- [5] De Bernardo M, Rosa N. Clarification on using ultrasonography to detect intracranial pressure. *JAMA Ophthalmol* 2017;135(9):1004–5.
- [6] De Bernardo M, Marotta G, Rosa N. Sonography of the optic nerve sheath diameter. *J Ultrasound Med* 2018;37(7):1845.
- [7] Iaconetta G, De Bernardo M, Rosa N. Coronal axis measurements of the optic nerve sheath diameter. *J Ultrasound Med* 2017;36(5):1073.
- [8] Rosa N, De Bernardo M. Ultrasound assessment of optic nerve sheath diameter in healthy volunteers. *J Crit Care* 2017;40:279.
- [9] De Bernardo M, Rosa N. Measuring optic nerve sheath diameter as a proxy for intracranial pressure. *JAMA Ophthalmol* 2018;136(11):1309–10.
- [10] Ossoinig KC. Standardized echography of the optic nerve. In: Till P, editor. *Documenta ophthalmologica proceedings series. Ophthalmic echography* 13Dordrecht: Springer Netherlands; 1990. p. 3–99.
- [11] De Bernardo M, Rosa N. Optic nerve sheath diameter measurement in patients with idiopathic normal-pressure hydrocephalus. *Eur J Neurol* 2018;25(2):e24.
- [12] De Bernardo M, Rosa N. Comment on 'invasive and noninvasive means of measuring intracranial pressure: a review'. *Physiol Meas* 2018;39(5):058001.
- [13] De Bernardo M, Rosa N. Comment on "optic nerve sheath diameter ultrasound evaluation in intensive care unit: possible role and clinical aspects in neurological critical patients' daily monitoring". *Biomed Res Int* 2018;2018(6154357).
- [14] Tenuta M, De Bernardo M, Rosa N. Comments on "neuromuscular ultrasonography of cranial nerves". *J Clin Neurol* 2017;13(2):212–3.
- [15] De Bernardo M, Rosa N. Transorbital sonography to evaluate optic nerve in hypertensive encephalopathy. *J Stroke Cerebrovasc Dis* 2018;27(4):1124.
- [16] De Bernardo M, Rosa N. Transbulbar B-mode sonography in multiple sclerosis: clinical and biological relevance. *Ultrasound Med Biol* 2018;44(2):508.

Carotid intima-media thickening predicts negative stress test in chest pain patients in an emergency department observation unit



Chest pain is a common presentation in the Emergency Department (ED) [1]. While most patients with chest pain are ultimately determined to have non-cardiac pain, a proportion of patients will have cardiogenic chest pain with no acute ECG changes and negative initial cardiac biomarkers.

The American Heart Association (AHA) recommends that patients in the ED with non-ultra low risk chest pain and the possibility for ACS be evaluated with serial ECGs, cardiac troponins and stress testing before discharge or within 72 h after discharge [2]. Thus, patients are frequently admitted to Observation Units (OU) for evaluation. The goal of stress testing is to evaluate for provoked ischemia requiring coronary intervention.

Ultrasound of Carotid Intima-Media Thickness (CIMT) has been proposed as a tool to non-invasively identify patients with increased atherosclerotic burden [3–6]. Studies have shown that CIMT is associated with cardiovascular disease [7–11]. The value of CIMT in the ED evaluation of chest pain is unknown. The goal of this study is to evaluate if CIMT in an ED OU population can be used as a tool to predict which patients will have a positive stress test.

This is an IRB approved, prospective, convenience sample of patients admitted to our ED OU with chest pain, dyspnea or coronary equivalents who received a stress test during their observation stay. The study was performed in a large, urban, academic ED with an annual volume of over 100,000 patients per year, an emergency medicine residency and emergency ultrasound fellowship.

CIMT scanning was performed with a Philips Sparq (Bothell WA) ultrasound machine using the L12–4 MHz linear transducer. The carotid artery was scanned at the level of the carotid bulb (Fig. 1). A registered vascular sonographer and ultrasound fellow performed scanning and data collection.

Three CIMT measurements were taken on each side 1 cm proximal to the carotid bulb (Fig. 2). The averages for each side were compared to a known standard for age and ethnicity [3]. The composite average of both sides was compared with the composite average of same reference standards. CIMT greater than the 75th percentile for age, gender and ethnicity was considered positive. Data was considered for 2 scenarios: average CIMT on *either* side or composite average CIMT of *both* sides. The reference standard differentiated patient ethnicity as black or white. For this study we used the categorization of white for all non-African ethnicities. Stress test results, interpreted by attending cardiologists, were considered the comparative standard. Results were categorized as positive or negative with indeterminate stress tests categorized as negative. Data was analyzed using Stata v 14.0 (College Station, TX).

The final analysis included 57 patients. Demographic and stress test information is shown in Table 1. Five patients had a positive stress test (8.8%).

When consideration was made for average CIMT of *either* side, 30 (52.6%) patients had a positive CIMT (Table 2). The sensitivity, specificity and negative predictive value (NPV) of CIMT for positive stress was 80% (0.95 CI 28.4–99.5%), 50% (0.95 CI 35.8–64.2%) and 96.3% (0.95 CI 81–99.9%) respectively with an odds ratio (OR) of 4 (0.95 CI 0.4–38.2 $p = 0.229$) using a univariate logistic regression analysis.

When consideration was made for average CIMT of *both* sides, 20 (35.1%) patients had a positive CIMT (Table 3). The sensitivity, specificity and NPV of CIMT for positive stress was 40% (0.95 CI 5.27–85.3%), 65.4% (0.95 CI 50.9–78%) and 91.9% (0.95 CI 78.1–98.3%) respectively with an OR of 1.3 (0.95 CI 0.2–8.2, $p = 0.81$) using a univariate logistic regression analysis.

Multiple studies have demonstrated a correlation between CIMT and increasing risk of cardiovascular disease. Chambless et al. [7] demonstrated a correlation between CIMT measurement > 1 mm and increased risk for coronary heart disease. O'Leary et al. [8] showed the risk of myocardial infarction or stroke increased with increasing CIMT. Coskun et al. [11] demonstrated that patients with CAD on angiography had a significantly thicker CIMT (1.48 ± 0.2 mm vs 0.78 ± 0.21 mm).

To our knowledge, this is the first study to evaluate CIMT in the ED evaluation of patients with chest pain. In our study we demonstrated a trend toward positive stress test with a positive CIMT. We showed that an abnormal unilateral average CIMT demonstrated a modest sensitivity for positive stress (80%) but high NPV (96.3%) and odds ratio 4 suggesting that patients with normal bilateral CIMT measurements are not likely to have a positive stress test. Interestingly this result less predictive for the composite average of both carotids (sensitivity 40%, NPV 91.9%, OR 1.3).

We found that a normal CIMT measurement for both carotid arteries correlates with a negative stress test with an extremely high negative predictive value. Our data suggest that given further study, CIMT measurement may be a reasonable determinate of disposition in ED patients with chest pain.




Fig. 1. Long axis of the common carotid artery at the level of the carotid bulb (*) with hyperechoic intima (arrow) and hypoechoic media (arrowhead).




Fig. 2. CIMT measurements of the common carotid artery within 1 cm of the carotid bulb.

Matthew P. Tabbut, MD, FACEP*

Jon W. Schrock, MD FACEP

Charles L. Emerman MD, FACEP, FAAEM

Robert A. Jones DO, FACEP

Department of Emergency Medicine, Division of Emergency Ultrasound,
MetroHealth Medical Center, Cleveland, OH 44109, United States of
America

Department of Emergency Medicine, Case Western Reserve
University School of Medicine, Cleveland, OH 44106, United States
of America

*Corresponding author at: 2500 MetroHealth Dr. BG-3, Cleveland, OH
44109, United States of America.

E-mail address: mtabbut@metrohealth.org.

Diane Gramer, RDMS, RVT

Department of Emergency Medicine, Division of Emergency Ultrasound,
MetroHealth Medical Center, Cleveland, OH 44109, United States of America

9 November 2018

<https://doi.org/10.1016/j.ajem.2018.12.053>

Table 1
Patient demographic and stress tests information.

Patient demographics		
	Frequency (n = 57)	Percent
Gender		
Male	20	35%
Female	37	65%
Ethnicity		
White	24	42.1%
Black	21	36.8%
Hispanic	10	17.5%
American Indian	1	1.8%
Asian	1	1.8%
Age		
Mean Age	53	
Min Age	33	
Max Age	79	
Stress test		
	Frequency (n = 57)	Percent
Exercise stress ECHO	33	57.9%
Pharmacologic stress ECHO	22	38.6%
Pharmacologic nuclear stress	2	3.5%

Table 2
2 × 2 table for positive CIMT on either side.

	Stress Pos	Stress Neg	
CIMT Pos	4	26	30
CIMT Neg	1	26	27
	5	52	57

Table 3
2 × 2 table for positive composite average CIMT.

	Stress Pos	Stress Neg	
CIMT Pos	4	16	20
CIMT Neg	1	36	37
	5	52	57

References

- [1] Foy AJ, Liu G, Davidson WR, Sciamanna C, Leslie DL. Comparative effectiveness of diagnostic testing strategies in emergency department patients with chest pain: an analysis of downstream testing, interventions and outcomes. *JAMA Intern Med* 2015;175:428–36.
- [2] Amsterdam EA, et al. AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes. *Circulation* 2014;2014(130):2354–95.
- [3] Howard G, Sharrett AR, Heiss G, Evans GW, Chambless LE, Riley WA, et al. Carotid artery intimal-medial thickness distribution in general populations as evaluated by B-mode ultrasound. ARIC investigators. *Stroke* 1993;24:1297–304.
- [4] Mookadam F, Moustafa SE, Lester SJ, Warsame T. Subclinical atherosclerosis: evolving role of carotid intima-media thickness. *Prev Cardiol* 2010;13:186–97.
- [5] Stein JH, Korcarz CE, Hurst RT, Lonn E, Kendall CB, Mohler ER, et al. Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: a consensus statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force endorsed by the Society for Vascular Medicine. *J Am Soc Echocardiogr* 2008;21:93–111.
- [6] Eleid MF, Lester SJ, Wiedenbeck TL, Patel SD, Appleton CP, Nelson MR, et al. Carotid ultrasound identifies high risk subclinical atherosclerosis in adults with low Framingham risk scores. *J Am Soc Echocardiogr* 2010;23:802–8.
- [7] Chambless LE, Heiss G, Folsom AR, Rosamond W, Szklo M, Sharrett AR, et al. Association of Coronary Heart Disease incidence with carotid arterial wall thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) study, 1987–1993. *Am J Epidemiol* 1997;146:483–94.
- [8] O'Leary DH, Polak JF, Kronmal RA, Manolio TA, Burke GL, Wolfson Jr SI. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older patients. *N Engl J Med* 1999;340:14–22.

- [9] Lorenz MW, Von Kegler S, Steinmetz H, Markus HS, Sitzer M. Carotid intima-media thickening indicates a higher vascular risk across a wide age range: prospective data from the carotid atherosclerosis progression study (CAPS). *Stroke* 2006;37:87–92.
- [10] Lorenz MW, Markus HS, Bots ML, Rosvall M, Sitzer M. Prediction of clinical cardiovascular events with carotid-intima media thickness: a systematic review and meta-analysis. *Circulation* 2007;115:459–67.
- [11] Coskun U, Jildiz A, Esen OB, Baskurt M, Cakar MA, Kilikesmez KO, et al. Relationships between carotid intima-media thickness and coronary angiographic findings: a prospective study. *Cardiovasc Ultrasound* 2009;7:59.

Consent for emergency treatment: Emergency department patient recall and understanding



Informed consent is a crucial component of patient autonomy and shared decision-making. Previous studies have shown that comprehension of a variety of types of informed consent by patients is poor [1–3]. This study was undertaken to identify patient recall and understanding of the emergency department (ED) Consent for Treatment document.

In this prospective survey study design, eligible participants included ED patients age 18 and older, who were able to communicate, not in distress, and consented to participate. After signing informed consent document per ED registration protocol, a convenience sample of patients meeting the inclusion criteria were invited to participate.

A total of 293 patients consented to participate (95% participation rate). The mean age was 52 and the majority of participants were female (58%) and White (62%). The majority of participants stated that they had signed a consent document ($N = 272$; 93%). A minority of patients read the entire document (7%) or read part of the document (11%). Most patients did not read the document (36%) or received only a verbal explanation (45%) (Table 1). Many patients did not recall anything about what they signed ($N = 107$; 39%). The most frequently recalled elements of consent included consent for treatment ($N = 144$; 52%), information regarding finances and billing ($N = 36$; 13%), and privacy rights ($N = 12$; 4%) (Table 2).

Reading the document prior to signing was associated with African American ethnicity ($p = 0.01$). Age, gender, mode of arrival, and triage level were not associated with reading the document (Table 3). Respondents who indicated they didn't know what they had consented to were significantly older (median 56 years) than respondents who remembered something from the consent form (median 47; $p = 0.01$). A

Table 1
Did you read the document prior to signing?

Yes, I read the entire document	20 (7%)
Yes, I read part of the document	31 (11%)
No, I did not read the document	101 (36%)
No, I did not read the document but I received a verbal explanation	126 (45%)

Table 2
What did you consent to?^a

Don't know	107 (39%)
Treatment	144 (52%)
Attending physician	0 (0%)
Privacy/HIPAA	12 (4%)
Photography	0 (0%)
Finances, billing	36 (13%)
Personal property	0 (0%)
Patient rights	9 (3%)

^a More than 1 response possible. Percentages are calculated based on $n = 276$.