




ELSEVIER

Contents lists available at ScienceDirect

# Photodiagnosis and Photodynamic Therapy

journal homepage: [www.elsevier.com/locate/pdpdt](http://www.elsevier.com/locate/pdpdt)

Letter to the Editor

## The current status of phototherapy in multimodal anti-cancer nanomedicines



Dear Editor,

We have read with great interest, the recently published article by Vivek et al. (2018), describing the authors' research on the therapeutic applications of a multifunctional nanomedicine [1]. This paper discusses the combined efficacy of photodynamic therapy (PDT), photothermal therapy (PTT) and carboplatin (CBP) chemotherapy, using CBP and indocyanine green (ICG) loaded nanoparticles (NPs), functionalised with polyethylene glycol and folic acid. Significant uptake and retention of NPs *in vitro*, was observed in the folate receptor-overexpressing breast cancer cell line, MCF7. The intracellular acidic pH environment of cancer cells induced the release of therapeutic agents into the cytoplasm. Upon near-infrared light irradiation, the increase in intracellular temperature and simultaneous increasing production of reactive oxygen species, was attributed to the photothermal conversion and photodynamic properties of ICG respectively [2]. The dual synergistic effect of PDT and PTT resulted in an apoptotic cell death. Furthermore, significant regression in tumour growth with no noticeable adverse effects, was observed in animal models treated with NPs and near-infrared light irradiation.

This study is a good example of the rapid growing interest in multifunctional anti-cancer nanomedicines, combining PDT, PTT and conventional chemotherapy. We recently reviewed the scope of photothermal therapy-based nanomedicines, in colorectal cancers and highlight the diverse combinations of phototherapeutic, chemotherapeutic, optical imaging and cancer-targeting nanomedicines that have been investigated in the past decade [3]. These nanomedicines have shown to significantly improve the management of solid colorectal cancers, by efficiently packaging different treatment methods into a single entity.

Unlike the study by Vivek et al. (2018) which demonstrated both PTT and PDT through ICG [1]. Seo et al. (2014) synthesised and evaluated PTT and PDT using NPs comprised of gold nanorods (to induce PTT) and methylene blue (to induce PDT) [4]. The authors successfully show synergy between PDT and PTT, in comparison to PDT or PTT alone. The aforementioned studies, demonstrate the flexibility in customising PTT and PDT inducing NPs, using different types of photosensitising agents.

Thus far, many novel multimodal PDT and/or PTT nanomedicines have been the subject of 'proof-of-principle' pre-clinical studies, showing potential clinical applications. Currently, PDT and PTT (through thermal ablation) are utilised in clinics to treat cancers and both techniques have independently proven to be successful as stand-alone methods and in combination with more conventional means of

treatment [5,6]. However, due to the apparent lack of controlled and randomised clinical trials investigating phototherapy nanomedicines, it is difficult to design appropriate clinical protocols and identify key candidates for further evaluation and clinical translation. Nevertheless, the remarkable progression in this field with a focus on precision medicine means that the clinical application of phototherapy nanomedicines is feasible in the near future.

### Funding sources

The National Institute for Health Research (NIHR) infrastructure at Leeds. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

### Conflicts of interest

None.

### References

- [1] R. Vivek, N. Varukattu, R. Chandrababu, S. Alok, P. Thondhi, V. Alagarsamy, S. Kannan, Multifunctional nanoparticles for trimodal photodynamic therapy-mediated photothermal and chemotherapeutic effects, *Photodiagn. Photodyn. Ther.* 23 (2018) 244–253, <https://doi.org/10.1016/j.pdpdt.2018.06.025>.
- [2] J.T. Alander, I. Kaartinen, A. Laakso, T. Pättilä, T. Spillmann, V.V. Tuchin, M. Venermo, P. Välisuo, A review of indocyanine green fluorescent imaging in surgery, *Int. J. Biomed. Imaging* 2012 (2012) 1–26, <https://doi.org/10.1155/2012/940585>.
- [3] M.I. Khot, H. Andrew, H.S. Svavarsdottir, G. Armstrong, A.J. Quyn, D.G. Jayne, A review on the scope of photothermal therapy-based nanomedicines in preclinical models of colorectal cancer, *Clin. Colorectal Cancer* (2019), <https://doi.org/10.1016/j.clcc.2019.02.001>.
- [4] S.H. Seo, B.M. Kim, A. Joe, H.W. Han, X. Chen, Z. Cheng, E.S. Jang, NIR-light-induced surface-enhanced Raman scattering for detection and photothermal/photodynamic therapy of cancer cells using methylene blue-embedded gold nanorod@SiO<sub>2</sub> nanocomposites, *Biomaterials* 35 (2014) 3309–3318, <https://doi.org/10.1016/j.biomaterials.2013.12.066>.
- [5] K.F. Chu, D.E. Dupuy, Thermal ablation of tumours: biological mechanisms and advances in therapy, *Nat. Rev. Cancer* 14 (2014) 199–208, <https://doi.org/10.1038/nrc3672>.
- [6] K. Moghissi, K. Dixon, S. Gibbins, A surgical view of photodynamic therapy in oncology: a review, *Surg. J. (New York, N.Y.)* 1 (2015) e1–e15, <https://doi.org/10.1055/s-0035-1565246>.

M. Ibrahim Khot\*, Gemma Armstrong, Helen Andrew,  
Candice L. Downey, David G. Jayne  
*School of Medicine, St James's University Hospital, University of Leeds,  
Leeds, UK*  
E-mail address: [M.I.Khot@leeds.ac.uk](mailto:M.I.Khot@leeds.ac.uk) (M.I. Khot).

\* Corresponding author at: Section of Translational Anaesthesia and Surgical Sciences, Level 9, Wellcome Trust Brenner Building, St James's University Hospital, Beckett St, Leeds, LS9 7TF, UK.

<https://doi.org/10.1016/j.pdpdt.2019.04.024>

Received 25 April 2019

Available online 28 April 2019

1572-1000/© 2019 Elsevier B.V. All rights reserved.