

## Original Article

## The international St-aging system as a prognostic marker in general senior population: Findings from the EPIDOS cohort study

Marie Pouzoullic<sup>a</sup>, Anne-Marie Schott<sup>b,g,h</sup>, Dolores Sánchez-Rodríguez<sup>c,i,j</sup>, Regis Bataille<sup>d,e</sup>, Cedric Annweiler<sup>a,f,\*</sup>, On behalf of the SOCOS group

<sup>a</sup> Department of Geriatric Medicine, Angers University Hospital, Angers University Memory Clinic, Research Center on Autonomy and Longevity, UPRES EA 4638, University of Angers, Angers, France

<sup>b</sup> Pôle Information Médicale Evaluation Recherche, Hospices Civils de Lyon, F-69003 Lyon, France

<sup>c</sup> Geriatrics Department, Parc de Salut Mar (Centre Fórum-Hospital del Mar), Barcelona, Spain

<sup>d</sup> Institut de Cancérologie de l'Ouest, Angers, France

<sup>e</sup> INSERM UMR 892, Institut de Recherche Thérapeutique de l'Université de Nantes, France

<sup>f</sup> Roberts Research Institute, Schulich School of Medicine and Dentistry, the University of Western Ontario, London, Ontario, Canada

<sup>g</sup> INSERM U1033, F-69000 Lyon, France

<sup>h</sup> Université Lyon 1, F-69000 Lyon, France

<sup>i</sup> Rehabilitation Research Group, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM), Barcelona, Spain

<sup>j</sup> School of Medicine, Universitat Autònoma de Barcelona, Spain

## ARTICLE INFO

## Keywords:

Albumin  
Beta-2 microglobulin  
Multiple myeloma international staging system  
Prognosis  
Survival  
Older adults

## ABSTRACT

The International Staging System (ISS) -calculated from serum albumin and beta-2 microglobulin ( $\beta 2m$ ) - is an established prognostic marker in multiple myeloma (MM), which has also been suggested to account for survival among general senior population. Our objective was to examine long-term survival of older women free of MM according to baseline ISS. The study included 230 community-dwelling healthy older women without known MM from the EPIDOS cohort (mean  $\pm$  SD, 80.4  $\pm$  3.4 years). Serum albumin and  $\beta 2m$  were measured at baseline, and used to calculate the ISS *a posteriori*. Abnormal ISS was defined as ISS = 2 or ISS = 3, although ISS = 1 was considered normal. The vital status was sought after a mean follow-up of 17.6  $\pm$  0.2 years (range, 16.8–18.3). Age, body mass index, mean arterial pressure, diabetes mellitus, hypertension, coronary heart disease, stroke, use corticosteroids, number of drugs daily taken, smoking, physical activity, fall history, bone mineral density, and creatinine clearance were used as potential confounders. All participants died during the 17-year follow-up. Compared to women with normal ISS, those with abnormal ISS ( $n = 24$ ) had shorter survival time (4.9  $\pm$  4.3 versus 8.7  $\pm$  5.2 years,  $P = .001$ ) and died earlier (85.6  $\pm$  4.8 versus 89.1  $\pm$  5.6 years old,  $P = .003$ ). Survival time after blood test correlated with both serum albumin ( $r = 0.16$ ,  $P = .015$ ) and  $\beta 2m$  ( $r = -0.27$ ,  $P < .001$ ). Cox regression revealed that abnormal ISS was associated with mortality (adjusted HR = 3.03,  $P < .001$ ). Kaplan-Meier distributions showed that participants with abnormal ISS died earlier than those with normal ISS (log-rank  $P < .001$ ). In conclusion, community-dwelling older women with abnormal ISS had shorter survival time than the others, suggesting that ISS could be considered as a universal prognostic “aging system” rather than a specific MM “staging system”.

## 1. Introduction

The first step of any efficient geriatric care relies on the early identification of older adults most vulnerable to stress, who are at greater risk of failure of usual coping mechanisms, with consequent lower survival chance. This screening is yet limited by the difficulty of implementation into clinical practice due to the time consumption of standard validated tools [1,2]. A simple and easily accessible blood test

is thus desirable to improve the feasibility of the screening for older adults with limited life expectancy in daily clinical practice.

There is ample evidence that serum albumin (SA) and beta-2 microglobulin ( $\beta 2m$ ) are both biological prognostic biomarkers of survival [3–5]. The combination of SA and  $\beta 2m$  constitutes a potent staging system in multiple myeloma (MM) [6–10], with a higher prognostic value compared to all other markers including gene expression profiles [9]. In 2005, this SA/ $\beta 2m$  combination was internationally recognized

\* Corresponding author at: Department of Geriatrics, Angers University Hospital, 49933 Angers cedex 9, France.  
E-mail address: [Cedric.Annweiler@chu-angers.fr](mailto:Cedric.Annweiler@chu-angers.fr) (C. Annweiler).

<https://doi.org/10.1016/j.ejim.2019.07.031>

Received 10 April 2019; Received in revised form 25 July 2019; Accepted 30 July 2019

Available online 03 August 2019

0953-6205/ © 2019 European Federation of Internal Medicine. Published by Elsevier B.V. All rights reserved.

**Table 1**  
Characteristics and comparison of participants separated into two groups based on the International Staging System\* (n = 230).

	Whole sample (n = 230)	International staging system		P-value <sup>b</sup>
		Normal <sup>a</sup>	Abnormal <sup>a</sup>	
		(n = 206)	(n = 24)	
Age, years	80.4 ± 3.4	80.3 ± 3.5	80.6 ± 2.7	0.711
Body mass index, kg/m <sup>2</sup>	25.5 ± 4.4	25.5 ± 4.5	25.7 ± 3.6	0.848
Mean arterial pressure, mmHg	106.4 ± 12.4	106.8 ± 12.4	102.9 ± 12.3	0.144
History of diabetes mellitus, n (%)	15 (6.5)	15 (7.3)	0 (0.0)	0.170
History of arterial hypertension, n (%)	107 (46.5)	94 (46.1)	13 (54.2)	0.453
History of coronary heart disease, n (%)	48 (20.9)	39 (19.0)	9 (37.5)	<b>0.035</b>
History of stroke, n (%)	12 (5.2)	9 (4.4)	3 (12.5)	0.092
Use corticosteroids, n (%)	8 (3.5)	8 (3.9)	0 (0.0)	0.325
Number of drugs taken per day	5.1 ± 3.0	5.0 ± 2.9	6.5 ± 3.9	<b>0.022</b>
Smoking, n (%)	28 (12.2)	26 (12.7)	2 (8.3)	0.538
Regular physical activity <sup>c</sup> , n (%)	116 (50.4)	105 (51.2)	11 (45.8)	0.618
History of falls in preceding 6 months, n (%)	57 (24.8)	50 (24.4)	7 (29.2)	0.609
Whole-body mineral density, g/cm <sup>2</sup>	0.98 ± 0.09	0.98 ± 0.09	0.97 ± 0.08	0.560
Serum albumin concentration, g/L	45.1 ± 4.5	45.5 ± 4.3	42.2 ± 5.2	<b>0.001</b>
Serum beta-2 microglobulin concentration, mg/L	2.49 ± 0.96	2.26 ± 0.50	4.46 ± 1.55	< <b>0.001</b>
Estimated glomerular filtration rate <sup>d</sup> , mL/min	51.3 ± 15.9	52.3 ± 15.8	41.3 ± 14.1	<b>0.006</b>
Survival time after blood test, years	8.3 ± 5.2	8.7 ± 5.2	4.9 ± 4.3	<b>0.001</b>
Age of death, years	88.8 ± 5.6	89.1 ± 5.6	85.6 ± 4.8	<b>0.003</b>

Data presented as mean ± standard deviation where applicable.

P-value significant (i.e., < 0.05) indicated in bold.

<sup>a</sup> : normal ISS defined as ISS = 1; abnormal ISS defined as ISS = 2 or ISS = 3.

<sup>b</sup> : comparisons between participants with normal ISS and participants with abnormal ISS based on independent samples t-test or Chi-square test, as appropriate.

<sup>c</sup> : at least one hour recreational physical activity (walking, gymnastics, cycling, swimming or gardening) per week for the past month or more.

<sup>d</sup> : calculated from the Cockcroft formula.

as the MM international staging system (ISS) [10]. However and importantly, it was proposed that the ISS may actually take into account more the age-related comorbidity burden of MM older patients than the intrinsic malignancy of the MM clone [11]. Decreased SA and increased  $\beta$ 2m reflect declines across multiple physiologic systems and may account for the comorbidity burden among older adults [11]. This suggests that the ISS should not be considered as a specific MM staging system, but rather as an “aging system”, i.e. a universal prognostic biomarker in older adults.

We hypothesized that the ISS could predict survival time in general senior population. We had the opportunity to examine the association of ISS with survival time in a sample of healthy women aged  $\geq 75$  years from the large representative community-based EPIDémiologie de l’OStéoporose (EPIDOS) cohort. The objective of this historical, longitudinal cohort study was to examine long-term survival of older women according to baseline ISS.

## 2. Materials and methods

We studied 230 participants without known MM, a subset recruited from 1992 to 1994 in the EPIDOS study, a French large observational prospective multicenter cohort study designed to evaluate risk factors for hip fractures among healthy community-dwelling women aged 75 years and older [12]. The present analysis considered data from all women who had: i) full data on serum measures; and ii) full data on mortality up to October 1, 2010.

All study participants had a blood test at the time of inclusion. Sera were stored at  $-100\text{ }^{\circ}\text{C}$  until analyses were performed in a sample of women drawn using a random number table that generated in an unpredictable, haphazard a sequence of number corresponding to the number of participants included in the study [13]. SA (g/L) and  $\beta$ 2m (mg/L) concentrations were determined using automated standard laboratory methods at the University Hospital of Lyon, France. The ISS was scored in 3 stages as previously described [10]: stage 1 defined as  $\beta$ 2m < 3.5 mg/L with SA  $\geq 3.5$  g/dL; stage 2 defined as  $\beta$ 2m between 3.5 and 5.5 mg/L with any SA level, or  $\beta$ 2m < 3.5 mg/L with SA <

3.5 g/L; stage 3 defined as  $\beta$ 2m  $\geq 5.5$  mg/L. Abnormal ISS was defined as ISS = 2 or ISS = 3, and ISS = 1 was considered normal.


Mortality data were collected either via telephone or through a search of the French national death registry CépiDC [Centre d’épidémiologie sur les causes médicales de décès] up to October 1, 2010 [14]. The survival time was calculated between the date of recruitment in the EPIDOS study and the date of death or end of the follow-up.

All participants had a full examination consisting of clinical examination and structured questionnaires on socio-demographic data, health status, and physical functioning. Age, body mass index, mean arterial pressure, diabetes mellitus, hypertension, coronary heart disease, history of stroke, current use of corticosteroids, number of drugs taken per day, smoking, regular practice of physical activity, history of falls in preceding 6 months, whole-body bone mineral density, and estimated glomerular filtration rate (estimated from the Cockcroft-Gault formula:  $[(140 - \text{age}_{\text{years}}) \times \text{weight}] / \text{creatinine}_{\text{mol/L}} \times 1.04$ ) were measured and used as potential confounders.

The study was conducted in accordance with the ethical standards set by Helsinki declaration (1983). The EPIDOS study was approved by the local ethics committee. All participants gave written informed consents.

### 2.1. Statistical analysis

The participants' baseline characteristics were summarized using means and standard deviations or frequencies and percentages, as appropriate. Participants were separated into 2 groups according to ISS at baseline (i.e., normal ISS versus abnormal ISS). Firstly, between-group comparisons were performed using the independent samples t-test or Chi-square test, as appropriate. Secondly, three consecutive Cox regression models were used to examine the associations of mortality (dependent variable) with SA,  $\beta$ 2m and abnormal ISS adjusted on baseline characteristics (independent variables). The models produce a survival function that provides the probability of death at a given time for the characteristics supplied for the independent variables. Third, the



**Fig. 1.** Relationships between survival time and serum concentrations of albumin and beta-2 microglobulin.


The tick lines are the best-fit linear regression lines, and the thin lines at the top and bottom are the limits of the 95% confidence intervals.

- Survival time on serum albumin.
- Survival time on serum beta-2 microglobulin.
- Survival time on serum albumin.
- - - Survival time on serum beta-2 microglobulin.

elapsed time to death was studied by survival curves computed according to the Kaplan-Meier method and compared by the log-rank test. *P*-values < .05 were considered statistically significant. All statistics were performed using SPSS (version 19.0; SPSS, Inc., Chicago, IL).

**3. Results**

All 230 participants (mean age at baseline, 80.4 ± 3.4 years) died during the follow-up (mean, 17.6 ± 0.2 years; range, 16.8–18.3 years). As shown in Table 1, women with abnormal ISS (*n* = 24) had shorter survival time than those with normal ISS (4.9 ± 4.3 versus 8.7 ± 5.2 years; *P* = .001) and died earlier (mean age of death, 85.6 ± 4.8 versus 89.1 ± 5.6 years; *P* = .003). Survival time after blood test correlated positively with SA concentration (*r* = 0.16, *P* = .015), and negatively with serum β2m concentration (*r* = -0.27, *P* < .001) (Fig. 1). Cox regression models revealed an inverse association between SA and mortality (fully adjusted hazard ratio (HR) = 0.96, *P* = .036) and a positive association between β2m and mortality (fully adjusted HR = 1.30, *P* = .004) (Table 2). Abnormal ISS was also associated positively with mortality (HR = 2.29, *P* < .001), even after adjustment for potential confounders (fully adjusted



**Fig. 2.** Kaplan-Meier estimates of the cumulative probability of participants' survival according to the International Staging System (ISS) (*n* = 230).

HR = 3.03, *P* < .001) (Table 2). Consistently, Kaplan-Meier distributions showed in Fig. 2 that participants with abnormal ISS had shorter survival time than those with normal ISS (log-rank *P* < .001).

**4. Discussion**

Our results show that abnormal ISS predicted, irrespective of studied potential confounders, shorter long-term survival time in community-dwelling healthy older women.

To the best of our knowledge, we provide here the first evidence of the prognostic value of the ISS in general senior population. This finding is consistent with previous studies reporting that SA and β2m are both biological prognostic factors of survival [3–5]. Numerous epidemiological studies have showed an association between lower SA concentrations and increased morbid-mortality [3,4,15,16]. The estimated increase in the odds of death ranges from 24% to 56% for each 2.5 g/L decrement in SA concentration [3]. This association predicts overall and cause-specific mortality including cardiovascular mortality. Several biological mechanisms have been suggested to explain the association of lower SA concentrations with increased mortality [3]. First, a protective effect of SA is likely based on its potential nitrovasodilator effect, which could modulate cardiovascular diseases processes. Another protective mechanism may be related to a significant antioxidant effect of SA in blood and extracellular fluids. SA enhances the removal of reactive oxygen species involved in the pathogenesis of numerous diseases such as atherosclerosis. SA may also be directly beneficial by increasing the ability of serum to bind potential toxins. Finally, SA is an

**Table 2**

Hazard ratios of mortality (dependent variable) according to serum albumin concentration, serum beta-2 microglobulin concentration, and abnormal International Staging System at baseline (independent variables)<sup>a</sup> (*n* = 230).

	Mortality								
	Model 1			Model 2			Model 3		
	HR	95%CI	P-value	HR	95%CI	P-value	HR	95%CI	P-value
Serum albumin concentration	0.97	[0.94–1.00]	0.063	0.97	[0.93–1.00]	0.069	0.96	[0.92–0.99]	<b>0.036</b>
Serum beta-2 microglobulin concentration	1.30	[1.15–1.48]	< <b>0.001</b>	1.35	[1.17–1.55]	< <b>0.001</b>	1.30	[1.09–1.55]	<b>0.004</b>
Abnormal international staging system	2.29	[1.49–3.53]	< <b>0.001</b>	2.87	[1.79–4.59]	< <b>0.001</b>	3.03	[1.72–5.33]	< <b>0.001</b>

HR: hazard ratio; CI: Confidence interval; Model 1: unadjusted Cox model; Model 2: Model 1 + adjustment for age, body mass index, mean arterial pressure, diabetes mellitus, hypertension, coronary heart disease, stroke, use corticosteroids, number of drugs per day, smoking, regular physical activity, history of falls in preceding 6 months, whole-body bone mineral density; Model 3: Model 2 + adjustment for estimated glomerular filtration rate.

HR significant (i.e. *P* < .05) indicated in bold.

<sup>a</sup> : separate regression analyses were conducted for each model.

inhibitor of human endothelial apoptosis and lower SA concentrations may result in a general increased vascular permeability leading to tissue damage [3,4,16]. Like SA, higher  $\beta$ 2m has been associated with increased morbi-mortality while aging, independent of any history of MM [5,17–21]. Firstly, greater levels of serum  $\beta$ 2m have been reported in healthy older adults compared to younger ones [18,21]. Secondly,  $\beta$ 2m is an established biomarker of disease activity not only in malignancies and chronic renal dysfunction, but also in autoimmune affections and infections [17,19–21]. Thus, an increase in serum  $\beta$ 2m concentration reflects pathological aging across multiple physiologic systems. In line with this, increased  $\beta$ 2m concentrations were used as a marker of frailty phenotype in geriatric inpatients [20]. Thirdly, a strong association has been reported among older adults between increased serum  $\beta$ 2m concentration and higher rate of all-cause mortality, specifically cardiovascular mortality [5]. All these findings suggest that the combination of SA with  $\beta$ 2m adequately explores the balance of inflammation proteins and constitutive proteins, and may reflect the physiological reserves available to the senior, making it an “aging system”.

Some limitations of our study should be considered. Firstly, the study cohort was restricted to relatively vigorous older women with interest in personal health issues who may be unrepresentative of older adults in general. Secondly, although we were able to control for important characteristics that could modify the association between ISS and mortality, residual potential confounders such as the presence of inflammatory systemic diseases or the causes of deaths might still be present. Thirdly, data were recorded before 2000. It is possible, though unproven, that the profiles and the risks of the participants have changed since then, as well as the implementation and enforcement of various prevention programs. The present results show thus trends that would better be confirmed by an independent survey in another cohort.

In conclusion, abnormal ISS predicted shorter survival time in community-dwelling healthy older women. Further studies with different cohorts are warranted to corroborate this finding, and assess the value of the ISS as an “aging system” and a universal marker of frailty phenotype in older adults.

### Study funding

This work was supported by the French Ministry of Health. The sponsor had no role in the design and conduct of the study, in the collection, management, analysis, and interpretation of the data, or in the preparation, review, or approval of the manuscript.

### Contributions of the authors

Annweiler has access to all the study data, takes responsibility for the accuracy of the analysis, and has authority over manuscript preparation and the decision to submit the manuscript for publication.

- Study concept and design: Annweiler.
- Acquisition of data: Schott.
- Analysis and interpretation of data: Annweiler, Pouzoulic.
- Drafting of the manuscript: Annweiler, Pouzoulic.
- Critical revision of the manuscript for important intellectual content: Schott, Sánchez-Rodríguez, Bataille.
- Statistical analysis: Annweiler.
- Obtaining funding: Schott.
- Administrative, technical, or material support: Schott.
- Supervision: Annweiler.

### Declaration of Competing Interest

All other authors state that they have no conflicts of interest with

this paper.

### Acknowledgments

Investigators of EPIDOS study. Coordinators: Breart, Dargent-Molina, Meunier, Schott, Hans, and Delmas. Principal investigators: Baudoin and Sebert (Amiens); Chapuy and Schott (Lyon); Favier and Marcelli (Montpellier); Hausherr, Menkes and Cormier (Paris); Grandjean and Ribot (Toulouse).

### References

- [1] Bergman H, Ferrucci L, Guralnik J, Hogan DB, Hummel S, Karunanathan S, et al. Frailty: an emerging research and clinical paradigm issues and controversies. *J Gerontol A Biol Sci Med Sci* 2007;62:731–7.
- [2] Rodrigues-Manas L, Feart C, Mann G, Viña J, Chatterji S, Chodzko-Zajko W, et al. Searching for an operational definition of frailty: a delphi method based consensus statement. The frailty operative definition-consensus conference project. *J Gerontol A Biol Sci Med Sci* 2013;68:62–7.
- [3] Goldwasser P, Feldman J. Association of serum albumin and mortality risk. *J Clin Epidemiol* 1997;50:693–703.
- [4] Corti MC, Guralnik JM, Salive ME, Sorokin JD. Serum albumin level and physical disability as predictors of mortality in older persons. *JAMA* 1994;272:1036–42.
- [5] Shinkai S, Chaves PH, Fujiwara Y, Watanabe S, Shibata H, Yoshida H, et al. Beta2-microglobulin for risk stratification of total mortality in the elderly population: comparison with cystatin C and C-reactive protein. *Arch Intern Med* 2008;168:200–6.
- [6] Bataille R, Durie BGM, Grenier J, Sany J. Prognostic factors and staging in multiple myeloma. *J Clin Oncol* 1986;4:80–7.
- [7] Durie BGM, Salmon SE. A clinical staging system for multiple myeloma. *Cancer* 1975;36:842–54.
- [8] Durie BGM, Stock-Novack D, Salmon SE, Finley P, Beckord J, Crowley J, et al. Prognostic value of pre-treatment serum B2 microglobulin in myeloma: a Southwest Oncology Group study. *Blood* 1990;75:823–30.
- [9] Jacobson JL, Hussein MA, Barlogie B, Durie BG, Crowley JJ, Southwest Oncology Group. A new staging system for multiple myeloma patients based on the Southwest Oncology Group (SWOG) experience. *Br J Haematol* 2003;122:441–50.
- [10] Greipp Ph R, San Miguel J, Durie BGM, Crowley JJ, Barlogie B, Bladé J, et al. International staging system for multiple myeloma. *J Clin Oncol* 2005;23:3414–20.
- [11] Bataille R, Annweiler C, Beauchet O. Multiple myeloma international staging system: “staging” or simply “aging” system?. *Clin Lymphoma Myeloma Leuk* 2013;13:635–7.
- [12] Dargent-Molina P, Favier F, Grandjean H, Baudoin C, Schott AM, Hausherr E, et al. Fall-related factors and risk of hip fracture: the EPIDOS prospective study. *Lancet* 1996;348:145–9.
- [13] Annweiler C, Schott AM, Montero-Odasso M, Berrut G, Fantino B, Herrmann FR, et al. Cross-sectional association between serum vitamin D concentration and walking speed measured at usual and fast pace among older women: the EPIDOS study. *J Bone Miner Res* 2010;25:1858–66.
- [14] Bailly S, Haesebaert J, Decullier E, Dargent-Molina P, Annweiler C, Beauchet O, et al. Mortality and profiles of community-dwelling fallers. Results from the EPIDOS cohort. *Maturitas* 2014;79:334–9.
- [15] Gom I, Fukushima H, Shiraki M, Miwa Y, Ando T, Takai K, et al. Relationship between serum albumin level and aging in community-dwelling self-supported elderly population. *J Nutr Sci Vitaminol* 2007;53:37–42.
- [16] Herrmann FR, Safran C, Levkoff SE, Minaker KL. Serum albumin level on admission as a predictor of death, length of stay, and readmission. *Arch Intern Med* 1992;152:125–30.
- [17] Bataille R, Grenier J, Sany J. Beta-2-microglobulin in myeloma: optimal use for staging, prognosis, and treatment. A prospective study of 160 patients. *Blood* 1984;63:468–76.
- [18] Vincent C, Revillard JP, Galland M, Traeger J. Serum beta2-microglobulin in hemodialyzed patients. *Nephron* 1978;21:260–8.
- [19] Erez E, Aravot D, Erman A, Sharoni E, van Oyk DJ, Raanani E, et al. Increased production of B2-microglobulin after heart transplantation. *J Heart Lung Transplant* 1998;17:538–41.
- [20] Annweiler C, Bataille R, Ferrière N, Douillet D, Fantino B, Beauchet O. Plasma beta-2 microglobulin as a marker of frailty in older adults: a pilot study. *J Gerontol A Biol Sci Med Sci* 2011;66:1077–9.
- [21] Shinkai S, Kumagai S, Fujiwara Y, Amano H, Yoshida Y, Watanabe S, et al. Predictors for the onset of functional decline among initially non-disabled older people living in a community during a 6-year follow-up. *Geriatr Gerontol Int* 2003;3:S31–9.