

The UK's pandemic influenza research portfolio: a model for future research on emerging infections



Colin R Simpson, Dan Beever, Kirsty Challen, Daniela De Angelis, Ellen Fragaszy, Steve Goodacre, Andrew Hayward, Wei Shen Lim, G James Rubin, Malcolm G Semple, Marian Knight, on behalf of the NIHR hibernated influenza studies collaborative group*

The 2009 influenza A H1N1 pandemic was responsible for considerable global morbidity and mortality. In 2009, several research studies in the UK were rapidly funded and activated for clinical and public health actions. However, some studies were too late for their results to have an early and substantial effect on clinical care, because of the time required to call for research proposals, assess, fund, and set up the projects. In recognition of these inherent delays, a portfolio of projects was funded by the National Institute for Health Research in 2012. These studies have now been set up (ie, with relevant permissions and arrangements made for data collection) and pilot tested where relevant. All studies are now on standby awaiting activation in the event of a pandemic being declared. In this Personal View, we describe the projects that were set up, the challenges of putting these projects into a maintenance-only state, and ongoing activities to maintain readiness for activation, and discuss how to plan research for a range of major incidents.

Introduction

In response to the 2009 influenza A H1N1 pandemic, UK research funders, including the National Institute for Health Research (NIHR), rapidly sponsored and activated several research studies for clinical and public health actions (table). The programme showed the ability of the NIHR to commission, fund, and deliver clinically relevant research under challenging circumstances.⁹ However, even with accelerated processes, some studies were completed too late for their results to have an immediate substantial effect on clinical care.^{10–12} This experience was shared by major research networks in other countries.¹³ By contrast, studies which had been funded separately, either before the emergence of the pandemic,¹⁴ or via the Department of Health in England, were able to provide timely reports on their results.¹⁵

In recognition of the effect of the NIHR-funded work on the 2009 influenza A H1N1, and following reflection on the inherent delays in calling for research proposals, assessing, funding, and setting up the subsequent projects, including obtaining relevant ethical and regulatory approvals, the NIHR funded a second set of studies in 2012. These projects were set up (relevant permissions put in place and arrangements made for data collection) and pilot tested where relevant. All the studies are being maintained in a state of readiness or hibernation, in that they are on standby mode in a maintenance-only state to await activation in the event of a pandemic being declared. Other organisations have subsequently recognised the need for better planning of research for a future pandemic.¹⁶

The NIHR Clinical Research Network portfolio of pandemic influenza studies

The portfolio of studies (table) cover key pathways of health care for influenza illness, including: surveillance, primary prevention (vaccination), triage (both in the community and at the hospital front door), and clinical management (therapeutics). The population-level susceptibility, severity,

and spread of pandemic influenza study (PIPS) has developed the Health Survey for England as a tool for rapid population based serological surveys of influenza infection and disease.¹⁷ During the first two outbreaks of a pandemic, the study will assess the public's susceptibility to and spread of the novel virus both geographically and by age. By combining our estimated number of infections in the population with estimates of the numbers of hospitalisations and deaths, the proportion of infections leading to these outcomes can be quantified—a key measure of pandemic severity.

The Flu Telephone Survey Template study (FluTEST) has been funded to develop rapid turn-around telephone surveys, for use in monitoring behaviour across the general population and identifying ways of improving how public health advice is communicated.¹⁸ How the public behave during any future pandemic will have a crucial role in how quickly influenza spreads in the population. Understanding whether the public are adhering to public health recommendations (eg, washing hands, being vaccinated, and using health services appropriately), and if not, why not, is essential information.^{14,18,19}

The Early estimation of pandemic influenza Antiviral and Vaccine Effectiveness (EAVE) project has created a new sentinel system linking primary care data to RT-PCR swabs, stored serology, and hospital and mortality outcomes. The EAVE sentinel system, which builds on the seasonal influenza vaccination study called Seasonal Influenza Vaccine Effectiveness,²⁰ allows rapid evaluation of any new vaccination (and antivirals). The EAVE project also provides information to help with targeting any available pandemic influenza vaccine at those who are considered to be at increased risk of serious illness or death from pandemic influenza infection—eg, those with any underlying medical conditions; those who might not have cross-reactivity from exposure to previous pandemics or vaccinations; and novel risk groups that are uniquely at risk because of a tropism exhibited by the virus.² For any new pandemic influenza vaccine, it might

Lancet Infect Dis 2019;
19: e295–300

Published Online
April 18, 2019
[http://dx.doi.org/10.1016/S1473-3099\(18\)30786-2](http://dx.doi.org/10.1016/S1473-3099(18)30786-2)

*Members listed at the end of the paper

School of Health, Faculty of Health, Victoria University of Wellington, Wellington, New Zealand (Prof C R Simpson PhD); Usher Institute, The University of Edinburgh, Edinburgh, UK (Prof C R Simpson); Clinical Trials Research Unit, School of Health and Related Research, University of Sheffield, UK (D Beever MPH, Prof S Goodacre PhD); Lancashire Teaching Hospitals National Health Service Trust, Preston, UK (K Challen PhD); Medical Research Council Biostatistics Unit, University of Cambridge, Cambridge, UK (D De Angelis PhD); Centre for Public Health Data Science, Institute of Health Informatics (E Fragaszy MSc, Prof A Hayward MD), and Institute of Epidemiology and Health Care (Prof A Hayward), University College London, London, UK; Department of Psychological Medicine, Weston Education Centre, King's College London, London, UK (G J Rubin PhD); Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK (E Fragaszy); Nottingham University Hospitals NHS Trust, Nottingham, UK (Prof W S Lim DM); Institute of Translational Medicine, University of Liverpool, UK (Prof M G Semple PhD); and National Perinatal Epidemiology Unit, University of Oxford, Oxford, UK (Prof M Knight DPhil)

Correspondence to: Prof Colin R Simpson, School of Health, Faculty of Health, Victoria University of Wellington, Wellington 6140, New Zealand colin.simpson@vuw.ac.nz

take some time until matched vaccines are available. Stockpiled pre-pandemic vaccine (eg, H5N1, H7N9) for the UK population, recommended to be of sufficient quantities for those aged 16 or younger and aged 65 or older, could be used for early testing.^{21,22}

The real-time refinement and validation of criteria and tools used in primary care (FLU-CATs) study captures the criteria that allows validation of several triage tools in the early stage of an outbreak in readiness for deployment of these tools should surge occur.²³ During pandemics, surge in health-care demand can exceed the capacity to provide normal standards of care. Health-care workers who are unfamiliar with clinical assessment and admission decision-making might be asked to fulfil gatekeeper roles. Triage tools might aid decisions in identifying people who are most likely to benefit from the higher levels of care that can only be provided in hospital. Anonymised details of patients with influenza-like illness attending a study general practice will be shared with the study team. These

details include: assessment, management, selected medical record excerpts, record hospital discharge data, and death notices.²³ The FLU-CATs study activates during winter influenza seasons at a limited number of sites (five to 20) to maintain study processes. Annual winter activation of the FLU-CATs study has allowed validation of the Department of Health Community Assessment Tools for people presenting with seasonal influenza-like illness.²³

The PANdemic INfluenza Triage in the Emergency Department (PAINTED) study addresses the issue of needing to prioritise a potentially large number of patients with limited resources in the event of a pandemic.²⁴ Clinicians will need to identify patients at higher risk of poor outcomes, ideally with minimal or no supplementary testing. This study will use a standardised clinical assessment form to collect data on adults and children presenting to UK emergency departments in the event of a pandemic. Patients will be followed up at 30 days to

	Lead applicant and institute	Purpose of study
The population-level susceptibility, severity, and spread of pandemic influenza study (PIPS) ¹	Andrew Hayward, University College London	The PIPS study developed a system to rapidly assess real-time community-level susceptibility and spread of infection and illness in the event of a pandemic. It achieved this by adding additional questions and specimen collection to the Health Survey for England, an annual, nationally representative survey that recruits participants throughout the year.
Flu Telephone Survey Template study (FluTEST) ²	James Rubin, King's College London	The preliminary work of FluTEST involved developing, testing, and publishing a broad range of survey questions that can be used with confidence in the next pandemic to identify public knowledge, attitudes, and behaviour. During a pandemic, the study team will support the UK Department of Health and Social Care in deploying these items and interpreting the results.
Early estimation of pandemic influenza Antiviral and Vaccine Effectiveness (EAVE): use of a unique community and laboratory national linked dataset ³	Colin Simpson, Victoria University of Wellington and University of Edinburgh	The EAVE project created a new sentinel system that links primary care data to RT-PCR swabs, stored serology, and hospital and mortality outcome data. The project allows for a rapid evaluation of any new vaccination (and antivirals), and provides information to help with targeting any available pandemic influenza vaccine at those who are considered to be at increased risk of serious illness or death from pandemic influenza infection.
Real-time refinement and validation of criteria and tools used in primary care to aid hospital referral decisions for patients of all ages in the event of surge during an influenza pandemic—FLU-CATs (Influenza Community Assessment Tools) ⁴	Malcolm Semple, University of Liverpool	FLU-CATs is a study that checks if decision tools can be used by general practitioners and other health-care professionals to help them to choose who can be cared for safely in the community, and who needs urgent referral to hospital. FLU-CATs runs each winter influenza season in a small number of general practices to keep the study processes running smoothly and ready to react should there be a new influenza outbreak. In the event of a new influenza outbreak, FLU-CATs will quickly identify which problems that patients have that will best predict the level of care that they need.
PANdemic INfluenza Triage in the Emergency Department (The PAINTED study) ⁵	Steve Goodacre, University of Sheffield	PAINTED aims to identify the most accurate triage method for predicting severe illness among patients attending the emergency department with suspected pandemic influenza.
Maternal and perinatal outcomes of pandemic influenza in pregnancy (UKOSS) ⁶	Marian Knight, University of Oxford	The UKOSS influenza in pregnancy study will use an existing research platform to collect information on pregnant and post-partum women admitted to hospital with confirmed influenza infection. The management of women will be described, focusing particularly on the role of extracorporeal membrane oxygenation (ECMO), and women will be followed up to pregnancy completion to collect information on both maternal and perinatal outcomes.
Multi-centre Adjuvant Steroids in Adults with Pandemic Influenza (ASAP) Trial ⁷	Wei Shen Lim, Nottingham University Hospitals	The ASAP trial is a multicentre (>40 sites), blinded, randomised controlled clinical trial to determine if low-dose corticosteroids (dexamethasone 6 mg started within 24 h of admission once a day for 5 days), in addition to standard care, is associated with a lower risk of death or admission to intensive care, compared with placebo.
Real-time Modelling of a Pandemic Influenza Outbreak (RTM) ⁸	Daniela De Angelis, University of Cambridge	The RTM project advances the state of art for real-time pandemic modelling by developing an existing model used to reconstruct the 2009 H1N1 pandemic on the basis of realistic epidemic surveillance data collected. The work for the RTM has already been done and has resulted in a new monitoring tool that allows: capture of spatial variation in influenza transmission; use of efficient computational algorithms for the provision of timely statistical estimates and predictions; and incorporation of the above into freely available software. The tool is now available to Public Health England for use in the advent of a pandemic, and key staff has been trained in its use, supported by collaborators at the University of Cambridge to deal with workforce shortage during a pandemic. The real-time model has been tested in the monitoring of the 2017–18 seasonal influenza, and the model is ready to be used in the current season to estimate infection and clinical attack rates and to predict timing and magnitude of the peak influenza activity.

Table: Projects in the National Institute of Health Research's pandemic influenza preparedness portfolio

identify those who have died or needed life-saving interventions, and the information collected will be used to evaluate current, and develop new, triage methods.

The UK Obstetric Surveillance System (UKOSS) influenza in pregnancy study will collect information on pregnant and post-partum women admitted to hospital with confirmed influenza infection.²⁵ Pregnant women are known to be at high risk of severe outcomes for several infections, and appeared to be disproportionately affected in the 2009 influenza A H1N1 pandemic.²⁵ UKOSS will collect information on the management of women, focusing particularly on the role of extracorporeal membrane oxygenation, and women will be followed up to pregnancy completion to collect information on both maternal and perinatal outcomes.

The multicentre Adjuvant Steroids in Adults with Pandemic Influenza (ASAP) trial will determine the role of corticosteroids in adults admitted to hospital with influenza infection. Corticosteroids were used widely during the 2009 influenza A H1N1 pandemic, particularly for severely ill patients. However, increased mortality was found in observational studies.²⁶ The ASAP trial will recruit 2200 participants by the end of the first part of the next pandemic with trial results available to inform clinical practice before the start of the second part.⁷

The real-time modelling of a pandemic influenza outbreak (RTM) study has been funded to advance the state of the art of real-time modelling of influenza epidemics, and to provide a tool to monitor and predict the development of an ongoing pandemic outbreak in the UK. Outputs from the RTM study include: models to produce epidemic forecasts specific to age and region;²⁷ algorithms, building on the latest developments in statistical computation, to allow epidemic analyses to be updated in a timely fashion as the epidemic unfolds;³ and bespoke software and relevant training of Public Health England staff.

The need to put studies on standby while awaiting a pandemic is a feature of the pandemic influenza portfolio studies that has raised several issues. This situation reflects both the need to maintain interest and engagement of stakeholders (including those who initially agreed to host the research, policy makers, and others), and changes to the landscape in terms of research regulations, changes in science, and social changes.

Maintaining engagement of research sites to ensure rapid activation

Reduced research nurse availability in the event of a pandemic, both because of sickness absence and indirectly because of reallocation to clinical duties, is a particular issue for several of the studies.²⁴ Furthermore, for the ASAP trial, it is crucial that hospital sites are in a simmering state of readiness for an extended period, and potential obstacles must be minimised that might arise following activation and might require time-costly solutions.⁷ For instance, there will be no opportunity to

conduct and evaluate a pilot study once a pandemic arises. Instead, an activation simulation exercise was required which enabled the identification of stumbling-blocks and refinement of trial processes, adding to trial readiness. One important issue highlighted was the need for backup plans and pre-assigned deputy roles (including for the chief investigator), as part of trial resilience.²⁸

The PIPS study was specifically designed to support existing infrastructure provided by the Health Survey for England,²⁹ an annual, nationally representative, household survey which collects health information and blood specimens throughout the year.¹⁷ This study is therefore dependent on long-term continuation and its continuation during a pandemic.

The RTM study's prompt activation in the event of a pandemic depends on the continued commitment of relevant public bodies (eg, Public Health England and Department of Health) and collaborators to provide ready access to the agreed available data.

Changing regulatory environment and technologies

Studies report that research sites' research administration staff regularly attempt to close hibernated studies, with researchers needing to repeatedly explain the hibernation status to National Health Service (NHS) organisations that are unfamiliar with the concept and do not have systems in place to incorporate this form of research into their regulatory frameworks. The term hibernated thus possibly represents a misnomer; such studies are perhaps better thought of as in progress but awaiting activation of data collection. There might be a need for new processes designed specifically for studies such as these.

The PAINTED study was designed to recruit patients on the basis of clinical suspicion of pandemic influenza.²⁴ Advances in point of care technology mean that suspected seasonal influenza can be rapidly diagnosed or ruled out in the emergency department.³⁰ If applicable in a pandemic, this practice would change the study population but would not undermine the need for the study or change the study design. An observational cohort study would still be needed to identify predictors of adverse outcome in diagnosed pandemic influenza (rather than suspected pandemic influenza), and therefore a modified PAINTED study with a different population would continue. Influenza serological assays require specialist expertise only available in selected laboratories that will have limited research capacity during a pandemic because of a substantial increase in workload. Maintaining expertise outside of these centres is necessary to ensure that researchers can rapidly initiate testing of samples.¹ Pilot work carried out in the PIPS study¹ identified that establishing de-novo serological testing capacity led to unacceptable delays. In response to this finding, the NIHR is funding short laboratory placements in WHO's Collaborating Centre for Reference and

Research on Influenza for laboratory staff to maintain a pool of expertise that can be rapidly deployed. WHO's collaborating centre will also have access to the novel virus and the first available serological assay so these links will support the study's rapid assay set up and deployment. Consideration is being given in how to best run rapid turn-around surveys. For example, the number of households that have solely mobile phones is increasing. Such households might differ from landline households on several demographic and psychological levels. The FluTEST team is monitoring changes in the science of polling methods and exploring the possible cost-effectiveness of alternative approaches such as social media monitoring.⁸

Maintaining readiness of electronic platforms

Rapid evolution of the IT landscape is inevitable so adequate resources for maintenance of such studies is imperative. The FLU-CATS study was established on an electronic health record (eHR) system (Vision, In Practice Systems Ltd) in partnership with the Clinical Practice Research Datalink.²³ The use of this platform has gradually decreased within the primary care community as web-based and cloud-based systems have become more popular. To retain representative coverage in the UK, the FLU-CATS study has been obliged to add an additional eHR solution which will sit on the EMIS Health eHR (Egton Medical Information Systems Ltd, Leeds, UK). This solution will be rolled out in collaboration with the Royal College of General Practitioners Research and Surveillance Centre.

The PAINTED study has been designed to use paper records, in accordance with typical NHS emergency department practice at the time.²⁴ However, increasing numbers of NHS emergency departments are developing electronic data systems. The new Emergency Care Data Set has a research data item that allows identification of cases recruited to research and could be used to electronically flag cases included in the PAINTED study. Further pilot work is required to determine whether and how the new Emergency Care Data Set and new emergency department data systems can be used to optimise data collection.

The UKOSS research platform switched from a paper-based reporting system to an electronic system during the hibernation period. Additional funding to activate the study to collect information on seasonal influenza management and outcomes in pregnancy has enabled the required programming to be undertaken to maintain the hibernated pandemic influenza study in readiness.²⁵

Workforce issues during a pandemic

The PIPS study is dependent on the continuation of the Health Survey for England during a pandemic and its ability to either maintain laboratory expertise in influenza serology or recruit staff with relevant experience. The data analysis protocols have been automated and could be run by alternative staff. A similar approach has been taken by

the EAVE project, whereby systems have been automated sufficiently for non-core staff to run the relevant syntax to extract and analyse data. For FLU-CATS, the study process relies on both continued activity by general practitioners in primary care to continue their usual consultation activity for influenza-like illness using their usual eHR. The study runs on background processes that harvest information from these consultations. Substitution by locums will not inherently affect the study provided consultations continue and the eHR is used. The study research team is small; however, the processes for data analysis have been archived and test activated against the current eHR systems, so the processes could be run by substitute staff. For FluTEST, the open access publication of the protocol and survey questions allows anyone to use the FluTEST items and analyse the resulting data. For ASAP, all sites and participating research partners (including manufacturing units) have been requested to identify persons who can deputise for key research staff in the event of absence from work. There is also an internal pandemic plan to ensure the project can continue to deliver the trial, including a co-chief investigator. PAINTED is working on creating an electronic triage form for emergency department systems that could be used to collect electronic data for the study as part of routine clinical care. Furthermore, using record linkage to NHS Digital will allow the study to be undertaken with minimal research nurse support.

Influenza pandemic preparedness and emerging infective and major incident challenges

This national portfolio of hibernated pandemic influenza studies is now mature, and, we believe, clearly illustrates the value of the UK's clinical research system and the potential for rapid research, and clinical and public health response, in a future pandemic. However, when and whether a future influenza pandemic occurs is unpredictable. What is clearer, is that other emerging infective challenges will continue to occur. Can these studies act as a model for research in a future pandemic or emerging infection? Experience suggests they can as UKOSS was used to study Zika virus exposure during pregnancy.³¹

Other hibernated studies could equally address research questions likely to be essential in any future epidemic or pandemic, and not just those specific to influenza. For instance, the FLU-CATS response to changes in the architecture of eHRs—ie, the study tools could be used in either primary or secondary care settings for a variety of emerging diseases, and deliberately or accidentally released agent (chemical, biological, radiological, and nuclear agent)—although the performance of the tool will vary with the population being triaged.²³ PAINTED is focused on triage, given its importance in ensuring patients with suspected influenza are managed appropriately in a busy emergency department setting.²⁴ However, a robust triage process is likely to be important if any other emerging infection results in large numbers of patients seeking

emergency care, to ensure that the health system can function. As departments continue to face capacity issues, exploring the role of triage in the event of other epidemics or pandemics would be valuable in assessing its effectiveness in times of severe stress on health systems.

The EAVE project has created a new sentinel system linking primary care data to RT-PCR swabs, stored serology, and hospital and mortality outcomes. The project will allow for a rapid evaluation of any new vaccination irrespective of the infection it is developed to test against. Also, as noted previously, serological analysis of stored samples on susceptibility is dependent on the provision of an assay that will allow this work to be undertaken. Experience of the last pandemic would indicate that there will likely be some delay in the development and deployment of such an assay. Prior agreement for access to the first assay approved for use would be important to maximise the study's benefit.

The PIPS project specimen and data collection could also be used for other epidemic or pandemic scenarios provided a serological marker of susceptibility or infection is available. Similarly, surveys using the FluTEST survey method could be used to monitor behaviour whatever the public health emergency (including chemical, biological, radiological, and nuclear risks).

The RTM system could also be used in other outbreaks and could be regularly used outside a pandemic to anticipate the burden of seasonal influenza. This practice would also test the ease of its implementation and give Public Health England staff the opportunity to develop their expertise. However, use on seasonal data would pose challenges around release of relevant data (eg, on general practice consultations), whose access should be instead guaranteed during a pandemic.

Irrespective of the emerging infection, trials will be needed—ie, a network set up and ready to test one intervention versus standard care will be invaluable to fast track the evidence needed for clinical practice, and the periodic activation exercise undertaken by the ASAP trial provides a model for this. Work has already begun to develop the ASAP trial network for the broader purpose of determining the role of adjuvant corticosteroids in severe acute respiratory viral infections other than pandemic influenza.

Lessons learned and next steps

Although more information on project performance will be available once the projects have been brought out of hibernation for testing, there have already been several lessons learned during the set-up phase. There might be a need for incremental modification in the event of changes in the architecture of eHRs. Approaches are needed to mitigate for reduced staff availability in the event of a pandemic because of sickness absence and indirectly because of reallocation to clinical duties—eg, pre-assigned deputy roles and automation of processes. Commitment of relevant public bodies and

collaborators—eg, to provide ready access to agreed available data should be maintained. A pool of expertise (beyond selected laboratories) that can be rapidly deployed is needed to carry out de-novo serological testing during a pandemic. Organisational regulatory frameworks need to be flexible for studies that require long periods before activation. With technological advances for running rapid turn-around surveys, consideration should be given to alternative approaches that might be faster and cheaper.

Should the UK's portfolio of hibernated pandemic influenza studies provide a model for planning research in public health emergencies, including emerging infections, going forward? All too often researchers are in competition when trying to answer research questions rapidly in an emergency situation. The research response to the Ebola outbreak in parts of Africa was late and then inappropriately competitive, resulting in several underpowered and unsuccessful studies.³² By contrast, the UK model allows for advance funding and planning of a complementary suite of studies, system testing (a test has been carried out in 2018 with reports compiled by the NIHR), and developing a collaborative network of researchers. Existing international networks—eg, the International Severe Acute Respiratory and Emerging Infection Consortium and the International Network of Obstetric Survey Systems either are or could be primed and ready to do similar studies. However, the challenge going forward will be for commercially funded studies not to compete for scarce patient resources and to fit within this framework to ensure the highest quality studies are done most expediently. With this in mind, is now the time for a national or international register of planned pandemic and emerging infection studies with agreements over priorities and co-enrolment and collaboration?

While an influenza pandemic heads the UK National Risk Register of Civil Emergencies,³³ large infectious outbreaks do not constitute the only public health risk. Research to improve care is also essential in other public health emergencies, such as natural disasters and the deliberate release of biochemical weapons as part of terrorist attacks.^{16,34} Coordination across research and funding bodies (commercial, charity, and government) will be required if expectations demanded by the public or the research community are to be met in response to public health emergencies. Is this the time to consider a new way of working together as part of emergency preparedness, including public engagement on the ethical questions that arise?

Contributors

MK and CRS led the writing of this Personal View. All authors made contributions to the writing and discussions on the scope of this Personal View, and critically reviewed revisions and the final paper.

Declaration of interest

SG is Chair of the National Institute for Health Research (NIHR) Health Technology Assessment Clinical Evaluation and Trials Board, and chaired the NIHR Pandemic Influenza Themed Call Board. All authors reports grants from NIHR, during the conduct of the study. DDA reports grants from Public Health England, outside the submitted works.

For more on the **International Severe Acute Respiratory and Emerging Infection Consortium** see <https://isaric.tghn.org>

For more on the **International Network of Obstetric Survey Systems** see <https://www.npeu.ox.ac.uk>

WSL's institution has received unrestricted investigator-initiated research funding from Pfizer for an unrelated research study in pneumonia where WSL is the chief investigator.

Acknowledgments

The authors would like to thank all the collaborators of this work. Helen Walters (National Institute for Health Research Evaluation, Trials and Studies Coordinating Centre, Southampton, UK) and Garry Meakin (University of Nottingham, UK) contributed to the collaborative group meeting at which the concept of this Personal View was first discussed. These studies were funded by grants from the National Institute for Health Research Health under its Pandemic Influenza Preparedness call. Professor Hayward is a National Institute for Health Research Senior Investigator. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the HS&DR, HTA and Public Health programmes, the National Institute for Health Research, the National Health Service, or the Department of Health and Social Care, UK.

References

- 1 Fragaszy EB, Quinlivan M, Breuer J, et al. Population-level susceptibility, severity and spread of pandemic influenza: design of, and initial results from, a pre-pandemic and hibernating pandemic phase study using cross-sectional data from the Health Survey for England (HSE). *Public Health Res* 2015; **3**: 1–24.
- 2 Rubin GJ, Bakhshi S, Amlôt R, Fear N, Potts H, Michie S. The design of a survey questionnaire to measure perceptions and behaviour during an influenza pandemic: the Flu Telephone Survey Template (FluTeST). *Health Serv Deliv Res* 2014; **2**: 41.
- 3 Simpson CR, Lone N, McMenamin J, et al. Early estimation of pandemic influenza Antiviral and Vaccine Effectiveness (EAVE): use of a unique community and laboratory national data-linked cohort study. *Health Technol Assess* 2015; **19**: 1–32.
- 4 Health Protection and Influenza Research. <https://www.nottingham.ac.uk/research/groups/healthprotection/projects/flu-cats.aspx> (accessed April 3, 2019).
- 5 The PAINTED Study. <https://www.sheffield.ac.uk/scharr/painted> (accessed Sept 3, 2018).
- 6 Seasonal Influenza in Pregnancy. <https://www.npeu.ox.ac.uk/ukoss/current-surveillance/flu> (accessed Nov 5, 2018).
- 7 Lim WS, Brittain C, Duley L, et al. Blinded randomised controlled trial of low-dose Adjuvant Steroids in Adults admitted to hospital with Pandemic influenza (ASAP): a trial 'in hibernation', ready for rapid activation. *Health Technol Assess* 2015; **19**: 1–78.
- 8 Birrell PJ, Pebody RG, Charlett A, Zhang X-S, De Angelis D. Real-time modelling of a pandemic influenza outbreak. *Health Technol Assess* 2017; **21**: 1–118.
- 9 Walley T, Davidson P. Research funding in a pandemic. *Lancet* 2010; **375**: 1063–65.
- 10 Simpson CR, Ritchie LD, Robertson C, Sheikh A, McMenamin J. Effectiveness of H1N1 vaccine for the prevention of pandemic influenza in Scotland, UK: a retrospective observational cohort study. *Lancet Infect Dis* 2012; **12**: 696–702.
- 11 Goodacre S, Challen K, Wilson R, Campbell M. Evaluation of triage methods used to select patients with suspected pandemic influenza for hospital admission: cohort study. *Health Technol Assess*. 2010; **14**: 173–236.
- 12 Killingley B, Greatorex J, Cauchemez S, et al. Virus shedding and environmental deposition of novel A (H1N1) pandemic influenza virus: interim findings. *Health Technol Assess*. 2010; **14**: 237–354.
- 13 Annane D, Antona M, Lehmann B, et al. Designing and conducting a randomized trial for pandemic critical illness: the 2009 H1N1 influenza pandemic. *Intensive Care Med*. 2012; **38**: 29–39.
- 14 Rubin GJ, Amlôt R, Page L, Wessely S. Public perceptions, anxiety and behavioural change in relation to the swine flu outbreak: a cross-sectional telephone survey. *BMJ* 2009; **339**: b2651.
- 15 Nguyen-Van-Tam JS, Openshaw PJ, Hashim A, et al. Risk factors for hospitalisation and poor outcome with pandemic A/H1N1 influenza: United Kingdom first wave (May–September 2009). *Thorax* 2010; **65**: 645–51.
- 16 Lurie N, Manolio T, Patterson AP, Collins F, Frieden T. Research as a part of public health emergency response. *N Engl J Med* 2013; **368**: 1251–55.
- 17 Fragaszy EB, Quinlivan M, Breuer J, et al. Population-level susceptibility, severity and spread of pandemic influenza: design of, and initial results from, a pre-pandemic and hibernating pandemic phase study using cross-sectional data from the Health Survey for England (HSE). *Public Health Res* 2015; **3**: 1–24.
- 18 Rubin GJ, Potts HWW, Michie S. Likely uptake of swine and seasonal flu vaccines among healthcare workers. A cross-sectional analysis of UK telephone survey data. *Vaccine* 2011; **29**: 2421–28.
- 19 Rubin GJ, Potts HWW, Michie S. The impact of communications about swine flu (influenza A H1N1v) on public responses to the outbreak: results from 36 national telephone surveys in the UK. *Health Technol Assess* 2010; **14**: 183–266.
- 20 Simpson CR, Lone N, Kavanagh K, et al. Seasonal influenza vaccine effectiveness (SIVE): an observational retrospective cohort study—exploitation of a unique community-based national-linked database to determine the effectiveness of the seasonal trivalent influenza vaccine. *Health Serv Deliv Res* 2013; **1**: 10.
- 21 Scientific Pandemic Influenza Group. Modelling SPI-M Modelling Summary. <https://www.gov.uk/government/publications/spi-m-publish-updated-modelling-summary> (accessed Nov 21, 2018).
- 22 Baguelin M, Flasche S, Camacho A, et al. Assessing optimal target populations for influenza vaccination programmes: an evidence synthesis and modelling study. *PLoS Med* 2013; **10**: e1001527.
- 23 Venkatesan S, Myles PR, McCann G, et al. Development of processes allowing near real-time refinement and validation of triage tools during the early stage of an outbreak in readiness for surge: the FLU-CATs Study. *Health Technol Assess* 2015; **19**: 1–132.
- 24 Goodacre S, Irving A, Wilson R, Beever D, Challen K. The PAndemic INfluenza Triage in the Emergency Department (PAINTED) pilot cohort study. *Health Technol Assess*, 2015; **19**: 1–69.
- 25 Knight M, Brocklehurst P, O'Brien P, Quigley MA, Kurinczuk JJ. Planning for a cohort study to investigate the impact and management of influenza in pregnancy in a future pandemic. *Health Serv Deliv Res* 2015; **3**: 6.
- 26 Rodrigo C, Leonardi-Bee J, Nguyen-Van-Tam JS, Lim WS. Effect of corticosteroid therapy on influenza-related mortality: a systematic review and meta-analysis. *J Infect Dis* 2015; **212**: 183–89.
- 27 Birrell PJ, Zhang X-S, Pebody RG, Gay N, De Angelis D. Reconstructing a spatially heterogeneous epidemic: characterising the geographic spread of 2009 A/H1N1 infection in England. *Sci Rep* 2016; **11**: 29004.
- 28 Lim WS, Meakin G, Brittain C, Bewick T, Duley L. Improving readiness for recruitment through simulated trial activation: the Adjuvant Steroids in Adults with Pandemic influenza (ASAP) trial. *Trials* 2017; **18**: 546.
- 29 Fuller E, Mindell J, Prior G, eds. Health Survey for England, 2016. London: NHS Digital, 2017.
- 30 Davis S, Allen AJ, O'Leary R, et al. Diagnostic accuracy and cost analysis of the Alere i Influenza A&B near-patient test using throat swabs. *J Hosp Infect* 2017; **97**: 301–09.
- 31 NPEU. Pregnancy outcomes in mothers with a history of travel to a country with active Zika transmission. <https://www.npeu.ox.ac.uk/ukoss/current-surveillance/zika-virus> (accessed Sept 4, 2018).
- 32 Thielman NM, Cunningham CK, Woods C, Petzold E, Sprentz M, Russell J. Ebola clinical trials: five lessons learned and a way forward. *Clin Trials* 2016; **13**: 83–86.
- 33 UK National Risk Register of Civil Emergencies, Cabinet Office, 2015. <https://www.gov.uk/government/publications/national-risk-register-of-civil-emergencies-2017-edition> (accessed Nov 12, 2018).
- 34 Levine AC, Teicher C, Aluisio AR, et al. Regional Anesthesia for Painful Injuries after Disasters (RAPID): study protocol for a randomized controlled trial. *Trials* 2016; **17**: 542.

© 2019 Elsevier Ltd. All rights reserved.