



## Original article

Isolation and characterization of a *Rickettsia* from the ovary of a Western black-legged tick, *Ixodes pacificus*

Maryam Alowaysi<sup>a,1</sup>, Junyan Chen<sup>a,1</sup>, Sierra Stark<sup>a</sup>, Kristine Teague<sup>a</sup>, Monique LaCourse<sup>a</sup>, Joanna Proctor<sup>a</sup>, Katie Vigil<sup>a</sup>, Jeremy Corrigan<sup>a</sup>, Aja Harding<sup>a</sup>, Jinze Li<sup>a</sup>, Timothy Kurtti<sup>b</sup>, Jianmin Zhong<sup>a,\*</sup>

<sup>a</sup> Department of Biological Sciences, Humboldt State University, Arcata, CA, 95521, USA

<sup>b</sup> Department of Entomology, University of Minnesota, St. Paul, MN, 55108, USA

## ARTICLE INFO

## Keywords:

'*Rickettsia monacensis*' str. Humboldt  
*Ixodes pacificus*  
Phylogenetic analysis  
Multilocus sequence typing  
Multi-spacer sequence typing

## ABSTRACT

A rickettsial isolate was obtained from a partially engorged *Ixodes pacificus* female, which was collected from Humboldt County, California. The isolate was provisionally named *Rickettsia* endosymbiont *Ixodes pacificus* (REIP). The REIP isolate displayed the highest nucleotide sequence identity to *Rickettsia* species phylotype G021 in *I. pacificus* (99%, 99%, and 100% for *ompA*, 16S rRNA, and *gltA*, respectively), a bacterium that was previously identified in *I. pacificus* by PCR. Analysis of sequences from complete opening frames of five genes, 16S rRNA, *gltA*, *ompA*, *ompB*, and *sca4*, provided inference to the bacteria's classification among other *Rickettsia* species. The REIP isolate displayed 99.8%, 99.4%, 99.2%, 99.5%, and 99.6% nucleotide sequence identity for 16S rRNA, *gltA*, *ompA*, *ompB*, and *sca4* gene, respectively, with genes of '*R. monacensis*' str. IrR/Munich, indicating the REIP isolate is closely related to '*R. monacensis*'. Our suggestion was further supported by phylogenetic analysis using concatenated sequences of 16S rRNA, *gltA*, *ompA*, *ompB*, and *sca4* genes, concatenated sequences of *dksA-xerC*, *mppA-purC*, and *rpmE-tRNA<sup>Met</sup>* intergenic spacer regions. Both phylogenetic trees implied that the REIP isolate is most closely related to '*R. monacensis*' str. IrR/Munich. We propose the bacterium be considered as '*Rickettsia monacensis*' str. Humboldt for its closest phylogenetic relative (= DSM 103975 T = ATCC TSD-94 T).

## 1. Introduction

*Ixodes pacificus*, the Western black-legged tick, is a predominant vector of *Borrelia burgdorferi* sensu stricto and *Anaplasma phagocytophilum*, the etiologic agents of Lyme borreliosis and anaplasmosis, respectively, in the Pacific West Coast region of the United States (Burgdorfer et al., 1985; Foley et al., 2008; Piesman et al., 1999). Ixodid ticks, as well as other arthropods, often harbor rickettsiae that are known as obligate intracellular bacteria and arthropod-borne pathogens of vertebrate hosts (Hardstone Yoshimizu and Billeter, 2018). Presently, 32 *Rickettsia* species within the spotted fever, typhus, *Rickettsia bellii*, and *Rickettsia canadensis* groups have been validated (Abdad et al., 2017; <http://www.bacterio.net/rickettsia.html>). Utilization of molecular techniques has facilitated the continual detection and isolation of rickettsiae from ticks around the world (Parola et al., 2013; Cazorla et al., 2008; Richards, 2012). Several serological studies have noted that *I. pacificus* harbors a spotted fever group *Rickettsia* that is

antigenically related to pathogenic species (Hughes et al., 1976; Philip et al., 1978, 1981). We have previously identified *Rickettsia* species phylotype G021 and *Rickettsia* species phylotype G022 via PCR amplification using genomic DNA extracted from *I. pacificus* ticks; of which, the bacterial prevalence was found to be 100% and 2%, respectively (Phan et al., 2011; Cheng et al., 2013b). Transmission electron microscope and Fluorescence in situ hybridization assays detected the presence of *Rickettsia* species in midgut and ovaries of *I. pacificus* (Phan et al., 2011; Bagheri et al., 2017). It was later noted that *Rickettsia* species phylotype G021 was found to possess 100% transovarial and transstadial efficiency as well as the genetic capacity to synthesize tetrahydrofolate, indicating a possible nutritional relationship between the host and endosymbiont (Bodnar et al., 2018; Cheng et al., 2013a; Kurlovs et al., 2014; Hunter et al., 2015).

\* Corresponding author.

E-mail address: [jz15@humboldt.edu](mailto:jz15@humboldt.edu) (J. Zhong).

<sup>1</sup> These authors contributed equally to this work.

## 2. Material and methods

### 2.1. Source of ticks and tick cell culture

To isolate *Rickettsia* phylotype G021 from a mixed population of microorganisms in *I. pacificus*, we collected a partially engorged *I. pacificus* female that was donated from McKinleyville Animal Care Center, McKinleyville, CA, in March 2013. The standard Universal Transverse Mercator Grid System coordinate for the tick collection site is 10 N 429550 4530570. The isolation of rickettsiae from the engorged tick followed methods reported previously (Kurtti et al., 1996, 2015). In brief, ovaries were extirpated from a surface disinfected, partially engorged, female. Ovarian fragments were cultured with tick cell line IRE11 and incubated in a candle jar at 25 °C. Once established, the isolate was maintained in tick cell line ISE6 at 34 °C (Simser et al., 2002; Munderloh et al., 1994). In the sixth serial transfer, genomic DNA was prepared and submitted to BioProject PRJNA232537 for genome sequencing (SRX476329; SRX476328 and SRX476327).

The isolate was initially named *Rickettsia* endosymbiont *Ixodes pacificus* (REIP). Infection of IRE11 and ISE6 cell lines was conducted by mixing  $10^6$  cells with 0.3–0.5 mL of REIP-infected IRE11 lysate, resulting in 20–100 rickettsiae per tick cell, in 25 cm<sup>2</sup> flasks (Fisher Scientific, USA). Cells of rickettsiae and IRE11 were counted using a Petroff Hauser counting chamber under a Leica DM750 phase-contrast microscope with Leica high-resolution camera module (Leica Microsystems, USA). Infected cells were cultured for three weeks at 25 °C in L-15B300 complete medium supplemented with 10% fetal bovine serum, 5% tryptose phosphate broth, 0.1% bovine lipoprotein cholesterol concentrate, and 10 mM HEPES (Kurtti et al., 2005; Munderloh and Kurtti, 1989). Giemsa stain (KaryoMAX®, Thermo Fisher Scientific, USA) was performed on a weekly basis to monitor infection status. Giemsa stained REIP-infected ISE6 cells (or IRE11) were observed under the Leica DM750 microscope.

### 2.2. DNA extraction and PCR amplification

To classify REIP into genus and species, genomic DNA was extracted from  $5 \times 10^5$  REIP-infected IRE11 cells using DNeasy Blood & Tissue Kit for both MLST and MST (Qiagen, Germany). Amplification of 16S rRNA, *gltA*, *ompA*, *dksA-xerC*, *mppA-purC*, and *rpmE-tRNA<sup>fMet</sup>* was conducted using PCR using primers specified in Table 1 (Simser et al., 2002; Fournier and Raoult, 2007; Cheng et al., 2013b). Specifically, 2 µL genomic DNA (50 ng/µL), 1 µL 5 mM forward and reverse primers, 10 µL GoTaq® Green Master Mix (Promega, USA), and 6 µL PCR grade water were added in PCR reactions. Negative controls contained all components with the exception of REIP genomic DNA. The β-actin-coding gene was amplified simultaneously to serve as positive control using β-actin specific primers and field-collected *I. pacificus* (Cheng

et al., 2013b). PCR reactions were performed in an Applied Biosystems 2720 Thermal Cycler (Applied Biosystems, USA) with an initial denaturation at 95 °C for 5 min; this was followed by 30–45 cycles of denaturation at 95 °C for 30 s, annealing at 50–60 °C for 30 s, and extension at 72 °C for 30 s. The final extension was completed at 72 °C for ten minutes. Following amplification, PCR products were run on 2% agarose gel with 1X TAE buffer and stained in 10 µg/mL ethidium bromide. Alphamager® HP was used to visualize and photograph gels (ProteinSimple, USA).

### 2.3. Cloning and DNA sequencing


PCR amplicons were ligated into StrataClone's pSC-A-amp/kan cloning vector (Agilent Technologies, USA). Transformants were plated on LB agar containing 100 µg/mL ampicillin and 40 µl of 2% X-gal. White colonies were then restreaked on LB/AMP/X-gal plates. Following restreaking, white colonies were grown in LB broth containing 100 mg/mL ampicillin overnight at 37 °C. Plasmids were purified by E.Z.N.A.® Plasmid Mini Kit (Omega, USA) and constructs were confirmed by running EcoRI restriction enzyme digests (Promega, USA) on 1% agarose gel with 1X TAE buffer and stained in ethidium bromide. Sequencing of the PCR amplicons was conducted by Elim Biopharmaceuticals, Inc. using M13 reverse primer (5'-GGAAACAGCT ATGACCATG-3'). The GenBank accession numbers for the 16S rRNA, *gltA*, and *ompA* genes, *dksA-xerC*, *mppA-purC*, and *rpmE-tRNA<sup>fMet</sup>* regions for determining the numbers of species and genotypes of the REIP isolate are KX505845, KX505846, KX505847, KX505844, KX505842, KX505843, respectively.

### 2.4. Phylogenetic analysis

We utilized the completed genome sequence of REIP, which is available in NCBI (accession number NZ\_LAOP01000001). Homologous sequences of the whole open reading frame of each of the five genes of REIP, 16S rRNA (NZ\_LAOP01000001), *gltA* (KJW02430), *ompA* (KJW02278), *ompB* (KJW03404), and *sca4* (KJW03052), were obtained by BLAST analysis. For phylogenetic reconstructions, homologous sequences for each gene (Supplemental Table) were aligned using Clustal Omega at EMBL-EBI. The resulting aligned sequences of 16S rRNA (1,440 bp), *gltA* (1,233 bp), *ompA* (1,330 bp), *ompB* (2,847 bp) and *sca4* (2,691 bp) genes of REIP and 21 validated *Rickettsia* species were concatenated to generate a sequence of 9,829 characters for each species in Mesquite (version 3.04). Sequences from other validated *Rickettsia* species were not included in this study since their *ompA* gene is either truncated or lacking. The concatenated nucleotide sequences were translated to amino acid sequences, with a purpose of aligning the nucleotide sequences to match the amino acid alignments in Mesquite. Phylogeny of the concatenated sequence was constructed by SeaView

**Table 1**  
PCR reaction primers, annealing temperatures, and amplicon sizes.

Gene name	Primer (5'-3')	Annealing temperature	Amplicon size (bp)	Reference
16S rRNA	F-AGAGTTTGATCCTGGCTCAG R-AACGTCATTATCTTCCTTGC	58 °C	436	Simser et al., 2002
<i>gltA</i>	F-GGGGGCCTGCTCACGGCGG R-ATTGCAAAAAGTACAGTGAACA	50 °C	382	Simser et al., 2002
<i>ompA</i>	F-ATGGCGAATATTTCTCCAAAAA R-AGTGCAGCATTGCTCCCCCT	60 °C	530	Simser et al., 2002
<i>dksA-xerC</i>	F-TTTTCATGACGCTCTTGAGC R-GTAAAGAAAAGATAATTCCGTGGTT	58 °C	165	Fournier and Raoult, 2007
<i>mppA-purC</i>	F-GCATATGCRGTRGGTAGTTAT R-CACAGCCTCAATTCTAATT	59 °C	1000	Fournier and Raoult, 2007
<i>rpmE-tRNA<sup>fMet</sup></i>	F-TTCCGAAATGTAGTAAATCAATC R-TCAGGTTATGAG CCTGACGA	57 °C	368	Fournier and Raoult, 2007
β-actin	F-TTGTCCGCGACATCAAGGA R-CGGGAAGCTCGTAGGACTTCT	57 °C	110	Cheng et al., 2013b



**Fig. 1.** Giemsa stain of *Rickettsia endosymbiont Ixodes pacificus* (REIP)-infected ISE6 cells propagated for three weeks at 26 °C. (a) Uninfected ISE6 control without bacteria residing in the cytoplasm. (b) Heavily infected ISE6 cells three weeks post infection with REIP. Original magnification:  $\times 1000$ . Bar, the scale bar in  $\mu\text{m}$ .

(version 4.4.3) (<http://pbil.univlyon1.fr/software/seaview.html>). Gouy and Guindon S. & Gascuels' method was used to determine evolutionary distance values, which were then used to construct phylograms by neighbor-joining method and maximal parsimony method (Gouy et al., 2010). The pairwise nucleotide sequence identity of each gene between REIP and validated *Rickettsia* species was calculated in SIAS (<http://imed.med.ucm.es/Tools/sias.html>) using aligned sequences from Clustal Omega.

### 3. Results and discussion

We successfully propagated the REIP isolate using embryonic tick cell lines IRE11 and ISE6 (Simser et al., 2002; Munderloh and Kurtti, 1989), in L15B300 media formulated for tick cell culture (Munderloh and Kurtti, 1989; Oliver et al., 2014). Giemsa stained REIP-infected ISE6 cells were observed under light microscope. Cocci-shaped bacteria were viewed residing in the cytoplasm or outside lysed cells; uninfected controls had no visible signs of bacterial inhabitation (Fig. 1).

To confirm that there is only one bacterial species and/or genotype in the REIP isolate, we utilized multilocus sequence typing (MLST) using three genes—16S rRNA, *gltA*, and *ompA*—allowing for the detection of single nucleotide polymorphisms that distinguish *Rickettsia* species (Fournier et al., 2003; Zhu et al., 2005a). However, because coding DNA may not provide enough nucleotide sequence variation to differentiate within some rickettsial species, we also used multi-spacer sequence typing (MST)—utilizing intergenic spacers such as *dksA-xerC*, *mppA-purC*, and *rpmE-tRNA<sup>Met</sup>*—as this method is capable of differentiating at the intraspecies level (Fournier et al., 2004; Zhu et al., 2005b). The three genes and three intergenic spacers were amplified via PCR and amplicons were ligated into StrataClone's pSC-A-amp/kan cloning vector. Sequencing data determined the insert size of clones for *gltA*, *ompA*, 16S rRNA genes of REIP to be 382, 530 and 436 base pair (bp)-long, respectively. PCR amplification and cloning of intergenic spacers determined the insert size of *dksA-xerC*, *mppA-purC*, and *rpmE-tRNA<sup>Met</sup>* regions were 165, 121, 134 and 368 bp, respectively (data not shown). Additional, sequences from 30 clones of 16S rRNA, *gltA*, *ompA*, *dksA-xerC*, *mppA-purC*, and *rpmE-tRNA<sup>Met</sup>* genes of REIP were aligned in ClustalX version 2.1. Nucleotide sequence alignments of thirty clones for each of the genes and intergenic spacers determined that the nucleotide sequences were identical to one another in the amplified regions. Additionally, no difference was noted when comparing amplified 16S rRNA, *gltA*, and *ompA* sequences to sequences obtained from the genome sequencing project of REIP (Supplemental Table). Considering REIP's nucleotide homology among all thirty clones, we assumed the REIP isolate possessed one genotype.

We utilized complete open reading frames of 16S rRNA, *gltA*, *ompA*, *ompB*, and *sca4* nucleotide sequences of REIP obtained from the genome sequencing project in NCBI (accession number NZ\_LAOP01000001) to classify the REIP isolate. The pairwise nucleotide sequence identity comparison was conducted between 1,440 bp 16S rRNA, 1,233 bp *gltA*, 1,330 bp *ompA*, 2,847 bp *ompB*, and 2,691 bp *sca4* genes of REIP and homologous sequences of 21 validated *Rickettsia* species in GenBank. The nucleotide sequence identities of 16S rRNA, *gltA*, *ompA*, *ompB*, and *sca4* genes of REIP with homologous sequences of *R. buchneri* str. ISO7 (Kurtti et al., 2015) was 99.7%, 99.5%, 91.7%, 76.4%, and 47.3%, respectively (Table 2). Based on Fournier et al., 2003 Fournier et al.'s (2003) classification scheme, these values confirmed the isolate is a member of the *Rickettsia* genus. For all genes, REIP shared the highest nucleotide sequence identity with '*R. monacensis*' str. IrR/Munich (99.8, 99.4, 99.2, 99.5, and 99.6% for 16S rRNA, *gltA*, *ompA*, *ompB*, and *sca4*, respectively). The identities between the two species for all genes indicate that the REIP isolate is closely related to '*R. monacensis*'. Moreover, the isolate also displayed similarity to *Rickettsia* species phylotype G021. For instance, the isolate shared 99%, 99%, and 100% nucleotide identity to *Rickettsia* species phylotype G021's *ompA*, 16S rRNA, and *gltA* sequences, respectively (Phan et al., 2011). Considering the nucleotide sequences of 30 clones for each gene were identical, we ruled out the possibility that the discrepancies observed between nucleotide sequences for REIP and phylotype G021's genes could have resulted from random errors during PCR. The incongruity is more likely due to bacterial genetic diversity within species or an error when sequencing phylotype G021. Therefore, we are in favor of the conclusion that REIP is the same species as phylotype G021, however possesses a different genotype.

Sequence similarities present between REIP, *R. buchneri*, '*R. monacensis*', and *Rickettsia* species phylotype G021 were also reflected via phylogenetic analysis. Phylogenetic analysis was conducted by comparing the REIP isolate's concatenated *gltA*, *ompA*, *ompB*, *sca4*, and 16S rRNA nucleotide sequences to homologous sequences of rickettsiae available in GenBank. Trees generated from the concatenated sequences reflected those previously generated by our lab for *Rickettsia* species

**Table 2**

The nucleotide sequence identity of the *gltA*, *ompA*, *ompB*, *sca4*, and 16S rRNA genes between '*Rickettsia monacensis*' str. Humboldt and *Rickettsia buchneri* str. ISO7 and *R. monacensis* str. IrR/Munich.

Strains	' <i>R. monacensis</i> ' str. Humboldt				
	16S rRNA	<i>gltA</i>	<i>ompA</i>	<i>ompB</i>	<i>sca4</i>
<i>R. buchneri</i> str. ISO7	99.65	99.51	91.71	76.41	47.32
<i>R. monacensis</i> str. IrR/Munich	99.79	99.35	99.17	99.50	99.55




Fig. 2. Phylogenetic tree of concatenated *gltA*, *ompA*, *ompB*, *sca4*, and 16S rRNA gene sequences of '*Rickettsia monacensis*' str. Humboldt. The tree was generated using neighbor-joining distance method and 1000 bootstrap replicates. Bootstrap values > 50% are shown at the nodes. Bar, nucleotide distance. '*R. monacensis*' str. Humboldt (or *Rickettsia* endosymbiont *Ixodes pacificus*) is represented at the top of the tree. Other selected sequences on the tree include: *R. aeschlimannii*, *R. africae*, *R. akari*, *R. australis*, *R. buchneri*, *R. conorii*, *R. felis*, *R. heilongjiangensis*, *R. honei*, *R. japonica*, *R. massiliae*, '*R. monacensis*' str. IrR/Munich, *R. montanensis*, *R. parkeri*, *R. peacockii*, *R. philipii*, *R. raoultii*, *R. rhipicephali*, *R. rickettsii*, *R. sibirica*, and *R. slovaca*. The accession numbers of genes of all *Rickettsia* species are listed in Supplemental Table.


phylotype G021 (Phan et al., 2011). The tree topology indicated that REIP was clustered with '*R. monacensis*' str. IrR/Munich and that the two bacteria were placed in the same clade as *R. buchneri* str. ISO7. This monophyletic group forms a sister group to the rest of a clade that contains 16 spotted fever group rickettsiae. The concatenated sequences of the five genes of the spotted fever group rickettsiae are distant from the sequences of *R. akari*, *R. australis*, and *R. felis* (Fig. 2). The neighbor joining tree indicated that the monophyletic groups, in which REIP resides, are distinct from the rest of the spotted fever group rickettsiae. We inferred that the REIP isolate is closely related to '*R. monacensis*' and shares similar evolutionary history with *R. buchneri* and '*R. monacensis*' str. IrR/Munich. As such, based on our nucleotide sequence identities and phylogenetic analyses, we propose that the REIP isolate constitutes a new strain of '*R. monacensis*' different from '*R. monacensis*' strain IrR/Munich. Therefore, we propose the name '*Rickettsia monacensis*' str. Humboldt, for the bacterium.

Previous studies have demonstrated the ability of MST to differentiate between genotypes of some *Rickettsia* species by at least one nucleotide mutation (Fournier and Raoult, 2007; Zhu et al., 2005b; Znazen et al., 2013). As such, MST was used to further understand the evolutionary relationship between '*R. monacensis*' str. Humboldt and other *Rickettsia* species. A neighbor joining tree was generated based on the concatenated nucleotide sequence of *dkxA-xerC*, *mppA-purC*, and *rpmE*-tRNA<sup>Met</sup> regions and their homologous sequences from 22 validated *Rickettsia* species in GenBank (Supplemental Table). There was a limit to what rickettsial strains could be used to generate the tree as some strains don't have all three spacer sequences and these regions for other strains have yet to be sequenced and made available in GenBank. As shown in Fig. 3, the phylogram indicated that '*R. monacensis*' str. Humboldt and '*R. monacensis*' str. IrR/Munich form a unique clade. The phylogram also confirmed that '*R. monacensis*' str. Humboldt is closely related to '*R. monacensis*' str. IrR/Munich and that '*R. monacensis*' str. Humboldt, '*R. monacensis*' str. IrR/Munich, *R. buchneri* str. ISO7, and *R. massiliae* str. Mtut1T form a monophyletic group that is distinct from other spotted fever group rickettsiae (Fig. 3).

Based on MLST, MST, and phylogenetic tree constructions of this study, we have drawn a conclusion that the REIP isolate from *I. pacificus*

is '*R. monacensis*'. Our conclusion is supported by a recent genome-wide comparison among genomes of *Rickettsia* species using core genome alignment sequence identity (CGASI), a method using both length and sequence identity of genome for species classification. Using a CGASI cutoff of equal or greater than 96.8%, '*R. monacensis*' str. Humboldt from *I. pacificus* and '*R. monacensis*' str. IrR/Munich are classified as the same species. There are only two *Rickettsia* species in the spotted fever group based on CGASI reclassification. In addition to '*R. monacensis*' species, all other *Rickettsia* species in the spotted fever group are classified as the second *Rickettsia* species (Chung et al., 2018). Our conclusion that '*R. monacensis*' str. Humboldt and '*R. monacensis*' str. IrR/Munich are two different strains within the same species is also supported by CGASI since the core genome alignment sequence identity between '*R. monacensis*' str. Humboldt and '*R. monacensis*' str. IrR/Munich is 99.6% (Chung et al., 2018). In addition, the different genome content between the two strains is reflected by unique plasmids that exist in their genomes. Pulse-field gel electrophoresis and Southern blot identified pRM plasmid of '*R. monacensis*' str. IrR/Munich (Baldrige et al., 2007) (GenBank Accession Number EF564599), whereas genome sequencing of '*R. monacensis*' str. Humboldt revealed the presence of pREIP plasmid (KR611317). Although the size of pREIP (102 kb) differs dramatically with that of pRM (23.5 kb), parts of the pREIP share 99% nucleotide sequence identity with pRM (T. Kurtti, personal communication). However, the pREIP and pRM plasmids have low nucleotide homology with other rickettsial plasmids from other spotted fever group rickettsia, including *R. amblyommatis*, *R. buchneri*, *R. helvetica*, *R. hoogstraalii*, *R. massiliae*, and *R. peacockii* (Baldrige et al., 2008; Gillespie et al., 2012; T. Kurtti, personal communication).

'*R. monacensis*' was originally isolated in an *I. ricinus* tick in Munich, Germany (Simser et al., 2002). Later, '*R. monacensis*' infection in *I. ricinus* was identified all along Europe, including Hungary (Sreter-Lancz et al., 2006), Portugal (de Carvalho et al., 2008), Switzerland (Boretti et al., 2009), Sweden (Elfving et al., 2010), Italy (Corrain et al., 2012), The Netherlands (Koetsveld et al., 2016), Turkey (Gargili et al., 2012), Spain (Palomar et al., 2012), Poland (Rymaszewska and Piotrowski, 2013), Romania (Ionita et al., 2013), Croatia (Tijssen-Klasen et al., 2013), Slovakia (Spitalska et al., 2014), and Estonia (Katargina et al.,



**Fig. 3. Phylogenetic tree of concatenated nucleotide sequences of *dksA-xerC*, *mppA-purC*, and *rpmE-tRNA<sup>Met</sup>* intergenic spacers of ‘*Rickettsia monacensis*’ str. Humboldt.** The tree was generated using neighbor-joining distance method and 1000 bootstrap replicates. Bootstrap values > 50% are shown at the nodes. Bar, nucleotide distance. ‘*R. monacensis*’ str. Humboldt (or *Rickettsia* endosymbiont *Ixodes pacificus*) is represented at the top of the tree. Other selected sequences on the tree include: *R. aeschlimanii*, *R. africae*, *R. akari*, *R. australis*, *R. buchneri*, *R. conorii*, *R. felis*, *R. heilongjiangensis*, *R. helvetica*, *R. japonica*, *R. massiliae*, ‘*R. monacensis*’ str. IrR/Munich, *R. montanensis*, *R. parkeri*, *R. prowazekii*, *R. raoultii*, *R. rhipicephali*, *R. rickettsii*, *R. sibirica*, *R. slovacica*, and *R. typhi*. The accession numbers of genes of all *Rickettsia* species are listed in Supplemental Table.

2015). In addition to Europe, ‘*R. monacensis*’ has been reported in Africa, Asia, and South and Central America. In Africa, ‘*R. monacensis*’ was isolated from *I. ricinus* in Tunisia (Sfar et al., 2008) and Algeria (Dib et al., 2009). Also, the bacterium infection was documented in *I. nipponensis* ticks in Korea (Lee et al., 2013) and in *I. sinensis* ticks in China (Ye et al., 2014). In South and Central America, *I. boliviensis* and *Rhipicephalus sanguineus* sensu lato were tested positive for ‘*R. monacensis*’ in Nicaragua and Costa Rica (Springer et al., 2018; Troyo et al., 2014). The global distribution of ‘*R. monacensis*’ is, at least partially, attributed to migrating birds, which serve as possible dispersers of the bacterium (Elfving et al., 2010; Biernat et al., 2016). Many not yet isolated rickettsia in *Ixodes* ticks have been reported. Some of them are genetically close to ‘*R. monacensis*’ str. Humboldt and ‘*R. monacensis*’. For example, the *Rickettsia* species detected in *Amblyomma ovale* (Springer et al., 2018). Future isolation and characterization of the bacteria, especially genomic sequencing and CGASI analysis, are required to confirm that they belong to ‘*R. monacensis*’.

#### Declaration of interest

None.

#### Acknowledgements

This research was supported by National Institutes of Health grant 1R15AI099902-01. We would like to express our great appreciation to the veterinarian and staff at McKinleyville Animal Care Center, McKinleyville, CA, for donating partially engorged *Ixodes pacificus* females for our research.

#### Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.ttbdis.2019.04.017>.

## References

- Abdad, M.Y., Abdallah, R.A., Karkouri, K.E., Beye, M., Stenos, J., Owen, H., Unsworth, N., Robertson, I., Blacksell, S.D., Nguyen, T.T., Nappez, C., Raoult, D., Fenwick, S., Fournier, P.E., 2017. *Rickettsia gravesii* sp. nov.: a novel spotted fever group rickettsia in Western Australian *Amblyomma triguttatum triguttatum* ticks. *Int. J. Syst. Evol. Microbiol.* 67, 3156–3161.
- Bagheri, G., Lehner, J.D., Zhong, J., 2017. Enhanced detection of *Rickettsia* species in *Ixodes pacificus* using highly sensitive fluorescence in situ hybridization coupled with Tyramide Signal Amplification. *Ticks Tick. Dis.* 8, 915–921.
- Baldrige, G.D., Burkhardt, N.Y., Felsheim, R.F., Kurtti, T.J., Munderloh, U.G., 2007. Transposon insertion reveals pRM, a plasmid of *Rickettsia monacensis*. *Appl. Environ. Microbiol.* 73, 4984–4995.
- Baldrige, G.D., Burkhardt, N.Y., Felsheim, R.F., Kurtti, T.J., Munderloh, U.G., 2008. Plasmids of the pRM/pRF family occur in diverse *Rickettsia* species. *Appl. Environ. Microbiol.* 74, 645–652.
- Biernat, B., Stanczak, J., Michalik, J., Sikora, B., Cieniuch, S., 2016. *Rickettsia helvetica* and *R. monacensis* infections in immature *Ixodes ricinus* ticks derived from sylvatic passerine birds in west-central Poland. *Parasitol. Res.* 115, 3469–3477.
- Bodnar, J.L., Fitch, S., Rosati, A., Zhong, J., 2018. The *foIA* gene from the *Rickettsia* endosymbiont of *Ixodes pacificus* encodes a functional dihydrofolate reductase enzyme. *Ticks Tick. Dis.* 9, 443–449.
- Boretti, F.S., Perreten, A., Meli, M.L., Cattori, V., Willi, B., Wengi, N., Hornok, S., Honegger, H., Hegglin, D., Woelfel, R., Reusch, C.E., Lutz, H., Hofmann-Lehmann, R., 2009. Molecular investigations of *Rickettsia helvetica* infection in dogs, foxes, humans, and *Ixodes* ticks. *Appl. Environ. Microbiol.* 75, 3230–3237.
- Burgdorfer, W., Lane, R.S., Barbour, A.G., Gresbrink, R.A., Anderson, J.R., 1985. The western black-legged tick, *Ixodes pacificus*: a vector of *Borrelia burgdorferi*. *Am. J. Trop. Med. Hyg.* 34, 925–930.
- Cazorla, C., Socolovschi, C., Jensenius, M., Parola, P., 2008. Tick-borne diseases: tick-borne spotted fever rickettsioses in Africa. *Infect. Dis. Clin. North Am.* 22, 531–544 ix-x.
- Cheng, D., Lane, R.S., Moore, B.D., Zhong, J., 2013a. Host blood meal-dependent growth ensures transovarial transmission and transstadial passage of *Rickettsia* sp. phylotype G021 in the western black-legged tick (*Ixodes pacificus*). *Ticks Tick. Dis.* 4, 421–426.
- Cheng, D., Vigil, K., Schanes, P., Brown, R.N., Zhong, J., 2013b. Prevalence and burden of two rickettsial phylotypes (G021 and G022) in *Ixodes pacificus* from California by real-time quantitative PCR. *Ticks Tick. Dis.* 4, 280–287.
- Chung, M., Munro, J.B., Dunning Hotopp, J.C., 2018. Using core genome alignments to assign bacterial species. *bioRxiv*. <https://doi.org/10.1101/328021>.
- Corrain, R., Drigo, M., Fenati, M., Menandro, M.L., Mondin, A., Pasotto, D., Martini, M., 2012. Study on ticks and tick-borne zoonoses in public parks in Italy. *Zoonoses Public Health* 59, 468–476.
- de Carvalho, I.L., Milhano, N., Santos, A.S., Almeida, V., Barros, S.C., De Sousa, R., Nuncio, M.S., 2008. Detection of *Borrelia lusitanae*, *Rickettsia* sp. IRS3, *Rickettsia monacensis*, and *Anaplasma phagocytophilum* in *Ixodes ricinus* collected in Madeira Island, Portugal. *Vector Borne Zoonotic Dis.* 8, 575–579.
- Dib, L., Bitam, I., Bensouilah, M., Parola, P., Raoult, D., 2009. First description of *Rickettsia monacensis* in *Ixodes ricinus* Algeria. *Clin. Microbiol. Infect.* 15 (Suppl 2), 261–262.
- Elfving, K., Olsen, B., Bergstrom, S., Waldenstrom, J., Lundkvist, A., Sjostedt, A., Mejlom, H., Nilsson, K., 2010. Dissemination of spotted fever rickettsia agents in Europe by migrating birds. *PLoS One* 5, e8572.
- Foley, J.E., Nieto, N.C., Adjemian, J., Dabritz, H., Brown, R.N., 2008. *Anaplasma phagocytophilum* infection in small mammal hosts of *Ixodes* ticks, western United States. *Emerg. Infect. Dis.* 14, 1147–1150.
- Fournier, P.E., Raoult, D., 2007. Identification of rickettsial isolates at the species level using multi-spacer typing. *BMC Microbiol.* 7, 72.
- Fournier, P.E., Dumler, J.S., Greub, G., Zhang, J., Wu, Y., Raoult, D., 2003. Gene sequence-based criteria for identification of new rickettsia isolates and description of *Rickettsia heilongjiangensis* sp. nov. *J. Clin. Microbiol.* 41, 5456–5465.
- Fournier, P.E., Zhu, Y., Ogata, H., Raoult, D., 2004. Use of highly variable intergenic spacer sequences for multispacer typing of *Rickettsia conorii* strains. *J. Clin. Microbiol.* 42, 5757–5766.
- Gargili, A., Palomar, A.M., Midilli, K., Portillo, A., Kar, S., Oteo, J.A., 2012. *Rickettsia* species in ticks removed from humans in Istanbul, Turkey. *Vector Borne Zoonotic Dis.* 12, 938–941.
- Gillespie, J.J., Joardar, V., Williams, K.P., Driscoll, T., Hostetler, J.B., Nordberg, E., Shukla, M., Walenz, B., Hill, C.A., Nene, V.M., Azad, A.F., Sobral, B.W., Caler, E., 2012. A *Rickettsia* genome overrun by mobile genetic elements provides insight into the acquisition of genes characteristic of an obligate intracellular lifestyle. *J. Bacteriol.* 194, 376–394.
- Gouy, M., Guindon, S., Gascuel, O., 2010. SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* 27, 221–224.
- Hardstone Yoshimizu, M., Billeter, S.A., 2018. Suspected and confirmed vector-borne rickettsioses of north america associated with human diseases. *Trop. Med. Infect. Dis.* 3 (1), 2.
- Hughes, L.E., Clifford, C.M., Gresbrink, R., Thomas, L.A., Keirans, J.E., 1976. Isolation of a spotted fever group rickettsia from the Pacific Coast tick, *Ixodes pacificus*, in Oregon. *Am. J. Trop. Med. Hyg.* 25, 513–516.
- Hunter, D.J., Torkelson, J.L., Bodnar, J., Mortazavi, B., Laurent, T., Deason, J., Thephavongsa, K., Zhong, J., 2015. The *Rickettsia* endosymbiont of *Ixodes pacificus* contains all the genes of de novo folate biosynthesis. *PLoS One* 10, e0144552.
- Ionita, M., Mitrea, I.L., Pfister, K., Hamel, D., Silaghi, C., 2013. Molecular evidence for bacterial and protozoan pathogens in hard ticks from Romania. *Vet. Parasitol.* 196, 71–76.
- Katargina, O., Geller, J., Ivanova, A., Varv, K., Tefanova, V., Vene, S., Lundkvist, Å., Golovljova, I., 2015. Detection and identification of *Rickettsia* species in *Ixodes* tick populations from Estonia. *Ticks Tick. Dis.* 6, 689–694.
- Koetsveld, J., Tijssse-Klasen, E., Herremans, T., Hovius, J.W., Sprong, H., 2016. Serological and molecular evidence for spotted fever group *Rickettsia* and *Borrelia burgdorferi* sensu lato co-infections in the Netherlands. *Ticks Tick. Dis.* 7, 371–377.
- Kurlovs, A.H., Li, J., Cheng, D., Zhong, J., 2014. *Ixodes pacificus* ticks maintain embryogenesis and egg hatching after antibiotic treatment of *Rickettsia* endosymbiont. *PLoS One* 9, e104815.
- Kurtti, T.J., Munderloh, U.G., Andreadis, T.G., Magnarelli, L.A., Mather, T.N., 1996. Tick cell culture isolation of an intracellular prokaryote from the tick *Ixodes scapularis*. *J. Invertebr. Pathol.* 67, 318–321.
- Kurtti, T.J., Simser, J.A., Baldrige, G.D., Palmer, A.T., Munderloh, U.G., 2005. Factors influencing in vitro infectivity and growth of *Rickettsia peacockii* (Rickettsiales: Rickettsiaceae), an endosymbiont of the Rocky Mountain wood tick, *Dermacentor andersoni* (Acari, Ixodidae). *J. Invertebr. Pathol.* 90, 177–186.
- Kurtti, T.J., Felsheim, R.F., Burkhardt, N.Y., Oliver, J.D., Heu, C.C., Munderloh, U.G., 2015. *Rickettsia buchneri* sp. nov., a rickettsial endosymbiont of the blacklegged tick *Ixodes scapularis*. *Int. J. Syst. Evol. Microbiol.* 65, 965–970.
- Lee, K.M., Choi, Y.J., Shin, S.H., Choi, M.K., Song, H.J., Kim, H.C., Klein, T.A., Richards, A.L., Park, K.H., Jang, W.J., 2013. Spotted fever group rickettsia closely related to *Rickettsia monacensis* isolated from ticks in South Jeolla province, Korea. *Microbiol. Immunol.* 57, 487–495.
- Munderloh, U.G., Kurtti, T.J., 1989. Formulation of medium for tick cell culture. *Exp. Appl. Acarol.* 7, 219–229.
- Munderloh, U.G., Liu, Y., Wang, M., Chen, C., Kurtti, T.J., 1994. Establishment, maintenance and description of cell lines from the tick *Ixodes scapularis*. *J. Parasitol.* 80, 533–543.
- Oliver, J.D., Burkhardt, N.Y., Felsheim, R.F., Kurtti, T.J., Munderloh, U.G., 2014. Motility characteristics are altered for *Rickettsia bellii* transferred to overexpress a heterologous *rickA* gene. *Appl. Environ. Microbiol.* 80, 1170–1176.
- Palomar, A.M., Santibanez, P., Mazuelas, D., Roncero, L., Santibanez, S., Portillo, A., Oteo, J.A., 2012. Role of birds in dispersal of etiologic agents of tick-borne zoonoses, Spain, 2009. *Emerg. Infect. Dis.* 18, 1188–1191.
- Parola, P., Paddock, C.D., Socolovschi, C., Labruna, M.B., Mediannikov, O., Kernif, T., Abdad, M.Y., Stenos, J., Bitam, I., Fournier, P.E., Raoult, D., 2013. Update on tick-borne rickettsioses around the world: a geographic approach. *Clin. Microbiol. Rev.* 26, 657–702.
- Phan, J.N., Lu, C.R., Bender, W.G., Smoak 3rd, R.M., Zhong, J., 2011. Molecular detection and identification of *Rickettsia* species in *Ixodes pacificus* in California. *Vector Borne Zoonotic Dis.* 11, 957–961.
- Phillip, R.N., Casper, E.A., Burgdorfer, W., Gerloff, R.K., Hughes, L.E., Bell, E.J., 1978. Serologic typing of rickettsiae of the spotted fever group by microimmunofluorescence. *J. Immunol.* 121, 1961–1968.
- Phillip, R.N., Lane, R.S., Casper, E.A., 1981. Serotypes of tick-borne spotted fever group rickettsiae from western California. *Am. J. Trop. Med. Hyg.* 30, 722–727.
- Piesman, J., Clark, K.L., Dolan, M.C., Happ, C.M., Burkot, T.R., 1999. Geographic survey of vector ticks (*Ixodes scapularis* and *Ixodes pacificus*) for infection with the Lyme disease spirochete, *Borrelia burgdorferi*. *J. Vector Ecol.* 24, 91–98.
- Richards, A.L., 2012. Worldwide detection and identification of new and old rickettsiae and rickettsial diseases. *FEMS Immunol. Med. Microbiol.* 64, 107–110.
- Rymaszewska, A., Piotrowski, M., 2013. Use of DNA sequences for *Rickettsia* identification in *Ixodes ricinus* ticks: the first detection of *Rickettsia monacensis* in Poland. *Microbes Infect.* 15, 140–146.
- Sfar, N., M'Ghirbi, Y., Letaief, A., Parola, P., Bouattour, A., Raoult, D., 2008. First report of *Rickettsia monacensis* and *Rickettsia helvetica* from Tunisia. *Ann. Trop. Med. Parasitol.* 102, 561–564.
- Simser, J.A., Palmer, A.T., Fingerle, V., Wilske, B., Kurtti, T.J., Munderloh, U.G., 2002. *Rickettsia monacensis* sp. nov., a spotted fever group *Rickettsia*, from ticks (*Ixodes ricinus*) collected in a European city park. *Appl. Environ. Microbiol.* 68, 4559–4566.
- Spitalska, E., Boldis, V., Verdakova, M., Selyemova, D., Rusnakova Taragelova, V., 2014. Rickettsial infection in *Ixodes ricinus* ticks in urban and natural habitats of Slovakia. *Ticks Tick. Dis.* 5, 161–165.
- Springer, A., Montenegro, V.M., Schicht, S., Wölfel, S., Schaper, S.R., Chitimia-Dobler, L., Siebert, S., Strube, C., 2018. Detection of *Rickettsia monacensis* and *Rickettsia amblyommatis* in ticks collected from dogs in Costa Rica and Nicaragua. *Ticks Tick. Dis.* 9, 1565–1572.
- Sreter-Lancz, Z., Szell, Z., Kovacs, G., Egyed, L., Marialigeti, K., Sreter, T., 2006. Rickettsiae of the spotted-fever group in ixodid ticks from Hungary: identification of a new genotype (*Candidatus Rickettsia kotlaniai*). *Ann. Trop. Med. Parasitol.* 100, 229–236.
- Tijssse-Klasen, E., Sprong, H., Pandak, N., 2013. Co-infection of *Borrelia burgdorferi* sensu lato and *Rickettsia* species in ticks and in an erythema migrans patient. *Parasit. Vectors* 6, 347.
- Troyo, A., Moreira-Soto, A., Carranza, M., Calderón-Arguedas, O., Hun, L., Taylor, L., 2014. Detection of an undescribed *Rickettsia* sp. in *Ixodes boliviensis* from Costa Rica. *Ticks Tick. Borne Dis.* 5, 883–886.
- Ye, X., Sun, Y., Ju, W., Wang, X., Cao, W., Wu, M., 2014. Vector competence of the tick *Ixodes sinensis* (Acari: Ixodidae) for *Rickettsia monacensis*. *Parasit. Vectors* 7, 512.
- Zhu, Y., Fournier, P.E., Ereemeeva, M., Raoult, D., 2005a. Proposal to create subspecies of *Rickettsia conorii* based on multi-locus sequence typing and an emended description of *Rickettsia conorii*. *BMC Microbiol.* 5, 11.
- Zhu, Y., Fournier, P.E., Ogata, H., Raoult, D., 2005b. Multispacer typing of *Rickettsia prowazekii* enabling epidemiological studies of epidemic typhus. *J. Clin. Microbiol.* 43, 4708–4712.
- Znazen, A., Khrouf, F., Elleuch, N., Lahiani, D., Marrekchi, C., M'Ghirbi, Y., Ben Jemaa, M., Bouattour, A., Hammami, A., 2013. Multispacer typing of *Rickettsia* isolates from humans and ticks in Tunisia revealing new genotypes. *Parasit. Vectors* 6, 367.