




ELSEVIER

Contents lists available at ScienceDirect

Ticks and Tick-borne Diseases

journal homepage: www.elsevier.com/locate/ttbdis

Original article

First detection of tick-borne encephalitis virus in *Ixodes ricinus* ticks and their rodent hosts in Moscow, Russia

Marat Makenov^{a,*}, Lyudmila Karan^a, Natalia Shashina^b, Marina Akhmetshina^b, Olga Zhurenkova^a, Ivan Kholodilov^c, Galina Karganova^{c,d}, Nina Smirnova^{a,e}, Yana Grigoreva^a, Yanina Yankovskaya^f, Marina Fyodorova^a

^a Central Research Institute of Epidemiology, Novogireevskaya st 3-A, 415, Moscow, 111123, Russia

^b Scientific Research Disinfectology Institute, Nauchnyy proezd st. 18, Moscow, 117246, Russia

^c Chumakov Institute of Poliomyelitis and Viral Encephalites (FSBSI "Chumakov FSC R&D IBP RAS), prem. 8, k.17, pos. Institut Poliomyelita, poselenie Moskovskiy, Moscow, 108819, Russia

^d Institute for Translational Medicine and Biotechnology, Sechenov University, Bolshaya Pirogovskaya st, 2, page 4, room 106, Moscow, 119991, Russia

^e Lomonosov Moscow State University, Leninskie Gory st. 1-12, MSU, Faculty of Biology, Moscow, 119991, Russia

^f Pirogov Russian National Research Medical University, Ostrovityanova st. 1, Moscow, 117997, Russia

ARTICLE INFO

Keywords:

Tick-borne encephalitis virus
Ixodes ricinus
Borrelia burgdorferi sensu lato
Vector-borne diseases
Small mammals

ABSTRACT

Here, we report the first confirmed autochthonous tick-borne encephalitis case diagnosed in Moscow in 2016 and describe the detection of tick-borne encephalitis virus (TBEV) in ticks and small mammals in a Moscow park.

The paper includes data from two patients who were bitten by TBEV-infected ticks in Moscow city; one of these cases led to the development of the meningeal form of TBE. Both TBEV-infected ticks attacked patients in the same area. We collected ticks and trapped small mammals in this area in 2017. All samples were screened for the presence of pathogens causing tick-borne diseases by PCR. The TBEV-positive ticks and small mammals' tissue samples were subjected to virus isolation. The sequencing of the complete polyprotein gene of the positive samples was performed.

A total of 227 questing ticks were collected. TBEV was detected in five specimens of *Ixodes ricinus*. We trapped 44 small mammals, mainly bank voles (*Myodes glareolus*) and pygmy field mice (*Apodemus uralensis*). Two samples of brain tissue from bank voles yielded a positive signal in RT-PCR for TBEV. We obtained six virus isolates from the ticks and brain tissue of a bank vole. Complete genome sequencing showed that the obtained isolates belong to the European subtype and have low diversity with sequence identities as high as 99.9%. GPS tracking showed that the maximum distance between the exact locations where the TBEV-positive ticks were collected was 185 m. We assume that the forest park had been free of TBEV and that the virus was recently introduced.

1. Introduction

Tick-borne encephalitis virus (TBEV) is a flavivirus that is transmitted by the bite of infected hard-bodied ticks. The geographic range of TBEV is mainly in Eurasia spanning Japan to the Russian far east towards Central Europe (Lindquist and Vapalahti, 2008; Süss, 2011; Zlobin, 2010). The prevalence of the virus in ticks, as well as its distribution, can increase based on the different yearly factors governing the different parts of the vectors' and hosts' ranges (Korenberg, 2009). In particular, changes in the boundaries of the TBE risk areas were described in Germany (Süss et al., 2004) and Russia (Tokarevich et al., 2011). Furthermore, the detection of new TBEV foci has been reported

in Sweden (Haglund, 2002; Jaenson et al., 2012), Norway (Csango et al., 2004; Haglund, 2002; Skarpaas et al., 2006, 2004), The Netherlands (De Graaf et al., 2016; Dekker et al., 2019), and Austria (Heinz et al., 2015). There is evidence regarding the reactivation of old foci of TBEV, which for a long time (decades) have been inactive, thus turning the places into a risk area (Frimmel et al., 2014; Jääskeläinen et al., 2006; Kristiansen, 2002).

In Russia, out of 85 regions, 47 belong to TBE risk areas ("On the list of endemic territories for viral tick-borne encephalitis in 2015," 2015 On the list of endemic territories for viral tick-borne encephalitis, 2015 On the list of endemic territories for viral tick-borne encephalitis in 2015," 2015), but Moscow and most districts of the Moscow Oblast

* Corresponding author.

E-mail address: makenov@cmd.su (M. Makenov).

<https://doi.org/10.1016/j.ttbdis.2019.101265>

Received 21 December 2018; Received in revised form 10 July 2019; Accepted 28 July 2019

Available online 30 July 2019

1877-959X/© 2019 Elsevier GmbH. All rights reserved.




Fig. 1. Tick-borne encephalitis-endemic areas in Russia (Based on (On the list of endemic territories for viral tick-borne encephalitis in 2015, 2015On the list of endemic territories for viral tick-borne encephalitis, 2015On the list of endemic territories for viral tick-borne encephalitis in 2015, 2015)).

are not endemic for TBEV. Only two districts in the north of the Moscow Oblast are included in the TBE risk area (“On the list of endemic territories for viral tick-borne encephalitis in 2015,” 2015On the list of endemic territories for viral tick-borne encephalitis, 2015On the list of endemic territories for viral tick-borne encephalitis in 2015,” 2015; Shevtsova et al., 2008), but local cases of the disease have not been recorded there for several decades. The Moscow Oblast borders two TBE endemic regions in the north and northwest with the Tver Oblast and Yaroslavl Oblast, respectively (Fig. 1). The most recent, confirmed cases of TBE in the Moscow Oblast are from 1948–1954. At the time of the outbreak, at least 150 patients were documented in two regions lying to the northwest of the Moscow Oblast (Drozdzov, 1956). In the city of Moscow, local cases of the disease were not registered. The city has experienced a significant population increase in the past, growing due to the expansion of the surrounding territories. Some of the new urban areas were not developed, and they were preserved as forest parks. The local fauna of such forest parks has remained and includes small mammals and ticks (Uspensky, 2017). Surveys of the forest parks of Moscow showed the presence of *Ixodes ricinus* in the majority of the city’s parks (Shashina and Germant, 2015). A recent study of the ticks collected in Moscow’s parks revealed the prevalence of *Borrelia burgdorferi* s.l., *Anaplasma phagocytophilum* and *Ehrlichia* sp., but TBEV was not detected (Shashina and Germant, 2015). In Moscow, 5–14 clinical TBE cases have been recorded annually, but none of them have been autochthonous (On the incidence of tick-borne borreliosis in Moscow in 2014, 2015On the incidence of tick-borne borreliosis in Moscow, 2015On the incidence of tick-borne borreliosis in Moscow in 2014, 2015; Yankovskaya et al., 2017).

In 2016, TBE was diagnosed in a patient who was bitten by a tick in a Moscow park. It was the first documented autochthonous TBE case diagnosed in Moscow. Therefore, we aimed to report this case and to

study the prevalence of TBEV among ixodid ticks and small mammals in an urban forest park. In addition, this paper seeks to identify the origin of the discovered TBE foci and to determine whether the described cases are due to the reactivation of formerly active foci or if there is evidence of introduction of TBEV from a new territory.

2. Methods

2.1. Patients

The paper includes data from two patients who were bitten by TBEV-infected ticks within the Moscow city limits. We interviewed both patients and considered their medical records. For the patient described in Case-2, blood samples were collected on the 17th and 41st days after the tick bite to test for anti-TBEV IgM and IgG.

2.2. Research area

We obtained first two TBEV-infected ticks from the same area in the western part of the city. Hence, we focused on forest parks from this area. These parks adjoin the forests of the Moscow Oblast and the Moscow Ring Road (a large highway with busy round-the-clock traffic) separates them. We divided the research area into three study sites (Fig. 2). Site-1 was the main site used for field research since in this forest park, a TBEV-infected tick attacked a patient in July 2017 (see Case-2 below). Additionally, we selected Site-2 and Site-3 north and south of Site-1, respectively, to check the areas in the city boundaries. The forest parks on Site-1, Site-2, and Site-3 are free of regulation and management services (such as grass mowing, deratization, spraying acaricides, etc.). The areas of these sites are 104 ha, 218 ha and 179 ha, respectively. The local people actively use these parks for recreational



Fig. 2. Study area with depicted locations of tick collection and small mammal trapping (A). The inserted maps represent the location of the study sites (B) and the administrative boundary of Moscow city (C).

activities.

2.3. Tick collection

Ticks were collected through flagging on regular transects in May 2017 (one sample session) and in August–September 2017 (2–3 times per week). Any tick collected was placed in a separate tube. We recorded the GPS coordinates of each tick collection site. Our sampling efforts were focused on Site-1, where we conducted nine flagging assays in August covering a total distance of 43.3 km during flagging. At Site-2 and Site-3, we carried out only one flagging assay. Due to a lack of sampling efforts at Site-2 and Site-3, we estimated tick abundance only for Site-1. Each flagging assay was performed by 2–4 collectors. Based on their GPS tracks, we calculated the number of ticks per kilometer for each collector. Subsequently, we calculated the mean number of ticks collected per kilometer of flagging for a ten-day period. All tick specimens were identified to stage and species by morphological characteristics (Filippova, 1977). Additionally, we performed molecular species identification (see below for details).

2.4. Small mammals

Small mammals were captured in October 2017 using live traps (Shchipanov, 1987) baited with sunflower oil and oat. The traps were set up for four nights placed at 5 m intervals along two linear transects. The total capture effort was 100 traps per night. We chose the location for small mammals trapping based on the GPS tracking of the TBEV-positive tick collection sites. Small mammals capture and handling procedures were approved by the Ethical Committee at Scientific Research Disinfectology Institute (Moscow, Russia).

2.5. Extraction, Molecular detection and genotyping of pathogens

All the collected ticks were tested for the presence of tick-borne pathogens by a real-time PCR assay. Initially, the ticks were washed consistently in 70% alcohol and 0.15 M NaCl solution, and then individual ticks were homogenized with TissueLyser LT (Qiagen, Hilden, Germany) in 300 μ l of medium 199 with Earle's salts (Chumakov Institute of Poliomyelitis and Viral Encephalitides, Moscow, Russia). Thereafter, DNA/RNA was extracted from 100 μ l of the tick suspension using a commercial kit AmpliSens RIBO-prep kit (Central Research Institute of Epidemiology, Moscow, Russia) following the manufacturer's instructions. PCR-based detection of TBEV, and *B. burgdorferi* s.l., *A. phagocytophilum*, *E. chaffeensis*, and *E. muris* was performed using a commercial multiplex PCR kit (AmpliSens TBEV, *B. burgdorferi* s.l., *A. phagocytophilum*, *E. chaffeensis*/*E. muris*-FL; Central Research Institute of Epidemiology, Moscow, Russia) according to the manufacturer's instructions. Each run included a negative control and the mix of the five positive recombinant DNA controls of the TBEV 5'UTR-C gene fragment; *B. burgdorferi* s.l. and *E. chaffeensis*, *E. muris* 16S gene fragments, *A. phagocytophilum* msp2 gene fragment and RNA internal control with 10^5 copies/ml. The RNA template was reverse transcribed using Thermo Scientific Maxima Reverse Transcriptase (Thermo Scientific, Schwerte, Germany).

To check the species diagnoses based on the morphological characteristics, we performed qPCR (Table A1). Furthermore, for tick identification, we sequenced the *ITS-2* gene fragments and *COI* gene fragments (Table A1). PCR products of the appropriate length were separated by electrophoresis on a 1.5% agarose gel supplemented with ethidium bromide. Amplicon purification was performed using the RIBO-prep extraction kit (Central Research Institute of Epidemiology, Moscow, Russia).

2.6. Virus isolation

The TBEV-positive ticks and small mammals' tissue samples were subjected to virus isolation. For this, we used the pig embryo kidney (PEK) cell line and suckling mice. The PEK cell line was maintained at 37 °C in medium 199 (Chumakov Institute of Poliomyelitis and Viral Encephalitis Moscow, Russia,) supplemented with 5% fetal bovine serum (FBS) (Gibco/Thermo Fisher Scientific, Waltham, MA, USA). The suckling mice (FSBI, Scientific Center of Biomedical Technologies, Stolbovaya Branch, Russia) were infected intracerebrally with the supernatant obtained from the PCR-positive tick and tissue sample suspensions. The mice were euthanized at the appearance of signs of the disease or on the 14th day of the experiment. The mice were maintained according to the international guidelines for animal husbandry, including the recommendations of CIOMS, 1985 and the FELASA Working Group Report, 1996–1997.

2.7. 50% plaque reduction neutralization test (PRNT50)

We tested rodents' serum with neutralizing antibody for TBEV using a 50% plaque reduction neutralization test (PRNT50). The analysis was performed as described earlier (Pripuzova et al., 2013). For this test, we used strain MOS-152-T-2017, isolated from the tick that bit the patient described in Case-2. Briefly, sequential dilutions of the sera samples were prepared in 199 medium on Earle solution with the addition of 2% FBS (Gibco/Thermo Fisher Scientific, Waltham, MA, USA). An equal volume of viral suspension containing 40–70 plaque forming units was added to the sera dilution. Virus-positive sera were incubated at 37 °C for 1 h, added to PEK cell monolayers, and incubated for one hour for virus adsorption. Then, each well was overlaid with 1% Bacto-agar (Difco Laboratories, Detroit, MI, USA) on Earle solution containing 7.5% FBS and 0.015% neutral red. Every experiment included appropriate controls, i.e., negative and positive murine sera with known titers. The antibody titer was calculated according to the modified Reed and Muench method (Reed and Muench, 1938).

2.8. The sequencing

We used RNA isolated from tick suspensions and rodents' brain tissue for sequencing. The purified PCR products were sequenced bidirectionally using BigDye Terminator v1.1 Cycle Sequencing kit (Thermo Fisher Scientific, Austin, TX, USA) on an Applied Biosystems 3500xL Genetic Analyzer (Applied Biosystems, Foster City, CA, USA). The primers used in sequencing are shown in Table A1. The sequences obtained were deposited in NCBI GenBank under the following accession numbers: TBEV MH663426-MH663428 (complete polyprotein gene, 9521 bp) and MH663429-MH663430 (NS1 protein gene, 1054 bp). The partial sequence alignment and phylogenetic tree construction were performed using the software MEGA7.

2.9. Data analysis

We conducted the data analysis in R using the package PropCIs (Scherer, 2014). The sites where the questing ticks were collected and the small mammals were trapped were mapped using Quantum GIS free software. The nucleotide sequences were aligned, compared, and analyzed using MEGA7, ClustalW, and BLAST. The phylogenetic reconstruction was inferred using the neighbor-joining method with 1000 bootstrap replications. The evolutionary distances were computed using the Kimura 2-parameter method and are presented in units of the number of base substitutions per site.

3. Results

3.1. Patient cases

3.1.1. Case 1

A 43-year-old man was bitten by a tick in a forest park in the west of Moscow on August 8, 2016. The patient stated that he had not had any other tick bites and had not visited TBE-endemic areas. The patient also declined TBE vaccination. On August 13, fever, headache, and myalgia developed in the patient. Erythema migrans were absent at the site of the tick bite. The patient started antibiotic treatment (doxycycline) on an outpatient basis after consulting a doctor. On August 29, the patient was hospitalized with high fever (temperature 40.0 °C). ELISA tests on anti-TBE IgM and IgG showed positive results. The diagnosis was Tick-borne encephalitis, meningeal form.

3.1.2. Case 2

On July 25, 2017, a 27-year-old woman reported a tick bite after walking in a forest park in the west of Moscow. The patient brought the tick to a commercial laboratory to test. RT-PCR yielded a positive result for TBEV. The patient had not been previously vaccinated against TBE. On July 27, the patient received prophylactic immunoglobulin therapy (1:80, 6 ml). Two ELISAs of serum samples for anti-TBE IgM and IgG conducted on the 17th and 41st days after the tick bite showed negative results. No symptoms of acute infection manifested.

3.2. Questing ticks

The questing ticks field assays resulted in the collection of 227 ticks from vegetation in the study area, and 225 specimens out of the total collection were identified morphologically and through qPCR to be *I. ricinus* (Table 1). Only one tick was identified as *Ixodes persulcatus* (female), and one was identified as *Dermacentor reticulatus* (male). The mean abundance of adult *I. ricinus* at Site-1 varied in August from 1.6 to 3.0 ticks/km (Table 2). We did not estimate the abundance of ticks at Site-2 and Site-3.

TBEV was detected in five specimens, including two adult males, two adult females, and one nymph of *I. ricinus* (Table 3). All TBEV-positive ticks were collected at Site-1. We determined the exact coordinates of the collection site for the three TBEV-positive ticks. Mapping showed that all the TBEV-positive ticks were obtained at the same location along the small river on the western part of the park (Fig. 2). The maximum distance between the TBEV-positive ticks was 185 m. The mean prevalence of *B. burgdorferi* s.l. in ticks at all the studied sites was 32.4%, the difference between Site-1 and Site-2 was not significant ($\chi^2 = 1.58$, $df = 1$, $p = 0.21$).

3.3. Small mammals

Out of the 44 rodents that were live-trapped, 14 of these were released since they were not fully mature (individuals without previous breeding experience) and had no ticks on them (Table 4.). At Site-1, we trapped small mammals of four species; the sample included mainly bank voles (*Myodes glareolus*) and pygmy field mice (*Apodemus uralensis*). From the trapped animals, we collected larvae of *I. ricinus* and larvae and adult females of *Ixodes trianguliceps* (Table 4). In addition to

Table 1
Questing ixodid ticks collected in the study area.

Location	<i>Ixodes ricinus</i>		<i>Ixodes persulcatus</i>	<i>Dermacentor reticulatus</i>	Total
	Adults	Nymphs			
Site-1	106	42	1	0	149
Site-2	47	10	0	1	58
Site-3	17	3	0	0	20

Table 2
The abundance of *Ixodes ricinus* at Site-1, August 2017.

Ten-day periods	Number of flagging assays	Total distance of flagging assays, km	Tick abundance, ticks/km (mean ± SD)		
			Adults	Nymphs	Total
1	3	16.0	1.6 ± 1.1	0.2 ± 0.3	1.8 ± 1.4
2	3	14.2	1.7 ± 0.8	1.2 ± 0.7	3.0 ± 0.6
3	3	13.1	3.0 ± 2.2	1.4 ± 1.3	4.4 ± 2.8

Table 3
Prevalence of TBEV and *Borrelia burgdorferi* s.l in collected *Ixodes ricinus* ticks.

Stage	Total	Number of PCR-positive samples, Frequency (% – CI 95%)	
		TBEV	<i>B. burgdorferi</i> s.l
Site-1	149	5 (3.4% – 1.1–7.7%)	54 (36.2% – 28.5–44.5%)
Adults:	107	4 (3.7% – 1.0–9.3%)	42 (39.3% – 30.0–49.2%)
Female	52	2 (3.8% – 0.5–13.2%)	22 (42.3% – 28.8–56.8%)
Male	55	2 (3.6% – 0.4–12.5%)	20 (36.4% – 23.8–50.4%)
Nymph	42	1 (2.4% – 0.1–12.6%)	12 (28.6% – 15.7–44.6%)
Site-2	58	0 (0.0%)	15 (25.9% – 15.3–39.0%)
Adults:	48	0 (0.0%)	15 (31.3% – 18.7–46.3%)
Female	23	0 (0.0%)	7 (30.4% – 13.2–52.9%)
Male	25	0 (0.0%)	8 (32.0% – 14.9–53.5%)
Nymph	10	0 (0.0%)	0 (0.0%)
Site-3	20	0 (0.0%)	7 (35.0% – 15.4–59.2%)
Adults:	17	0 (0.0%)	4 (23.5% – 6.8–49.9%)
Female	14	0 (0.0%)	3 (21.4% – 4.7–50.8%)
Male	3	0 (0.0%)	1 (33.3% – 0.8–90.6%)
Nymph	3	0 (0.0%)	3 (100.0%)

the ixodid ticks, we found a large number of chiggers identified as *Hirsutiella zachvatkini*.

We tested samples of small mammalian tissues for the presence of TBEV RNA and DNA of *B. burgdorferi* s.l.. Two specimens of brain tissue of bank vole (*M. glareolus*) yielded a positive signal in RT-PCR for TBEV (Table 6). PRNT50 showed that 24 of the 28 studied rodents had neutralizing antibodies for TBEV (Table 6). The distance between TBEV-positive rodent trapping locations was 245 m (Fig. 2). Additionally, we found DNA of *B. burgdorferi* s.l. in 15 rodents (Table 5).

3.4. Virus characteristics

Six isolates of TBEV were obtained, including an isolate from the tick that bit the patient described in Case-2, four isolates from questing ticks, and one isolate from the brain tissue of a bank vole. We genetically characterized three TBEV-positive tick samples by sequencing their complete polyprotein gene. For another two tick samples, we sequenced the NS1 protein gene. The obtained isolates belong to the European subtype (Fig. 3). The sequences are most similar to the strain Ljubljana I (U27494), which was isolated from a patient in Slovenia in 1992 (Wallner et al., 1995). The TBEV complete polyprotein gene sequences at Moscow Forest Park had low diversity with sequence identities as high as 99.9%; the genetic distance with other strains of

Table 4
The number of live-trapped small mammals and the number of ixodid ticks collected from the investigated animals.

	Total	<i>Myodes glareolus</i>	<i>Apodemus uralensis</i>	<i>Apodemus agrarius</i>	<i>Sorex aureus</i>
Small mammals					
Trapped	44	20	21	1	2
Collected for further work	30	20	9	1	0
Prevalence of ticks collected from small mammals, the number and % (CI95%)					
<i>Ixodes ricinus</i> larvae	3 10.0% (2.1–26.5%)	1 5.0% (0.1–24.9%)	1 11.1% (0.3–48.2%)	1 100.0%	–
<i>Ixodes trianguliceps</i> larvae	11 36.7% (19.9–56.1%)	3 15.0% (3.2–37.9%)	8 88.9% (51.8–99.7%)	0 0.0%	–
<i>Ixodes trianguliceps</i> female	1 3.3% (0.1–17.2%)	0 0.0%	1 11.9% (0.3–48.2%)	0 0.0%	–

Table 5
Prevalence of vector-borne and zoonotic pathogens in the tissue samples of the investigated small mammals.

Pathogens	Number of PCR-positive samples, Frequency (% – CI 95%)	
	<i>Myodes glareolus</i>	<i>Apodemus uralensis</i>
The total number of tested animals	20	9
TBEV	2 (10.0% – 1.2–31.7%)	0
<i>Borrelia burgdorferi</i> s.l.	8 (40.0% – 19.1–63.9%)	7 (77.8% – 40.0–97.2%)

Table 6
The number of animals with the corresponding titers of neutralizing antibodies for TBEV (based on a 50% plaque reduction neutralization test of the rodents' serum).

Species / stage classes	Total number	The titer of neutralizing antibodies for TBEV				
		< 1:10*	1:10 – 1:50	1:50 – 1:100	1:100 – 1:150	> 1:150
<i>Myodes glareolus</i>	20	1	4	5	2	8
Juvenile	2	0	0	1	0	1
Subadult	9	1	2	2	0	4
Adult	9	0	2	2	2	3
<i>Apodemus uralensis</i>	8	3	3	0	1	1
Subadult	6	2	3	0	1	0
Adult	2	1	0	0	0	1
Total	28	4	7	5	3	9

* We consider titers less than 1:10 as negative results.

European subtype was approximately 2.0–2.9% (Fig. 3).

4. Discussion

One important research question concerns the origin of the virus in the urban forest park. The answer to this question could shed light on the possibility of introduction of the virus to other parks, cities, regions, etc. As a result, we considered two hypotheses. First, the virus was introduced in the urban forest park with infected host animals (birds and/or small mammals) or with infected feeding ticks transporting by migrating birds/mammals, and secondly, the virus has been circulating here before but just had not been detected.

I. ricinus and *D. reticulatus* are common species for Moscow's forest parks (Shashina and Germant, 2015; Yankovskaya et al., 2017). In Moscow Oblast the abundance of adult *I. ricinus* in August was up to 46.5 ticks per kilometer of flagging (Kislenko and Korotkov, 2002), the




Fig. 3. Neighbor-joining phylogenetic tree of the complete polyprotein gene (11 148 bp). The bar below indicates the number of base substitutions per site.

mean prevalence of *B. burgdorferi* s.l. in adult unfed ticks was 19.5% (Kislenko and Korotkov, 2002; Korotkov et al., 2008). The incidence of Lyme borreliosis in Moscow in 2015 was 9.5 cases per 100 000 residents (Yankovskaya et al., 2017).

Our primary hypothesis is that the forest park was free of TBEV, and the virus was recently introduced. Consequently, three scenarios of this process are possible: first the virus was introduced with an engorged tick female on a migrating bird or small mammal. Secondly, the virus was brought into the forest park by an infected host animal with high viremia at the time of high questing activity of ticks. Alternatively, the third possibility was both of these scenarios simultaneously.

The hypothesis of recent TBEV introduction is supported by different arguments.

Previously, TBEV has not been detected in ticks or small mammals in Moscow and Moscow Oblast (“On the list of endemic territories for viral tick-borne encephalitis in 2015,” 2015; “On the list of endemic territories for viral tick-borne encephalitis, 2015” 2015). The exception is an outbreak in two districts in the northeastern part of Moscow Oblast in 1948–1957 (Drozdov, 1956). During the outbreak,

about 150 cases of TBE were confirmed. Most of them were caused by consumption of raw goat or cow milk. At the end of the 1950s the local government had taken active measures with large-scale acaricidal spraying, which allowed to eliminate the foci. Subsequently, the TBEV was not registered in the region. It is worth mention that in 2007, one PCR-positive TBEV tick was found north of Moscow Oblast (Shevtsova et al., 2008). However, a follow-up study showed no TBEV-positive ticks in this area (Shevtsova et al., 2009).

Second, epidemiological studies of TBE in Moscow and Moscow Oblast showed that all cases were non-autochthonous, and ticks outside of Moscow Oblast had bitten all patients with TBE (On the incidence of tick-borne borreliosis in Moscow in 2014, 2015; On the incidence of tick-borne borreliosis in Moscow, 2015; On the incidence of tick-borne borreliosis in Moscow in 2014, 2015; Yankovskaya et al., 2017). Autochthonous TBE cases have not been detected since the 1950s.

Third, we found TBEV-positive ticks only at an exact location of Site-1. Mapping of the exact coordinates of the ticks’ collection sites showed the close localization of TBEV-positive ticks (Fig. 2). This evidence also supports the hypothesis of the virus’s recent introduction. If the virus had been introduced by an infected female tick, we can expect

that its offspring (larvae) had been dispersed in the vicinity of the place of oviposition. The distance of the dispersion of questing nymphs and adult ticks depends on the home range size of hosts. The primary hosts for larvae of *I. ricinus* are rodents (Korenberg et al., 2013; Randolph and Craine, 1995). In the forest parks of Moscow, the most common small mammals are the bank vole (*M. glareolus*) and the pygmy field mouse (*A. uralensis*) (Tikhonova et al., 2012). We obtained the same results, and these two species also dominated in our sample. The home range of the bank vole in the breeding season varies up to 3260 m² and for pygmy field mice — 1100–1800 m² (Ilchenko and Zubchaninova, 1963; Kikkawa, 1964; Naumov, 1951; Zhigarev, 2004). In our study, the maximum distance between locations where the infected ticks were collected is 185 m. This is consistent with the home range size of the main hosts of immature ticks — bank voles and pygmy field mice. Therefore, the spatial distribution of the infected ticks does not contradict the hypothesis that they are the offspring of one infected female. In addition to the transovarial transmission route, TBEV can be transmitted during blood meals from a host with high viremia to feeding ticks or between cofeeding ticks (Alekseev and Chunikhin, 1990; Labuda et al., 1993). The dense occurrence of infected ticks in Site-1 does not contradict these two scenarios as well. It is important to note that our sample is biased in favor of a more thorough study of Site-1, so we can only ascertain the presence of TBEV in the ticks at Site-1. Our sample does not allow us to draw conclusions about the presence/absence of TBEV in ticks at Site-2 or Site-3 or in other city parks.

Fourth, only single specimens of *I. persulcatus* and *D. reticulatus* were identified in the study area. Moscow and Moscow Oblast are within the ranges of *I. ricinus*, *I. persulcatus* and *D. reticulatus* (Filippova, 1997, 1977). Each of these species has a mosaic distribution in the region (Korotkov et al., 2008). *I. persulcatus* in Moscow Oblast has a unimodal seasonal activity pattern with the peak in May–June that declines through August (Filippova, 1985). Therefore, we found only one female *I. persulcatus*, probably because of low activity. *D. reticulatus* has a bimodal pattern of seasonal activity, with the first peak in May and the second peak in August. We collected ticks during August–September and found only a single specimen of *D. reticulatus*. Consequently, we can conclude that this species does not form a permanent population in the forest park and that the detected specimen was accidentally found here, arriving here, apparently, with migrating birds or mammals. This evidence also highlights the hypothesis regarding the introduction of the virus.

The fifth assertion is that in the European part of Russia, three subtypes of TBEV are spread, with the Siberian subtype being predominant (Demina et al., 2009; Zlobin et al., 2003). Our results revealed that the identified virus belongs to the European subtype. The sequences of Moscow's TBEV are most similar to the strain isolated from a patient in Slovenia in 1992 (Wallner et al., 1995). This finding can also serve as an indirect confirmation of the hypothesis that the virus was introduced from the outside; the most likely scenario is that the virus was introduced by migrating birds. The spread of TBEV with migrating birds was shown in Tomsk city (Russia), in Sweden, Norway, and Estonia (Geller et al., 2013a,b; Hasle, 2013, 2010; Moskvitina et al., 2014; Waldenström et al., 2007). The avifauna of the studied forest park includes 83 species, including the fieldfare (*Turdus pilaris*), common blackbird (*Turdus merula*), redwing (*Turdus iliacus*), song thrush (*Turdus philomelos*), thrush nightingale (*Luscinia luscinia*), European robin (*Erithacus rubecula*) and common chaffinch (*Fringilla coelebs*) (Kalyakin et al., 2014). These birds can act as tick hosts (Korenberg, 1966, 1962; Korenberg et al., 1964) and as tick vectors (Dubska et al., 2009; Hasle, 2013, 2010; Humair et al., 1993; Marsot et al., 2012; Michalik et al., 2008). The overwintering areas of the mentioned thrushes are located mainly in Italy and France (Bolshakov et al., 2009). These countries are considered low-risk areas for TBEV (European Center for Disease Prevention and Control, 2016). However, it is known that during their migration, birds use various stopover sites, where they feed and rest and where ticks may attach (Sándor et al.,

2014). Therefore, birds could acquire the virus or infected ticks in Eastern Europe on the way from the overwintering grounds. Based on the arguments above, we are inclined to think that the TBEV found in this study was introduced to the forest park from other regions. Further research is needed to provide an unambiguous answer to this question.

The alternative hypothesis is that TBEV was present in this territory previously and has circulated for a long time but was not previously detected. The natural foci of TBEV exhibited a mosaic spatial structure (Korenberg et al., 2013). In particular, it was shown that natural foci of TBEV could be divided into a core zone and an expansion zone (Korenberg et al., 2013; Kucheruk, 1972). The TBEV core zone was the most stable and most epizootically active component of the foci (Korenberg et al., 2013). The TBEV expansion zone was an area with less intensive virus circulation surrounding the core zone (Kucheruk, 1972). We might assume that the location at Site-1 where we found TBEV-positive ticks is the TBEV core zone (a place where the virus has persisted for many years in ticks and their hosts). However, this hypothesis does not seem very likely for the following reason: the location where we found TBEV-infected ticks is crossed by two footpaths that are actively used by locals, and if we were to assume that the core of the infection is located in this place, then autochthonous cases of TBE would have been recorded among the local population even earlier. However, no cases of autochthonous TBE have been recorded in Moscow until 2016, and we consider this to be a strong argument against the alternative hypothesis.

Another important issue raised by this work is how many viral copies need to be introduced into the region for the establishment of new foci. For such tasks, the concept of R_0 – the basic reproduction number is useful. R_0 is the estimate of the average number of secondary cases produced by one primary case in a completely susceptible population (Diekmann et al., 1990). For example, in a host population of capybaras *Hydrochoerus hydrochaeris* susceptible to the tick *Amblyomma sculptum*, it was shown that the introduction of a single infected capybara or a single infected tick was unable to trigger Brazilian spotted fever (caused by the bacterium *Rickettsia rickettsii*) in a nonendemic area (Polo et al., 2017). Only the introduction of a single infected capybara with at least one infected attached tick was enough to trigger the disease in a nonendemic area (Polo et al., 2017). R_0 was defined for TBEV, but only using a model small mammal host and *I. ricinus* (Hartemink et al., 2008). The contribution of other vectors and hosts to the R_0 of TBEV has not been studied. In the forest park, we found not only *I. ricinus* but also *I. trianguliceps*. Ticks of *I. trianguliceps* species do not attack humans but take part in the circulation of the TBEV (Malyushina and Katin, 1965). Thus, our work highlights the need to construct more complex models for estimating R_0 with the inclusion of several groups of hosts (rodents and birds) and at least two vectors (*I. ricinus* and *I. trianguliceps*).

Our findings also contribute to the discussion about the role of urban areas in the development of new foci of vector-borne diseases. It is known that urban areas differ from rural areas in abiotic factors as well as in the fauna of small mammals and birds and their respective behaviors (home-range size, wintering, migration, etc.) (Forman, 2014; Klausnitzer, 1993). Therefore, the role of urban areas in the emergence of new foci requires special attention.

Declaration of Competing Interest

None.

Acknowledgments

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

We thank Novik, I.V. and Midyanik, G.A. for assistance with tick collection, and Uspensky, I.V. for comments that greatly improved the manuscript. We would also like to thank Antonovskaya, A.A.

(Lomonosov Moscow State University) for her help in identifying chigger species.

Appendix A

Table A1
Oligonucleotide sequences of the primers and probes used in PCR amplification and sequencing.

Name	Nucleotide sequence (5'-3')	PCR conditions	References	
qPCR for tick species identification				
Ir_F	CAGCTATCCATATTTGGTCAACAT	95 °C for 15 min	The primers and probes of this assay are self-designed	
Ip_F	AGGAGCAACAGTAATCACAAAC GGGTGAAAGGGATTTATCAAT			
Ixod_R	CAGTTGATAATAACTACTCAAC			
Ir-F2_FAM	CAGTTGATAATAACCCCTAACTCAAC			
Ip-J2_JOE		95 °C for 10 s 60 °C for 20 s	x45	
ITS-2 fragment amplification and sequencing for tick species identification				
58S F3/1	GGGTCGATGAAGAACGCAGCCAGC	95 °C for 15 min	(Fukunaga et al., 2000)	
28S R1/1	TTCTATGCTTAAATTCAGGGGGTTGTC	95 °C for 10 s 63 °C for 15 s 72 °C for 15 s 72 °C for 2 min		x40
COI gene fragment amplification and sequencing for tick and mammal species identification				
ST-COI-F2	CTCYACYAAWCAYAAAGACATTGGAAC	95 °C for 15 min 95 °C for 10 s 60 °C for 15 s 72 °C for 15 s 72 °C for 2 min		x40 self-designed (Geller et al., 2013a,b)
TBEV sequencing ^a				
TBE83s	AGTCGTGAACGTGAGAAAAAGA			
TBE595a	CCATGTC(A/C)GT(A/G)GCCAGIATCACACAGGT			
TBE538s	GCGGCAAC(T/C)CAGGTGCG(T/C)GTGG			
TBE1130a	GGGTTCTCCTG(G/A)TA(G/A)ATGGAGTCAA			
TBEE1	TCA CGG TGC ACA CAT CTG GAA AA			
TBE6	CTT GGT TCC CTC AAT GTG (C/T)GC CAC AGG AA			
RTBE 1-4	GTT GAC YTK GCY CAG ACY GTC AT			
TBEE2	CAT CAG CTC CCA CTC CGA GTG TCA T			
TBE-3f	ATAGTGGGAGGAAGAGGCRTC			
TBE-3r	TCCGTGCCACCATTGAARCG			
TBE-4f	TGTCCATGAGCTTYCTCTTGGCTG			
TBE-4r	TATTTCCATGGCATACCAGCARTCAG			
TBE-5f	AACTGCTCATGGCCTGCGYAGC			
TBE-5r	CAGCCTCCAGACCGAGCAAG			
TBE-6f	TGCTCATGGCTCTYATGACACAGC			
TBE-6r	TGATGCACCTCATGGGCCYTC			
TBE-7f	ACACGATGTGGCAYGTGACKAGAG			
TBE-7r	TCCACTGTGATCATCATCACACTG0TC			
TBE-8f	AGCTMACCGGACCAGGCG			
TBE-8r	AAGCACCAGATGACCCAGGG			
TBE-9f	ATGCATGAGGAACCTGGCAGC			
TBE-9r	ACCGTTCKGTTTCATGACCCCG			
TBE-10f	ATGGTGGATGGGGAYGTCAATCAAC			
TBE-10r	TCCCGTGGCTTCCATGGCC			
TBE-11f	AACACCCGTACCGGACRTGGC			
TBE-11r	ACCACCTGACGAGCATCTTCCAAG			

* All listed primers for TBEV sequencing are self-designed.

References

- Alekseev, A.N., Chunikhin, S.P., 1990. The exchange of the tick-borne encephalitis virus between ixodid ticks feeding jointly on animals with a subthreshold level of viremia. *Med. Parazitol. (Mosk)*. 48–50.
- Bolshakov, C.V., Shapoval, A.P., Zelenova, N.P., 2009. Results of bird trapping and ringing by the Biological Station "Rybachy" on the Courish Spit in 2007. *Avian Ecol. Behav.* 16, 21–51.
- Csango, P.A., Blakstad, E., Kirtz, G.C., Pedersen, J.E., Czettel, B., 2004. Tick-borne encephalitis in Southern Norway. *Emerg. Infect. Dis.* 10, 533–534. <https://doi.org/10.3201/eid1003.020734>.
- De Graaf, J.A., Reimerink, J.H.J., Voorn, G.P., De Vaate, E.A.L.B., De Vries, A., Rockx, B., Schuitemaker, A., Hira, V., 2016. First human case of tick-borne encephalitis virus infection acquired in the Netherlands, July 2016. *Eurosurveillance* 21, 4–6. <https://doi.org/10.2807/1560-7917.ES.2016.21.33.30318>.
- Dekker, M., Laverman, G.D., de Vries, A., Reimerink, J., Geeraedts, F., 2019. Emergence of tick-borne encephalitis (TBE) in the Netherlands. *Ticks Tick. Dis.* 10, 176–179. <https://doi.org/10.1016/j.ttbdis.2018.10.008>.
- Demina, T.V., Dzhioev, Y.P., Verkhovina, M.M., Kozlova, I.V., Tkachev, S.E., Doroshchenko, E.K., Lisak, O.V., Zlobin, V.I., 2009. Molecular epidemiology of tick-borne encephalitis virus: geographic variability determined through the method of molecular hybridization. *Epidemiol. Vaccinal Prev.* 46, 27–39.
- Diekmann, O., Heesterbeek, J.A.P., Metz, J.A.J., 1990. On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations. *J. Math. Biol.* 28, 365–382. <https://doi.org/10.1007/BF00178324>.
- Drozdzov, S.G., 1956. Milk Double-wave Fever in the Moscow Region (materials of Etiological and Epidemiological Study of the Outbreak).
- Dubská, L., Literák, I., Kocianová, E., Taragelova, V., Sychra, O., 2009. Differential role of

- passerine birds in distribution of *Borrelia spirochetes*, based on data from ticks collected from birds during the postbreeding migration period in central Europe. *Appl. Environ. Microbiol.* 75, 596–602. <https://doi.org/10.1128/AEM.01674-08>.
- Annual Epidemiological Report 2016 – Tick-borne Encephalitis. European Centre for Disease Prevention and Control, Stockholm.
- Filippova, N.A., 1997. Ixodid Ticks of Subfamily Ambyomminae // Fauna of Russia and Neighbouring Countries. Arachnoidea. Nauka, Saint Petersburg.
- Filippova, N.A. (Ed.), 1985. Taiga Tick *Ixodes persulcatus* Schulze (Acarina, Ixodidae): Morphology, Systematics, Ecology, Medical Importance. Nauka, Leningrad.
- Filippova, N.A., 1977. Ixodid Ticks of the Subfamily Ixodinae // Fauna SSSR Paukoobraznye. Nauka, Leningrad.
- Forman, R.T., 2014. *Urban Ecology: Science of Cities*. Cambridge University Press.
- Frimmel, S., Krienen, A., Riebold, D., Loebermann, M., Littmann, M., Fiedler, K., Klaus, C., Süß, J., Steisinger, E.C., 2014. Tick-borne encephalitis virus habitats in North East Germany: reemergence of TBEV in ticks after 15 years of inactivity. *Biomed. Res. Int.* 2014. <https://doi.org/10.1155/2014/308371>.
- Fukunaga, M., Yabuki, M., Hamase, A., Oliver, J.H., Nakao, M., 2000. Molecular phylogenetic analysis of ixodid ticks based on the ribosomal DNA spacer, internal transcribed spacer 2, sequences molecular phylogenetic analysis of ixodid ticks based on the ribosomal DNA spacer, internal transcribed spacer 2. *Sequences* 86, 38–43.
- Geller, J., Meyer, C., Parker, M., Hawk, H., 2013a. Redesign of PCR primers for mitochondrial cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. *Mol. Ecol. Resour.* 13, 851–861. <https://doi.org/10.1111/1755-0998.12138>.
- Geller, J., Nazarova, L., Katargina, O., Levits, A., Järvekülg, L., Golovljova, I., 2013b. Tick-borne pathogens in ticks feeding on migratory passerines in western part of Estonia. *Vector-Borne Zoonotic Dis.* 13, 443–448. <https://doi.org/10.1089/vbz.2012.1054>.
- Haglund, M., 2002. Occurrence of TBE in areas previously considered being non-endemic: Scandinavian data generate an international study by the International Scientific Working Group for TBE (ISW-TBE). *Int. J. Med. Microbiol.* 291, 50–54.
- Hartemink, N.A., Randolph, S.E., Davis, S.A., Heesterbeek, J.A.P., 2008. The basic reproduction number for complex disease systems: defining R0 for tick-borne infections. *Am. Nat.* 171, 743–754. <https://doi.org/10.1086/587530>.
- Hasle, G., 2013. Transport of ixodid ticks and tick-borne pathogens by migratory birds. *Front. Cell. Infect. Microbiol.* 3, 1–6. <https://doi.org/10.3389/fcimb.2013.00048>.
- Hasle, G., 2010. *Dispersal of Ticks and Tick-borne Pathogens by Birds*. University of Oslo.
- Heinz, F.X., Stiasny, K., Holzmann, H., Kundli, M., Six, W., Wenk, M., Kainz, W., Essl, A., Kunz, C., 2015. Emergence of tick-borne encephalitis in new endemic areas in Austria: 42 years of surveillance. *Eurosurveillance* 20. <https://doi.org/10.2807/1560-7917.ES2015.20.13.21077>.
- Humair, P.F., Turrian, N., Aeschlimann, A., Gern, L., 1993. *Ixodes ricinus* immatures on birds in a focus of Lyme borreliosis. *Parasitologica* 40, 237–242.
- Ilchenko, A.I., Zubchaninova, E.V., 1963. Year-round observations of marked bank voles and woodland mice in Moscow region. *Zool. Zhurnal* 42, 609–617.
- Jääskeläinen, A.E., Tikkaoski, T., Uzcátegui, N.Y., Alekseev, A.N., Vaheri, A., Vapalahti, O., 2006. Siberian subtype tickborne encephalitis virus, Finland. *Emerg. Infect. Dis.* 12, 1568–1571. <https://doi.org/10.3201/eid1210.060320>.
- Jaenson, T.G., Hjertqvist, M., Bergström, T., Lundkvist, Å., 2012. Why is tick-borne encephalitis increasing? A review of the key factors causing the increasing incidence of human TBE in Sweden. *Parasit. Vectors* 5, 184. <https://doi.org/10.1186/1756-3305-5-184>.
- Kalyakin, M.V., Voltzit, O.V., Groot Kurkamp, H., Rezanov, A.G., 2014. *Atlas of the Birds of Moscow City*. Phiton XXI, Moscow.
- Kikkawa, J., 1964. Movement, activity, and distribution of small rodents *Clethrionomys glareolus* and *Apodemus sylvaticus* in woodland. *J. Anim. Ecol.* 33, 237–244.
- Kislenko, G.S., Korotkov, Y.S., 2002. The forest tick *Ixodes ricinus* (Ixodidae) in foci of tick-borne borrelioses in the north-west of Moscow province. *Parasitologia* 36, 447–456.
- Klausnitzer, B., 1993. *Ökologie Der Großstadtf fauna*. G. Fischer.
- Korenberg, E.I., 2009. Chapter 4 recent epidemiology of tick-borne encephalitis. An effect of climate change? *Advances in Virus Research*. Elsevier Inc., pp. 123–144. [https://doi.org/10.1016/S0065-3527\(09\)74004-7](https://doi.org/10.1016/S0065-3527(09)74004-7).
- Korenberg, E.I., 1966. Birds and the problem of natural foci of tick-borne encephalitis. *Zool. Zhurnal* 45, 245–260.
- Korenberg, E.I., 1962. The role of birds as hosts of ixodid ticks in the natural nidi of encephalitis in the forest zone. *Zool. Zhurnal* 41, 1220–1225.
- Korenberg, E.I., Pomelova, V., Osin, N., 2013. Infections with Natural Focality Transmitted by Ixodid Ticks. *Tipografiya Nauka, Moscow*.
- Korenberg, E.I., Surova, L.G., Kovalevsky, U.V., Tupikova, N.V., 1964. Birds as hosts of ticks in the European South taiga forests. *Zool. Zhurnal* 69, 16–29.
- Korotkov, Y.S., Kislenko, G.S., Burenkova, L.A., Rudnikova, N.A., Karan, L.S., 2008. Spatial and temporal variability of *Ixodes ricinus* and *Ixodes persulcatus* infection with the Lyme disease agent in Moscow region. *Parasitologia* 42, 441–451.
- Kristiansen, K., 2002. TBE in Denmark in particular on Bornholm. *Int. J. Med. Microbiol.* 291, 62–63.
- Kucheruk, V.V., 1972. Structure, Typology and Zoning of the Natural Foci of Human Diseases. The Results of the Development of the Doctrine of the Natural Focality and Further Tasks, Moscow, pp. 180–187.
- Labuda, M., Jones, L.D., Williams, T., Danielova, V., Nuttall, P.A., 1993. Efficient transmission of tick-borne encephalitis virus between co-feeding ticks. *J. Med. Entomol.* 30, 295–299.
- Lindquist, L., Vapalahti, O., 2008. Tick-borne encephalitis. *Lancet* 371, 1861–1871. [https://doi.org/10.1016/S0140-6736\(08\)60800-4](https://doi.org/10.1016/S0140-6736(08)60800-4).
- Malyushina, E.P., Katin, A.A., 1965. On the isolation of tick-borne encephalitis virus from ticks *Ixodes trianguliceps* Bir. *Actual issues viral Infect.* 135–136.
- Marsot, M., Henry, P.Y., Vourc'h, G., Gasqui, P., Ferquel, E., Laignel, J., Grysan, M., Chapuis, J.L., 2012. Which forest bird species are the main hosts of the tick, *Ixodes ricinus*, the vector of *Borrelia burgdorferi* sensu lato, during the breeding season? *Int. J. Parasitol.* 42, 781–788. <https://doi.org/10.1016/j.ijpara.2012.05.010>.
- Michalik, J., Wodecka, B., Skoracki, M., Sikora, B., Staficzak, J., 2008. Prevalence of avian-associated *Borrelia burgdorferi* s.l. Genospecies in *Ixodes ricinus* ticks collected from blackbirds (*Turdus merula*) and song thrushes (*T. philomelos*). *Int. J. Med. Microbiol.* 298, 129–138. <https://doi.org/10.1016/j.ijmm.2008.03.004>.
- Moskvitina, N.S., Korobitsyn, I.G., Tyuten'kov, O.Y., Gashkov, S.I., Kononova, Y.V., Moskvitin, S.S., Romanenko, V.N., Mikryukova, T.P., Protopopova, E.V., Kartashov, M.Y., Chausov, E.V., Konovalova, S.N., Tupota, N.L., Semestsova, A.O., Ternovoi, V.A., Loktev, V.B., 2014. The role of birds in the maintenance of tick-borne infections in the Tomsk anthropogenic foci. *Biol. Bull.* 41, 387–393. <https://doi.org/10.1134/S1062359014040086>.
- Naumov, N.P., 1951. Rodents, A new method for study of ecology of small. *Fauna Ecol. Rodents* 4, 3–21.
- On the Incidence of Tick-borne Borreliosis in Moscow in 2014. Moscow.
- On the List of Endemic Territories for Viral Tick-borne Encephalitis in 2015 [WWW Document]. URL. http://rospotrebnadzor.ru/deyatelnost/epidemiological-surveillance/?ELEMENT_ID=5771.
- Polo, G., Mera Acosta, C., Labruna, M.B., Ferreira, F., 2017. Transmission dynamics and control of *Rickettsia rickettsii* in populations of *Hydrochoerus hydrochaeris* and *Amphymma sculptum*. *PLoS Negl. Trop. Dis.* 11. <https://doi.org/10.1371/journal.pntd.0005613>.
- Pripuzova, N.S., Gmyl, L.V., Romanova, L.I., Tereshkina, N.V., Rogova, Y.V., Terekhina, L.L., Kozlovskaya, L.I., Vorovitch, M.F., Grishina, K.G., Timofeev, A.V., Karganova, G.G., 2013. Exploring of primate models of tick-borne Flaviviruses infection for evaluation of vaccines and drugs efficacy. *PLoS One* 8. <https://doi.org/10.1371/journal.pone.0061094>.
- Randolph, S.E., Crane, N.G., 1995. General framework for comparative quantitative studies on transmission of tick-borne diseases using Lyme borreliosis in Europe as an example. *J. Med. Entomol.* 32, 765–777. <https://doi.org/10.1093/jmedent/32.6.765>.
- Reed, L.J., Muench, H., 1938. A simple method of estimating fifty per cent endpoints. *Am. J. Epidemiol.* 27, 493–497.
- Sándor, A.D., Mărcuțan, D.I., D'Amico, G., Gherman, C.M., Dumitrache, M.O., Mihalca, A.D., 2014. Do the ticks of birds at an important migratory hotspot reflect the seasonal dynamics of *Ixodes ricinus* at the migration initiation site? A case study in the Danube delta. *PLoS One* 9, 1–7. <https://doi.org/10.1371/journal.pone.0089378>.
- Scherer, R., 2014. PropCIs: Various Confidence Interval Methods for Proportions. R Package Version 0.2-5. [WWW Document]. URL. <https://cran.r-project.org/package=PropCIs>.
- Shashina, N.I., Germant, O.M., 2015. The Problem of Nonspecific Prevention of Natural Focal Tick-borne Infections in the Moscow City Parklands. Results and Prospects of Studying the Problems of Infectious and Parasitic Diseases, Tyumen, pp. 220–223.
- Schichanov, N.A., 1987. Universal live trap for small mammals. *Zool. Zhurnal* 66, 759.
- Shevtsova, A.S., Motuzova, O.A., Belova, O.A., Burenkova, L.A., Gmyl, L.V., Kaira, A.N., Karganova, G.G., Kozlovskaya, L.I., Korotkov, Y.S., Pivanova, G.P., Rogova, Y.V., Romanova, L.Y., 2008. A Description of local foci of tick-borne diseases in Moscow Oblast. *Med. Virol.* 25, 127–132.
- Shevtsova, A.S., Romanova, L.Y., Motuzova, O.A., Belova, O.A., Burenkova, L.A., Gmyl, L.V., Kaira, A.N., Kozlovskaya, L.I., Korotkov, Y.S., Pivanova, G.P., Rogova, Y.V., Karganova, G.G., 2009. Characteristic of local foci of Lyme disease, Anaplasmosis, and Tick-borne encephalitis. *Sanit. Dr.* 1, 12–15.
- Skarpaas, T., Golovljova, I., Vene, S., Ljøstad, U., Sjørnsen, H., Plyusnin, A., Lundkvist, Å., 2006. Tickborne encephalitis virus, Norway and Denmark. *Emerg. Infect. Dis.* 12, 1136–1138. <https://doi.org/10.3201/eid1207.051567>.
- Skarpaas, T., Ljøstad, U., Sundøy, A., 2004. First human cases of tickborne encephalitis, Norway. *Emerg. Infect. Dis.* 10, 2241–2243. <https://doi.org/10.3201/eid1012.040598>.
- Süss, J., 2011. Tick-borne encephalitis 2010: epidemiology, risk areas, and virus strains in Europe and Asia-An overview. *Ticks Tick. Dis.* 2, 2–15. <https://doi.org/10.1016/j.ttbdis.2010.10.007>.
- Süss, J., Schrader, C., Falk, U., Wohanka, N., 2004. Tick-borne encephalitis (TBE) in Germany- Epidemiological data, development of risk areas and virus prevalence in field-collected ticks and in ticks removed from humans. *Int. J. Med. Microbiol.* 293, 69–79.
- Tikhonova, G.N., Tikhonov, I.A., Surov, A.V., Bogomolov, P.L., Kutenkova, E.V., 2012. Ecological Aspects of the Formation of the Fauna of Small Mammals in Urban Areas of the Middle Russia. KMK, Moscow.
- Tokarevich, N.K., Tronin, A.A., Blinova, O.V., Buzinov, R.V., Boltentkov, V.P., Yurasova, E.D., Nurse, J., 2011. The impact of climate change on the expansion of *Ixodes persulcatus* habitat and the incidence of tick-borne encephalitis in the north of European Russia. *Glob. Health Action* 4, 8448. <https://doi.org/10.3402/gha.v4i0.8448>.
- Uspensky, I.V., 2017. Blood-sucking ticks (Acarina, Ixodoidea) as an essential component of the urban environment. *Entomol. Rev.* 97, 941–969. <https://doi.org/10.1134/S0013873817070107>.
- Waldenström, J., Lundkvist, A., Falk, K.I., Garpmo, U., Bergström, S., Lindegren, G., Sjöstedt, A., Mejlum, H., Fransson, T., Haemig, P.D., Olsen, B., 2007. Migrating birds and tickborne encephalitis virus. *Emerg. Infect. Dis.* 13, 1215–1218. <https://doi.org/10.3201/eid1308.061416>.
- Wallner, G., Mandl, C.W., Kunz, C., Heinz, F.X., 1995. The flavivirus 3'-noncoding region: Extensive size heterogeneity independent of evolutionary relationships among strains

- of tick-borne encephalitis virus. *Virology* 213, 169–178. <https://doi.org/10.1006/viro.1995.1557>.
- Yankovskaya, Y.D., Shashina, N.I., Karan, L.S., Germant, O.M., Akimkin, V.G., 2017. Epidemiological and parasitological aspects and peculiarity prevention of infections transmitted by Ixodid ticks in Moscow. *Epidemiol. Vaccinal Prev.* 93, 80–85. <https://doi.org/10.1093/molbev/mst197>.
- Zhigarev, I.A., 2004. Small Mammals of Recreational and Natural Forests of the Moscow Region. Promitey, Moscow.
- Zlobin, V.I., 2010. Tick-borne encephalitis in Russia federation: etiology, epidemiology and prevention strategy. *Terra medica Nov.* 2, 13–21.
- Zlobin, V.I., Belikov, S.I., Dzhioev, Y.P., Demina, T.V., Kozlova, I.V., 2003. Molecular Epidemiology of Tick-borne Encephalitis. Irkutsk. RIO VSNC SO RAMN, Irkutsk.