

Trends in and Predictors of Depression Among Participants in the Collaborative Initial Glaucoma Treatment Study (CIGTS)



DAVID C. MUSCH, LESLIE M. NIZIOL, NANCY K. JANZ, AND BRENDA W. GILLESPIE

- **PURPOSE:** To characterize longitudinal trends and factors predictive of depressive symptoms following glaucoma diagnosis in Collaborative Initial Glaucoma Treatment Study (CIGTS) participants.
- **DESIGN:** Cohort study using follow-up data from a clinical trial.
- **METHODS:** A total of 607 participants with newly diagnosed open-angle glaucoma were enrolled at 14 clinical centers in the United States from October 1993 through April 1997, randomized to treatment with medication or surgery, and followed every 6 months through 2004. The 8-item Center for Epidemiologic Studies Depression Scale (CES-D) was administered at baseline and follow-up visits. Three outcome measures were investigated: overall CES-D depression score, presence of mild or worse depression (CES-D score ≥ 7), and number of depressive symptoms endorsed.
- **RESULTS:** The average baseline CES-D score was 2.4 (SD = 3.8), 12.5% of subjects reported symptoms associated with mild or worse depression, and 55.3% reported at least 1 depressive symptom. By 1 year posttreatment, depression measures decreased (1.5, 6.7%, and 38.4%, respectively), with modest decreases thereafter. Baseline factors predictive of mild or worse depression included worse vision-related quality of life (VRQOL) (odds ratio [OR] = 2.41), female sex (OR = 1.42), younger age (OR per 10 years younger = 1.24), and less than high school education (OR = 2.93); other outcomes showed similar results.
- **CONCLUSIONS:** Depressive symptoms decreased considerably during the first year after treatment initiation, but were elevated in those with impaired VRQOL. Given the potential of depression to reduce treatment adherence and thus increase the risk of glaucoma progression, eye care providers should ask patients about depressive symptoms, provide reassurance when appropriate, and make referrals as necessary. (*Am J Ophthalmol* 2019;197: 128–135. © 2018 Elsevier Inc. All rights reserved.)

Accepted for publication Sep 12, 2018.

From the Department of Ophthalmology and Visual Sciences, Medical School (D.C.M., L.M.N.); and Departments of Epidemiology (D.C.M.), Health Behavior and Health Education (N.K.J.), and Biostatistics (B.W.G.), School of Public Health, University of Michigan, Ann Arbor, Michigan, USA.

Inquiries to David C. Musch, Kellogg Eye Center, University of Michigan, 1000 Wall St, Ann Arbor, MI 48105, USA; e-mail: dmusch@umich.edu

NUMEROUS STUDIES HAVE REPORTED DEPRESSION and associated symptomatology to be frequently found among patients with glaucoma.^{1–5} Those with more advanced glaucoma had a greater frequency of depressive symptoms than those with little to no visual field loss.^{6–8} However, Wilson and associates found no difference in self-reported depressive symptoms between glaucoma patients and controls.⁹ Evidence is limited and mixed regarding the possibility that depression in glaucoma patients results from the topical medications that are used to treat the condition.^{10,11} The association of age and depression is unclear, but the 2011–2014 National Health and Nutrition Examination Survey (NHANES) reported past-month antidepressant use in 19% of those aged 60 or older.¹² Glaucoma patients with symptoms of depression have been reported to be less adherent to their prescribed medication(s).^{13–15}

In the Collaborative Initial Glaucoma Treatment Study (CIGTS), we administered an abbreviated 8-item version of the 20-item Center for Epidemiologic Studies Depression Scale (CES-D)¹⁶ to evaluate depressive symptomatology at the time of open-angle glaucoma (OAG) diagnosis among the 607 participants. We reported significant associations at baseline between CES-D responses and participants' self-reported perception of their visual function from the Visual Activities Questionnaire (VAQ).¹⁷ With control for age, sex, race, and comorbidities, we found more than a 3-fold increased odds (odds ratio [OR], 3.72) of a higher (worse) score on the CES-D with a 1-unit higher (worse) score on the VAQ.⁷ All associations indicated that more difficulty with vision-related tasks was associated with greater depressive symptomatology. We also corroborated the finding by Wang and associates⁴ of no significant association of depression with clinical measures of glaucoma severity.

Changes over time in depressive symptoms among people being treated for OAG are lacking in the peer-reviewed literature. We know from our study of CIGTS participants' fear of blindness¹⁸ that the percentage of participants who expressed this fear was initially substantial but diminished over time. Once diagnosed with glaucoma, patients are often treated for their remaining lifetime. They would benefit to know whether depressive symptoms, if evident initially, are likely to improve over time, and to know about factors that affect depressive symptoms.

Therefore, we evaluated longitudinal trends in depressive symptoms and the potential impact of demographic, clinical, and treatment-related factors on the prevalence of depressive symptoms in CIGTS participants.

METHODS

THIS COHORT STUDY INVOLVED FOLLOW-UP OF PARTICIPANTS involved in the CIGTS, a multicenter clinical trial in which a total of 607 participants with newly diagnosed OAG were enrolled at 14 clinical centers over a 3.5-year period from October 1993 through April 1997 (ClinicalTrials.gov Identifier #NCT00000149). Approval for this analysis of data collected on its participants was granted by the University of Michigan Investigational Review Board. For each patient, a “study eye” (usually the eye with more glaucomatous changes) was selected for initial treatment. Participants were randomized to receive either topical medications or surgery (trabeculectomy) to treat their glaucoma, and were followed after initial treatment at 3 months, 6 months, and every 6 months thereafter for up to 10 years. Fellow-eye treatment, when warranted, was specified by protocol to match the study-eye treatment. Baseline assessment and follow-up visits included both an in-person clinical examination and a quality-of-life (QOL) interview administered by telephone shortly after the clinic visit. The clinical examination measured the mean deviation (MD) from visual field testing, intraocular pressure (IOP), and visual acuity (VA), among other glaucoma indicators, to track disease progression. Telephone interviews included assessments of vision-specific QOL (33-item VAQ),¹⁷ glaucoma medication adherence (2 questions), and depressive symptomatology (8-item CES-D).¹⁶ Further details of the CIGTS are given in Musch and associates.¹⁹ This study was approved by the Investigative Review Board at the University of Michigan, and informed consent was obtained from all participants. In addition, the study was conducted in accordance with Health Insurance Portability and Accountability Act regulations and adhered to the tenets of the Declaration of Helsinki.

The 33-item VAQ was the best instrument available at the start of the CIGTS to assess self-reported visual function and impact of vision on daily life among those with glaucoma. Each item of the VAQ asks about the frequency (never, rarely, sometimes, often, always) of a specific vision-related problem. Items are scored from 1 to 5, with higher scores indicating worse visual function; a summary score is computed as the mean of all 33 items.

For each glaucoma medication a patient was taking, adherence was assessed by 2 questions: (1) Over the last 2 weeks, how closely did you follow the instructions of your physician for taking your glaucoma medication? (1-10 scale; 1 = not very closely at all, 10 = very closely),

TABLE 1. Baseline Patient Demographics and Clinical Measures of the Eye (N = 607 Subjects)

Continuous Variables	Mean (SD)	Min, Max	Median
Age (years)	58.0 (10.9)	28, 75	59.2
MD—study eye (dB)	−5.4 (4.3)	−23.5, 3.4	−4.4
IOP—study eye (mm Hg)	27.5 (5.6)	19, 50	27
VAQ	2.0 (0.7)	1.0, 4.4	1.9
Categorical Variables	Frequency, N (%)		
Male	334 (55%)		
Race			
White	337 (56%)		
Black	231 (38%)		
Asian	10 (2%)		
Other	29 (5%)		
Education			
≤6th grade	23 (4%)		
Grade 7-11	105 (17%)		
Grade 12	167 (28%)		
Some college	146 (24%)		
College degree	87 (14%)		
Graduate education	79 (13%)		
Marital status			
Never married	69 (11%)		
Married	365 (60%)		
Divorced/separated	113 (19%)		
Widowed	60 (10%)		
Diabetes	102 (17%)		
Hypertension	225 (37%)		

IOP = intraocular pressure; Max = maximum; MD = mean deviation; Min = minimum; VAQ = Visual Activities Questionnaire.

and (2) Did you miss any dose of your medication yesterday? (Yes, No). For each patient, a summary measure of adherence was calculated for question (1) by taking the minimum of the adherence scores over medications and for question (2) by counting any report of a missed dose on the previous day for any medication.

The 8-item CES-D asks about the presence (Yes, No) and frequency over the past week of specific symptoms of depression. Symptom frequency (0-3 scale) reflects the number of days per week that specific symptoms were experienced (0: less than 1 day; 1: 1-2 days; 2: 3-4 days; 3: 5-7 days). Participants who reported the presence of a symptom but at less than 1 day frequency were scored as 0.5 to distinguish from a report of no symptom.

Because CES-D depression symptom scores were highly skewed, we used 3 summary measures of depressive symptomatology to capture different characteristics of the distribution. First, an overall symptom score was computed as the sum of the 8 symptoms' frequencies over the past week, with scores ranging from 0 to 24. Second, an indicator for subjects with mild or worse depression was designated as an overall symptom score ≥ 7 . This threshold for mild or worse depression was approximated from the

TABLE 2. Frequency of Symptoms of Depression at Baseline (N = 606), Year 1 (N = 544), and Year 5 (N = 485)^a

CES-D Item	Baseline		Year 1		Year 5	
	Frequency	%	Frequency	%	Frequency	%
1. My sleep was restless	187	30.9	125	23.0	89	18.4
2. I felt sad	184	30.4	80	14.7	58	12.0
3. I felt depressed	168	27.7	77	14.2	54	11.1
4. I could not get going	131	21.6	97	17.8	45	9.3
5. I felt that everything I did was an effort	113	18.7	60	11.0	36	7.4
6. I felt that I could not shake off the blues even with the help of my family and friends	84	13.9	29	5.3	21	4.3
7. I did not feel like eating; my appetite was poor	74	12.2	41	7.5	25	5.2
8. I had crying spells	52	8.6	18	3.3	15	3.1

CES-D = Center for Epidemiologic Studies Depression Scale.

^aN = 605 for Baseline, items 1 and 6; N = 484 for Year 5, item 7.

relative position of the threshold used in the 20-item CES-D (overall symptom score ≥ 16 on a scale from 0-60).²⁰ Lastly, a count of reported symptoms, irrespective of the frequency over the past week, was computed, with a range from 0 to 8. Each summary measure of depressive symptoms was calculated from survey responses at the baseline assessment (postdiagnosis, pretreatment, prerandomization) and at follow-up visits (posttreatment).

• **STATISTICAL METHODS:** Characteristics of the CIGTS sample were summarized with means and standard deviations (SD) for continuous measures and frequencies and percentages for categorical measures. The frequency of each CES-D symptom was calculated at baseline, year 1, and year 5. The 3 summary measures of depressive symptoms are graphically presented at baseline and 9-year follow-up using box plots (for CES-D symptom score), a forest plot (for CES-D score ≥ 7), and stacked bar charts (for number of CES-D symptoms). Additionally, within-subject changes in CES-D symptom scores over time are illustrated using a lasagna plot.^{21,22} For easier visualization, CES-D scores were grouped as 0-6.9, 7-11.9, 12-17.9, and 18-24.

Linear mixed regression, repeated-measures logistic regression, and repeated-measures negative binomial regression models were used to investigate predictors of each of the 3 summary measures of depressive symptoms (overall CES-D symptom score, an indicator for mild or worse depression, and count of depressive symptoms, respectively). All models accounted for the correlation between repeated measures over time within a subject. Alternative strategies to address skewness in the data were explored, including log transformation. The log and raw data models gave similar results and thus we retained the linear mixed model on the raw scale for ease

of interpretation. Variables investigated in models included time since diagnosis, patient demographics, clinical measures of glaucoma severity (MD and IOP, both baseline and time-varying), treatment (medicine or surgery), type of medication or medication combination (carbonic anhydrase inhibitor [CAI], beta blocker [BB], prostaglandin analog [PGA]), and baseline VAQ. Variable selection used the method of best subsets.²³ Model results are reported as regression coefficient estimates, ORs, or incident rate ratios (IRR), with 95% confidence intervals (CI), for linear, logistic, and negative binomial models, respectively.

Secondary analyses explored glaucoma-related contributions to depression, whether clinically evaluated severity or self-reported visual function measures. We used linear regression to investigate the association of glaucoma progression (estimated for each subject by the slope of MD over the first 5 years) with depression at 5 years (measured by the CES-D score). This model was adjusted for variables found to be significantly associated with depression in our primary analysis (except for VAQ score because of confounding with MD slope). MD slope was entered as either a continuous variable or a dichotomous variable with cut-points of ≤ -0.5 dB/year or ≤ -1.0 dB/year. The associations between 5-year VAQ and both MD slope and MD at 5 years were assessed with Pearson correlations.

Lastly, we investigated the association between symptoms of depression and medication adherence for the 307 CIGTS subjects randomized to medication. Repeated-measures logistic regression was used to assess the association between CES-D score and probability of missing a dose of any glaucoma medication on the previous day (dichotomous measure). The between-subjects and within-subject correlations between CES-D score and the continuous medication adherence score, in the presence




FIGURE 1. Longitudinal trends in depression as measured by the 8-item Center for Epidemiologic Studies Depression Scale (CES-D) summary depression measures among participants of the Collaborative Initial Glaucoma Treatment Study (CIGTS), including box plots to display the overall symptom score (Top), a forest plot to display the prevalence of mild or worse depression over time from baseline (Middle), and stacked bar charts to display the percentage of subjects with 0-8 symptoms (Bottom; eg, at baseline, 45% of subjects have no symptoms, and the percentage is smaller at each higher category, with 2% having all 8 symptoms). Box plot displays the interquartile range (IQR), including the lower quartile/25th percentile (bottom of the box) and the upper quartile/75th percentile (top of the box), the median/50th percentile (line within the box; any median line not visible is equal to zero), the mean

of repeated observations, were estimated using the methods of Bland and Altman.^{24,25} SAS version 9.4 (SAS Institute, Cary, North Carolina, USA) was used for all statistical analyses.

RESULTS

CIGTS PARTICIPANTS HAD AN AVERAGE AGE OF 58.0 YEARS (SD = 10.9) at baseline; 55% were male and 56% white (Table 1). Baseline measures of the designated study eye showed an average mean deviation of -5.4 (SD = 4.3) dB and IOP of 27.5 (SD = 5.6) mm Hg. The average vision-related QOL at baseline, as measured by the VAQ total score, was 2.0 (SD = 0.7).

The frequency of each CES-D depression symptom at baseline and at 1 year and 5 years posttreatment is shown in Table 2. All symptoms were reported more frequently at baseline than at follow-up. Restless sleep was consistently the most commonly reported symptom over time (30.9% at baseline, 23.0% at year 1, and 18.4% at year 5), followed by feeling sad and feeling depressed.

Consistent with the symptom-specific trends described above, Figure 1 shows elevated depressive symptom measures at diagnosis, substantial decreases by 1 year, and further decreasing trends over follow-up in each of the 3 CES-D summary depressive symptom measures. The mean CES-D symptom score was 2.4 (SD = 3.8) at diagnosis and 1.5 (SD = 3.1) at 1 year (Figure 1, Top); 12.5% of subjects reported a frequency of depressive symptomology consistent with mild or worse depression at diagnosis, but only 6.7% at 1 year (Figure 1, Middle); 55.3% of subjects reported ≥ 1 depressive symptom at diagnosis and 38.4% at 1 year (Figure 1, Bottom). Further modest decreases in all 3 depression measures were observed between 1 and 9 years.

CES-D symptom scores over time for the same subject are shown in Figure 2 for each of the 607 CIGTS subjects. Each subject's score is displayed as 1 layer (row) of the plot. CIGTS subjects who reported the most depressive symptoms at baseline (darker gray sections at the top left corner of the plot) also reported more symptoms over follow-up, but intermittently and mostly to a decreasing extent. Subjects who reported less frequent or no symptoms of depression at baseline reported similar levels throughout their follow-up (center light gray section). Subjects with less frequent or no symptoms of depression at baseline,

(dot within the box), the lower fence/whisker (smallest observation within $1.5 \times IQR$), the upper fence/whisker (largest observation within $1.5 \times IQR$), and outliers (stars; observations located outside $1.5 \times IQR$). Outliers have been jittered to visualize overlying data points.




FIGURE 2. Display of the Center for Epidemiologic Studies Depression Scale (CES-D) overall symptom score within a subject over time for each of the 607 Collaborative Initial Glaucoma Treatment Study (CIGTS) subjects. In this “lasagna plot,” each row shows a subject’s symptom score history from enrollment (time = 0) to 108 months (9 years). Darker gray represents higher CES-D symptom score categories (see category intervals at right), and white areas represent missing data at that time point. At time = 0, subjects are ordered from lowest to highest depression score category, and for those within the same score category at time = 0, patients are ordered from lowest to highest depression score at 3 months, etc.

followed by missing data, were sorted to appear in the lower part of the plot.

Regression model results for each of the 3 depressive symptom measures are displayed in Table 3. Similar effects were observed in each of the 3 models, so for brevity we describe only results for the logistic regression model predicting mild or worse depression (CES-D score ≥ 7). As observed in the descriptive results and plots, there was a strong effect of follow-up time on risk for mild or worse depression. At initial diagnosis (baseline), subjects had a 41% increased odds of mild or worse depression (OR = 1.41; $P = .0072$) relative to year 1. For every additional year of follow-up, subjects had a 6% decrease in odds of mild or worse depression (OR = 0.94; $P = .0066$). Other factors investigated and found to be consistently associated with an increased odds of mild or worse depression included younger age (OR = 1.24 for every 10 year decrease in age; $P = .0031$), female sex (OR = 1.42 vs male; $P = .0726$), race (OR = 2.20 Other vs White race, $P = .0077$; OR = 0.08 Asian vs White race, $P = .0046$), less education (OR = 2.93 for < high school education vs graduate education; $P = .0001$), and worse baseline vision-related QOL (OR = 2.41 for a 1-unit increase [worsening] in VAQ score [1-5 scale]; OR = 5.79 for a 2-unit increase; $P = .0001$). Other variables investigated in adjusted models but not found to have significant associations with the summary depressive symptom measures included baseline or time-varying MD and IOP of the study

eye, initial treatment (medicine vs surgery), indicators for a visit within 1 year prior to cataract surgery or prior to argon laser trabeculoplasty, comorbidities (diabetes and hypertension), marital status, season in which the follow-up visit occurred, clinical center where treated, type of medication or medication combination (for CAI, BB, and PGA medications), and 2 interactions (center by season and age by sex).

Exploration of predictors of depressive symptoms related to glaucoma revealed that clinical measures of glaucoma (MD, IOP) showed little or no correlation with depressive symptoms, whereas self-reported visual function (VAQ) showed strong correlation with these symptom measures (Figure 3). Both the 5-year MD and glaucoma progression as measured by 0-5 year MD slope showed no significant correlation with CES-D score at 5 years ($r = -0.08$, $P = .0953$ and $r = -0.00$, $P = .9656$, respectively). Testing these associations in models adjusting for age, sex, race, and education yielded no significant association between 5-year CES-D score and either 0-5 year MD slope ($P = .6754$) or 5-year MD ($P = .0787$). Further, no significant effect on depressive symptom measures was seen when MD slope was dichotomized at -0.5 dB/year ($P = .0696$) or at -1.0 dB/year ($P = .4609$). In contrast, VAQ score was significantly associated with depressive symptom measures at 5 years ($r = 0.32$, $P < .0001$). The individual VAQ item most predictive of depressive symptoms was “I have problems carrying out activities that require a lot of visual concentration and attention” ($r = 0.34$, $P < .0001$).

The CES-D score was found to be significantly associated with medication adherence. For every 1-unit increase in the CES-D score, there was a 4% increase in the odds of missing a dose of medication during the previous day (OR = 1.04, 95% CI = 1.01, 1.08, $P = .0062$) and a 35% increased odds of missing a dose for every 7-unit increase in the CES-D score (OR = 1.35, 95% CI = 1.09–1.67). The continuous measures of adherence and depressive symptoms had an overall between-subjects correlation of -0.24 ($P < .0001$); that is, subjects with higher (worse) CES-D scores tended to have lower adherence. The within-subject correlations between adherence and CES-D score had an interquartile range from -0.29 to 0.23 , indicating a modest tendency for those with more depressive symptoms at a given time point to have more nonadherence at that time point.

DISCUSSION

CIGTS PARTICIPANTS REPORTED THE MOST DEPRESSIVE symptoms and were more likely to have a depression score indicative of mild or worse depression at or near the time of the clinic visit at which they were initially diagnosed with OAG and informed that they would likely need lifetime treatment for this condition. By 1 year posttreatment, these same participants reported fewer and less-frequent

TABLE 3. Multivariable Longitudinal Models of the 3 CES-D Summary Depression Measures (CES-D Symptom Score, CES-D Symptom Score ≥ 7 , Count of CES-D Symptoms)

Multivariable Models	Linear Mixed Model of CESD Total Score			Logistic Regression of CESD Total Score ≥ 7			Negative Binomial Regression of # CESD Symptoms		
	Estimate ^a	95% CI	P Value	OR ^a	95% CI	P Value	IRR ^a	95% CI	P Value
Time^b									
Baseline vs year 1	0.51	(0.29, 072)	<.0001	1.41	(1.10, 1.82)	.0072	1.45	(1.30, 1.61)	<.0001
1-8 years (per 1 year)	-0.09	(-0.12, -0.06)	<.0001	0.94	(0.91, 0.98)	.0066	0.92	(0.90, 0.94)	<.0001
Age (per 10 years younger)	0.21	(0.39, 0.04)	.0159	1.24	(1.08, 1.44)	.0031	1.12	(1.04, 1.22)	.0031
Sex (female vs male)	0.55	(0.17, 0.94)	.0052	1.42	(0.97, 2.09)	.0726	1.47	(1.20, 1.80)	.0002
Race (vs white)									
Black	0.29	(-0.13, 0.71)	.1769	1.21	(0.83, 1.76)	.3177	1.16	(0.94, 1.44)	.1633
Asian	-0.96	(-2.43, 0.51)	.2019	0.08	(0.01, 0.45)	.0046	0.46	(0.25, 0.85)	.0130
Other	1.51	(0.59, 2.42)	.0012	2.20	(1.23, 3.92)	.0077	2.11	(1.45, 3.06)	<.0001
Education (vs graduate education)									
< High school	0.90	(0.20, 1.61)	.0125	2.93	(1.69, 5.10)	.0001	1.41	(1.01, 1.98)	.0442
High school	-0.12	(-0.76, 0.51)	.7080	1.50	(0.89, 2.54)	.1304	0.90	(0.66, 1.21)	.4756
Some college	0.40	(-0.24, 1.05)	.2167	2.30	(1.29, 4.10)	.0045	1.14	(0.83, 1.56)	.4247
College degree	0.10	(-0.61, 0.80)	.7854	1.64	(0.85, 3.17)	.1381	1.11	(0.77, 1.60)	.5726
Baseline VAQ (per 1-unit increase on 1-5 scale; higher is worse)	1.26	(0.99, 1.54)	<.0001	2.41	(1.91, 3.04)	<.0001	1.89	(1.67, 2.15)	<.0001

CESD = Center for Epidemiologic Studies Depression Scale; CI = confidence interval; IRR = incident rate ratio; OR = odds ratio; VAQ = Visual Activities Questionnaire.

Note: Linear mixed regression used the compound symmetry covariance structure to adjust for the correlation between repeated measures. Repeated-measures logistic and negative binomial regressions used generalized estimating equations to adjust for the correlation between repeated measures.

^aIncreased risk of depression is indicated by Estimate > 0, OR > 1, or IRR > 1.

^bBecause the drop in depression scores from baseline to 1 year was larger than predicted by a linear relationship between 1 and 8 years in all 3 models, time since diagnosis was modeled with a linear term for all time points as well as an indicator variable for deviation from the line at the baseline time point.

depressive symptoms and lower prevalence of mild or worse depression. These findings closely parallel the decline in fear of blindness over time since diagnosis that we reported previously,¹⁸ as well as the responses over time of patients informed of other potentially serious conditions.²⁶ This trend likely relates to an initially elevated concern when patients are informed of having a chronic, vision-threatening condition. The level of concern and associated depressive symptoms in glaucoma are substantially ameliorated with extended time under treatment, during which most patients were told that the treatment they had been receiving was effective in reducing IOP, stabilizing visual field loss, and controlling their OAG. However, there is evidence that poorer glaucoma control is associated with more depressive symptoms, while depressive symptoms are in turn related to nonadherence and possibly poorer glaucoma control. Diniz-Filho and associates²⁷ reported that fast visual field progression was associated with a greater occurrence of depressive symptoms in 102 patients with glaucoma, and Freeman and associates²⁸ found that a maladaptive coping strategy (denial) led to a higher risk of glaucomatous progression.

Personal factors associated with a higher frequency of depressive symptoms included younger age, lower educational attainment, female sex, race, and baseline VAQ. The literature documents a greater tendency of female than male subjects to report depression.^{29,30} Pratt and associates reported that 24.4% of women and 12.6% of men aged 60 or older in NHANES household interviews had taken an antidepressant medication in the past month.¹² Zhou and associates found that both younger age and female sex were independent predictors of anxiety but not depression in their study of 506 Chinese patients with glaucoma.⁵ Our finding of a 42% increased odds of mild or worse depression in female vs male CIGTS participants is consistent with, although somewhat less than, that reported in the general medical literature. The fact that those with an educational attainment less than high school had higher rates of depression may relate to unmeasured factors, such as the extent to which the patient understood glaucoma and its treatment and lower health literacy. The latter has been associated with perceived risk that exceeds actual risk, leading to undue worry despite reassurances of minimal risk.³¹ Race effects showed no difference between blacks and




FIGURE 3. Relationships among clinical measures (mean deviation [MD] and 0-5 year MD slope), self-reported Visual Activities Questionnaire (VAQ), and self-reported Center for Epidemiologic Studies Depression Scale (CES-D) scores. These correlations show that the strongest glaucoma-related predictor of depression was self-assessment of difficulty with visual activities, not clinical measures of visual function or progression.

whites. The significant effects seen in Asians ($n = 10$) and “Other” races ($n = 29$) compared to whites were based on sample sizes too small for meaningful inference. Participants

who reported worse vision-related QOL at baseline were more likely to report depressive symptoms, which fits well with past reports on this association.^{5,7,32,33}

While these findings are based on a large group of newly diagnosed glaucoma patients who were evaluated in a standardized manner over extended follow-up during treatment for glaucoma, we lack information on the participants’ depressive symptoms prior to being informed of their OAG diagnosis. Further, we did not collect information on medications or other treatment for depression. Some participants may have had preexisting depressive symptoms, and some may have required treatment for depression during follow-up, and so our findings need to be considered with these caveats in mind.

Patterns of depressive symptoms observed over follow-up of CIGTS participants are encouraging, as the higher symptom frequency at diagnosis decreased to a lower, stable level over time. However, 11% of patients continued to affirm the CES-D item indicating that they felt depressed even at 5 years after diagnosis. The association of such symptoms with lack of adherence to medications previously reported^{13–15} and validated herein may be attributable to maladaptive strategies of coping, such as denial.²⁸ The possible consequence of faster glaucomatous progression²⁷ warrants attention to the glaucoma patient’s psychological status and better risk communication by eye care providers. Clinicians might consider using the VAQ or a single VAQ item to assess the patient’s glaucoma burden and possible related depression. In keeping with Rabins’ advice,³⁴ continued assessment and management of patients’ anxiety about glaucoma is a necessary addition to the standard clinical reminders about the importance of treatment adherence. Signs of depression, even if unrelated to glaucoma, can lead to low medication adherence and warrant referral for care by a qualified health care professional.

FUNDING/SUPPORT: THIS STUDY WAS SUPPORTED BY A RESEARCH GRANT FROM THE NATIONAL INSTITUTES OF HEALTH/National Eye Institute (grant number EY025719), Bethesda, Maryland, USA, and by a departmental grant from Research to Prevent Blindness (RPB), New York, New York, USA. Dr Musch is a recipient of RPB’s Lew R. Wasserman Merit Award. The sponsors had no role in the study design; in the collection, analysis, and interpretation of the data; in the writing of the report; or in the decision to submit the article for publication. **Financial Disclosures:** David C. Musch is a member of independent data and safety monitoring boards for Glaukos, Inc (San Clemente, California, USA), InnFocus, Inc. (Miami, Florida, USA), and Ivantis, Inc (Irvine, California, USA) and a consultant to Editas Medicine (Boston, Massachusetts, USA), Iridex, Inc (Mountainview, California, USA), and Notal Vision, Inc. (Manassas, Virginia, USA). The following authors have no financial disclosures: Leslie M. Niziol, Nancy K. Janz, and Brenda W. Gillespie. All authors attest that they meet the current ICMJE criteria for authorship.

REFERENCES

- Mabuchi F, Yoshimura K, Kashiwagi K, et al. High prevalence of anxiety and depression in patients with primary open-angle glaucoma. *J Glaucoma* 2008;17(7): 552–557.
- Lin H-C, Chien C-W, Hu C-C, Ho J-D. Comparison of comorbid conditions between open-angle glaucoma patients and a control cohort: a case-control study. *Ophthalmology* 2010;117(11):2088–2095.
- Popescu ML, Boisjoly H, Schmaltz H, et al. Explaining the relationship between three eye diseases and depressive symptoms in older adults. *Invest Ophthalmol Vis Sci* 2012; 53(4):2308–2313.
- Wang SY, Singh K, Lin SC. Prevalence and predictors of depression among participants with glaucoma in a nationally-representative population sample. *Am J Ophthalmol* 2012;154(3):436–444.e2.
- Zhou C, Qian S, Wu P, Qiu C. Anxiety and depression in Chinese patients with glaucoma: sociodemographic, clinical,

- and self-reported correlates. *J Psychosom Res* 2013;75(1): 75–82.
6. Cumurcu T, Cumurcu BE, Celikel FC, Etikan I. Depression and anxiety in patients with pseudoexfoliative glaucoma. *Gen Hosp Psychiatr* 2006;28(6):509–515.
 7. Jampel HD, Frick KD, Janz NK, et al. Depression and mood indicators in newly diagnosed glaucoma patients. *Am J Ophthalmol* 2007;144(2):238–244.
 8. Skalicky S, Goldberg I. Depression and quality of life in patients with glaucoma: a cross-sectional analysis using the Geriatric Depression Scale-15, assessment of function related to vision, and the Glaucoma Quality of Life-15. *J Glaucoma* 2008;17(7):546–551.
 9. Wilson MR, Coleman AL, Yu F, et al. Depression in patients with glaucoma as measured by self-report surveys. *Ophthalmology* 2002;109(5):1018–1022.
 10. Schweitzer I, Maguire K, Tuckwell V. Antiglaucoma medication and clinical depression. *Austr NZ J Psychiatry* 2001; 35(5):569–571.
 11. Kaiserman I, Kaiserman N, Elhayany A, Vinker S. Topical beta-blockers are not associated with an increased risk of treatment for depression. *Ophthalmology* 2006;113(7):1077–1080.
 12. Pratt LA, Brody DJ, Gu Q. Antidepressant use among persons aged 12 and over: United States, 2011–2014. *NCHS Data Brief* 2017;283:1–8.
 13. Pappa C, Hyphantis T, Pappa S, et al. Psychiatric manifestations and personality traits associated with compliance with glaucoma treatment. *J Psychosom Res* 2006;61(5):609–617.
 14. Jayawant SS, Bhosle MJ, Anderson RT, Balkrishnan R. Depressive symptomatology, medication persistence, and associated healthcare costs in older adults with glaucoma. *J Glaucoma* 2007;16(6):513–520.
 15. Lim MC, Watnik MB, Imson KR, et al. Adherence to glaucoma medication: the effect of interventions and association with personality type. *J Glaucoma* 2013;22(6):439–446.
 16. Radloff LS. The CES-D scale: a self report depression scale for research in the general population. *Appl Psychol Meas* 1977; 1(3):385–401.
 17. Sloane ME, Ball K, Owsley C, et al. The Visual Activities Questionnaire: developing an instrument for assessing problems in everyday visual tasks. *Tech Dig Noninvasive Assess Vis Syst* 1992;1:26–29.
 18. Janz NK, Wren PA, Guire KE, et al. Fear of blindness in the Collaborative Initial Glaucoma Treatment Study: patterns and correlates over time. *Ophthalmology* 2007;114(12): 2213–2220.
 19. Musch DC, Lichter PR, Guire KE, et al. The Collaborative Initial Glaucoma Treatment Study (CIGTS): study design, methods, and baseline characteristics of enrolled patients. *Ophthalmology* 1999;106(4):653–662.
 20. Lewinsohn PM, Seeley JR, Roberts RE, Allen NB. Center for Epidemiological Studies-Depression Scale (CES-D) as a screening instrument for depression among community-residing older adults. *Psychol Aging* 1997; 12(2):277–287.
 21. Wilkinson L, Friendly M. The history of the cluster heat map. *Am Stat* 2009;63(2):179–184.
 22. Swihart BJ, Caffo B, James BD, et al. Lasagna plots: a saucy alternative to spaghetti plots. *Epidemiology* 2010;21(5): 621–625.
 23. Hosmer DW, Jovanovic B, Lemeshow S. Best subsets logistic regression. *Biometrics* 1989;45(4):1265–1270.
 24. Bland JM, Altman DG. Calculating correlation coefficients with repeated observations: Part 1—Correlation within subjects. *BMJ* 1995;310(6977):446.
 25. Bland JM, Altman DG. Calculating correlation coefficients with repeated observations: Part 2—Correlation between subjects. *BMJ* 1995;310(6980):633.
 26. Maunsell E, Brisson J, Deschenes L. Psychological distress after initial treatment of breast cancer. Assessment of potential risk factors. *Cancer* 1992;70(1):120–125.
 27. Diniz-Filho A, Abe RY, Cho HJ, et al. Fast visual field progression is associated with depressive symptoms in patients with glaucoma. *Ophthalmology* 2016;123(4): 754–759.
 28. Freeman EE, Lesk MR, Harasymowycz P, et al. Maladaptive coping strategies and glaucoma progression. *Medicine (Baltimore)* 2016;95(35):e4761.
 29. Djernes JK. Prevalence and predictors of depression in populations of elderly: a review. *Acta Psychiatr Scand* 2006;113(5): 372–387.
 30. Vink D, Aartsen MJ, Schoevers RA. Risk factors for anxiety and depression in the older: a review. *J Affect Disorder* 2008; 106(1-2):29–44.
 31. Hawley ST, Janz NK, Griffith KA, et al. Recurrence risk perception and quality of life following treatment of breast cancer. *Breast Cancer Res Treat* 2017;161(3):557–565.
 32. Brody BL, Garnst AC, Williams RA, et al. Depression, visual acuity, comorbidity, and disability associated with age-related macular degeneration. *Ophthalmology* 2001;108(10): 1893–1901.
 33. Hahm B-J, Shin Y-W, Shim E-J, et al. Depression and the vision-related quality of life in patients with retinitis pigmentosa. *Br J Ophthalmol* 2008;92(5):650–654.
 34. Rabins PV. Depressive symptoms in ophthalmology patients. *JAMA Ophthalmol* 2016;134(9):1015.