

Review

Advances in the Development of Anti-*Toxoplasma gondii* Vaccines: Challenges, Opportunities, and Perspectives

Jin-Lei Wang,¹ Nian-Zhang Zhang,¹ Ting-Ting Li,¹ Jun-Jun He,¹
Hany M. Elsheikha,^{2,*} and Xing-Quan Zhu^{1,*}

Important progress has been made in understanding how immunity is elicited against *Toxoplasma gondii* – a complex pathogen with multiple mechanisms of immune evasion. Many vaccine candidates have been tested using various strategies in animal models. However, none of these strategies has delivered as yet, and important challenges remain in the development of vaccines that can eliminate the tissue cysts and/or fully block vertical transmission. In this review, we provide an overview of the current understanding of the host immune response to *T. gondii* infection and summarize the key limitations for the development of an effective, safe, and durable toxoplasmosis vaccine. We also discuss how the successes and failures in developing and testing vaccine candidates have provided a roadmap for future vaccine development.

Why Do We Need a Toxoplasmosis Vaccine?

T. gondii is an obligate intracellular protozoan of global importance, causing toxoplasmosis in humans and animals [1,2]. This parasite places a major health and socioeconomic burden on society, with more than one-third of the global population at risk of infection [1,2]. The complex life cycle of *T. gondii* involves sexual reproduction in the intestine of members of the family Felidae, the only definitive host, and asexual propagation in a broad spectrum of intermediate hosts (Figure 1). *T. gondii* has the ability to infect all warm-blooded animals and humans, using multiple routes of transmission (Figure 1). Three developmental stages – **tachyzoites** (see Glossary), tissue **cysts** containing **bradyzoites**, and **oocysts** – are infectious to hosts, and *T. gondii* can be transmitted horizontally or vertically [1,2].

Most infections in immunocompetent people are asymptomatic and remain undetected with only mild flu-like, myalgia, or other nonspecific symptoms. However, when primary or reactivated infection occurs during pregnancy, or around the time of conception, serious health consequences may occur, such as abortion or severe congenital defects in the affected fetus, including mental retardation, blindness, and hydrocephaly [3]. Reactivation or primary infection in immunocompromised patients (e.g., HIV-infected patients, organ transplant recipients, cancer patients) may cause severe complications, such as encephalitis or pneumonitis [4]. In addition to causing disease in humans, *T. gondii* may induce abortion and stillbirth in livestock, especially in sheep and goats, posing a serious threat to sustainable agricultural economies [2].

In humans, the primary therapeutic regimen involves the combined use of the antifolate drugs pyrimethamine and sulfadiazine; however, failure rates remain significant [5]. Other

Highlights

There is a compelling need to develop a safe and effective toxoplasmosis vaccine.

Successful vaccination of domestic cats is the key step in reducing *T. gondii* transmission to humans and food-producing animals.

An effective toxoplasmosis vaccine must be able to induce both humoral and cellular immune responses, directed against multiple different proteins, at different stages of the parasite life cycle.


Live attenuated *T. gondii* strains offer good protection against toxoplasmosis, but the possibility of reversion to the virulent type remains possible.

There is a need to identify more immunogenic antigens, adjuvants, and antigen-delivery systems together with defining robust immunocorrelates of protection.

Having a standardized protocol for assessment of vaccine efficacy can facilitate the synergy between the results obtained by various research groups.

'Omics' technologies have revolutionized our understanding of the pathophysiology of toxoplasmosis, paving the way for development of a safe and effective vaccine.

¹State Key Laboratory of Veterinary Etiological Biology, Key Laboratory of Veterinary Parasitology of Gansu



Province, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu Province 730046, PR China
²Faculty of Medicine and Health Sciences, School of Veterinary Medicine and Science, University of Nottingham, Sutton Bonington Campus, Loughborough, LE12 5RD, UK

*Correspondence:
hany.elsheikha@nottingham.ac.uk
 (H.M. Elsheikha) and
xingquanzhu1@hotmail.com
 (X.-Q. Zhu).

Trends in Parasitology

Figure 1. The Two-Host, Predator-Prey, Life Cycle of *Toxoplasma gondii*. (A) Members of the family Felidae (domestic and wild cats) are the only known definitive hosts of *T. gondii*. (B) Cats become infected after ingestion of viable tissue cysts. (C) Cats shed unsporulated oocysts in their feces into the environment. (D) After shedding, oocysts mature in the environment into infective (sporulated) oocysts. Oocysts are able to survive in the environment for long periods of time, and the sporulated oocysts can contaminate soil, water, vegetables, and fruits (E), providing a route of infection to a wide range of intermediate hosts (F) and cats (G). *T. gondii* infection is initiated in humans by ingestion of oocysts in contaminated food or water (H) or ingestion of viable tissue cysts in raw or undercooked meat (I). Other means of infection include contact with cats (J), mother-to-child transmission (K), or by organ transplantation (L). Shortly after
 (Figure legend continued on the bottom of the next page.)

regimens – including pyrimethamine together with clindamycin, clarithromycin, azithromycin, or atovaquone, or monotherapy with trimethoprim–sulfamethoxazole (TMP-SMX) – have been used for treating toxoplasmosis; however, none have been found to be superior to the pyrimethamine–sulfadiazine combination [5]. All treatment regimens used in clinical practice are active against the rapidly dividing tachyzoite stage of the parasite, but do not exert any significant activity against the slow-dividing bradyzoites within the tissue cysts, and cannot cure persistent infections [5]. Also, of concern with current therapeutics are the side-effects that may occur in the treated host [5]. Vaccines offer better alternatives for efficient long-term disease control, while reducing the reliance on, and consequences of, chemical therapeutics. Currently, only one commercially available vaccine (Toxovax[®], MSD Animal health) is licensed for use in sheep to prevent abortion, but this vaccine has several drawbacks (see below) [6]. The development of an effective, safe, and durable vaccine against *T. gondii* infection remains a necessity to facilitate the control of toxoplasmosis.

Mechanisms of Immunity to Toxoplasmosis

Protective immunity to *T. gondii* is complex, involves the innate and adaptive immune response, and is primarily dependent on T helper 1 (Th1) cell-mediated immunity driven by production of high levels of interleukin-12 (IL-12) and interferon- γ (IFN- γ) [7–9]. Following infection with *T. gondii*, innate immune cells – including dendritic cells (DCs), macrophages and neutrophils – migrate to the site of infection where they detect the parasite, mainly via Toll-like receptors (TLRs), and secrete IL-12, to stimulate CD4⁺ and CD8⁺ T cells, and natural killer (NK) cells to produce IFN- γ (Figure 2). IFN- γ limits the parasite's proliferation and the progression of infection through multiple intracellular mechanisms, such as stimulating the expression of **guanylate-binding proteins (GBPs)** and **immunity-related GTPases (IRGs)** to degrade the **parasitophorous vacuole (PV)**, increasing the expression of major histocompatibility complex (MHC) and antigen-presentation genes, and upregulation of other antiparasitic factors such as reactive oxygen species (ROS) and reactive nitrogen species (RNS) [7–9]. IL-12 and IFN- γ are critical cytokines for clearance of tachyzoites during acute infection; they are also essential for inducing tissue cyst formation and sustaining latent infection [7–9]. In addition, IL-1, IL-2, IL-4, IL-6, and tumor necrosis factor alpha (TNF- α) play a protective role in *T. gondii* infection. The anti-inflammatory cytokines IL-4, IL-10, IL-13, IL-27, and transforming growth factor (TGF- β) are responsible for minimizing damage caused by an excessive immune-inflammatory response [7–9].

Adaptive immunity depends largely on antigen-presenting cells (APCs), such as DCs and macrophages, and their ability to present *T. gondii* antigen and activate B cells, and CD4⁺ and CD8⁺ T cells, in secondary lymphoid organs. Activation of B cells, and CD4⁺ and CD8⁺ T cells, requires both the recruitment of APCs from the infection site to secondary lymphoid organs and the activation of resident APCs through exposure to *T. gondii* antigens. Once activated, B cells serve as APCs to stimulate CD4⁺ T cells, and they secrete cytokines such as IFN- γ and IL-12 [10]. Activation of T cells elicits immunity to *T. gondii*, and adoptive transfer of activated T cells can confer resistance to naive hosts [11]. CD4⁺ and CD8⁺ T cells are essential for long-term survival of the host, acting synergistically to prevent reactivation of cysts, and are important for the development of protective immunity against reinfection [12].

infection, infective oocysts and tissue cysts transform to fast-replicating tachyzoites, the parasite stage responsible for acute infection. As the infection progresses, tachyzoites differentiate into slow-replicating bradyzoites enclosed within tissue cysts, which mark the latent form of infection and can persist throughout the lifetime of the affected host.

Glossary

Apicomplexan AP2 (ApiAP2): a family of apicomplexan proteins with a 60 amino acid DNA-binding domain related to the Apetala-2 (AP2) family in plants that are linked to stress responses.

Bradyzoite: a slowly replicating form of *T. gondii*, found within tissue cysts, which is associated with latent infection.

Cysts: thick-walled intracellular vacuolar structures that contain quiescent bradyzoites and can persist for the life of the host, predominantly in muscle and neuronal tissues.

Guanylate-binding proteins

(GBPs): a family of large (65–75 kDa) GTPases induced by interferon; they contribute to immune control of *T. gondii*.

Immunity-related GTPases (IRGs):

a family of GTPases, about 47 kDa, stimulated by interferon gamma. They are important in the control of *T. gondii* in mice. Only one is present in humans, and it is not responsive to interferon.

Immunomics: an integrated immunology, genomics, proteomics, transcriptomics, and bioinformatics approach aimed at the discovery of immunogenic antigens by studying the global interaction between the host immune system and pathogens.

Oocysts: thick-walled environmental forms of *T. gondii* that are excreted in the feces of cats and can survive for lengthy periods outside the host under various ecological conditions.


Parasitophorous vacuole (PV): a special membrane-bound intracellular vacuolar compartment formed during host cell invasion by the parasite; it serves as a surrogate niche where the parasites divide through a special process known as endodyogeny.

Parasitophorous vacuole

membrane (PVM): a physical membrane that surrounds the parasitophorous vacuole and is modified by the parasite to facilitate nutrient acquisition and avoid host defenses.

Reverse vaccinology: an approach used for the discovery of candidate vaccine antigens from genomic information.

Structural vaccinology: the use of knowledge of protein molecular



structure to identify protective antigens.

Tachyzoite: a fast-replicating form of *T. gondii* associated with the acute phase of infection.

Figure 2. Host Immune Responses against *Toxoplasma gondii*. Chemokines secreted from cells at the site of infection trigger the recruitment of neutrophils, dendritic cells (DCs), monocytes, and macrophages to the site of infection.

(Figure legend continued on the bottom of the next page.)

Humoral immunity is also essential for the control of toxoplasmosis [2,13]. Infection with *T. gondii* increases the levels of circulating immunoglobulins – IgA, IgE, IgM, and IgG – which exert their protective effects through multiple mechanisms, such as opsonization of the parasites by phagocytosis; inhibiting the parasite's attachment to host cells; blocking parasite invasion; and activating the classical complement pathway [10].

Historical Account of Toxoplasmosis Vaccines

Considerable efforts have been made to develop a vaccine against toxoplasmosis over the past 30 years. Toxoplasmosis vaccine research has advanced considerably in recent times, with a large number of vaccine candidates having been tested using various vaccination strategies. Among these, inactivated vaccines, live-attenuated vaccines, DNA vaccines, protein vaccines, epitope vaccines, and live vector-based vaccines have been investigated in animal models to determine their ability to induce protective immune responses against *T. gondii* infection. However, none of these vaccination approaches have been able to eliminate the tissue cyst and/or block vertical transmission [14–16].

Vaccines based on killed organisms, lysate antigen, excretory–secretory (ES) products, subunit proteins, and DNAs, are particularly effective against *T. gondii* infection as they can induce strong cytotoxic T lymphocytes and activate a number of lymphocytes. Among the tested antigens, studies have focused on the surface antigens (SAGs), rhoptry antigens (ROPs), microneme antigens (MICs), dense granule antigens (GRAs), and other antigens secreted by tachyzoites and bradyzoites, which play critical roles in *T. gondii* motility, attachment, invasion, replication, and manipulation of the host's immune response [14–16]. Multiantigen vaccines, expressing several proteins or peptides representing different stages of the parasite, seem more powerful and efficient than single-antigen vaccines [14–16]. The components of the vaccine (e.g., peptides and adjuvants) seem to play roles in the efficacy of the vaccine. Several adjuvants, such as Freund's adjuvant, interleukins, CpG DNA, monophosphoryl lipid A, MHC-restricted adjuvant, have been incorporated into vaccines to enhance immune responses [14–16]. For example, coadministration of IL-15 and IL-21 with MIC8 stimulated specific humoral and cellular immune responses to acute and chronic *T. gondii* infection [17].

Live nonpathogenic microbes (bacteria, protozoa, and virus) are the preferred vectors for delivering *T. gondii* antigen due to their ability to mimic the intracellular niche of *T. gondii* and induce an immune response without an additional adjuvant [18–20]. Immunization using a combined virus-like particle (VLP) vaccines expressing ROP18 and MIC8 induced robust humoral and cellular immune responses, and reduced parasite burden [19]. Also, heterologous

The interaction of *T. gondii*-derived profilin protein with Toll-like receptor 11 (TLR11) and TLR12 on DCs is required for the secretion of interleukin-12 (IL-12) in mice. In addition to stimulating IL-12 secretion, tumor necrosis factor alpha (TNF- α) is also produced by macrophages, in response to TLR2- and TLR4-mediated sensing of *T. gondii* glycosylphosphatidylinositol (GPI)-anchored proteins. IL-12 stimulates natural killer (NK) cells, and CD4⁺ and CD8⁺ T cells to produce IFN- γ . Neutrophils are an important source of IFN- γ ; however, the mechanisms that regulate neutrophil-derived IFN- γ remain incompletely known. The regulatory IL-10 cytokine plays an important role in the modulation of immune pathways and can prevent the lethal overproduction of T helper 1 type proinflammatory cytokines. IFN- γ has pleiotropic effects on the course of infection. IFN- γ amplifies a signal through a surface IFN- γ receptor to activate signal transducer and activator of transcription 1 (STAT1). In response to activation of STAT1, monocytes and macrophages upregulate their expression of the indoleamine 2,3-dioxygenase (IDO) enzyme and inducible nitric oxide synthase (iNOS), and reactive oxygen species (ROS). IDO depletes tryptophan, an essential amino acid for *T. gondii* growth. iNOS, in addition to production of nitric oxide (NO), controls parasite replication by depleting arginine, another essential amino acid for *T. gondii* growth. ROS has been implicated in the IFN- γ -mediated control of *T. gondii* in a cell type- and species-specific manner. Both hematopoietic and nonhematopoietic cells upregulate two families of defense proteins called guanylate-binding proteins (GBPs) and immunity-related GTPases (IRGs), which accumulate at the parasitophorous vacuole (PV) and contribute to the parasite clearance.

prime-boost immunization strategies could induce high levels of specific humoral and cellular immune responses, providing a basis for the development of preventative and therapeutic interventions against *T. gondii* infection. For example, prime immunization using recombinant influenza viruses expressing SAG2 antigen, followed by a booster vaccination using recombinant adenovirus encoding the same antigen, induced strong immune responses and protected mice with an 85% reduction in parasite burden [20].

Live-attenuated vaccines are among the most efficient approaches that confer immune protection because they mimic natural infection, providing an approximation of the microenvironment for antigen processing and presentation. Live-attenuated vaccines based on mutants, such as T-263, TS-4, and S48, have shown high efficacy in the prevention of shedding oocysts from cats, reduction of the incidence of abortion in sheep, and reduction of tissue cyst formation in food-producing animals [14–16]. However, only one commercial vaccine, Toxovax[®], based on the live-attenuated tachyzoite S48 strain, is licensed in New Zealand, the UK, and other European countries, and is used to reduce the incidence of abortion in sheep [6]. The S48 tachyzoites in this live-attenuated commercial vaccine (Toxovax[®]) were originally isolated from an aborted lamb in New Zealand, and with repeated passage in excess of 3000 times in mice have lost their capacity to form tissue cysts or produce oocysts [21]. However, Toxovax[®] has a relatively short shelf life and cannot fully eliminate the parasite. Furthermore, because the molecular basis of this live-attenuated tachyzoite S48 strain is not fully understood, spontaneous mutations may raise concerns due to the possibility of reversion to a pathogenic phenotype, which can pose a high risk to vulnerable individuals [21].

New Trends in Toxoplasmosis Vaccine Development

Live-Attenuated Vaccines Based on Gene Editing

In recent years, with the advances in genetic engineering methods, various approaches have been used to develop an attenuated toxoplasmosis vaccine [22]. Several genetically attenuated parasite strains, generated by deletion of certain genes, have shown good efficacy against toxoplasmosis (Table 1). The protective efficacy of genetically attenuated parasites depends on the challenge dose and challenge time, as well as the parasite background and host susceptibility. Not all genetically attenuated parasites have the potential to induce a strong immunity to wild-type *T. gondii* challenge. There is evidence that the apical secretion of some rhoptry proteins is a key element for the induction of long-lasting sterile protective immunity [23]. On the other hand, *T. gondii* has been under selection pressure due to its interaction with its hosts' immune response; and it is possible to genetically alter the parasite in a manner that prevents inhibition of, and maybe even boosts, immune responses. This may be achieved by overexpression of some vital protective

Table 1. Selected List of Attenuated Live Vaccines Tested against Experimental *Toxoplasma gondii* Infection

Targeted gene	Parasite strain	Dosage	Animal model	Spectrum of protective efficacy	Refs
CPSII	RH	10 ⁷ tachyzoites	BALB/c, C57BL/6 mice	Acute and chronic infection	[11,77]
MIC1-3	RH	20 tachyzoites	Swiss OF1 mice	Acute, chronic, and congenital infection	[78]
OMPDC	RH, Pru	10 ⁷ tachyzoites	C57BL/6 mice	Acute and chronic infection	[79,80]
AMA1	RH	10 ⁶ tachyzoites	BALB/c, C57BL/6, CD-1 mice	Acute and chronic infection	[81]
PTS	RH	5 × 10 ³ tachyzoites	C57BL/6 mice	Acute and chronic infection	[82]
GRA17	RH	5 × 10 ⁴ tachyzoites	Kunming mice	Acute, chronic, and congenital infection	[83]
CDPK2	Pru	5 × 10 ² tachyzoites	Kunming mice	Acute, chronic, and congenital infection	[84]
MIC1-3	RH	10 ⁵ tachyzoites	Sheep	Congenital infection	[85]

antigens, or by deletion of some important genes that the parasite uses to manipulate its host cell, or deletion of some key genes involved in controlling bradyzoite differentiation. For example, tachyzoites engineered to constitutively express bradyzoite-specific SRS9 protein trigger a stronger SRS9-specific immune response than the wild-type tachyzoite [24]. Live-attenuated toxoplasmosis vaccines can provide more protection than vaccines prepared by other approaches. However, before a licensure of an efficacious live-attenuated vaccine can be sought, there is a need for evaluation systems to validate vaccine safety and efficacy with higher predictive values in the preclinical and clinical studies in order to prioritize promising candidate strains and guide the development of the much needed vaccines.

Nanoparticle-Based Vaccines

Recombinant subunit vaccines, containing one or multiple target antigens from a pathogen, are a safer alternative to live-attenuated vaccines, but they are less immunogenic and thus less effective at eliciting protective immunity [14–16]. Recent advances in nanomaterial engineering have opened up new avenues for the development of novel vaccines [25]. Nanoparticles (NPs) have significant potential as a delivery system. First, antigen encapsulation within NPs, or via covalent attachment to the NPs, protects the integrity of the antigen against enzymatic degradation, prolonging its systemic circulation time and enhancing the likelihood of presentation to immune cells [25]. Second, NP-based antigen systems can deliver antigens to certain cells, such as DCs, by coupling NPs with antibodies that bind with specific receptors on the surface of target cells [26]. Third, NPs use lower doses of antigens and adjuvant, thus reducing the risk of toxicity and side effects [27]. Some NPs are innately immunogenic and can serve as an adjuvant for activation of the immune system [28].

Many types of NP have been used as a potent immunostimulatory delivery system to improve the induction of an effective immune response against *T. gondii* infection [29]. For example, immunization of mice using a multiepitope recombinant *T. gondii* vaccine containing predicted T and B cell epitopes of SAG1, AMA1, ROP2, and GRA4 proteins, and encapsulation of the expressed protein in poly lactic-co-glycolic acid NPs as a delivery vehicle, enhanced Th1 immune response and significantly reduced the parasite load, prolonging the survival of infected mice [30]. Immunization of mice with cocktail DNA vaccines, containing ROM4 and GRA14 together with calcium phosphate nanoparticles (CaPNs), enhanced immune responses to acute toxoplasmosis [31]. A pioneering study used an adjuvant-free modified dendrimer nanoparticle vaccine, which simultaneously carries multiple replicon RNAs encoding GRA6, ROP2A, ROP18, SAG1, SAG2A, and AMA1, to protect mice against lethal *T. gondii* challenges [32]. Although the protective efficacy of NP vaccine is higher than conventional subunit vaccines, it is still less effective than live-attenuated vaccines.

Exosome-Based Vaccines

Exosomes are nano-sized (~30–150 nm in diameter) membrane-bound vesicles released by most eukaryotic cells through fusion of multivesicular bodies with the plasma membrane [33]. They can transfer mRNA, miRNA, and proteins between cells, representing a new way of intercellular communication which is crucial to the development of immune homeostasis [34]. The exosomes also contain pathogen-derived antigens, and they represent a novel tool for vaccination against infectious diseases [35]. Immunization of mice with exosomes derived from *T. gondii*-pulsed DCs induced a specific and protective T cell response to congenital toxoplasmosis [36]. Recent research has shown that exosomes isolated from *T. gondii* can induce humoral and cellular immune responses, which are protective against acute infection in mice [37]. These studies may motivate further exploration of the potential of exosome as a new vaccine delivery system to support research efforts in the quest for a toxoplasmosis vaccine.

Carbohydrate-Based Vaccines

The surface of many pathogens contains carbohydrates such as polysaccharides, oligosaccharides, and glycans, which are used by pathogens as receptors for attachment to, and invasion of, host cells [38]. Carbohydrates on the pathogen surface are recognized by the host innate immune receptors, leading to the production of antiglycan antibodies. This makes the surface carbohydrates of pathogens attractive candidates for vaccine development [39,40]. In *T. gondii*, glycosylphosphatidylinositols (GPIs) can be recognized by TLR-2 and TLR-4 to induce an inflammatory response, such as TNF- α production [41]. Two major *T. gondii* GPI glycoforms have been evaluated as potential vaccine candidates. Although antibodies were induced, immunization did not protect mice against challenge with the virulent *T. gondii* RH strain. Mouse serum antibodies were more specific to the linker used to attach the carbohydrate to the carrier protein than to the desired carbohydrate side-branch of the parasite GPI [42]. Given the proven success of carbohydrate vaccines to combat several microbial infections, further selection of more immunogenic carbohydrate antigens, and leveraging the potential of glycan array technologies, may facilitate the development of an effective carbohydrate-based anti-*T. gondii* vaccine.

Hurdles That May Limit Development of Effective Vaccines

Despite more than three decades of a critical mass of basic and preclinical research into toxoplasmosis vaccine development, and despite employing various approaches to test many antigens and various formulations, none of the tested vaccines can eliminate the tissue cysts and/or fully block vertical transmission. Several challenges for the development of an effective vaccine exist, such as the genetic diversity among *T. gondii* strains; the multistage life cycle of the parasite; the challenge of clinical translation; the remarkable ability of the parasite to evade and subvert host immunity; the ability of *T. gondii* to establish latent infection; the lack of a standardized vaccination protocol; and the influence of coinfection with other pathogens on the host response to vaccination.

The Impact of Strain Variations on Immunogenicity

Genetic diversity studies of *T. gondii* isolates from North America and Europe have revealed that *T. gondii* has a clonal structure in which the majority of *T. gondii* strains are classified into three predominant lineages: type 1, type 2, and type 3 [43]. These three lineages are all similar (only ~1–2% divergence at the nucleotide level) and probably originated from a common ancestor around 10 000 years ago [44]. Despite the low genetic diversity between these three lineages, they differ markedly in their pathogenicity. Type 1 strains are lethal in laboratory mice, with an infectious dose of a single viable organism, whereas types 2 and 3 strains are considerably less virulent. Type 1 strains also exhibit increased migratory capacity *in vivo* and *in vitro* and have a higher growth rate *in vitro* than the type 2 and type 3 strains [45]. Other differences between the three predominant lineages include the frequency of differentiation, induction of intestinal pathology during acute infection, development of central nervous system pathologies during chronic infection, and manipulation of host cell signaling (see below) in laboratory mice [45].

Further analysis of the global population structure of *T. gondii* strains has revealed that the majority of strains are divided into six major clades, which contain more than 16 haplo-groups with distinct geographical patterns [46,47]. Compared with strains from Europe and North America, *T. gondii* isolates from South America are more genetically diverse, with no predominant genotypes, and they show stronger evidence of genetic recombination. Like type 1 strains, the majority of *T. gondii* strains from South America are highly virulent to mice [46,47]. Genome analysis of 62 globally distributed *T. gondii* strains revealed a diversification of secretory

proteins that are important determinants in host–pathogen interactions, highlighting crucial differences between major clades of *T. gondii* [48]. Expansion of polymorphic genes encoding secretory proteins that interact with, and modulate, the host signaling pathways have made *T. gondii* strains efficient modulators of host immune responses [48]. Previous studies have shown that some genetically distinct strains may have the ability to superinfect the same host, suggesting insufficient cross protection from immunization with a single *T. gondii* strain [49,50].

A Complex, Multistage Life Cycle

T. gondii has a multistage life cycle, wherein it expresses many different proteins during the stage-to-stage progression [51]. Therefore, immune responses produced against a specific life cycle stage (i.e., tachyzoite, bradyzoite, or oocyst) of *T. gondii* may fail to protect against infection caused by other stages, making vaccine design for this parasite more challenging than for simpler organisms, such as most viruses and bacteria. Vaccines that target a stage-specific antigen, or are based on a single antigen or on two antigens, may elicit an immune response that lacks sufficient cross-protection. The complex life cycle of this parasite, coupled with significant genetic variations, as discussed above, raises doubts that single-antigen vaccines could confer the desired broad-spectrum protection. Combining immunogenic antigens from the different forms of *T. gondii* life-cycle in a multistage-specific cocktail may offer a more robust approach towards the development of a toxoplasmosis vaccine capable of inducing sterile immunity and effectively preventing infection.

Species Barriers to Clinical Translation

T. gondii can infect almost all warm-blooded animals, and its pathogenesis has been studied in different animal models. Sheep, goats, pigs, and even chickens have been used to model toxoplasmosis. However, the laboratory mouse is considered the most widely used model to study the pathogenesis and immunological events involved in the control or prevention of *T. gondii* infection [52]. Hundreds of vaccination studies over the past 30 years have been performed in laboratory mice, yielding important insights into toxoplasmosis vaccine development. However, the use of the mouse model to study the development of toxoplasmosis vaccine has several drawbacks.

Cross-species differences create barriers for translational research which may impede the success of clinical trials. Rodents are the natural intermediate host of *T. gondii*, and they frequently encountered this parasite over the course of their evolution, so this long-term interaction might have shaped the immune system of mice to handle *T. gondii* in a way different from that in an accidental intermediate host, such as humans and food-producing animals [52]. Therefore, mechanisms of innate response to *T. gondii* infection are expected to vary between rodents and humans, although both rely on IFN- γ and STAT1 signaling pathways [7–9]. In mice, TLR11 and TLR12 are the innate sensors of *T. gondii*, where they recognize parasite profilin, an actin-binding protein (Figure 2) [53]. TLR11 is a pseudogene, and TLR12 is absent from the human genome, and the sensors involved in *T. gondii* recognition by human innate immune cells remain unknown [7–9]. In addition, the IRGs, a family of IFN- γ -inducible proteins, are essential for the resistance of mice to *T. gondii* infection [54]. There are 23 members of this family in the mouse genome while humans lack most IRGs and only one IRG homologue, IRGM, is present in the human genome. Furthermore, IRGM is severely truncated and is not regulated by IFN- γ , so it possibly does not function in humans as it does in mice; this is another example of human/mouse differences in IFN- γ -mediated immunity to *T. gondii* [54]. Therefore, antigens identified in mice may not protect humans or food-producing animals, and more vaccine studies should be performed in food-producing animals, such as sheep and pigs.

Manipulation of Host Cell Signaling Pathways

As an obligate intracellular parasite, *T. gondii* has evolved several strategies to fine-tune the cellular microenvironment to promote its own growth and persistence inside host cells, thereby increasing the spread of infection and the chance of transmission to new hosts. After invasion of the host cell, *T. gondii* employs a repertoire of effector proteins secreted from dense granules and rhoptry organelles to manipulate host gene expression and signaling pathways [55].

Some of the dense granule effector proteins traffic to the host cell nucleus (GRA16, GRA24 and TgIST), whereas others translocate to the **PV membrane (PVM)** (GRA15) or partially interact with the host cell cytosol while exhibiting a vacuole-restricted localization (GRA6). In all *T. gondii* strains, GRA16 upregulates the expression of host genes involved in cell cycle progression, metabolism, and the p53 tumor suppressor pathway [56]. GRA24 prolongs autophosphorylation and nuclear translocation of host cell p38 α , which correlates with the activation of transcription factors such as Egr-1 and c-Fos, and the production of proinflammatory cytokine IL-12 and chemokine monocyte chemoattractant protein 1 (MCP-1) [57]. TgIST blocks the IFN- γ response by blocking STAT1-dependent proinflammatory gene expression [58,59]. GRA6 activates the host transcription factor NFAT4 (nuclear factor of activated T cells 4) in a strain-specific manner, promoting the production of chemokines, such as Cxcl2 and Ccl2 [60]. GRA15 is another strain-specific effector that activates the NF- κ B pathway and induces more IL-12 secretion in type 2 strains, compared to type 1 or type 3 strains [61].

Some effector proteins secreted from the rhoptry organelle can also alter the host's transcriptional response. For example, rhoptry kinase ROP16 injected into the host cell nucleus can maintain constitutive activation of signal transducer and activator of transcription (STAT)-3 and STAT6 in a strain-specific manner; this results in decreased secretion of IL-12, skewing the response to Th2 [62,63]. In addition, polymorphic ROP5/17/18 complexes can phosphorylate IRGs and protect the parasite from clearance [64]. Whether the identified polymorphic effector proteins have similar or different effects across different host species remains unknown, and the diverse mechanisms used by *T. gondii* to evade and manipulate host immunity hinder the development of an effective vaccine.

Establishment of Latent Infection

A key factor in the pathogenesis and transmission of *T. gondii* is the ability to convert from a rapidly replicating stage (tachyzoite) to a slow-dividing stage (cyst-containing bradyzoites). This tachyzoite–bradyzoite transformation occurs in response to selective pressure imposed by the host immune system. The tachyzoite–bradyzoite transformation, and the protection offered by the highly glycosylated cyst wall, allows *T. gondii* to avoid immune-mediated destruction and persist, effectively invisible, for years within the infected host, increasing the chances of transmission [65]. Significant changes in bradyzoite gene expression are associated with tissue cyst development, and the **apicomplexan AP2 (ApiAP2)** transcription factors are a crucial mechanism regulating this developmental transcriptome. A total of 67 ApiAP2 repressors and activators located in the *T. gondii* genome are responsive to stress signals and are active in tachyzoite–bradyzoite differentiation [66–68]. For example, the *Api2IV-4* gene, normally expressed by tachyzoites, has a role in inhibiting bradyzoite gene expression. Deletion of *Api2IV-4* caused the expression of bradyzoite-specific proteins in tachyzoites, and the mistiming of bradyzoite antigens in Δ *Api2IV-4* tachyzoites induced a potent host immune response that effectively eliminated the parasite and prevented brain tissue cyst formation [68]. However, the mechanisms that allow *T. gondii* to adapt to different tissue environments and permit long-term persistence as dormant tissue cysts remain to be fully elucidated. Therefore, a better

understanding of the molecular mechanisms that drive the tachyzoite–bradyzoite interconversion is required to manage transmission and prevention of *T. gondii* infection.

The Lack of a Standardized Vaccination Protocol

Considerable efforts by many research groups have been put into designing and testing various vaccination strategies against toxoplasmosis. However, there has been a lack of synergy between toxoplasmosis vaccine studies because they vary in the parasite strain used, infective stage, infection dose or route of parasite infection, and host species [14–16]. *T. gondii* parasites have a diverse population structure; however, the strains frequently used to evaluate the protective efficacy of a toxoplasmosis vaccine are type 1 RH and type 2 (Pru and ME49); all of these strains have been maintained in laboratories for many years and may no longer represent the naive isolates; for example, RH strains have lost the ability to form oocysts or tissue cysts




Figure 3. An Integrated Strategy for the Discovery and Testing of Vaccine Candidates in the Postgenomics era. (A) Discovery of novel vaccine *Toxoplasma gondii* candidates. Vaccine candidates can be identified through analysis of the *T. gondii*'s genome (the complete genetic content of *T. gondii*), transcriptome (the complete set of RNA transcripts expressed by *T. gondii* under a certain condition), proteome (the complete set of proteins expressed by *T. gondii* under a certain condition), secretome (the complete set of proteins secreted by *T. gondii* under a certain condition), and surfome (the subset of proteins that are surface exposed). In parallel, the immunoproteome (the set of antigens that interact with the host immune system) can provide a panoramic view of the spectrum, intensity, and dynamics of antibodies binding to *T. gondii* proteins, revealing their immunogenicity potential. (B) Toxoplasmosis biomarkers and immune-correlates of protection across vulnerable host species. Further characterization of protective immune response is required in order to identify novel unambiguous markers that correlate with, and predict, vaccine-induced protective immunity against *T. gondii* infection. (C) Computational (*in silico*) analysis (e.g., comparative genomics, population genetics, stage-specific antigen, and prediction of B and T cell epitopes) can facilitate all aspects of preclinical and clinical vaccine development. (D) The structural vaccinology (the 3D structure of proteins, in particular the structural epitopes of immunogenic antigens) can be a useful aid in any future *T. gondii* vaccine development. (E) Experimental infection models are crucial for testing the clinical efficacy of *T. gondii* vaccine. (F) Animal-free (*in vitro*) systems can provide preclinical proof-of-concept in support of vaccine development.

[69]. In the natural setting, toxoplasmosis is transmitted by oral or congenital routes. However, intraperitoneal challenge has been frequently used in vaccine experiments, and may not reflect a natural infection route. Although oral challenge has been used to mimic infection, most studies have focused on the oral administration of tissue cysts, with few studies using infective oocysts. These inconsistencies in vaccination regimens could hinder the accurate evaluation of the vaccination efficacy. The genetic background of experimental animals also affects the outcome of any vaccination. Immunity conferred by vaccination in mouse models may not reflect the mechanisms implicated in the protective immune response in food-producing animals or humans. Therefore, future vaccination studies should use different *T. gondii* strains, administered through a natural infection route, in different animal models, particularly domestic cats and food-producing animals.

Coinfection with Other Pathogens

In the natural setting, a host may be simultaneously infected with several pathogens (parasites, viruses, and bacteria). Coinfections alter the host's immune response compared with a single-pathogen infection [70]. *T. gondii* infection induces a Th1-type immune response; whereas other pathogens, such as helminths, induce a Th2-type immune response [71]. Whether or not Th2 type immunity induced by prior or concurrent helminth infection can antagonize a protective Th1 type immune response of toxoplasmosis vaccines remains to be determined. Understanding the immune response to mixed infections should be a part of future vaccine development.

Perspectives for Toxoplasmosis Vaccines

Although substantial progress has been made in the search for, and the development of, vaccines to prevent *T. gondii* infection, there is still no effective and safe toxoplasmosis vaccine to eliminate the tissue cysts and/or fully block vertical transmission. Technology-oriented research is the way forward to identify and isolate more immunogenic vaccine candidates. For example, **reverse vaccinology** and **immunomics** offer enormous opportunities to enable targeted antigen discovery, and rational vaccine design (Figure 3) [72–74]. Reverse vaccinology aims to identify the complete repertoire of antigens that a pathogen secretes or expresses on its surface; while immunomics aims to elucidate sets of proteins or peptides that interact with the host immune system and the potential mechanisms involved in these interactions [72–74]. Recent technological advances in computational analysis, and the availability of genomics resources of both host and *T. gondii* have facilitated the identification of novel antigen vaccine candidates [75]. **Structural vaccinology**, which focuses on understanding the structure of immune silent and immune-dominant antigens, is increasingly applied to the rational structure-guided design of immunogens to optimize their immunogenicity, stability, and safety [76]. The application of these genome-based reverse vaccinology techniques to the development of toxoplasmosis vaccines may increase the chance of success.

Concluding Remarks

Decades of toxoplasmosis vaccine development have resulted in the identification and testing of various candidate antigens; however, none have shown durable and adequate protective efficacy. Potential impediments and solutions to the development of an effective toxoplasmosis vaccine have been discussed. There is still much to be learned about the immunopathogenesis of toxoplasmosis and the identification of better immune-correlates and vaccination candidates (see Outstanding Questions). Successful vaccine development depends on a full understanding of the immunopathogenesis of *T. gondii* infection, choosing appropriate candidate antigens, and selecting the right adjuvant or delivery vehicle, as well as testing the efficacy in

Outstanding Questions

Can candidate vaccine antigens identified in mice protect humans or food-producing animals?

Can a humanized animal model predict the protective efficacy of a toxoplasmosis vaccine to reflect human responses?

What can be done to improve the efficacy and safety of live-attenuated vaccines?

How many immunogenic candidates remain to be discovered, and how can we modify existing antigens to make them more immunogenic?

How do coinfections with other pathogens influence the efficacy of vaccination against toxoplasmosis?

How many mechanisms are used by *T. gondii* to evade host immune responses and establish persistent infection?

What new technologies are having the greatest impact on *T. gondii* immunobiology research, and why?

suitable animal models in order to enable a transition from preclinical testing to food-producing animal and domestic cat trials. The future of toxoplasmosis vaccines appears bright, and the clinical data and resources generated by genomics, transcriptomics, proteomics, metabolomics, and other technologies are likely to build substantially on, and invigorate, basic research to develop effective and safe vaccines against toxoplasmosis.

Acknowledgments

Project support was kindly provided by the National Natural Science Foundation of China (Grant Nos. 31802180, 31230073), the National Key Research and Development Program of China (Grant No. 2017YFD0501304), the International Science and Technology Cooperation Project of Gansu Provincial Key Research and Development Program (Grant No. 17JR7WA031), the Elite Program of Chinese Academy of Agricultural Sciences, and the Agricultural Science and Technology Innovation Program (ASTIP) (Grant No. CAAS-ASTIP-2016-LVRI-03).

Disclaimer Statement

The authors declare that they have no conflicts of interest.

References

- Montoya, J.G. and Liesenfeld, O. (2004) Toxoplasmosis. *Lancet* 363, 1965–1976
- Robert-Gangneux, F. and Dardé, M.L. (2012) Epidemiology of and diagnostic strategies for toxoplasmosis. *Clin. Microbiol. Rev.* 25, 264–296
- Elsheikha, H.M. (2008) Congenital toxoplasmosis: priorities for further health promotion action. *Public Health* 122, 335–353
- Wang, Z.D. *et al.* (2017) Prevalence and burden of *Toxoplasma gondii* infection in HIV-infected people: a systematic review and meta-analysis. *Lancet HIV* 4, e177–e188
- Dunay, I.R. *et al.* (2018) Treatment of toxoplasmosis: historical perspective, animal models, and current clinical practice. *Clin. Microbiol. Rev.* 31, e00057–17
- Buxton, D. and Innes, E.A. (1995) A commercial vaccine for ovine toxoplasmosis. *Parasitology* 110, S11–S16
- Yarovinsky, F. (2014) Innate immunity to *Toxoplasma gondii* infection. *Nat. Rev. Immunol.* 14, 109–121
- Pittman, K.J. and Knoll, L.J. (2015) Long-term relationships: the complicated interplay between the host and the developmental stages of *Toxoplasma gondii* during acute and chronic infections. *Microbiol. Mol. Biol. Rev.* 79, 387–401
- Hunter, C.A. and Sibley, L.D. (2012) Modulation of innate immunity by *Toxoplasma gondii* virulence effectors. *Nat. Rev. Microbiol.* 10, 766–778
- Sayles, P.C. *et al.* (2000) B cells are essential for vaccination-induced resistance to virulent *Toxoplasma gondii*. *Infect. Immun.* 68, 1026–1033
- Gigley, J.P. *et al.* (2009) Long-term immunity to lethal acute or chronic type II *Toxoplasma gondii* infection is effectively induced in genetically susceptible C57BL/6 mice by immunization with an attenuated type I vaccine strain. *Infect. Immun.* 77, 5380–5388
- Hwang, S. and Khan, I.A. (2015) CD8⁺ T cell immunity in an encephalitis model of *Toxoplasma gondii* infection. *Semin. Immunopathol.* 37, 271–279
- Dard, C. *et al.* (2016) Relevance of and new developments in serology for toxoplasmosis. *Trends Parasitol.* 32, 492–506
- Li, Y. and Zhou, H. (2018) Moving towards improved vaccines for *Toxoplasma gondii*. *Expert Opin. Biol. Ther.* 18, 273–280
- Zhang, N.Z. *et al.* (2015) Recent advances in developing vaccines against *Toxoplasma gondii*: an update. *Expert Rev. Vaccines* 14, 1609–1621
- Hiszczyska-Sawicka, E. *et al.* (2014) Veterinary vaccines against toxoplasmosis. *Parasitology* 141, 1365–1378
- Li, Z.Y. *et al.* (2014) Synergy of mL-21 and mL-15 in enhancing DNA vaccine efficacy against acute and chronic *Toxoplasma gondii* infection in mice. *Vaccine* 32, 3058–3065
- Wang, L.J. *et al.* (2018) Protective immune response against *Toxoplasma gondii* elicited by a novel yeast-based vaccine with microneme protein 16. *Vaccine* 36, 3943–3948
- Lee, S.H. *et al.* (2018) Virus-like particle vaccines expressing *Toxoplasma gondii* rhoptry protein 18 and microneme protein 8 provide enhanced protection. *Vaccine* 36, 5692–5700
- Machado, A.V. *et al.* (2010) Prime and boost immunization with influenza and adenovirus encoding the *Toxoplasma gondii* surface antigen 2 (SAG2) induces strong protective immunity. *Vaccine* 28, 3247–3256
- Buxton, D. *et al.* (1991) Vaccination of sheep with a live incomplete strain (S48) of *Toxoplasma gondii* and their immunity to challenge when pregnant. *Vet. Rec.* 129, 89–93
- Wang, J.L. *et al.* (2016) The past, present, and future of genetic manipulation in *Toxoplasma gondii*. *Trends Parasitol.* 32, 542–553
- Sloves, P.J. *et al.* (2015) Apical organelle secretion by *Toxoplasma* controls innate and adaptive immunity and mediates long-term protection. *J. Infect. Dis.* 212, 1449–1458
- Kim, S.K. and Boothroyd, J.C. (2005) Stage-specific expression of surface antigens by *Toxoplasma gondii* as a mechanism to facilitate parasite persistence. *J. Immunol.* 174, 8038–8048
- Fang, R.H. and Zhang, L. (2016) Nanoparticle-based modulation of the immune system. *Annu. Rev. Chem. Biomol. Eng.* 7, 305–326
- Mintern, J.D. *et al.* (2013) Targeting dendritic cells: the role of specific receptors in the internalization of polymer capsules. *Adv. Healthc. Mater.* 2, 940–944
- Skwarczynski, M. *et al.* (2010) Polyacrylate dendrimer nanoparticles: a self-adjuncting vaccine delivery system. *Angew. Chem. Int. Ed. Engl.* 49, 5742–5745
- Dacoba, T.G. *et al.* (2017) Modulating the immune system through nanotechnology. *Semin. Immunol.* 34, 78–102
- Assolini, J.P. *et al.* (2017) Nanomedicine advances in toxoplasmosis: diagnostic, treatment, and vaccine applications. *Parasitol. Res.* 116, 1603–1615
- Roozbehani, M. *et al.* (2018) Characterization of a multi-epitope peptide with selective MHC-binding capabilities encapsulated in PLGA nanoparticles as a novel vaccine candidate against *Toxoplasma gondii* infection. *Vaccine* 36, 6124–6132
- Rahimi, M.T. *et al.* (2017) Immunological evaluation of a DNA cocktail vaccine with co-delivery of calcium phosphate nanoparticles (CaPNs) against the *Toxoplasma gondii* RH strain in BALB/c mice. *Parasitol. Res.* 116, 609–616
- Chahal, J.S. *et al.* (2016) Dendrimer-RNA nanoparticles generate protective immunity against lethal Ebola, H1N1 influenza, and *Toxoplasma gondii* challenges with a single dose. *Proc. Natl. Acad. Sci. U. S. A.* 113, E4133–E4142

33. Théry, C. *et al.* (2002) Exosomes: composition, biogenesis and function. *Nat. Rev. Immunol.* 2, 569–579
34. Raposo, G. and Stoorvogel, W. (2013) Extracellular vesicles: exosomes, microvesicles, and friends. *J. Cell Biol.* 200, 373–383
35. Coakley, G. *et al.* (2015) Exosomes and other extracellular vesicles: the new communicators in parasite infections. *Trends Parasitol.* 31, 477–489
36. Beauvillain, C. *et al.* (2009) Exosomes are an effective vaccine against congenital toxoplasmosis in mice. *Vaccine* 27, 1750–1757
37. Li, Y. *et al.* (2018) Characterization of exosomes derived from *Toxoplasma gondii* and their functions in modulating immune responses. *Int. J. Nanomed.* 13, 467–477
38. Astronomo, R.D. and Burton, D.R. (2010) Carbohydrate vaccines: developing sweet solutions to sticky situations? *Nat. Rev. Drug. Discov.* 9, 308–324
39. Rodrigues, J.A. *et al.* (2015) Parasite glycobiology: a bittersweet symphony. *PLoS Pathog.* 11, e1005169
40. Jaurigue, J.A. and Seeberger, P.H. (2017) Parasite carbohydrate vaccines. *Front. Cell. Infect. Microbiol.* 7, 248
41. Debierre-Grockiego, F. *et al.* (2007) Activation of TLR2 and TLR4 by glycosylphosphatidylinositols derived from *Toxoplasma gondii*. *J. Immunol.* 179, 1129–1137
42. Götz, S. *et al.* (2015) Investigation of the protective properties of glycosylphosphatidylinositol-based vaccine candidates in a *Toxoplasma gondii* mouse challenge model. *Glycobiology* 25, 984–991
43. Howe, D.K. and Sibley, L.D. (1995) *Toxoplasma gondii* comprises three clonal lineages: correlation of parasite genotype with human disease. *J. Infect. Dis.* 172, 1561–1566
44. Khan, A. *et al.* (2007) Recent transcontinental sweep of *Toxoplasma gondii* driven by a single monomorphic chromosome. *Proc. Natl. Acad. Sci. U. S. A.* 104, 14872–14877
45. Saeij, J.P. *et al.* (2005) Differences among the three major strains of *Toxoplasma gondii* and their specific interactions with the infected host. *Trends Parasitol.* 21, 476–481
46. Su, C. *et al.* (2012) Globally diverse *Toxoplasma gondii* isolates comprise six major clades originating from a small number of distinct ancestral lineages. *Proc. Natl. Acad. Sci. U. S. A.* 109, 5844–5849
47. Shwab, E.K. *et al.* (2018) Human impact on the diversity and virulence of the ubiquitous zoonotic parasite *Toxoplasma gondii*. *Proc. Natl. Acad. Sci. U. S. A.* 115, E6956–E6963
48. Lorenzi, H. *et al.* (2016) Local admixture of amplified and diversified secreted pathogenesis determinants shapes mosaic *Toxoplasma gondii* genomes. *Nat. Commun.* 7, 10147
49. Jensen, K.D. *et al.* (2015) *Toxoplasma gondii* superinfection and virulence during secondary infection correlate with the exact ROP5/ROP18 allelic combination. *mBio* 6, e02280
50. Dao, A. *et al.* (2001) Successful reinfection of chronically infected mice by a different *Toxoplasma gondii* genotype. *Int. J. Parasitol.* 31, 63–65
51. Wang, Z.X. *et al.* (2017) Proteomic differences between developmental stages of *Toxoplasma gondii* revealed by iTRAQ-based quantitative proteomics. *Front. Microbiol.* 8, 985
52. Szabo, E.K. and Finney, C.A.M. (2017) *Toxoplasma gondii*: one organism, multiple models. *Trends Parasitol.* 33, 113–127
53. Koblansky, A.A. *et al.* (2013) Recognition of profilin by Toll-like receptor 12 is critical for host resistance to *Toxoplasma gondii*. *Immunity* 38, 119–130
54. Howard, J.C. *et al.* (2011) The IRG protein-based resistance mechanism in mice and its relation to virulence in *Toxoplasma gondii*. *Curr. Opin. Microbiol.* 14, 414–421
55. Hakimi, M.A. *et al.* (2017) *Toxoplasma* effectors targeting host signaling and transcription. *Clin. Microbiol. Rev.* 30, 615–645
56. Bougdour, A. *et al.* (2013) Host cell subversion by *Toxoplasma* GRA16, an exported dense granule protein that targets the host cell nucleus and alters gene expression. *Cell Host Microbe* 13, 489–500
57. Braun, L. *et al.* (2013) A *Toxoplasma* dense granule protein, GRA24, modulates the early immune response to infection by promoting a direct and sustained host p38 MAPK activation. *J. Exp. Med.* 210, 2071–2086
58. Olias, P. *et al.* (2016) *Toxoplasma* effector recruits the Mi-2/NuRD complex to repress STAT1 transcription and block IFN- γ -dependent gene expression. *Cell Host Microbe* 20, 72–82
59. Gay, G. *et al.* (2016) *Toxoplasma gondii* TgIST co-opts host chromatin repressors dampening STAT1-dependent gene regulation and IFN- γ -mediated host defenses. *J. Exp. Med.* 213, 1779–1798
60. Ma, J.S. *et al.* (2014) Selective and strain-specific NFAT4 activation by the *Toxoplasma gondii* polymorphic dense granule protein GRA6. *J. Exp. Med.* 211, 2013–2032
61. Rosowski, E.E. *et al.* (2011) Strain-specific activation of the NF- κ B pathway by GRA15, a novel *Toxoplasma gondii* dense granule protein. *J. Exp. Med.* 208, 195–212
62. Butcher, B.A. *et al.* (2011) *Toxoplasma gondii* rhoptry kinase ROP16 activates STAT3 and STAT6 resulting in cytokine inhibition and arginase-1-dependent growth control. *PLoS Pathog.* 7, e1002236
63. Jensen, K.D. *et al.* (2011) *Toxoplasma* polymorphic effectors determine macrophage polarization and intestinal inflammation. *Cell Host Microbe* 9, 472–483
64. Etheridge, R.D. *et al.* (2014) The *Toxoplasma* pseudokinase ROP5 forms complexes with ROP18 and ROP17 kinases that synergize to control acute virulence in mice. *Cell Host Microbe* 15, 537–550
65. Jeffers, V. *et al.* (2018) A latent ability to persist: differentiation in *Toxoplasma gondii*. *Cell. Mol. Life Sci.* 75, 2355–2373
66. Holmes, M.J. *et al.* (2017) Translational control in the latency of apicomplexan parasites. *Trends Parasitol.* 33, 947–960
67. Lesage, K.M. *et al.* (2018) Cooperative binding of ApiAP2 transcription factors is crucial for the expression of virulence genes in *Toxoplasma gondii*. *Nucleic Acids Res.* 46, 6057–6068
68. Radke, J.B. *et al.* (2018) Transcriptional repression by ApiAP2 factors is central to chronic toxoplasmosis. *PLoS Pathog.* 14, e1007035
69. Dubey, J.P. *et al.* (1999) Infection and immunity with the RH strain of *Toxoplasma gondii* in rats and mice. *J. Parasitol.* 85, 657–662
70. Damania, B. and Dittmer, D.P. (2014) What lies within: coinfections and immunity. *Cell Host Microbe* 16, 145–147
71. Inclán-Rico, J.M. and Siracusa, M.C. (2018) First responders: innate immunity to helminths. *Trends Parasitol.* 34, 861–880
72. Sette, A. and Rappuoli, R. (2010) Reverse vaccinology: developing vaccines in the era of genomics. *Immunity* 33, 530–541
73. Rinaudo, C.D. *et al.* (2009) Vaccinology in the genome era. *J. Clin. Invest.* 119, 2515–2525
74. Kassegne, K. *et al.* (2016) Immunomic approaches for antigen discovery of human parasites. *Expert Rev. Proteomics* 13, 1091–1101
75. Goodswen, S.J. *et al.* (2018) A gene-based positive selection detection approach to identify vaccine candidates using *Toxoplasma gondii* as a test case protozoan pathogen. *Front. Genet.* 9, 332
76. Gourlay, L. *et al.* (2017) Structure and computation in immunoreagent design: from diagnostics to vaccines. *Trends Biotechnol.* 35, 1208–1220
77. Fox, B.A. and Bzik, D.J. (2002) *De novo* pyrimidine biosynthesis is required for virulence of *Toxoplasma gondii*. *Nature* 415, 926–929
78. Ismael, A.B. (2006) Mic1-3 knockout of *Toxoplasma gondii* is a successful vaccine against chronic and congenital toxoplasmosis in mice. *J. Infect. Dis.* 194, 1176–1183
79. Fox, B.A. and Bzik, D.J. (2010) Avirulent uracil auxotrophs based on disruption of orotidine-5'-monophosphate decarboxylase elicit protective immunity to *Toxoplasma gondii*. *Infect. Immun.* 78, 3744–3752
80. Fox, B.A. and Bzik, D.J. (2015) Nonreplicating, cyst-defective type II *Toxoplasma gondii* vaccine strains stimulate protective immunity against acute and chronic infection. *Infect. Immun.* 83, 2148–2155

81. Lagal, V. *et al.* (2015) AMA1-deficient *Toxoplasma gondii* parasites transiently colonize mice and trigger an innate immune response that leads to long-lasting protective immunity. *Infect. Immun.* 83, 2475–2486
82. Arroyo-Olarte, R.D. *et al.* (2015) Phosphatidylthreonine and lipid-mediated control of parasite virulence. *PLoS Biol.* 13, e1002288
83. Wang, J.L. *et al.* (2017) Immunization with *Toxoplasma gondii* GRA17 deletion mutant induces partial protection and survival in challenged mice. *Front. Immunol.* 8, 730
84. Wang, J.L. *et al.* (2018) Live attenuated Pru:Ncdpk2 strain of *Toxoplasma gondii* protects against acute, chronic, and congenital toxoplasmosis. *J. Infect. Dis.* 218, 768–777
85. Mévélec, M.N. *et al.* (2010) Mic1-3 knockout *Toxoplasma gondii* is a good candidate for a vaccine against *T. gondii*-induced abortion in sheep. *Vet. Res.* 41, 49