



True Lumen Stabilization to Overcome Malperfusion in Acute Type I Aortic Dissection

Konstantinos Tsagakis, MD, PhD,* Rolf A. Jánosi, MD, PhD,[†] Ulrich H. Frey, MD, PhD,[‡] Thomas Schlosser, MD, PhD,[§] Roberto Chiesa, MD, PhD,^{||} Tienush Rassaf, MD, PhD,[†] and Heinz Jakob, MD, PhD*

Acute type I aortic dissection (AD) complicated by true lumen (TL) collapse and malperfusion downstream is associated with devastating prognosis. The study reports an institutional mid-term experience with TL stabilization by uncovered stents to restore perfusion as a supplement to proximal thoracic aortic surgery. Between January 2007 and May 2017, 181 out of 270 acute type A AD patients were operated on type I AD. Eighteen uncovered stents (10%) were used to expand the aortic TL in presence of visceral and/or peripheral malperfusion. The procedures took place in a hybrid operating room and were combined with proximal aortic surgery. During follow-up (mean \pm standard deviation 3.44 ± 2.1 years), the fate of AD was evaluated by computed tomography. Indication for TL stenting included visceral (44%) or peripheral malperfusion (11%) or both (45%). Stenting of aortic branches followed in 33%. All patients underwent proximal repair and were combined with frozen elephant trunk (67%) or retrograde descending aorta stent grafting (11%). Thirty-day mortality was 16.7%. Two-year survival was 71.8%. The false lumen around the uncovered stents remained patent in 89% and the aortic diameter increased 0.1 cm/y. No intimal rupture or occlusion of arteries occurred. In 1 patient, the stented aortic lumen was visualized after 6.3 years and neointima ingrowth covering the nitinol frame was found. In acute type I AD, combined endovascular-surgical procedures in a hybrid operation room setting can be used safely to resolve distal malperfusion. Encapsulation of uncovered stents within the intimal wall provides a stable fundament for endovascular techniques to close entry tears and false lumen.

Semin Thoracic Surg 31:740–748 © 2018 The Author. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Keywords: acute aortic dissection, malperfusion, frozen elephant trunk, stenting

Abbreviations: AD, aortic dissection; FL, false lumen; FET, frozen elephant trunk; TL, true lumen; SD, standard deviation

*Department of Thoracic and Cardiovascular Surgery, West German Heart and Vascular Centre Essen, University Hospital of Essen, University of Duisburg-Essen, Essen, Germany

[†]Department of Cardiology and Vascular Medicine, West German Heart and Vascular Centre Essen, University Hospital of Essen, University of Duisburg-Essen, Essen, Germany

[‡]Clinic for Anesthesiology and Intensive Care Unit, University Hospital of Essen, University of Duisburg-Essen, Essen, Germany

[§]Institute for Diagnostic and Interventional Radiology and Neuroradiology, University Hospital of Essen, University of Duisburg-Essen, Essen, Germany

^{||}Division of Vascular Surgery, IRCCS San Raffaele Scientific Institute, Milano, Italy

Conflicts of Interest: Heinz Jakob is consultant to JOTEC GmbH.

Funding: None.

Address reprint requests to Konstantinos Tsagakis, MD, Department of Thoracic and Cardiovascular Surgery, West German Heart and Vascular Centre, University Hospital of Essen, Hufeland Str. 55, Essen 45122, Germany. E-mail: konstantinos.tsagakis@uk-essen.de



Combined surgical and endovascular treatment in acute type I aortic dissection.

Central Message

Treatment of true lumen collapse and malperfusion downstream by uncovered aortic stents combined with proximal repair in acute type I dissection is a safe approach in hybrid operating room setting.

Perspective Statement

Uncovered stents allow stabilization of a collapsed true lumen and became covered by neointima without impairment of end-organ perfusion in mid-term. The reinforced dissection's septum by the encapsulated stent may allow selective endovascular closure of residual entries. In case of complete false lumen exclusion, neointima ingrowth may induce complete healing of aortic dissection.

INTRODUCTION

Acute type I aortic dissection (AD) complicated by severe true lumen (TL) collapse and malperfusion has a devastating prognosis.¹ The recommended “straight forward” surgical treatment consisting of immediate sternotomy and proximal aorta replacement enables the release of tamponade and circulatory stabilization and in many cases restoration of TL perfusion in the thoracic aorta, simply by exclusion of the proximal entry tear.² Depressurization of the false lumen (FL), however, does not regularly result in TL expansion, since re-entries in the descending and abdominal aorta may maintain increased pressure in the FL, thus sustaining the distal malperfusion. Furthermore, narrowing of abdominal or peripheral aortic branches as a result of thrombus formation into the FL leads to static malperfusion. Malperfusion may persist throughout the operative treatment despite FL depressurization, increasing the risk for an end-organ ischemia and thus resulting in necrosis, which is associated with a high mortality rate of 40–70%.^{3–5}

To reduce the duration of end-organ ischemia, in 2007, we have started to restore abdominal organ perfusion endovascularly prior to surgery in a hybrid operating room setting using self-expandable uncovered stents.⁶ The study reports our experience with this kind of malperfusion management in acute type I AD patients and presents the durability of the treatment and the fate of residual AD in mid-term follow-up.

PATIENTS AND METHODS

From January 2007 to May 2017, 270 patients underwent emergency surgery for acute type A AD, out of whom 181 had type I AD. Eighteen of them (10%) underwent endovascular TL stabilization along the thoracoabdominal aorta by placing self-expandable uncovered stents. The study was approved from the institutional review board of University of Duisburg-Essen and no informed consent was required. Patients' characteristics are presented in Table 1. Malperfusion was obvious according to clinical symptoms at the time of admission and in association with the computed tomography (CT) findings. Visceral malperfusion was documented in case of abdominal pain, vomiting, hematemesis, or hematochezia. Peripheral malperfusion presented with peripheral pain, pulselessness, or paresis. In intubated and sedated patients, clinical symptoms were documented according to the external clinical report. According to our “Hybrid-Concept” in acute AD,⁷ admission and treatment took place in a hybrid operating room, which incorporates invasive angiography and surgery on the same operating table. Invasive angiography was used to confirm the severity of TL collapse and malperfusion, and to differentiate between static and dynamic malperfusion. In case of dynamic malperfusion, a threshold of 6 hours of nearly complete visceral malperfusion after onset of symptoms was chosen to restore distal perfusion prior to surgery. In case of a shorter than 6-hour period of dynamic malperfusion, surgery of the thoracic aorta was immediately performed followed by repeat angiography postoperatively, to detect potential persistence of malperfusion. In such a case, additional stenting was undertaken. In case of static

Table 1. Preoperative Characteristics

N (%), Mean ± SD	N = 18
Age	55.4 ± 30
Male	15 (83)
Intubated before admission	7 (39)
Tamponade	4 (22)
Penn classification*	
A	—
B	11 (61)
C	—
BC	7 (39)
Malperfusion related symptoms	
Neurologic deficit	9 (50)
cerebral	5/9
peripheral	6/9
Abdominal pain	11 (61)
Vomiting	10 (56)
Hematemesis	2 (11)
Hematochezia	1 (6)
One side limb ischemia	4 (22)
Both side limb ischemia	5 (28)
No symptoms/unknown	
Aortic valve regurgitation	
Mild	6 (33)
Moderate	5 (28)
Severe	5 (28)
Coronary artery disease	7 (39)
Creatinine >1.5 mg/dL	8 (44)

*Classification of acute AD according to University of Pennsylvania.²⁶

malperfusion, TL and abdominal aortic side branch perfusion was restored prior to surgery independently from malperfusion duration. In case of evidence for bowel necrosis, explorative laparotomy was performed and an interim stay in the ICU to overcome the metabolic disorder prior to proximal aortic repair in otherwise hemodynamically stable patients followed.⁶ Coronary angiography was performed in all patients over 50 years to exclude concomitant coronary artery disease.

HYBRID ROOM CONCEPT

All patients with suspected aortic syndrome are submitted to our hybrid room since 2005 with an aortic team awaiting the patient, consisting of a cardiac surgeon, cardiologist, and anesthesiologist. The patient was controlled hemodynamically by placing arterial lines into both radial arteries and 1 femoral artery as well as central venous catheters including a pulmonary artery catheter in sedation followed by intubation when the patient was admitted breathing spontaneously. Transesophageal echocardiography has been performed for confirmation of suspected AD and evaluation of tamponade and/or aortic valve regurgitation. In circulatory stable patients, angiography was performed first through the femoral artery, using an 8F sheath with a side-port for additional monitoring of the artery pressure. In circulatory unstable patients, immediate tamponade release via median sternotomy and cautious pericardial drainage placement took place. Angiography followed under stable

conditions. In case of cardiopulmonary resuscitation or free rupture, immediate surgery was started and invasive diagnostics followed postoperatively.

ENDOVASCULAR TL TREATMENT

Stenting was performed through a surgically exposed femoral artery and over a Back-up Meier Guide Wire (Boston Scientific, Marlborough, MA). The limb arising from the TL was chosen, even in case of malperfusion. Dynamic malperfusion was treated by TL expansion with the sinus-XL (Optimed, Ettlingen, Germany) or the E-XL (JOTEC, Hechingen, Germany) 100–130 mm in length stents. The first uncovered stent was deployed at the proximal part of TL collapse, commonly at the diaphragmatic level. A serial stent implantation distally was added, if required (Fig. 1A). Narrowed visceral arteries were stabilized by balloon-expandable stents through the frame of the aortic stents (Fig. 1B). Stent diameter was chosen according to the measurement of maximal TL

diameter perpendicular to the course of the aorta in axial CT image or according to intravascular ultrasound. Maximal TL diameter was defined as the distance between the anchor points of the dissected intimal wall. A one size smaller or equal stent diameter to TL maximal diameter was chosen. Oversizing was avoided in order to reduce the risk of stent infolding.

AORTIC SURGERY

Our surgical management in acute type I AD has been described previously.⁸ In brief, cannulation site for the extracorporeal circulation varied according to the extent of AD into the arch vessels. In case of absent brachiocephalic trunk dissection, cannulation of the right axillary artery was favored. Otherwise, central cannulation of the ascending aorta under direct vision during a short period of warm circulatory arrest (≤ 120 seconds) was preferred.⁹ Entries in the ascending aorta and the arch were resected and the distal anastomosis was performed




Figure 1. True lumen stabilization of the thoracoabdominal aorta by uncovered stents to resolve mesenteric malperfusion in acute type I aortic dissection combined with proximal descending aorta endografting (A) or frozen elephant trunk (B). True lumen collapse (C) was abolished by isolated stenting of the abdominal aorta (D). In static malperfusion (E), additional artery stenting was performed (F).

in open fashion. Patients with full circular arch AD and/or re-entries in the proximal descending aorta underwent frozen elephant trunk (FET) surgery using the E-vita open (JOTEC GmbH, Hechingen, Germany) prosthesis. FET was performed in combination with left subclavian artery debranching using an 8-mm prosthesis in order to facilitate the arch replacement and to perfuse all arch arteries during arch repair and to increase spinal cord perfusion via collaterals. In addition, selective descending aorta perfusion was re-established immediately in 7 patients after FET fixation using a Foley catheter as endo-clamp into the body of the stent graft in order to reduce visceral ischemia.¹⁰ Circulatory arrest distally was initiated at 28°C bladder temperature and arch repair was performed under bilateral or triple selective cerebral perfusion at 22°C blood temperature. Cerebral saturation was controlled by near infrared spectroscopy. The TL of the descending aorta was visualized endoscopically using an autoclavable flexible bronchovideoscope (BF type Q180-AC, Olympus), so called angioscope or aortoscope, to define re-entries and to guide the FET deployment.¹¹

DEFINITIONS AND STATISTICS

Malperfusion was confirmed and defined according to invasive angiography findings. Dynamic malperfusion was identified where the AD crossed the vessels' origin compromising perfusion.¹² Static malperfusion was identified where the dissection extended into the vessels' origin narrowing the TL. To evaluate the fate of the FL downstream, the aorta was divided into 3 segments. The proximal and mid part of the descending aorta down to TH 8 level was defined as segment A, distally down to coeliac trunk as segment B, and more distally down to bifurcation as segment C. Measurements of aortic diameter (cm) and area (cm²) of the aorta and TL were done at the level of pulmonary artery bifurcation, distal descending aorta, coeliac trunk, and inferior mesenteric artery. Measurements were undertaken in cross-sectional images after 3-dimensional multiplanar reconstruction. The SPSS Package 24 was used for statistical analysis. The Shapiro–Wilk test was used to evaluate data distribution. The paired-samples *t* test or the Wilcoxon signed-rank test was used to determine the differences of aortic diameter and area between the first and last CT examination, if




Figure 2. A 59-year-old patient presented with abdominal pain, hematemesis, and pulselessness distally. To reopen the collapsed true lumen (TL), an uncovered stent was introduced retrogradely. Wire and stent crossed over an entry at the superior mesenteric artery (SMA) level and the stent was landed partially within the false lumen (FL; arrow) resulting in TL occlusion proximally. To overcome this complication, the TL was expanded upstream by placement of 2 additional stents through the opened arch under angioscopic guidance. Proximal aorta repair and frozen elephant trunk (E-vita open plus) placement in zone 2 combined with subclavia artery debranching followed. The 18-month postoperative CT demonstrates the proximal part of the stent into the FL (A). The stented TL and visceral arteries remained open (B). (C) It demonstrates the angioscopic view into the entry tear in the abdominal aorta and the stent and pigtail catheter (*) crossing into the FL. A back-up guide wire (**) was placed through the channel of the angioscope beside the stent into the infrarenal TL. Two uncovered stents were guided over the wire downstream (D) and reopened the TL after deployment (E).

appropriate. Continuous variables are presented in mean ± standard deviation. Categorical variables are presented in percent. Survival and freedom from reintervention postoperatively were estimated by Kaplan–Meier analysis.

RESULTS

Malperfusion Treatment

The first diagnosis of AD was established elsewhere by CT in 14 patients (78%) and by echocardiography in 4 patients (22%). The delay between onset of symptoms and admission to the hybrid room (15 ± 33 hours) was <6 hours in 11 patients (61%) and >6 hours in 7 patients (39%). At the time of admission, 7 patients (39%) were hemodynamically compromised. Severe tamponade occurred in 4 patients. In 2 of them, invasive angiography was performed immediately after tamponade release. In the other 2, angiography followed the aortic surgery postoperatively.

Angiography demonstrated TL collapse inducing malperfusion in the mesenteric aortic part in 8 (44%), the infrarenal part in 2 (12%), and both in 8 (44%). Static malperfusion into the aortic branches was observed in 6 (33%): 2 in visceral arteries, 2 in visceral and iliac arteries, and 2 in iliac arteries only. TL stabilization by uncovered stents was achieved via the femoral artery in 16 and through the opened arch under angioscopic guidance in 2 patients. The reason to use the antegrade route was a persisting TL collapse after stent displacement in one (Fig. 2, Supplementary Video 1) and a large residual entry tear beyond the distal end of the FET in the other. In total, 35 uncovered stents, 7 sinus-XL and 28 E-XL stents, and 2 covered stents were placed. Stent deployment took place in segment A in 5 (28%), segment B in 14 (78%), and segment C in 15 (83%; Table 2). Static malperfusion could be resolved preoperatively by stenting of the aorta and the occluded arteries in all cases. In 4 of them, proximal aortic repair was performed after explorative laparotomy and an interim stay in ICU of 10.6 ± 29.7 hours to overcome severe metabolic disorder. Proximal repair consisted of ascending aorta ± arch replacement or FET surgery in 6 and 12 patients, respectively. In addition, aortic valve repair and/or replacement or coronary artery bypass grafting for coronary artery disease was required in 15 patients (83%). The time of cardiopulmonary bypass, cardioplegic arrest, visceral perfusion arrest, and selective cerebral perfusion was 261 ± 57, 142 ± 38, 48 ± 25 and 60 ± 21 minutes, respectively.

Postoperative Results

In-hospital and 30-day mortality was 16.7%. Cause of death was myocardial infarction in 1 and bowel necrosis in 2 patients after dynamic visceral malperfusion. In one of them, the delay to endovascular treatment was 14 hours. The other patient was admitted after 6 hours of delay but with Leriche syndrome. In addition, residual visceral and peripheral ischemia after the primary endovascular repair was observed in 2 (11%) and 2 (11%) patients, respectively. One patient underwent bowel resection prior to proximal aortic repair. The second patient

Table 2. Endovascular and Surgical Aortic Treatment

N (%) Mean ± SD	Malperfusion Site Downstream			
	Mesenteric N = 8	Peripheral N = 2	Both N = 8	Total N = 18
<i>Endovascular treatment</i>				
<i>Time of intervention</i>				
Preoperative	5 (63)	1 (50)	7 (88)	13 (72)
Intraoperative	1 (12)	—	1 (12)	2 (11)
Postoperative	2 (25)	1 (50)	1 (12)	4 (22)
<i>Aorta stenting</i>				
Segment A	1 (12)	—	4 (50)	5 (28)
Segment B	6 (75)	—	8 (100)	14 (78)
Segment C	5 (62)	2 (100)	8 (100)	15 (83)
<i>Arteries stenting</i>				
Visceral arteries	2 (25)	—	2 (25)	4 (22)
Iliac arteries	—	1 (50)	3 (37)	4 (22)
<i>Surgical treatment</i>				
<i>Aorta repair extension</i>				
Ascending aorta	—	1 (50)	2 (25)	3 (17)
Ascending aorta + arch	1 (12)	—	2 (25)	3 (17)
Ascending aorta + FET	7 (88)	1 (50)	4 (50)	12 (67)
<i>Perfusion management</i>				
Bilateral selective cerebral perfusion	4 (50)	1 (50)	5 (62)	10 (56)
Triple selective cerebral perfusion*	4 (50)	1 (50)	3 (37)	8 (44)
Selective descending aorta perfusion	2 (25)	1 (50)	4 (50)	7 (39)
<i>Aortic root replacement</i>				
Partial	2 (25)	1 (50)	2 (25)	5 (28)
Total	3 (37)	1 (50)	2 (25)	6 (33)
<i>Aortic valve</i>				
Repair	4 (50)	1 (50)	2 (25)	7 (39)
Replacement	3 (37)	1 (50)	2 (25)	6 (33)
CABG	5 (62)	—	1 (12)	6 (33)

*Both carotid arteries and left subclavia artery perfusion.

underwent endoscopy and medical treatment postoperatively. Peripheral interventions were performed in the distal extremities by thromboendarterectomy in one and transcatheter treatment in the second. No new stroke occurred. All patients with a preoperative cerebral neurologic deficit did recover but one. Eleven patients (61%) underwent temporary renal replacement therapy due to acute kidney injury. Of them, 10 had TL collapse at the visceral aortic part, 3 had additional renal artery dissection, and 1 had static renal artery malperfusion. No patient required dialysis after discharge. Hospital stay was 28.2 ± 21.5 days. All discharged patients except for 1—due to residual neurologic deficit—turned back to normal life.

Follow-up lasted 3.44 ± 2.1 years and estimated survival after 2 years was 71.8% (95% confidence interval 1.167–1.890). Four of 15 discharged patients died. Cause of death was secondary thoracoabdominal aortic surgery, unknown death at home, pneumonia after permanent stroke, and suicide. Three patients underwent secondary aortic intervention: 1 thoracoabdominal aortic replacement for infrarenal abdominal aneurysm as mentioned above, 1 proximal arch replacement due to pseudoaneurysm formation at the distal anastomosis after ascending aorta repair, and 1 FET for progressive distal arch enlargement after

2, 6, and 8 years, respectively. No patient underwent an endovascular reintervention.

Aortic Results

In the first postoperative CT examination, complete or partial FL thrombosis in segment A was documented in 61% and 28%, respectively (Table 3). Complete FL patency occurred in 11% and only in absence of FET treatment. In segments B and C, the FL remained completely or partially perfused in 83% and 89%, respectively. Infolding of the uncovered stents occurred in 3 patients (17%): 2 with sinus-XL and 1 with E-XL stents implanted. In the last CT examination, complete or partial FL patency in segments A, B, and C was 33%, 73%, and 87%, respectively. No occlusion of visceral arteries arising either from TL or FL occurred (Fig. 1C–F).

Between the first and last CT, the aortic diameter decreased in the mid-descending aorta 0.1 cm/y ($P = 0.005$) and increased in the distal descending aorta 0.1 cm/y ($P = 0.014$; Table 4). In the abdominal aorta, the aortic diameter increased

proximally ($P = 0.005$) and distally ($P = 0.027$) 0.08 cm/y. The TL/AL ratio increased in mid-descending aorta ($P = 0.005$) and remained stable distally.

Remodeling of Inner Aortic Surface—Case

One patient (55 years, male) was treated initially with uncovered stents in all 3 segments using 4 E-XL stents and ascending aorta repair (Fig. 3A). He had to undergo redo surgery after 6.3 years for pseudoaneurysm in the distal anastomosis. In absence of an entry tear in the aortic arch and proximal descending aorta, the AD across the arch had resolved. The FL in segment A was obliterated and remained perfused in segments B and C (Fig. 3B and D). During secondary surgery and open anastomosis in the proximal arch, the angioscope was used to visualize the aortic lumen downstream (Supplementary Video 2). All uncovered stents were endothelialized and encapsulated into the aortic wall (Fig. 3C). No lesion along the intimal surface was found apart from re-entries in the proximal and distal abdominal aorta. The overstented visceral orifices were open and the new intimal surface did not extend macroscopically into the vessels' offsprings (Fig. 3E).

Table 3. False Lumen in Postoperative and Follow-up Examinations

N (%)	First CT N = 18	Last CT N = 15
Segment A		
Complete thrombosis	11 (61)	10 (67)
Partial thrombosis	5 (28)	3 (20)
No thrombosis	2 (11)	2 (13)
Segment B		
Complete thrombosis	3 (17)	4 (27)
Partial thrombosis	1 (5)	5 (33)
No thrombosis	14 (78)	6 (40)
Segment C		
Complete thrombosis	2 (11)	2 (13)
Partial thrombosis	—	3 (20)
No thrombosis	16 (89)	10 (67)

DISCUSSION

Classic approach to treat accompanying visceral and/or peripheral malperfusion in acute type I AD is the fast replacement of the proximal thoracic aorta, thus achieving exclusion of the proximal entry tear and FL depressurization in many cases. However, the malperfusion associated with increased mortality demonstrates the necessity of alternative strategies to keep the patient alive.^{13,14} Deeb et al reported in 1997 the implementation of primary endovascular restoration of the abdominal perfusion followed by a surgical delay thereafter to overcome metabolic abnormalities and to reduce the surgical trauma in a hemodynamic and metabolic stable patient.¹ Although heavily criticized for this kind of “natural selection” approach, this concept revealed the shortcomings of classic open surgery to

Table 4. Computed Tomography Changes of Descending and Abdominal Aorta Between Postoperative and Last Follow-up Examination

Mean ± SD	Descending Aorta					
	Mid-level			Distal-level		
	Postop	FU	P	Postop	FU	P
Diameter (cm)	3.3 ± 0.5	3.0 ± 0.4	0.005	3.1 ± 0.4	3.4 ± 0.6	0.014
Ratio TL/AL (%)	57 ± 17	81 ± 27	0.005	58 ± 22	57 ± 27	0.792
Mean ± SD	Abdominal Aorta					
	Proximal-level			Distal-level		
	Postop	FU	P	Postop	FU	P
Diameter (cm)	2.9 ± 0.5	3.2 ± 0.6	0.005	2.4 ± 0.7	2.6 ± 0.8	0.027
Ratio TL/AL (%)	65 ± 19	63 ± 25	0.196	65 ± 23	61 ± 26	0.280

AL, aortic lumen; FU, follow-up; TL, true lumen.




Figure 3. Documentation of uncovered stents endothelialization in the thoracoabdominal aorta in 2 patients. Case 1 (A–E): Status after proximal aortic repair and thoracoabdominal true lumen stabilization with 3 E-XL uncovered stents for acute type I aortic dissection and open proximal arch resurgery for pseudoaneurysm after 6.3 years (A). The false lumen was excluded in the descending aorta (B) and remained perfused at the superior mesenteric artery level (C). Intraoperative angioscopy downstream during the opened arch resurgery demonstrated ingrowth of neointima along the entire stented aorta (D). Abdominal arteries were open without extension of the neointima into the orifices as visualized in superior mesenteric artery* (E). Case 2 (F) is provided from Roberto Chiesa (Milan, Italy) and confirms the coverage of uncovered stent by neointima in aortic dissection. Thoracoabdominal open aortic repair 18 months after PETTICOAT treatment for acute type B aortic dissection and progressive false lumen enlargement was performed. The uncovered stent was placed from diaphragm down to infrarenal aorta. The nitinol skeleton was covered by neointima-like tissue except of the space crossing the visceral arteries orifices. CT, coeliac trunk; LRA, left renal artery; RRA, right renal artery; SMA, superior mesenteric artery. Due to the encapsulation of the stent into the intimal wall endarterectomy-like manoeuvre to remove the stent became necessary.

deal with extensive mesenteric and peripheral malperfusion in emergency AD situations and introduced a new way to address the malperfusion sequelae. Modern hybrid operating room concepts like ours enable the combination of endovascular and surgical techniques and give the option for on-time diagnostics and treatment. Since no patient transport from a cath lab to an operating room and vice versa is needed, we accepted a short delay to sternotomy and cardiopulmonary bypass in order to evaluate the extent of AD and malperfusion severity by angiography and to decide for a specific treatment. Even in initially circulatory unstable patients due to tamponade invasive diagnostics and treatment is possible after cautious tamponade release and pericardial drainage under blood pressure control.⁷

Implantation of uncovered stents to further expand the TL is frequently applied in complicated acute type B AD and is known as the PETTICOAT concept.^{15,16} Uncovered stents are placed for extension of a proximally deployed stent graft in order to eliminate TL collapse and to improve perfusion distally. Similarly, we used this technique in acute type I AD in reverse fashion by first placing the uncovered stents into the abdominal aorta to resolve malperfusion and then by proceeding to the proximal aortic repair. In contrast to treatment by a stent graft, the uncovered stent allows deployment over the aortic vessels' orifices without need of precise landing except for the aortic bifurcation level. This simple technique justifies its use in emergency situation with minimal loss of time and

less endovascular jeopardies. However, placement of a guide wire within the TL must be secured in order to prevent misplacement, as described from our experience (Fig. 2). Thus, intravascular ultrasound is suggested to follow the wire along the TL of the abdominal aorta.

The stabilization of the TL by uncovered stents could be demonstrated throughout the investigation period but their use had less impact on FL exclusion, which is similar to the experience in type B AD.¹⁷ In contrast to the proximal descending aorta, in which shrinkage of FL and aortic diameter was demonstrated after FET, the FL distally around the uncovered stents remained completely or partially patent in 89%. Although less or no oversizing was applied in our series, we believe that even in case of excessive oversizing, exclusion of FL would fail due to the stent's tendency to infold. The perfused FL resulted into enlargement of the distal aortic segment 0.1 cm/y, which is similarly to other reports.^{18,19} However, the stable ratio of TL/AL area demonstrates that this enlargement refers to a dilatation of the entire aorta rather than to isolated FL progression.


The modest experience with this technique in acute AD raises concerns about the interaction between the nitinol skeleton and the fragile intimal wall as well as about the patency of the overstented abdominal orifices.²⁰ We applied this technique rarely in acute type I AD and only as a bail-out option in order to prevent fatal end-organ necrosis. Nevertheless, the encapsulation of the uncovered stents into the aortic wall 6.3 years after implantation demonstrates neointima-like ingrowth covering the nitinol frame and leaving the side branches untouched. No intimal rupture, no stent migration, and no occlusion of an overstented artery did occur in these patients. The quantitative influence of TL stabilization to blood supply of arteries arising from FL is unknown. However, the 100% patency of the overstented FL arteries and the residual FL perfusion around the uncovered stents supports our previous observation that FL arteries act as “exits” keeping the blood supply intact against thrombosis.²¹ The endothelialization of the stent can be considered as “reinforcement” of the dissected intimal wall. This observation was confirmed from Professor Roberto Chiesa, IRCCS San Raffaele Scientific Institute in Milano, Italy, by removal of an uncovered stent from the abdominal aorta 18 months after PETTICOAT treatment for acute type B AD (Fig. 3F). Similarly to our finding, a neointima-like tissue covered the entire nitinol skeleton but not the visceral arteries orifices. The stent was completely encapsulated into the intimal wall and its removal became possible only after an endarterectomy-like maneuver. This evolution to endothelialized stent and stable intimal wall opens up the option for specific closure of residual entries and FL by endovascular devices in the chronic phase of the disease with probably lesser risk for endoleak or intimal rupture.^{22,23} In addition, the endothelialization process may be supportive for the STABILISE concept,^{24,25} in which the entire aortic lumen is stabilized by uncovered stents after rupture of the dissection's septum by balloon and depressurization of the FL. In this case, the endothelialization process could be an important step toward complete healing of the entire aorta.

CONCLUSION

The application of endovascular diagnostics and treatment as a standard procedure for surgery of acute type I AD in our opinion is justified, when a hybrid operating room set-up is available, especially in patients suffering from visceral and peripheral malperfusion. The presented low mortality in this high-risk patient cohort demonstrates the safety of this hybrid concept in the presence of experienced cardiovascular surgeons and interventionalists. Although the endothelialization process over time of uncovered stent is not known in detail, the coverage of the fenestrated nitinol frame by “neo-intima” opens up opportunities for new endovascular strategies to exclude the residual FL or to heal the entire aorta.

SUPPLEMENTARY MATERIAL

The following is the supplementary data to this article:



Video 1. Endoscopically guided antegrade stenting to expand a collapsed true lumen in the abdominal aorta. The video refers to Figure 2 and demonstrates true lumen stabilization by stent placement downstream through the opened aortic arch in a bail-out situation. The endoscopically guided procedure was performed under selective cerebral perfusion and intermittent hypothermic circulatory arrest distally at 24°C bladder temperature. The endoscopy was performed by an autoclavable videoscope (BF type Q180-AC, Olympus), the so called angioscope. True lumen collapse and the entry tear in the mid-abdominal aortic level are demonstrated. The proximal part of the previously implanted uncovered stent had unintendedly crossed over a reentry tear into the false lumen resulting in collapse of the true lumen. The distal part of the stent was located correctly within the true lumen of the infrarenal aorta. Therefore, a stiff guide wire was advanced through the channel of the angioscope besides the stent skeleton into the true lumen. The angioscope was then removed and placed again to release the wire for stenting. An uncovered E-XL stent was introduced and deployed into the true lumen in over the wire technique thus compressing the first stent. This stenting was extended proximally into the thoracic aorta with an additional uncovered stent to secure the treatment. A complete expansion of the true lumen was achieved. Finally, an E-vita open plus hybrid stent graft was inserted and unfolded within the lumen of the uncovered stent, guided and controlled by angioscopy. The final control angioscopy demonstrates the satisfactory result with full restoration of the true lumen of the thoracoabdominal aorta. The time of selective cerebral perfusion and intermittent hypothermic circulatory arrest was 56 minutes and 16 minutes, respectively.



Video 2. Endoscopic visualization of stented thoracoabdominal aortic lumen. Presentation of the endoscopic visualization of the descending and abdominal aortic lumen 6.3 years (Fig. 3) after thoracoabdominal serial stenting with uncovered E-XL stents and proximal aorta replacement for acute type I AD. The procedure was performed through the opened arch under ongoing selective cerebral perfusion (50 mm Hg, 1000 mL/min) and hypothermic circulatory arrest at 28°C distally. Cause for the reoperation was a pseudoaneurysm formation in the distal ascending aorta involving the arch. A sump sucker placed into the descending aorta was used to create a bloodless field. Beginning from the proximal descending aorta, downstream endothelialization of the nitinol frame of the stents throughout the entire stented aorta is demonstrated. The markedly reduced blood flow indicates compromised collateralization. In addition, rise of spinal arteries from false lumen was detected in the CT examination. At the ending of the stents in the distal abdominal aorta, the true lumen is collapsed due to the circulatory arrest situation and the residual false lumen. Close to the distal end of the stent, an entry tear in clock position 5 was identified. Pulling the angioscope back to the level of the visceral arteries, the orifices were visualized. The artery orifice in clock position 2 is not affected from neointima ingrowth. This orifice refers to the superior mesenteric artery according to the CT imaging. Similar finding is presented in clock position 10 at the same video level. The orifice refers to a renal artery and the lumen is unaffected. Proximally, in clock position 10 and close to coeliac trunk, a bleeding entry tear is suggested.

REFERENCES

- Deeb GM, Williams DM, Bolling SF, et al: Surgical delay for acute type A dissection with malperfusion. *Ann Thorac Surg* 64:1669–1675, 1997
- Geirsson A, Szeto WY, Pochettino A, et al: Significance of malperfusion syndromes prior to contemporary surgical repair for acute type A dissection: Outcomes and need for additional revascularizations. *Eur J Cardiothorac Surg* 32:255–262, 2007
- Perera NK, Galvin SD, Seevanayagam S, et al: Optimal management of acute type A aortic dissection with mesenteric malperfusion. *Interact Cardiovasc Thorac Surg* 19:290–294, 2014
- Leontyev S, L egar e JF, Borger MA, et al: Creation of a scorecard to predict in-hospital death in patients undergoing operations for acute type A aortic dissection. *Ann Thorac Surg* 101:1700–1706, 2016
- Okita Y: Current surgical results of acute type A aortic dissection in Japan. *Ann Cardiothorac Surg* 5:368–376, 2016
- Tsagakis K, Pizanis N, Kamler M, et al: ICU controlled delay for acute type A aortic dissection repair after intervention for total visceral malperfusion: A way out of a dilemma? *Thorac Cardiovasc Surg* 56:298–300, 2008
- Tsagakis K, Konorza T, Dohle DS, et al: Hybrid operating room concept for combined diagnostics, intervention and surgery in acute type A dissection. *Eur J Cardiothorac Surg* 43:397–404, 2013
- Jakob H, Dohle D, Benedik J, et al: Long-term experience with the E-vita Open hybrid graft in complex thoracic aortic disease. *Eur J Cardiothorac Surg* 51:329–338, 2017
- Jakob H, Tsagakis K, Szabo A, et al: Rapid and safe direct cannulation of the true lumen of the ascending aorta in acute type A aortic dissection. *J Thorac Cardiovasc Surg* 134:244–245, 2007
- Tsagakis K, Dohle DS, Wendt D, et al: Left subclavian artery rerouting and selective perfusion management in frozen elephant trunk surgery. *Minim Invasive Ther Allied Technol* 24:311–316, 2015
- Tsagakis K, Kamler M, Benedik J, et al: Angioscopy—A valuable tool in guiding hybrid stent grafting and decision making during type A aortic dissection surgery. *Eur J Cardiothorac Surg* 38:507–509, 2010
- Erbel R, Alfonso F, Boileau C, et al: Diagnosis and management of aortic dissection. *Eur Heart J* 22:1642–1648, 2001
- Di Eusanio M, Trimarchi S, Patel HJ, et al: Clinical presentation, management, and short-term outcome of patients with type A acute dissection complicated by mesenteric malperfusion: Observations from the International Registry of Acute Aortic Dissection. *J Thorac Cardiovasc Surg* 145:385–390, 2013
- Czerny M, Schoenhoff F, Etz C, et al: The impact of pre-operative malperfusion on outcome in acute type A aortic dissection: Results from the GER-AADA Registry. *J Am Coll Cardiol* 65:2628–2635, 2015
- Nienaber CA, Kische S, Zeller T, et al: Provisional extension to induce complete attachment after stent-graft placement in type B aortic dissection: The PETTICOAT concept. *J Endovasc Ther* 13:738–746, 2006
- He H, Yao K, Nie WP, et al: Modified Petticoat technique with pre-placement of a distal bare stent improves early aortic remodeling after complicated acute Stanford type B aortic dissection. *Eur J Vasc Endovasc Surg* 50:450–459, 2015
- Melissano G, Bertoglio L, Rinaldi E, et al: Volume changes in aortic true and false lumen after the “PETTICOAT” procedure for type B aortic dissection. *J Vasc Surg* 55:641–651, 2012
- Halstead JC, Meier M, Etz C, et al: The fate of the distal aorta after repair of acute type A aortic dissection. *J Thorac Cardiovasc Surg* 133:127–135, 2007
- Kimura N, Tanaka M, Kawahito K, et al: Influence of patent false lumen on long-term outcome after surgery for acute type A aortic dissection. *J Thorac Cardiovasc Surg* 136:1160–1166, 2008
- Kische S, D’Ancona G, Belu IC, et al: Perioperative and mid-term results of endovascular management of complicated type B aortic dissection using a proximal thoracic endoprosthesis and selective distal bare stenting. *Eur J Cardiothorac Surg* 48:77–84, 2015
- Dohle DS, Jakob H, Schucht R, et al: The impact of entries and exits on false lumen thrombosis and aortic remodeling. *Eur J Cardiothorac Surg* 52:508–515, 2017
- K lbel T, Carpenter SW, Lohrenz C, et al: Addressing persistent false lumen flow in chronic aortic dissection: The knickerbocker technique. *J Endovasc Ther* 21:117–122, 2014
- Falkenberg M, Roos H, Lepore V, et al: Endovascular closure of chronic dissection entries in the aortic arch using the Amplatzer vascular plug II as a sealing button. *J Endovasc Ther* 23:378–383, 2016
- Hofferberth SC, Nixon IK, Boston RC, et al: Stent-assisted balloon-induced intimal disruption and relamination in aortic dissection repair: The STABILISE concept. *J Thorac Cardiovasc Surg* 147:1240–1245, 2014
- Matalanis G, Perera NK, Galvin SD: Total aortic repair: The new paradigm in the treatment of acute type A aortic dissection. *Ann Cardiothorac Surg* 5:216–221, 2016
- Augoustides JG, Szeto WY, Desai ND, et al: Classification of acute type A dissection: Focus on clinical presentation and extent. *Eur J Cardiothorac Surg* 39:519–522, 2011