



Review article

TSPO in diverse CNS pathologies and psychiatric disease: A critical review and a way forward



Tomás R. Guilarte

Robert Stempel College of Public Health & Social Work, Environmental Health Sciences, Cognitive Neuroscience & Imaging Florida International University, 11200 SW 8th St, AHC5-507, Miami, FL 33199, USA

ARTICLE INFO

Available online 4 September 2018

Keywords:

Translocator protein 18 kDa (TSPO) - peripheral benzodiazepine receptor (PBR)
Neuroinflammation
Neurodegenerative disease
Positron Emission Tomography (PET)
Biomarker
Schizophrenia
Microglia
Astrocytes

ABSTRACT

The use of Translocator Protein 18 kDa (TSPO) as a clinical neuroimaging biomarker of brain injury and neuroinflammation has increased exponentially in the last decade. There has been a furious pace in the development of new radiotracers for TSPO positron emission tomography (PET) imaging and its use has now been extensively described in many neurological and mental disorders. This fast pace of research and the ever-increasing number of new laboratories entering the field often times lack an appreciation of the historical perspective of the field and introduce dogmatic, but unproven facts, related to the underlying neurobiology of the TSPO response to brain injury and neuroinflammation. Paradoxically, while in neurodegenerative disorders and in all types of CNS pathologies brain TSPO levels increase, a new observation in psychiatric disorders such as schizophrenia is decreased brain levels of TSPO measured by PET. The neurobiological bases for this new finding is currently not known, but rigorous experimental design using multiple experimental approaches and careful interpretation of results is critically important to provide the methodological and/or biological underpinnings to this new observation. This review provides a perspective of the early history of validating TSPO as a biomarker of brain injury and neuroinflammation and a critical analysis of controversial topics in the literature related to the cellular sources of the TSPO response. The latter is important in order to provide the correct interpretation of PET studies in neurodegenerative and psychiatric disorders. Furthermore, this review proposes some yet to be explored explanations to new findings in psychiatric disorders and new approaches to quantitatively assess the glial sources of the TSPO response in order to move the field forward.

© 2018 Elsevier Inc. All rights reserved.

Contents

1. The peripheral benzodiazepine receptor: early history of validation as a biomarker of brain injury and neuroinflammation	45
2. A critical analysis of the glial sources of the TSPO response	45
3. Longitudinal studies demonstrate different temporal patterns of TSPO expression in activated microglia and astrocytes	47
4. Exploration of new findings in psychiatric disorders	53
5. Concluding remarks	55
Conflict of interest statement	55
Acknowledgments	55
References	55

Abbreviations: Bmax, number of binding sites; CBR, central benzodiazepine receptor; CNS, central nervous system; CNTF, cytokine ciliary neurotrophic factor; EAE, experimental autoimmune encephalomyelitis; GFAP, glial fibrillary acidic protein; Kd, affinity constant; LPS, lipopolysaccharide; PBR, peripheral benzodiazepine receptor; PET, positron emission tomography; poly:IC, polyribinosinic-polyribocytidylic acid; SIV, simian immunodeficiency virus; SIVE, simian immunodeficiency virus encephalopathy; TBI, traumatic brain injury; TMT, trimethyltin; TSPO, translocator protein 18 kDa.

E-mail address: tguilart@fiu.edu.

1. The peripheral benzodiazepine receptor: early history of validation as a biomarker of brain injury and neuroinflammation

1.1. From PBR to TSPO and beyond

The peripheral benzodiazepine receptor (PBR) was first described as a binding site for diazepam (valium) in the quest to identify the endogenous receptor for benzodiazepines in the brain (Baestrup & Squires, 1977). Studies using tritium-labeled diazepam discovered that brain tissue expressed [³H]diazepam specific binding sites, that could be displaced by the benzodiazepine clonazepam (Baestrup & Squires, 1977). It was also discovered that kidney, liver, and lungs also expressed a specific binding site for [³H]diazepam that was not displaced by clonazepam but was sensitive to Ro5–4864, a benzodiazepine derivative of diazepam that lacks affinity for the GABA_A receptor (Baestrup & Squires, 1977). Furthermore, [³H]diazepam binding in kidney was associated with the mitochondrial fraction, while in the brain it was associated with the plasma membrane fraction (Baestrup & Squires, 1977). These findings lead to the identification of the peripheral benzodiazepine receptor binding site (also previously known as mitochondria benzodiazepine receptor, and omega-3 receptor), because it bound the benzodiazepine [³H]diazepam and it was highly expressed in peripheral tissue; thus, the original name peripheral benzodiazepine receptor. This was a fortuitous finding because diazepam is a drug that has high binding affinity for both the central benzodiazepine receptor (CBR) associated with the GABA_A receptor, and the PBR (Maragos et al., 1982). Subsequently, detailed studies showed that in fact the CBR and PBR were distinct proteins that differed in tissue distribution, pharmacology, and cellular and subcellular localization (Anholt et al., 1986; Benavides et al., 1983; Patel & Maragos, 1982) with CBR primarily expressed in neurons and PBR in glial cells (Junck et al., 1989; Schoemaker et al., 1982; Starosta-Rubenstein et al., 1987).

The use of the non-benzodiazepine isoquinoline carboxamide PK11195 (initial studies used the R,S-racemic form and later studies used the active R-enantiomer) and the GABA_A receptor non-active derivative of diazepam Ro5–4864 (4'-chlorodiazepam) as selective ligands for PBR with no affinity for the CBR facilitated the characterization of brain PBR binding sites. Early studies using different models of brain injury and neurodegeneration indicated that PBR levels measured by [³H]PK11195 and [³H]Ro5–4864 specific binding increased markedly at primary and secondary sites of brain injury (Benavides et al., 1988; Benavides et al., 1990; Benavides et al., 1987; Diorio et al., 1991; Doble et al., 1987; Dubois et al., 1988; Gehlert et al., 1985; Guilarte et al., 1995; Miyazawa et al., 1995) and these studies provided the first description of PBR's association with gliosis in animal models of brain injury (Benavides et al., 1990; Diorio et al., 1991; Dubois et al., 1988; Guilarte et al., 1995; Miyazawa et al., 1995). Despite the fact that the name PBR was used for many years, it was misleading since it was not an exclusively "peripheral" organ protein nor was it a receptor in the classical sense of neurotransmitter receptors. At the time, PBR was proposed to "transport" or "translocate" cholesterol from the outer to the inner mitochondria membrane for pregnenolone synthesis; the rate-limiting step in the production of neurosteroid (Krueger & Papadopoulos, 1990). It is ironic that while the name PBR was changed to translocator protein 18 kDa (TSPO) in 2006 based on its proposed function and molecular weight (Papadopoulos et al., 2006), recent studies using conditional and global TSPO knockout mice have questioned the cholesterol "translocator" function of TSPO (Morohaku et al., 2014; Tu et al., 2014; Tu et al., 2015).

1.2. Early validation of TSPO as a biomarker of gliosis

The ability to image and quantitatively measure TSPO levels in the living human brain using Positron Emission Tomography (PET) has significant advantages over other methodologies because it provides a direct dynamic view of glial responses to ongoing brain injury and

inflammation. TSPO can also be used *ex vivo* to assess regional brain injury with exquisite spatial resolution and sensitivity using quantitative receptor autoradiography and more recently immunofluorescence confocal imaging. Many studies during the last three decades have demonstrated that TSPO is a sensitive biomarker that is able to detect neuronal injury ranging from subtle neuronal terminal damage (Chen & Guilarte, 2008; Guilarte et al., 2003), to frank neuronal loss (Guilarte et al., 1995; Kuhlmann & Guilarte, 1997, 1999, 2000), demyelination (Chen et al., 2004; Chen & Guilarte, 2006), virus-induced inflammation (Cagnin et al., 2001; Coughlin et al., 2014; Hammoud et al., 2005; Mankowski et al., 2003), and neurodegeneration in animal models (Domene et al., 2016; Ji et al., 2008; Loth et al., 2016) and in humans with neurological disease (Banati et al., 2000; Coughlin et al., 2015; Coughlin et al., 2017; Turner et al., 2004). The cellular basis of TSPO as a biomarker of brain injury is due to the fact that the hallmark response of the brain to injury is the activation of microglia and astrocytes, the glial cell types that express and upregulate TSPO (Chen & Guilarte, 2008). In the normal brain neuropil, TSPO is expressed at very low levels, but TSPO levels increase rapidly as a result of brain injury anywhere in the brain and from any etiology (Chen & Guilarte, 2008; Venneti et al., 2006). TSPO levels increase in glial cells prior to behavioral manifestation of disease or histological detection of neurodegeneration, making it an early biomarker of brain injury and inflammation (Domene et al., 2016; Kuhlmann & Guilarte, 1997; Loth et al., 2016). TSPO levels not only increase at primary sites of brain injury and inflammation, but also at secondary sites (Cagnin et al., 2001; Kuhlmann & Guilarte, 1999; Turner et al., 2004; Venneti et al., 2006). These characteristics make TSPO an exceptional biomarker that can be used to assess recovery from injury (Chen & Guilarte, 2006) and monitor the effectiveness of therapeutic strategies (Doorduyn et al., 2008).

The early studies of TSPO as a biomarker of brain injury and inflammation was primarily driven by the use of radiolabeled ligand binding studies using receptor binding techniques ("grind and bind filtration assay") or quantitative receptor autoradiography in *ex vivo* brain tissue. As noted above, these studies provided early evidence that TSPO levels measured by radioligand binding were associated with gliosis in areas of the brain undergoing neurodegeneration. However, the cellular sources of the TSPO response in injured brain tissue was not clearly defined, although some of the early studies described TSPO as being associated with microglia activation and thus, neuroinflammation (Banati et al., 1997; Conway et al., 1998; Myers et al., 1991; Stephenson et al., 1995; Vowinckel et al., 1997). The paucity of information on the cellular sources of the TSPO response was largely due to the lack of molecular and cellular biology tools including the lack of well-characterized and validated TSPO antibodies. Furthermore, despite significant efforts from several laboratories including our own, to develop *in situ* hybridization techniques for mapping the spatial distribution of TSPO gene expression in healthy and diseased brain tissue, no such method was ever developed or validated.

2. A critical analysis of the glial sources of the TSPO response

2.1. TSPO in microglia and/or astrocytes?—a controversy that remains embedded in the literature today

Early studies from a number of laboratories using quantitative receptor autoradiography demonstrated that the levels of radioligand binding to TSPO was very low in the normal brain neuropil but increased dramatically as a result of different types of brain pathology (Banati et al., 1997; Benavides et al., 1983; Benavides et al., 1987; Benavides et al., 1988; Benavides et al., 1990; Chen et al., 2004; Chen & Guilarte, 2006; Conway et al., 1998; Dubois et al., 1988; Gehlert et al., 1985; Guilarte et al., 1995; Kuhlmann & Guilarte, 1997, 1999, 2000; Myers et al., 1991; Stephenson et al., 1995; Vowinckel et al., 1997). At this time, autoradiography (film and/or emulsion) and receptor binding studies were the only methods available to identify and

assess the cellular sources of the TSPO response. Film autoradiography has tissue level resolution while emulsion autoradiography has resolution at the cellular level. Using [^3H]PK11195 quantitative receptor autoradiography and emulsion autoradiography, several laboratories including our own examined the TSPO response to brain injury using a variety of different animal models of CNS pathologies.

The approach taken in our laboratory was to use well-characterized neurotoxicant-based animal models of neurodegeneration that would target different brain regions as primary sites of injury and would produce different degrees of injuries to examine the sensitivity of TSPO in detecting a signal from the injured brain. We used neurotoxicants that had not been used previously in the TSPO literature in order to complement other models being used at the time. Combined, these different models would provide face-validity to TSPO as a general biomarker of brain injury and inflammation based on the glial response (Chen & Guilarte, 2008). The first neurotoxicant used in our laboratory was trimethyltin (TMT) which has a well-characterized regional and temporal pattern of neuropathology in the limbic system (Guilarte et al., 1995). In the initial study, we found that increased levels of [^3H]PK11195 specific binding to TSPO faithfully tracked the regional and temporal pattern of TMT-induced neuropathology. Furthermore, the increased temporal pattern of [^3H]PK11195 binding in the hippocampus followed the levels of glial cell activation based on immunostaining and astrocytes based on glial fibrillary acidic protein (GFAP) ELISA (Guilarte et al., 1995). However, these studies were not as methodologically advanced as to what is possible and available today [see Section 3.4 below].

At approximately the same time, other laboratories using transient global forebrain ischemia, facial nerve axotomy, and experimental autoimmune encephalomyelitis, attributed the TSPO response to brain injury exclusively to microglia and/or macrophage infiltration at the sites of injury with no apparent association to astrocytes (Banati et al., 1997; Conway et al., 1998; Myers et al., 1991; Stephenson et al., 1995; Vowinckel et al., 1997). However, careful examination of these early (and later) studies that attributed the cellular sources of the TSPO response exclusively to microglia reveals several limitations including: 1) visual inspection of the spatial pattern of TSPO autoradiography and immunohistochemistry of glial markers was only done at one time point in the progression of the injured brain (Mirzaei et al., 2016; Stephenson et al., 1995); 2) astrocyte staining was noted as being done but images were not provided (Banati et al., 1997); 3) immunohistochemistry for astrocytes was not performed at all (Arlicot et al., 2008; Harhausen et al., 2013; Kannan et al., 2007; Miller et al., 2013; Toyama et al., 2008; Vowinckel et al., 1997); 4) microglia was noted in the title as being the source of the TSPO signal, but microglia staining was not performed (Yankam Njiwa et al., 2016); or 5) despite the fact that the spatial pattern of gene expression of the astrocytic marker GFAP was similar to the spatial distribution of [^3H]PK11195 binding (autoradiography) at later time points in the progression of the injured brain in a global forebrain ischemia model, the authors conclude that because the early temporal expression of the GFAP mRNA and [^3H]PK11195 binding were different, it supported the proposed localization of TSPO in activated microglia (Conway et al., 1998). This conclusion was provided despite the fact that the spatial distribution of the GFAP mRNA and [^3H]PK11195 binding closely overlapped (by visual inspection) at a later time point (8 days after the ischemia event-see Fig. 2 panels F and I in Conway et al., 1998) and microglia immunohistochemistry was not performed.

These initial studies were taken as experimental evidence of an exclusive localization of TSPO in microglia with no association with astrocytes. They formed the foundation for the current disregard of the significance of the astrocyte contribution to elevated TSPO levels in many neurological and neurodegenerative conditions. Another significant methodological pitfall of the early studies was that associations of the spatial pattern of [^3H]PK11195 binding using film or emulsion autoradiography as compared to the pattern of immunostaining of microglia

and astrocyte markers were performed by visual inspection and could be influenced by experimenter interpretation. To this point, it is relevant to note that when publications using the same approach and animal model of brain injury were compared, significantly different interpretations were provided. For example, Banati et al. (1997) examining the TSPO response in the facial nucleus at 1, 4, and 21 days after facial nerve axotomy indicated that [^3H]PK11195 emulsion autoradiography was restricted to microglia (based on immunostaining; no astrocyte staining was provided). However, another group using the same facial nerve axotomy model noted that [^3H]PK11195 binding was associated with the staining of both microglia and astrocytes during a similar time period following axotomy, i.e., 5–10 days (Gehlert et al., 1997). They indicate that following facial nerve transection, elevated levels of TSPO measured by increased [^3H]PK11195 binding colocalized based on visual inspection with the immunostaining of both microglia and astrocytes (Gehlert et al., 1997).

2.2. TSPO radioligand binding and correlation with glial cell number and immunofluorescent staining of glial markers

Another approach used in the early studies to assess the cellular sources of the TSPO response was to correlate the level of [^3H]PK11195 binding to TSPO with the number of cells immunostained with a microglia/macrophage and/or an astrocyte marker. In an animal model of traumatic brain injury (TBI), increased [^3H]PK11195 binding was correlated with the number of ED-1-positive cells for microglia/macrophages or GFAP-positive astrocytes (Rao et al., 2000). The number of ED-1 or GFAP positive cells were both shown to significantly correlate with [^3H]PK11195 binding but it was noted that there was a “better” correlation with ED-1 than with GFAP. The authors seem to disregard the astrocytic TSPO response with the statement that “microglia/macrophages are the major cell type associated with the increased TSPO expression following TBI” while considering that “the possibility of some astroglial contribution to the increased TSPO expression following TBI cannot be ruled out” (Rao et al., 2000). Regardless of their interpretation of the results, one experimental limitation of this study is that cell counting did not use unbiased stereological cell counting methods nor was a description of the morphological or cell size criteria used to count cell number was provided. The latter is important with ED-1 positive microglia/macrophages since in the injured brain there is clustering of microglia/macrophages (microglial nodule) and loss of ramifications with microglia activation (or with entry of peripheral macrophages) making it difficult to determine the actual number of cells in a cluster.

Another study of a macaque model of neuroAIDS, [^{11}C]PK11195 PET imaging was performed in 11 simian immunodeficiency (SIV) infected macaques and 2 non-infected controls (Venneti et al., 2004). The animals used had a wide range of age and the length of infection varied significantly from 56 to 1622 days. This PET study showed that [^{11}C]PK11195 brain uptake was increased in 6 animals that developed SIV encephalopathy (SIVE) compared to those that did not. Postmortem brain analysis of the 6 SIVE animals using [^3H]PK11195 autoradiography confirmed the increased TSPO levels in SIVE relative to SIV animals and controls. Immunofluorescence for a macrophage marker (CD68), astrocytes (GFAP), or neurons (MAP2) was performed in the same regions. The authors then correlated the levels of [^{11}C]PK11195 specific binding *in vivo* with the levels of astrocytes, macrophages, and neurons based on the immunofluorescent signal. It was noted that TSPO levels using [^3H]PK11195 binding in postmortem tissue or *in vivo* [^{11}C]PK11195 was correlated with the CD68 signal but not with the GFAP or MAP2 signal using confocal laser microscopy. In this study, the authors used an average pixel fluorescence for a particular cellular phenotype using confocal imaging and the latter determined the cellular phenotype that was associated with the *ex vivo* [^3H]PK11195 binding [Venneti et al., 2004]. The study concludes that the increased TSPO in the brain of SIVE animals correlated with the abundance of microglia/macrophages and not astrocytes. However, these two different endpoints, that is receptor binding

(postmortem) or TSPO radioligand uptake (*in vivo*) and fluorescent imaging cannot be associated unless it is demonstrated that both signals are originating in the same cell which is not possible based on the approach used.

In a later study, the same authors examined the glial sources of the TSPO response in post-mortem tissue from a variety of human neurological conditions such as cerebral infarct, multiple sclerosis, frontotemporal dementia, amyotrophic lateral sclerosis, and Alzheimer's disease [Venneti et al., 2008]. Despite the fact that analysis of CD68 (microglia/macrophage) and GFAP (astrocytes) were elevated in all neurological conditions, the authors conclude that the TSPO signal arises from microglia based on correlation of autoradiography area with immunofluorescence area. Further, in a rat model of TBI, a similar conclusion was provided despite significant astrocyte activation [Venneti et al., 2007]. Lastly, in a study using postmortem brain tissue from Alzheimer's disease subjects, in the abstract and title the authors note that PK11195 binding correlates with the extent of microglia activation despite the fact that an association was also found between [³H]PK11195 binding and GFAP immunostaining (Venneti et al., 2009). The latter study is indicative of an investigator-based selective bias to an association of the TSPO signal with microglia and disregard the association with astrocytes. A common error in the TSPO literature.

While these early studies were important contributions to the TSPO literature and used the best available methods at the time, they had significant methodological limitations that could lead to biased interpretations based on the fact that: 1) the cellular sources of the TSPO response were determined by examining the spatial pattern of TSPO radioligand binding using film or emulsion autoradiography with the pattern of immunostaining of microglia/macrophage or astrocyte markers which required visual inspection and potential experimenter bias, or 2) were based on correlation of the number of microglia/macrophage- or astrocyte-labeled cells or immunofluorescent signal intensity with [³H]PK11195 binding. However, the counting of labeled cells did not use state-of-the-art unbiased stereological cell counting methods [see Kanaan et al., 2010; Lemus et al., 2015] and were often based on gross analysis of immunofluorescence signal intensity. Therefore, there was a great need to use more rigorous experimental approaches in order to begin to provide a more quantitative, accurate, and direct assessment of the glial sources of the TSPO signal in the injured brain.

2.3. TSPO immunohistochemistry-A new era

As noted previously, one of the limiting factors of the early studies examining the cellular sources of the TSPO response was the lack of commercially available, validated TSPO-specific antibodies that could be used for experiments in which double label of TSPO with specific glial markers could be performed. In 2000, using the same TMT animal model that we first described in 1995 (single i.p. injection of 8 mg/kg TMT in rats), we performed studies on the cellular sources of the TSPO response (Kuhlmann & Guilarte, 2000). The temporal TSPO response to TMT-induced neurodegeneration was performed at 2 days, 14 days, and 6 weeks following a single 8.0 mg/kg TMT injection. Besides the typical methods used at this time (film and emulsion autoradiography), we also performed the first TSPO immunohistochemistry with a validated TSPO antibody provided by Dr. V. Papadopoulos. The results showed that the levels of [³H]PK11195 binding in various limbic brain regions, including the hippocampus, increased significantly as a function of time following TMT injection. Furthermore, [³H]PK11195 binding to TSPO was sustained in brain regions undergoing neurodegeneration. We also showed that the cellular sources of the TSPO response was driven by an early microglia response that resolved by 6 weeks after injury while the astrocytic response appeared at 14 days and remained present at 6 weeks following TMT administration. TSPO immunohistochemistry provided the first evidence that TSPO-labeled cells in the TMT-treated hippocampus, with active neurodegeneration based on silver staining, expressed morphology consistent with both activated

microglia and astrocytes. Importantly, the presence of TSPO in both glial cell types was confirmed with the first-ever double-label immunostaining of TSPO with either *Griffonia simplicifolia* isolectin B₄ for microglia or GFAP for astrocytes (Kuhlmann & Guilarte, 2000). This study provided the first direct evidence based on double-label immunohistochemistry of a distinct TSPO colocalization with both microglia and astrocytes. However, it did not perform analysis of TSPO signal cellular colocalization with glial markers as a function of brain region or time after brain injury [see Section 3.4 below]. Nevertheless, it was the first study to use a TSPO-specific antibody and showed that TSPO colocalization occurred with both microglia and astrocytes.

3. Longitudinal studies demonstrate different temporal patterns of TSPO expression in activated microglia and astrocytes

Glial cells have a dynamic response to different brain pathologies in their morphology and function. Therefore, performing studies at a single point in time does not provide an accurate representation of the dynamic nature of the glial response in the time-continuum of brain disease or injury (McNeela et al., 2017). To address this important aspect of the cellular sources of the TSPO response, we used another model of neurotoxicant-induced brain injury using cuprizone. Cuprizone is a copper chelator that damages oligodendrocytes and produces widespread demyelination of the brain (Praet et al., 2014). This neurotoxicant-induced animal model of demyelination was selected because cuprizone-induced demyelination exhibits a recovery phase with extensive remyelination once the animal is removed from the cuprizone exposure (Praet et al., 2014). Using this model, we assessed the glia cell TSPO response in a demyelination-remyelination continuum (Chen et al., 2004; Chen & Guilarte, 2006). This was an excellent animal model to examine two important questions in the early validation of TSPO as a biomarker of brain injury and inflammation: 1) Is TSPO able to track not only brain injury but also resolution of injury? and 2) what are the dynamics of the glial cell TSPO response during the demyelination-remyelination continuum?

Demyelination phase: The results of the demyelination study showed a dose-response relationship between cuprizone dose and [³H]R-PK11195 specific binding to TSPO based on autoradiography analysis of brain regions undergoing demyelination. Demyelinating brain regions were confirmed by myelin staining using Black Gold histochemistry (Chen et al., 2004; Chen & Guilarte, 2006). Examination of the temporal response of [³H]R-PK11195 binding, microglia staining with Mac-1 and astrocyte staining with GFAP, found that increased [³H]R-PK11195 binding at 2 weeks of cuprizone exposure was primarily driven by microglia since there was limited or no staining of astrocytes at this time point in brain regions exhibiting increased [³H]R-PK11195 binding to TSPO. However, at 3 and 4 weeks of cuprizone exposure when [³H]R-PK11195 increased dramatically compared to non-exposed control animals, there was markedly increased staining of both microglia and astrocytes supporting a role of both glial cell types in the TSPO response but with different temporal profiles. That is, an early microglia response followed by a later activation of astrocytes. [³H]R-PK11195 emulsion microautoradiography with glial marker immunostaining confirmed the colocalization of the TSPO signal with both microglia and astrocytes. These studies showed the importance of examining the course of brain injury using a longitudinal design because if we had only performed the study at 2 weeks after cuprizone exposure our conclusion would have been that the TSPO response is exclusively from microglia. However, if would fail to demonstrate that in this animal model of brain injury, astrocytes are also activated at later time points in the same brain regions undergoing demyelination. Furthermore, [³H]R-PK11195 emulsion autoradiography demonstrated TSPO cellular colocalization with both microglia and astrocytes at 3 and 4 weeks of exposure (Chen et al., 2004).

Remyelination phase: We also examined the remyelination phase of the cuprizone model using a similar approach of [³H]R-PK11195 quantitative autoradiography and immunostaining for glial markers and Black Gold histochemistry for myelin in the corpus callosum (Chen & Guilarte, 2006). The corpus callosum is a major white matter fiber track connecting the cerebral hemispheres. In this study, we also performed [¹¹C]R-PK11195 microPET in similarly treated animals. We found that as previously shown (Chen et al., 2004), 3 weeks of cuprizone exposure of the animals resulted in approximately a 5-fold increase in [³H]R-PK11195 binding to TSPO in the corpus callosum relative to non-treated controls. The increase in [³H]R-PK11195 binding measured at 3 weeks of cuprizone-induced demyelination decreased significantly at 3 and 6 weeks after removing animals from the cuprizone exposure and it was inversely associated with increased levels of myelin staining imaged by Black Gold histochemistry (Chen & Guilarte, 2006). By 6 weeks of remyelination there was approximately a 50% decrease in the maximal [³H]R-PK11195 specific binding measured at 3 weeks of cuprizone-induced demyelination. This decrease in [³H]R-PK11195 binding to TSPO during the remyelination phase was associated with increased myelin staining in the corpus callosum and with little or no labeled Mac-1 positive microglia by 6 weeks of remyelination. On the other hand, GFAP-labeled astrocytes also decreased but there were more labeled astrocytes in the same brain region remaining at 6 weeks of remyelination (Chen & Guilarte, 2006). Considering that [³H]R-PK11195 was still significantly elevated to approximately 50% of the maximal demyelination response at the 6 weeks remyelination time point, strongly suggests that astrocytes provide a significant, if not the majority of the TSPO signal at 6 weeks of remyelination. Finally, [¹¹C]R-PK11195 microPET confirmed a significantly increased TSPO signal during the demyelination phase with the TSPO signal achieving complete recovery to control levels by 10 weeks following removal from the cuprizone exposure (Chen & Guilarte, 2006).

Collectively, these two longitudinal studies demonstrated that the TSPO response can track not only disease onset and progression but also resolution of injury. Furthermore, it demonstrates a critical point on the importance of examining the TSPO response in a longitudinal fashion in the context of the glial sources. That is, if one examines the glial sources of the TSPO response at an early time point (2 weeks) after initiation of cuprizone-induced demyelination, one would conclude that the TSPO response is exclusively from microglia since no astrocyte staining was present at this time. Similarly, if one only examined the 6-week time point after removing the animals from the cuprizone exposure during the remyelination phase, one would conclude that the TSPO signal is exclusively from astrocytes since at 6 weeks there was no or minimal microglia staining. In both cases, the conclusions when utilizing a single time point would have been correct for those specific time points but would have been completely inaccurate when taken into consideration the progression of the disease as there is contribution of both microglia and astrocytes during the different phases of the demyelination-remyelination continuum.

3.1. Studies support an astrocyte contribution to the TSPO response in diverse pathologies

Following our TMT and cuprizone studies (Chen et al., 2004; Chen & Guilarte, 2006; Kuhlmann & Guilarte, 2000) and due to the biological importance of understanding the glial sources of the TSPO response to brain injury for proper interpretation of PET studies and for potential therapeutic approaches, other laboratories interrogated this question using TSPO immunohistochemistry. They confirmed that the TSPO signal is from both microglia and astrocytes using a variety of animal models of CNS pathologies and in postmortem brain tissue in human neurodegenerative disorders (Arlicot et al., 2014; Biesmans et al., 2015; Cosenza-Nashhat et al., 2009; Dickens et al., 2014; Domene et al., 2016; Israel et al., 2016; Janssen et al., 2017; Ji et al., 2008; Lavis-

se et al., 2012; Lavisse et al., 2015; Liu et al., 2015; Maeda et al., 2007; Maeda et al., 2011; Mattner et al., 2011; Nguyen et al., 2018; Rojas et al., 2007; Sérière et al., 2015; Wang et al., 2014). These series of studies showed that the dynamic nature of the TSPO glial cell response to brain injury represents a continuum of change based on the type of injury and whether the injury is acute or chronic. From this collection of papers, there are two studies that are notable for discussion. First, Ji et al. (2008) examined the cellular sources of the increased TSPO response in two mouse models of Alzheimer's disease (APP23 and PS19 tau transgenic mice). Using double-label immunostaining, they found that in APP23 (mutant APP) mice, there was predominant localization of the TSPO signal to astrocytes in the vicinity of β -amyloid plaques with no detectable TSPO signal in Iba-1 positive microglia. Strikingly, and in contrast to the TSPO astrocyte response in APP23 mice, the reverse was found in the PS19 tau transgenic mice. That is, the build-up of phospho-tau in the PS19 mouse brain was associated with a TSPO signal in microglia and not in astrocytes. This study clearly indicated different glial sources of the TSPO response based on the animal model used. Although they did not do a time-course profile in these two animal models of Alzheimer's disease that would have potentially discovered additional glial cell TSPO colocalization at different time points as in the cuprizone studies presented above.

In the same publication, the authors also examined a variety of neurotoxicant-induced animal models to examine the cellular sources of the TSPO response using double-label immunostaining. They found that similar to the APP23 and PS19 mouse models, the cellular sources of the TSPO response was specific to the neurotoxicant used. In particular, when they examined the TSPO glial cell response in the striatum and corpus callosum of cuprizone-exposed animals to induce demyelination, they found that the vast majority of astrocytes in the striatum exhibited high levels of TSPO staining. They then performed *in vivo* microPET with [¹⁸F]FE-DAA1106 and found increased uptake in the striatum of cuprizone-treated mice relative to non-treated mice. This finding was another demonstration that *in vivo* TSPO imaging is able to measure increased TSPO levels originating from astrocytes. In summary, this study provided evidence that TSPO increases in both microglia and astrocytes and may reflect deleterious and beneficial consequences of glial responses, respectively, in the injured brain.

The second notable study is related to the development of an animal model of selective astrocyte activation through lentiviral gene transfer of the cytokine ciliary neurotrophic factor (CNTF) into the striatum in the absence of neurodegeneration and microglia activation (Lavisse et al., 2012). This approach resulted in extensive astrocyte activation visualized by GFAP staining with minimal or no increase in microglia activation markers. The authors showed that using two different TSPO radioligands *i.e.*, [¹⁸F]DPA-713 and [¹¹C]SSR180575 there was significant uptake of these two TSPO radioligands in the CNTF-injected striatum relative to the contralateral control striatum. Furthermore, the radioligand uptake was displaced by non-radioactive PK11195, demonstrating the pharmacologically selective binding of the radiotracer to TSPO. Examination of brain tissue indicated significantly increased TSPO mRNA and protein levels that colocalized with CNTF-activated astrocytes. This study showed a selective increase of TSPO levels in reactive astrocytes that yields a selective and increased radioligand TSPO signal using *in vivo* microPET imaging in the absence of microglia activation. In summary, these studies provided strong and confirmatory evidence that the sources of the PET radioligand uptake is also derived from astrocytes. It is notable that the dogma that the cellular source of the TSPO response in various CNS pathologies is exclusively from microglia is so erroneously entrenched in the literature that despite the fact that some studies have clearly shown and articulated that astrocytes significantly contribute to the TSPO signal, the astrocyte contribution to the TSPO signal was not reflected in the title of the publication (Brackhan et al., 2016; Dickens et al., 2014; Israel et al., 2016).

3.2. TSPO-PET literature indicating TSPO as an exclusive microglia biomarker

Despite the fact that there are now many studies demonstrating an important contribution of astrocytes to the TSPO signal in brain injury and inflammation using both *ex vivo* and *in vivo* imaging methods, the TSPO-PET literature continues to have a significant number of studies taking the reductionist view that TSPO is a biomarker of neuroinflammation whose signal is originating from microglia (Table 1). This view is prevalent despite the fact that the TSPO response is defined by a complex cross-talk between microglia and astrocytes following brain injury (Masgrau et al., 2017; Schain & Kreisl, 2017; Streit et al., 2004), and neuroinflammation is not a selective microglia event, but it also involves other cells including astrocytes (Masgrau et al., 2017; Schain & Kreisl, 2017; Streit et al., 2004). While it is scientifically expedient to believe that TSPO is a biomarker of brain injury and neuroinflammation whose signal is exclusively attributed to microglia and/or infiltrating peripheral macrophages, it is an inaccurate statement unless there is direct demonstration that a selective “microglia” TSPO signal occurs during the course of a specific brain disorder or pathology. Often, studies note that the TSPO response is due to microglia but they either do not examine TSPO expression in astrocytes or the immunostaining is only done at one time point or in one brain region and are highly likely missing the presence of an astrocyte-mediated increase in the TSPO signal at a different time point or brain region in the progression of the neuropathology or disease (Table 1). Consistent with this point, a study in which TSPO immunohistochemistry was performed with microglia/macrophage and astrocyte markers in the spinal cord of experimental autoimmune encephalomyelitis (EAE) rats shows an apparent selective increase of TSPO only in microglia and not in astrocytes (Abourbeh et al., 2012). However, despite the fact that there was a longitudinal assessment of neurological score that ranged from 0 to 5 from approximately 8–15 days post-immunization, only animals that had the highest (peak) neurological score, *i.e.*, 4–5 were used for TSPO and glial markers immunohistochemistry. Therefore, the authors missed the opportunity to examine the temporal profile of the TSPO glial cell response during the induction phase and resolution of the EAE disease (Abourbeh et al., 2012). Based on a single time point in the continuum of the EAE, the authors conclude that the increased TSPO levels in EAE is from microglia/macrophages rather than astrocytes. While it is entirely possible that at the peak of the behavioral manifestation of the EAE the TSPO response originates from microglia, one cannot assume that the other phases of the disease have a similar TSPO glial cell response profile since signal colocalization and quantification were not performed. It is highly likely that the TSPO contribution from astrocytes was missed by the selection of a single time point in the disease continuum as it has been discussed with the cuprizone model above.

3.3. Endotoxin-induced neuroinflammation: TSPO in microglia and astrocytes?

Other studies have examined the TSPO glial cell response in endotoxin-induced neuroinflammation. Hannestad et al. (2012) examined the TSPO response in baboons injected with lipopolysaccharide (LPS; single injection) and TSPO brain levels imaged before and at 1 h, 4 h, or 22 h after *i.v.* injection of LPS using [¹¹C]PBR28-PET. Whole brain analysis of PET scans revealed an increase in radioligand uptake in the whole brain that was increased only at the 4 h time point and not at the 1 or 22 h time points after LPS. However, when they used the animal's own baseline PET scan, they found increased TSPO levels at 1 and 4 h suggestive of a transient increase since at the 22 h time point PET analysis showed a marked decrease or no TSPO increase relative to the individual animal's baseline PET (Hannestad et al., 2012). Out of the 6 animals available, the authors euthanized 1 animal at 6 h after LPS administration to perform immunohistochemistry for TSPO and CD68 for microglia/macrophages and GFAP staining for astrocytes. The

immunostaining was only performed in the frontal lobe despite widespread increased brain TSPO uptake in the PET scan. From these limited immunostaining experiments the authors concluded that “cells that were immunoreactive (IR) for the antibody for TSPO are almost exclusively small, rod-like cells with short or no visible processes”. “These cells were very similar in morphology, size, and distribution to CD68 IR cells”. They further state: “Conversely, GFAP IR cells (astrocytes) were larger, much more abundant, and had more numerous and longer processes”. From this limited immunostaining in the absence of double-label studies they conclude that “Thus, TSPO immunoreactivity occurred almost exclusively in microglia, but not in astrocytes...”. This conclusion was representative of a single small area in the frontal lobe, from 1 animal, at a single time point after LPS (6 h) in which double-label of TSPO with each individual glial marker was not performed. But yet, the conclusion based on visual inspection of presumed cellular morphology provides the evidence that the TSPO response from LPS administration is exclusively from microglia. Further, it is known that TSPO does not label the entire microglia or astrocyte but certain subcellular compartments; thus, the distinction cannot be made on presumed cellular size. Moreover, LPS not only activates microglia *via* Toll-Like Receptor 4, but it also activates the same receptors in astrocytes (Brahmachari et al., 2006; Gorina et al., 2011). Finally, it is interesting to note that in the discussion of the study, the authors contradict their main conclusion that “TSPO immunoreactivity is occurring exclusively in microglia” by the following statement: “For the postmortem data, because double immunostaining was not performed it is possible that TSPO and CD68 expression occurred in different cell types, or that TSPO expression occurred in astrocytes in addition to microglia”.

This oversimplification of the cellular sources of the TSPO response was translated to a subsequent human study using LPS administration and [¹¹C]PBR28-PET (Sandiego et al., 2015). Eight healthy control volunteers had two [¹¹C]PBR28-PET scans on the same day. They had a baseline scan and then a second scan in which LPS (10 ng/kg, *i.v.*) was administered 180 min before the second scan. They found a 30–60% increase in [¹¹C]-PBR28 binding in multiple brain regions. Indeed, examination of the PET scan images after LPS indicated increased TSPO radioligand uptake throughout the entire brain. In the discussion, the authors conclude: “In the current study, the systemic LPS markedly increased the [¹¹C]PBR28 binding in humans. [¹¹C]PBR28 is a radioligand sensitive to TSPO levels expressed on activated microglia and is a putative biomarker of neuroinflammation. TSPO is also expressed in astrocytes; however, immunohistochemistry performed in nonhuman primate brain after LPS administration confirmed that TSPO expression occurred mainly in activated microglia [32]”. This conclusion referenced their previous baboon study (Hannestad et al., 2012) in which 1 single animal, at a single time point after LPS (6 h), one brain area that was assessed by visual inspection of TSPO labeled cells with glial markers and in which double-label of TSPO with each individual glial marker and colocalization signal analysis was not performed. Further, as noted above, they acknowledge in the discussion of the baboon study, the uncertainty of the statement that the TSPO response is exclusively from microglia (Hannestad et al., 2012). Despite these limitations and uncertainties in the baboon study, they indicate these two statements in the discussion of the human study: “In the current study, systemic LPS markedly increased [¹¹C]PBR28 binding in humans. [¹¹C]PBR28 is a radioligand sensitive to TSPO levels expressed on activated microglia and is a putative biomarker of neuroinflammation. TSPO is also expressed in astrocytes; however, immunohistochemistry performed in a nonhuman primate brain after LPS administration occurred mainly on microglia [32]”. They conclude: “thus, we have successfully extended the main findings of the nonhuman primate study to human subjects and established that LPS administration in humans is associated with a marked and significant increase in activated microglia”. While it is entirely possible that the early TSPO signal following LPS may be from microglia, the LPS response was examined at a single time point and may have missed the astrocytic TSPO response at other time points. In fact, a study from another group in which LPS was directly injected

Table 1
Representative animal and human studies in which the TSPO response is attributed only to microglia.

Human disease or animal model	Experimental methods	Experimental shortcoming	Reference
Autoimmune Encephalomyelitis (rats)	PET imaging; western blot and immunohistochemistry	Single time point; no signal colocalization analysis performed.	Abourbeh et al. (2012)
Excitotoxicity using quinolinic acid (rats)	Microglia in title; SPECT imaging; autoradiography and immunohistochemistry	No GFAP immunohistochemistry performed.	Arlicot et al. (2008)
Facial nerve axotomy (rats)	Emulsion microautoradiography; microglia immunohistochemistry	GFAP immunohistochemistry noted but not shown.	Banati et al. (1997)
Rasmussen's encephalitis (human)	PET imaging; immunohistochemistry	Assumed astrocytes do not contribute to TSPO signal. GFAP immunostaining not performed. Emulsion autoradiography with GFAP staining not performed.	Banati et al. (1999)
Multiple Sclerosis (human, rats)	PET imaging; autoradiography and immunohistochemistry	GFAP immunohistochemistry noted but not shown.	Banati et al. (2000)
Status Epilepticus (rats)	PET imaging; autoradiography and immunohistochemistry	High degree of correlation of TSPO PET signal with both microglia and astrocytes. No double label of TSPO with glial markers. Only microglia correlation is noted in manuscript title.	Brackhan et al. (2016)
Alzheimer's disease dementia (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Cagnin et al. (2001)
Multiple Sclerosis (human)	PET imaging; no assessment of microglia	No postmortem tissue immunohistochemistry performed.	Debruyne et al. (2003)
Lipopolysaccharide (LPS) (rats)	PET imaging; autoradiography and immunohistochemistry	Increased microglia and astrocyte staining; no double label with TSPO performed; only microglia is noted in manuscript title.	Dickens et al. (2014)
Multiple system atrophy (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Gerhard et al. (2003)
Corticobasal degeneration (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Gerhard et al. (2004)
Ischemic stroke (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Gerhard et al. (2005)
Idiopathic Parkinson's disease (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Gerhard et al. (2006)
Progressive Supranuclear Palsy (humans)	PET imaging	No postmortem tissue immunohistochemistry performed.	Gerhard et al. (2006)
Parkinson's Disease (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Ghadery et al. (2017)
Alzheimer's Disease (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Groom et al. (1995)
Lipopolysaccharide (LPS) (non-human primates)	PET imaging; immunohistochemistry	Individual TSPO, microglia, and astrocyte imaging. No double label immunostaining performed.	Hannestad et al. (2012)
Corticobasal degeneration (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Henkel et al. (2004)
Microglia depletion and activation (non-human primates)	PET imaging	No postmortem tissue immunohistochemistry performed.	Hillmer et al. (2017)
Alcohol Dependence (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Hillmer et al. (2017)
Creutzfeldt-Jakob Disease (human)	PET imaging	No TSPO, microglia, or astrocyte postmortem tissue immunohistochemistry performed.	Iaccarino et al. (2017)
Fatal Familial Insomnia (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Iaccarino et al. (2018)
Head Injury (mice)	microPET imaging; autoradiography; immunohistochemistry	High degree of correlation of TSPO PET signal with both microglia and astrocytes. No double label of TSPO with glial markers. Only microglia correlation is noted in manuscript title.	Israel et al. (2016)
Lipopolysaccharide (LPS) (rabbit)	microPET imaging; immunohistochemistry	No astrocyte staining; no double label immunohistochemistry.	Kannan et al. (2007)
Amnesic mild cognitive impairment (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Knezevic et al. (2017)
Neuropathic Osteoarthritic Pain (rats)	Receptor binding; autoradiography; immunohistochemistry	No astrocyte staining performed; no TSPO double-label immunohistochemistry performed.	Miller et al. (2013)
Alzheimer's Disease (5XFAD mice); postmortem human tissue from sporadic Alzheimer's disease	microPET imaging; autoradiography; immunohistochemistry	One age of animals used. No signal colocalization and analysis performed.	Mirzaei et al. (2016)
Ischemic stroke (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Pappata et al. (2000)
Huntington's Disease (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Politis et al. (2015)
Lipopolysaccharide (LPS) (human)	PET imaging	No postmortem tissue immunohistochemistry performed. Referenced nonhuman primate study of a single animal (Hannestad et al., 2012).	Sandiego et al. (2015)
Global forebrain ischemia (rats)	Autoradiography; immunohistochemistry	One time point; no double label of TSPO with glial markers; visual inspection of TSPO binding with immunohistochemistry.	Stephenson et al. (1995)
Huntington's Disease (human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Tai et al. (2007)
Alzheimer's disease (human)	PET imaging	Presumed TSPO staining in arteries and veins in brain. No microglia or	Tomasi et

Table 1 (continued)

Human disease or animal model	Experimental methods	Experimental shortcoming	Reference
Alcohol (rats)	microPET imaging; histological staining of microglia	astrocyte immunohistochemistry. No staining of astrocytes.	al. (2008) Toyama et al., 2008
Amyotrophic Lateral Sclerosis (ALS; human)	PET imaging	No postmortem tissue immunohistochemistry performed.	Turner et al. (2004)
AIDS (non-human primates)	PET imaging; autoradiography and immunohistochemistry	No double label or quantitative analysis signal colocalization.	Venneti et al. (2004)
Traumatic Brain Injury (TBI) (rats)	Autoradiography (<i>in vitro</i> and <i>ex-vivo</i>) and immunohistochemistry	No quantitative analysis of signal colocalization.	Venneti et al. (2007)
Cerebral Infarcts, Amyotrophic Lateral Sclerosis, Alzheimer Disease, Frontotemporal Dementia, and Multiple Sclerosis (humans)	Autoradiography and immunohistochemistry	No quantitative analysis signal colocalization.	Venneti et al. (2008)
Alzheimer's disease postmortem tissue (human) APP/PS1 transgenic mice	MicroPET studies; autoradiography; immunohistochemistry	No quantitative analysis of signal colocalization.	Venneti et al. (2009)
Multiple Sclerosis (MS) (humans) and Experimental Autoimmune Encephalomyelitis (EAE) (mice)	PET imaging; autoradiography and immunohistochemistry	No astrocyte immunohistochemistry; no double label immunohistochemistry	Vowinckel et al. (1997)
Epilepsy (rats)	microPET imaging;	No postmortem tissue immunohistochemistry performed.	Yankam Njiwa et al. (2016)

into the rat brain and in which a time-course study was performed using TSPO double label immunofluorescent imaging with microglia and astrocyte markers confirms that TSPO colocalized with microglia at the early time points with no TSPO astrocyte labeling but TSPO colocalized to astrocytes at a later time point when microglia do not express TSPO (Ory et al., 2015). The study by Ory et al. (2015) support a significant body of evidence that the TSPO response in microglia and astrocytes have different temporal profiles. A limitation of the Ory et al., study was the lack of quantification of TSPO signal colocalization with the glial markers (see below).

3.4. TSPO signal colocalization with microglia and astrocytic markers and computer-based quantitative image analysis—a way forward

The last two decades of studies using TSPO as a biomarker of brain injury have witnessed a gradual improvement of methodological approaches used for assessing the cellular sources of the TSPO response. The current availability and use of state-of-the-art computer-based image analysis technology has facilitated the quantitative analysis of the colocalization of immunofluorescent signals from different proteins in the same cell with exquisite sensitivity and spatial resolution. An example of this new approach is depicted in a recent TSPO study using a mouse model of Sandhoff disease (Loth et al., 2016). Sandhoff disease is an autosomal recessive disorder characterized by a deficiency of the lysosomal enzyme β -hexosaminidase leading to the accumulation of gangliosides and glycolipids in neurons, specifically GM2 and GA2 (Jeyakumar et al., 2002; Mahuran, 1999; Sango et al., 1995). The accumulation and aggregation of gangliosides is the triggering event that leads to progressive and widespread neurodegeneration. The TSPO response was examined in this animal model as a function of age (neurodegeneration increases with age) using the second generation radioligand [^3H]DPA-713 (autoradiography) and [^{125}I]iodo-DPA-713 (microSPECT) in parallel with behavioral and pathological expression of the disease. It was found that the increased TSPO response as measured by [^3H]DPA-713 autoradiography occurred much earlier in time in the thalamus (1.5 months) than the behavioral manifestation of disease (3 months). The thalamus is the brain region that exhibits the first neurodegenerative changes in this disease. Furthermore, increased [^3H]DPA-713 specific binding to TSPO occurred simultaneously with the aggregation of GM2 gangliosides in neurons and before evidence of active neurodegeneration using Silver staining (Loth et al., 2016). The increase in brain TSPO in the living anesthetized animal was measured using [^{125}I]-

iodo-DPA-713 microPET and brain uptake was blocked by the administration of non-radioactive PK11195. Double-label immunofluorescence imaging of TSPO (using an extensively validated TSPO antibody) with the microglia marker Mac-1 and the astrocyte marker GFAP was also performed in an age-dependent fashion (See Fig. 1). Notably, this was the first study to provide a quantitative analysis of percent colocalization of the TSPO immunofluorescence signal with the Mac-1 or GFAP immunofluorescence signal using confocal imaging and computer-based image analysis. The results of this quantitative approach showed that there was a greater percent colocalization of the TSPO signal with GFAP (astrocyte) than with Mac-1 (microglia) and it changed with disease progression (Loth et al., 2016). This state-of-the-art quantitative approach is an example of what is possible today using computer-based microscopy and image analysis software. During the writing of this review, a new study was published in a murine model of mesial temporal lobe epilepsy in which [^{18}F]DPA-714 microPET/computed tomographic scanning, autoradiography, and double label immunofluorescence was performed using a longitudinal approach (Nguyen et al., 2018). This study confirmed that the glial sources of the TSPO response was due to an early microglia component which decayed with time, and a predominant TSPO colocalization in astrocytes at the later time point providing further support that both microglia and astrocytes contribute to the TSPO signal. However, this study did not use computer-based microscopy and image analysis to determine the percent signal colocalization as performed by Loth et al. (2016), a method that is currently routinely used in many laboratories.

Recommendation 1. Unlike in the early studies that examined the cellular sources of the TSPO response, today there are well-validated and specific antibodies that are commercially available for TSPO as well as glial markers. Furthermore, there are a number of computer-based microscopy and image analysis software that are readily available commercially and can provide unbiased quantitative TSPO immunofluorescence signal colocalization with microglia and astrocytes markers. Therefore, it is recommended and highly desirable that studies examining the cellular sources of the TSPO response should be done with the same level of experimental rigor and be standardized so that studies using different animal models of human disease with a longitudinal design can be compared. This approach can also assist with the interpretation of human PET studies.

In summary, a critical review of the literature indicates that the cellular sources of the TSPO response to brain injury is disease specific and has a temporal component. Temporal examination of the glial sources in




Fig. 1. Temporal colocalization of TSPO with glial markers in wildtype (WT) and Sandhoff disease (SD) mice in the thalamus. A) Representative triple labeled immunofluorescent confocal images in the thalamus of WT and SD mice at four different ages. Triple labeled immunofluorescent confocal imaging confirmed that TSPO colocalized with the microglial marker Mac-1 as indicated by the purple and magenta colors and astrocyte marker GFAP as indicated by the yellow color. B) In the thalamus, the percent colocalization of TSPO with Mac-1 follows a similar pattern in both the WT and SD mice, with a more pronounced effect in the SD mice. The percent colocalization between TSPO and Mac-1 increases with age in both WT and SD mice and peaks at the 2 months of age. C) The percent colocalization between TSPO and GFAP also differs significantly between WT and SD mice, due to the low signal of both in the WT mice. In the SD mice, we observed a high degree of colocalization beginning at 1.5 months and remaining at 3 months of age. Each value represents the mean \pm SEM. $n = 3-4$ animals and experiments. Reprinted from *Neurobiology of Disease*, 85, MK Loth, J Choi, JL McGlothlan, MV Pletnikov, MG Pomper, TR Guilarte, TSPO in a murine model of Sandhoff disease: presymptomatic marker of neurodegeneration and disease pathophysiology, 174–186, 2016, with permission from Elsevier.

any human disease of interest is critically important because glial cell dynamics are known to change as a function of age and disease progression (McNeela et al., 2017). What is clear, however, is that it is inaccurate and a simplification of the complexity of glial responses to brain injury to continue to state that TSPO is a biomarker of microglia activation when astrocytes also play an important role in upregulating the TSPO signal in the neuroinflammatory response in many neurological and neurodegenerative disorders. Even within a single disease entity, the temporal profile of the TSPO signal originating from microglia/macrophages and/or astrocytes may vary depending upon the stage of the disease and unless a time-course in postmortem tissue is performed it is scientifically prudent to take great caution in making a generalized assumption, especially in PET studies. The most prudent approach is to

note that TSPO is representative of a glial response rather than a specific glial type. The technological advances that exist today provide the appropriate means to rigorously examine this important question in brain tissue from any animal model or in human post-mortem brain tissue.

What has been extensively validated over the years is that the TSPO signal measured with [3 H]R-PK11195 binding studies is driven by an increase in the maximal number of binding sites (B_{max}) with no change in the affinity constant (K_d) of the protein-ligand interaction (Chen et al., 2004; Chen & Guilarte, 2008; Gehlert et al., 1997; Rao et al., 2000; Venneti et al., 2004). What has yet to be discovered is the functional significance of the increased TSPO response to brain injury and neuroinflammation and whether the increased TSPO response has the same or

different biological function(s) in microglia and astrocytes. A resolution to these important question(s) can bring about a new era in the development of novel TSPO radiotracers with a possible selectivity for microglia or astrocytes. If the latter is ever accomplished, it would be a major scientific breakthrough in the TSPO field.

4. Exploration of new findings in psychiatric disorders

4.1. TSPO in psychiatric disorders: an exception to the rule or alternative explanations?

The last decade has seen an exponential increase in the number of studies related to TSPO as a biomarker of brain injury and neuroinflammation. Technological advances in high resolution PET scanners and the development of second generation TSPO ligands with more favorable pharmacokinetics and lower non-specific binding than the prototypical TSPO ligand R-PK11195, has facilitated this work (Cumming et al., 2017; Dolle et al., 2009; Dupont et al., 2017). Further, TSPO as a biomarker of inflammation has been extended to study other organ systems (Fairweather et al., 2014; Hardwick et al., 2005). Psychiatric disorders are no exception in the quest to use TSPO as a biomarker of neuroinflammation and the last 5 years have seen a number of reports with mixed results [see review by De Picker et al., 2017]. Paradoxically, some studies have provided the first description of decreased rather than increased brain TSPO levels in schizophrenia (Benloch et al., 2018; Coliste et al., 2017) and in major depression (Li et al., 2018). Overall, studies have found increased, decreased, or no change in brain TSPO levels in schizophrenia (De Picker et al., 2017) and in depression (Hannestad et al., 2013; Holmes et al., 2017; Li et al., 2018; Richards et al., 2018; Setiawan et al., 2015; Setiawan et al., 2018) compared to age-matched controls using PET imaging with radiolabeled R-PK11195 or with second generation TSPO PET ligands. There are many factors in study design such as TSPO radioligand used (R-PK11195 or second generation), PET imaging quantification with and without radioligand plasma input data, medicated vs naïve subjects, as well as corrections for the human TSPO polymorphism and vascular radioligand binding that have been discussed in the literature (De Picker et al., 2017; Turkheimer et al., 2015; Veronese et al., 2017) and could potentially account for some of the different findings of TSPO-PET in these psychiatric disorders. However, other factors that could potentially influence the interpretation of the PET results have yet to be investigated. This is critically important before the TSPO biomarker field takes a turn as it has recently occurred with studies on the function of TSPO (Morohaku et al., 2014; Tu et al., 2014, 2015). Today, TSPO appears to be an orphan protein looking for a function because the long-held dogma of its principal role as the rate limiting step in the transfer of cholesterol from the outer to the inner mitochondria membrane for steroid synthesis has been questioned (Gut et al., 2015; Selvaraj & Stocco, 2015; Selvaraj & Tu, 2016). While there are studies which seem to support and some that do not support the function of TSPO in steroid synthesis; this discourse has not yet occurred with the use of TSPO as a biomarker of brain injury and neuroinflammation. However, a reconceptualization of TSPO as a biomarker of neuroinflammation in psychiatric disorders has been proposed (Notter et al., 2017).

4.2. TSPO studies in schizophrenia prior to PET imaging—a historical review

Before examining the TSPO-PET literature in schizophrenia, it is important to review the early literature in which TSPO levels were measured in cells or postmortem brain tissue from schizophrenia subjects. These studies may provide important information that may help the interpretation of current TSPO-PET findings. As far back as 1986, there were studies examining the levels of TSPO in platelets from schizophrenia subjects (Gavish et al., 1986; Weizman et al., 1986). Two different studies showed that the maximal number of [³H]PK11195 binding sites (B_{max}) in platelets from untreated schizophrenia subjects did

not differ from age-matched healthy controls. However, in one study, schizophrenia subjects with chronic neuroleptic treatment exhibited a 30% decrease in [³H]PK11195 B_{max} with no change in K_d (affinity constant) compared to untreated schizophrenics and healthy controls (Gavish et al., 1986). In the second study, medicated schizophrenia subjects with and without tardive dyskinesia had 31.3% and 21.1% decreased [³H]PK11195 B_{max} relative to unmedicated patients and health controls (Weizman et al., 1986). These studies showed no significant alterations in [³H]PK11195 binding parameters to TSPO in platelets from untreated schizophrenia subjects relative to healthy controls, but TSPO B_{max} levels decreased significantly with schizophrenia medication or disease classification.

Later studies using platelets and other peripheral cells to measure TSPO levels supported the findings that TSPO binding parameters could be altered based on specific classifications of the disease. Additionally, co-morbidity factors such as alcohol and substance abuse also influenced TSPO binding parameters. For example, using granulocytes, Wodarz et al. (1998) found a 38% decrease in [³H]PK11195 B_{max} in residual-type schizophrenics relative to controls. Similarly, Ritsner et al. (2003) found a 30% decrease in platelet [³H]PK11195 B_{max} in aggressive schizophrenia relative to controls. Studies in peripheral cells from alcoholics, cocaine dependence, and after suicide attempts in schizophrenia subjects indicate that these factors can decrease [³H]PK11195 B_{max} in peripheral cells (Javaid et al., 1994; Suranyi-Cadotte et al., 1988). Collectively, all of these studies provided evidence of no change in peripheral cell [³H]PK11195 B_{max} or K_d in unmedicated schizophrenia subjects relative to healthy controls but schizophrenia with specific classifications or co-morbidity factors decreased peripheral cell [³H]PK11195 B_{max} with no change in K_d. These studies indicate that factors from disease classification to co-morbidity need to be carefully controlled to provide clarity to current findings from TSPO-PET studies in schizophrenia.

4.3. Relevant TSPO studies in brain tissue from schizophrenia subjects

There is a paucity of studies examining TSPO levels in postmortem brain tissue from schizophrenia subjects. The only study found in PubMed was by Kurumaji et al. (1997) that examined postmortem brain tissue from schizophrenia subjects and controls. Analysis of 26 brain regions indicated decreased [³H]PK11195 specific binding in superior parietal cortex, occipital cortex-visual area 1, and putamen that ranged from 23.1 to 41.5%. Parenthetically, in this study they found decreases in both B_{max} and K_d. That is, decreased number of binding sites with increased affinity.

Although the current review is focused on TSPO as a biomarker of neuroinflammation in schizophrenia, it was important to examine studies in which microglia and astrocyte morphology and viability have been examined in postmortem tissue from schizophrenia subjects. Notably, Wierzbica-Bobrowicz et al. (2004) have described that in brain tissue from schizophrenia subjects there is evidence of degenerative changes in microglia including fragmentation of processes, apoptotic changes and, importantly, degenerated mitochondria. These findings are relevant because TSPO is highly localized in mitochondria and their degeneration can influence the available number of TSPO binding sites that can be measured in TSPO-PET studies. A PubMed search under “schizophrenia and microglia” resulted in 272 hits and “schizophrenia and astrocytes” resulted in 350 hits (January 27, 2018). Therefore, there are many studies, beyond the scope of this review, that may be relevant in understanding glial cell mitochondrial changes that may influence TSPO expression in schizophrenia.

4.4. What are potential but yet unexplored interpretations of TSPO PET findings in schizophrenia?

The current results of brain TSPO changes (increased, decreased, or no change) in schizophrenia relative to age-matched controls may

depend on other potentially important factors that have yet to be considered but may need to be experimentally controlled and/or examined. Several points to this effect and relevant recommendations are provided below:

POINT 1. Disease modifying effects on peripheral TSPO levels or TSPO ligand binding proteins may alter the availability of TSPO radioligands for brain uptake:

It is well recognized that TSPO levels in peripheral organs including peripheral circulating blood cells are much higher than those expressed in normal or even in pathological brain tissue. Schizophrenia is thought to have a peripheral inflammatory component. Thus, it is entirely plausible that in peripheral tissues TSPO levels could be differentially altered from those in the brain at different stages of the disease so that peripheral TSPO is markedly altered and may modulate the amount of radiotracer delivered to the brain. A possible effect of increasing or decreasing peripheral TSPO levels could be reflected as an “apparent” decrease or increase in TSPO radioligand brain uptake measured by PET. In fact, such a modulatory effect of a treatment on [¹¹C]PBR28 arterial plasma concentration and brain uptake has been reported with LPS and PLX3397 in nonhuman primates (Hillmer et al., 2017). PLX3397 is a selective colony-stimulating factor 1 receptor inhibitor used to deplete microglia in the brain.

Similarly, it is possible that in schizophrenia or in other disease conditions, there is induction of gene(s) that encode protein(s) in peripheral organs, blood, and possibly even in the brain neuropil that are able to bind R-PK11195 and/or second generation TSPO ligands; thus, potentially altering radiotracer delivery, uptake, or binding to TSPO in the brain. Such a protein exists and has already been described in the literature but has not been accounted for in TSPO-PET studies. The acute phase reactant α 1-acid glycoprotein (AAG), an abundant protein in plasma, has been shown to bind PK11195 with high affinity (Lockhart et al., 2003). While there is no current information on whether AAG also binds second generation TSPO ligands, it is highly likely that it does. Relevant to schizophrenia, plasma levels of AAG have been shown to be altered in schizophrenia subjects (Goodman et al., 1961) and AAG plasma concentrations are altered by drugs used for the treatment of schizophrenia (Levinson & Levine, 1995; Telford et al., 2012). Furthermore, AAG protein levels are markedly increased in plasma, and potentially in the brain neuropil, by pathophysiological conditions including inflammation, depression, cancer, and autoimmune deficiency syndrome, altering the pharmacokinetic disposition of drugs that bind to AAG (Garrido et al., 1999; Holladay et al., 1998). Notably, AAG genetic variants have been shown to alter the blood to brain transfer of AAG-binding drugs (Jolliet-Riant et al., 1998). Finally, relevant to psychiatric disorders, studies have found a positive association between serum AAG with age in depressed female subjects (Young et al., 1999). In summary, the above-mentioned studies indicate that it is critically important to determine and understand the potential for disease-modified levels of AAG in plasma, and potentially in the brain, on TSPO ligand uptake and binding in the brain and its implications in the interpretation of TSPO-PET studies.

Recommendation 2. determine if AAG binds second generation TSPO radioligands with high affinity.

Recommendation 3. Examine plasma levels of AAG in schizophrenia subjects at the time of the TSPO-PET study, and under medication treatments, to determine if differences in AAG occur between schizophrenia subjects and age-matched healthy controls at different stages of the disease.

Recommendation 4. Determine in post-mortem brain tissue if AAG levels are altered in psychiatric disorders, neuroinflammatory disease, and other neuropathological conditions.

POINT 2. Microglia and astrocyte number in schizophrenia: analysis using rigorous unbiased stereological cell counting: There is evidence of microglia number changes in schizophrenia (Frick et al., 2013; van Kesteren et al., 2017) as well as astrocytes (Kim et al., 2017; Williams

et al., 2013). Therefore, changes in microglia and/or astrocyte number in schizophrenia may influence TSPO-PET findings.

Recommendation 5. Determine microglia and astrocyte number in schizophrenia brain tissue and age-matched healthy controls using rigorous unbiased stereological cell counting methods.

POINT 3. Mitochondria number in microglia and astrocytes in schizophrenia: are they different? Microglia and astrocytes are the brain cells that express and increase TSPO levels in a variety of brain pathologies. Further, TSPO is a protein that is highly expressed in mitochondria. Therefore, one must consider the possibility that TSPO levels in schizophrenia subjects could be influenced by disease-modifying changes in glial cell mitochondria number; thus, altering the putative number of TSPO binding sites even under conditions where no glial cell activation or glia cell number change is observed. There is evidence of decreased mitochondria number in glial cells in the brain of schizophrenia subjects (Ben-Scachar, 2002; Roberts, 2017; Somerville et al., 2012) as well as changes in mitochondria genes that could modulate TSPO expression (Iwamoto et al., 2005).

Recommendation 6. Determine mitochondria integrity and number in microglia and astrocytes in postmortem brain tissue in schizophrenia using 3-dimensional scanning electron microscopy.

POINT 4. Modeling schizophrenia in preclinical animal models and their effect on brain TSPO levels - what do they tell us? Schizophrenia is a uniquely human disease and it is not possible to capture its complexity in an animal model. However, animal models of schizophrenia do provide a presumed behavioral and neuropathology phenotype and mimic certain aspects of the human disease. There is a paucity of studies using appropriate animal models of schizophrenia that also assess brain TSPO levels. A recent study by Notter et al. (2018) examined the relationship of the brain TSPO response in an inflammatory developmental animal model of schizophrenia; that is, the injection of polyriboinosinic-polyribocytidylic acid (poly:IC) in time-pregnant mice. The offspring of these poly:IC exposed dams express deficits in prepulse inhibition of the acoustic startle response and social interaction similar to what has been described in schizophrenia subjects (Notter et al., 2017). In the same study, the authors also performed TSPO-PET using [¹¹C]DPA-713 in recent-onset schizophrenia subjects and matched healthy controls. They found a near ($p = 0.051$), but non-significant decrease in TSPO levels by PET in the middle frontal gyrus of schizophrenic subject relative to controls.

In the animal portion of this study, the authors perform TSPO immunohistochemistry in frontal cortex and hippocampus of poly:IC exposed animals and controls. They found an apparent decrease in brain TSPO levels using immunofluorescent labeling that appears to be selective to the frontal cortex with no change in the hippocampus (Notter et al., 2018). TSPO immunofluorescence signal was measured using relative optical intensity. However, receptor autoradiography provided no evidence of a change in TSPO specific binding in the frontal cortex or hippocampus of poly:IC exposed animals compared to non-exposed. The authors recognized the discrepancy in results using TSPO immunohistochemistry and receptor autoradiography but they argue that the differences using these two methods are explained by the lower resolution of *ex vivo* radiotracer autoradiography compared to immunohistochemistry which can provide analysis at the cellular and subcellular level. They state that the immunohistochemistry technique is likely to be more suitable in detecting relatively small changes in TSPO expression, and thus confirmation of decreased TSPO levels in the frontal cortex of poly:IC exposed animals. While the authors make an argument for a significant and selective decrease in TSPO levels in the frontal cortex in the *in utero* exposed poly:IC animals based on the immunohistochemistry results, this argument needs further validation as they did not measure the TSPO signal specifically at the cellular and/or subcellular level. That is, they performed optical density readings at the gross anatomical level which not only includes a signal from TSPO labeled cells but also

microvasculature and specific and non-specific primary and/or secondary antibody staining. It is likely that the receptor autoradiography data is providing the most accurate results. That is, no change in TSPO levels in frontal cortex or hippocampus of poly:IC animals compared to non-exposed controls as has been found in several of the TSPO-PET studies in schizophrenia subjects including results from the same investigators (Notter et al., 2018).

Recommendation 7. In animal studies it is important to use multiple methods to confirm a finding, including: 1) radioligand binding such as film and emulsion microautoradiography with the latter having resolution at the cellular level similar to confocal imaging immunofluorescence, 2) radioligand saturation isotherms with Scatchard analysis of brain tissue to determine Kd and Bmax values which is a more accurate representation of the radioligand interaction than a single radioligand concentration as it is typically used in autoradiography studies, and 3) protein analysis by quantitative Western blot.

POINT 5. Maintain independent variables the same in comparison groups: In the same study, a potentially important experimental design problem is that the animals used for TSPO immunohistochemistry received a behavioral experience, *i.e.*, assessment of pre-pulse inhibition and social interaction (Notter et al., 2018-supplementary materials) while the animals used for TSPO autoradiography did not experience behavioral tests. Thus, it is possible that the behavioral experience, along with the handling of animals for behavior—a form of environmental enrichment—may differentially affect TSPO brain levels in poly:IC animals used for immunohistochemistry relative to those used for autoradiography. Although the authors waited a week prior to euthanizing the animals that experienced the behavioral protocol, the effect of the behavior on regional TSPO levels is not known and was not studied.

Recommendation 8. All animals in both treatment groups should have experienced an identical behavioral protocol in order to eliminate any potential modulatory effects from the behavior and manipulation which only one set of animals experienced. In future studies, it is possible to use one hemisphere for autoradiography studies and the other hemisphere for immunohistochemistry so that there is a direct comparison of the two different methods in the same animal. Alternatively, an entirely behaviorally naïve set of animals should have been used for both the immunohistochemistry and autoradiography studies.

5. Concluding remarks

It is clear from the limited but increasing number of human TSPO studies in schizophrenia and other psychiatric disorders such as depression that the findings are equivocal in regards to TSPO as a biomarker of neuroinflammation. During the writing of this review, a report appeared of a meta-analysis in individual participant data of TSPO-PET studies in first-episode schizophrenia patients with psychosis and healthy controls (Plaven-Sigra et al., 2018). The meta-analysis concludes that overall, brain TSPO levels are decreased in 75 schizophrenia participants relative to 77 healthy controls from five independent studies. This meta-analysis provides evidence of a decrease in TSPO levels in the schizophrenia brain relative to aged-matched healthy controls using PET imaging and provides a compelling reason to study the recommendations outlined in this review.

While there are differences amongst the TSPO-PET studies in schizophrenia in regards to TSPO radioligand used, PET imaging quantification with and without radioligand plasma input data and analytical methods used, medicated vs naïve subjects, as well as corrections for the human TSPO polymorphism and vascular radioligand binding; there are other important factors that have not been considered and are delineated in this review. From the perspective of using preclinical animal models, it is critically important that rigorous experimental design and

interpretation of results be provided so that potential confounding factors are eliminated to reduce paradoxical findings and increase the reliability and reproducibility of the results. These considerations of rigor and authentication will go a long way to bring clarity so that the TSPO biomarker field is not turned up-side-down as has recently happened with studies on the function of TSPO. To this end, a remaining point of contention in the literature is the cellular sources of the TSPO response to brain injury and neuroinflammation which is often and still described by a significant number of publications in the literature as an exclusively microglial response. The current availability of well-validated and specific TSPO antibodies and computer-based software for microscopy and image analysis of confocal images now make it possible to provide accurate, reliable, and direct results on the cellular sources of the TSPO response with quantitative analysis of signal colocalization with glial markers in any human brain disease or preclinical model in which brain tissue samples are available. Using this type of approach, there is evidence that the TSPO response to different brain pathologies is derived from both microglia and astrocytes, although they may have different temporal profiles and degree of induction. The ready dismissal of the nature of the TSPO response to be less than a complex interaction of microglia and astrocytes, “a glial response”, in the context of diverse brain pathologies or neurodegenerative/mental disorders and their progression is no longer tenable and will not provide a near term solution, but it will delay the inevitable reality of how nature works. The progression of science will eventually provide the correct answer since “*in science, truth always wins*” (Nobel Prize winner Max Perutz; <http://www.vega.org.uk/video/programme/1>).

Conflict of interest statement

The author declares that there are no conflicts of interest.

Acknowledgments

This review is dedicated to the many doctoral students, postdoctoral fellows, research staff, and collaborators that have contributed significantly to this work in my laboratory over the last 20 plus years. I am especially indebted to my senior research scientist Ms. Jennifer Dziedzic that has been involved in these studies over the last 20 years. The TSPO work from our laboratory has been continuously funded from the National Institute of Environmental Health Sciences grant number ES007062-19. I acknowledge that there are many more publications that could have been cited, but the ones cited in this review were selected to support a specific comment or scientific point of view.

References

- Abourbeh, G., Theze, B., Maroy, R., Dubois, A., Brulon, V., Fontyn, Y., ... Boisgard, R. (2012). Imaging microglial/macrophage activation in spinal cords of experimental autoimmune encephalomyelitis rats by positron emission tomography using the mitochondrial 18 kDa translocator protein radioligand [¹⁸F]-DPA-714. *The Journal of Neuroscience* 32, 5728–5736.
- Anholt, R.R.H., Pedersen, P.L., De Suetz, E.B., & Snyder, S.H. (1986). The peripheral-type benzodiazepine receptor-localization to the mitochondrial outer membrane. *The Journal of Biological Chemistry* 261, 576–583.
- Arlicot, N., Katsifis, A., Garreau, L., Mattner, F., Vergote, J., Duval, S., ... Chalou, S. (2008). Evaluation of CLINDE as potent translocator protein (18 kDa) SPECT radiotracer reflecting the degree of neuroinflammation in a rat model of microglial activation. *European Journal of Nuclear Medicine and Molecular Imaging* 35, 2203–2211.
- Arlicot, N., Tronel, C., Bodard, S., Garreau, L., de la Crompe, B., Vandeveld, I., ... Chalou, S. (2014). Translocator protein (18 kDa) mapping with [¹²⁵I]-CLINDE in the quinolinic acid rat model of excitotoxicity: a longitudinal comparison with microglial activation, astrogliosis, and neuronal death. *Molecular Imaging* 13, 1–11.
- Baestrup, C., & Squires, R.F. (1977). Specific benzodiazepine receptors in rat brain characterized by high-affinity [³H]diazepam binding. *Proceedings of the National Academy Science* 74, 3805–3809.
- Banati, R.B., Goerres, G.W., Myers, R., Gunn, R.N., Turkheimer, F.E., Kreutzberg, G.W., ... Duncan, J.S. (1999). [¹¹C]-R-PK11195 positron emission tomography imaging of activated microglia in Rasmussen's encephalitis. *Neurology* 53, 2199–2203.

- Banati, R.B., Myers, R., & Kreutzberg, G.W. (1997). PK ('peripheral benzodiazepine') – binding sites in the CNS indicate early and discrete brain lesions: microautoradiography detection of [³H]PK11195 binding to activated microglia. *Journal of Neurocytology*, 26, 77–82.
- Banati, R.B., Newcombe, J., Gunn, R.N., Cagnin, A., Turkheimer, F., Heppner, F., ... Myers, R. (2000). The peripheral benzodiazepine binding site in the brain in multiple sclerosis: quantitative *in vivo* imaging of microglia as measure of disease activity. *Brain*, 123, 2321–2337.
- Benavides, J., Cornu, P., Dennis, T., Dubois, A., Hauw, J.J., MacKenzie, E.T., ... Scatton, B. (1988). Imaging of human brain lesions with an omega 3 site radioligand. *Annals of Neurology*, 24, 708–712.
- Benavides, J., Dubois, A., Gotti, B., Buordiol, F., & Scatton, B. (1990). Cellular distribution of omega 3 (peripheral type benzodiazepine) binding sites in the normal and ischaemic rat brain: an autoradiographic study with the photoaffinity ligand [³H]PK14105. *Neuroscience Letters*, 114, 32–38.
- Benavides, J., Fage, D., Carter, C., & Scatton, B. (1987). Peripheral type benzodiazepine binding sites are a sensitive indirect index of neuronal damage. *Brain Research*, 421, 167–172.
- Benavides, J., Quarteronet, D., Imbault, F., Malgouris, C., Uzan, A., Renault, C., ... Lefleur, G. (1983). Labelling of "peripheral-type" benzodiazepine binding sites in the rat brain by using [³H]PK11195, an isoquinoline carboxamide derivative: kinetic studies and autoradiographic localization. *Journal of Neurochemistry*, 41, 1744–1750.
- Benloch, J.M., Gonzalez, A.J., Pani, R., Preziosi, E., Jackson, C., Murphy, J., ... Schwaiger, M. (2018). The MINDVIEW project: First results. *Eur Psych*, 50, 21–27.
- Ben-Scabar, D. (2002). Mitochondria dysfunction in schizophrenia: a possible linkage to dopamine. *Journal of Neurochemistry*, 83, 1241–1251.
- Biesmans, S., Acton, P.D., Cotto, C., Langlois, X., Ver Donck, L., Bouwknecht, J.A., ... Nuydens, R. (2015). Effect of stress and peripheral immune activation on astrocyte activation in transgenic bioluminescent Cgfp-luc mice. *Glia*, 63, 1126–1137.
- Brackhan, M., Bascunana, P., Postema, J.M., Ross, T.L., Bengel, F.M., Bankstahl, M., & Bankstahl, J.P. (2016). Serial quantitative TSPO-targeted PET reveals peak microglia activation up to 2 weeks after an epileptogenic brain insult. *Journal of Nuclear Medicine*, 57, 1302–1308.
- Brahmachari, S., Fung, Y.K., & Pahan, K. (2006). Induction of glial fibrillary acidic protein expression in astrocytes by nitric oxide. *The Journal of Neuroscience*, 26, 4930–4939.
- Cagnin, A., Brooks, D.J., Kennedy, A.M., Gunn, R.N., Myers, R., & Turkheimer, F.E. (2001). *In vivo* measurement of activated microglia in dementia. *The Lancet*, 358, 461–467.
- Cagnin, A., Myers, R., Gunn, R.N., Lawrence, A.D., Stevens, T., Kreutzberg, G.W., ... Banati, R. B. (2001). *In vivo* visualization of activated glia by [¹¹C]-(R)-PK11195-PET following herpes encephalitis reveals projected neuronal damage beyond the primary focal lesion. *Brain*, 124, 2014–2027.
- Chen, M.K., Baidoo, K., Verina, T., & Guilarte, T.R. (2004). Peripheral benzodiazepine receptor imaging in CNS demyelination: functional implications of anatomical and cellular localization. *Brain*, 127, 1379–1392.
- Chen, M.K., & Guilarte, T.R. (2006). Imaging the peripheral benzodiazepine receptor response in central nervous system demyelination and remyelination. *Toxicological Sciences*, 91, 532–539.
- Chen, M.K., & Guilarte, T.R. (2008). Translocator protein 18 kDa (TSPO): molecular sensor of brain injury and repair. *Pharmacological Therapy*, 118, 1–17.
- Coliste, K., Plaven-Sigra, P., Fatouros-Bergman, H., Victorsson, P., Schain, M., Forsberg, A., ... Cervenka, S. (2017). Lower levels of the glial marker TSPO in drug-naïve first-episode psychosis patients as measured using PET and [¹¹C]PBR28. *Molecular Psychiatry*, 22, 850–856.
- Conway, E.L., Gundlach, A.L., & Craven, J.A. (1998). Temporal changes in glial fibrillary acidic protein messenger RNA and [³H]PK11195 binding in relation to imidazoline-2-receptor and α 2-adrenoceptor binding to the hippocampus following transient global forebrain ischaemia in the rat. *Neuroscience*, 82, 805–817.
- Cosenza-Nashhat, M., Zhao, M., Suh, H., Morgan, J., Natividad, R., Morgello, S., & Lee, S.C. (2009). Expression of the translocator protein of 18 kDa by microglia, macrophages and astrocytes based on immunohistochemical localization in abnormal human brain. *Neuropath Applied Neurobiology*, 35, 306–328.
- Coughlin, J.M., Wang, Y., Ma, S., Yue, C., Kim, P.K., Adams, A.V., ... Pomper, M.G. (2014). *Journal of Neurovirology*, 20, 219–232 Regional distribution of translocator protein using [¹¹C]DPA-713 PET in individuals infected with HIV.
- Coughlin, J.M., Wang, Y., Minn, I., Bienko, N., Ambinder, E.B., Xu, X., ... Pomper, M.G. (2017). Imaging of glial cell activation and white matter integrity in brains of active and recently retired National Football League players. *JAMA Neurology*, 74, 67–74.
- Coughlin, J.M., Wang, Y., Munro, C.A., Ma, S., Yue, C., Chen, M.S., ... Pomper, M.G. (2015). Neuroinflammation and brain atrophy in former NFL players: an *in vivo* multimodal imaging pilot study. *Neurobiology of Disease*, 74, 58–65.
- Cumming, P., Burgher, B., Patkar, O., Breakspear, M., Vasdev, N., Thomas, P., ... Banati, R. (2017). Sifting through the surfeit of neuroinflammation tracers. *Journal of Cerebral Blood Flow and Metabolism*, 38, 204–224.
- De Picker, L.J., Morrens, M., Chance, S.A., & Boche, D. (2017). Microglia and brain plasticity in acute psychosis and schizophrenia illness course: a meta-review. *Frontiers Psychology*, 8, 238.
- Debruyne, J.C., Versijpt, J., Van Laere, K.J., De Vos, F., Keppens, J., Strijckmans, K., ... De Reuck, J.L. (2003). PET visualization of microglia in multiple sclerosis patients using [¹¹C]PK11195. *European Journal of Neurology*, 10, 257–264.
- Dickens, A.M., Vainio, S., Marjamaki, P., Johansson, J., Lehtiniemi, P., Rokka, J., ... Airas, L. (2014). Detection of microglial activation in an acute model of neuroinflammation using PET and radiotracers [¹¹C]-(R)-PK11195 and [¹⁸F]-GE-180. *Journal of Nuclear Medicine*, 55, 466–472.
- Diorio, D., Welner, S.A., Butterworth, R.F., Meaney, M.J., & Suranyi-Cadotte, B.E. (1991). Peripheral benzodiazepine binding sites in Alzheimer's disease frontal cortex and temporal cortex. *Neurobiology of Aging*, 12, 255–258.
- Doble, A., Malgouris, C., Daniel, M., Daniel, N., Imbault, F., Basbaum, A., ... Lefeur, G. (1987). Labelling of peripheral-type benzodiazepine binding sites in human brain using [³H]PK11195: anatomical and subcellular distribution. *Brain Research Bulletin*, 18, 49–61.
- Dolle, F., Luus, C., Reynolds, A., & Kassio, M. (2009). Radiolabelled molecules for imaging the translocator protein (18 kDa) using positron emission tomography. *Current Medicinal Chemistry*, 16, 2899–2923.
- Domene, A., Cavanagh, C., Page, G., Bodard, S., Klein, C., Delarasse, C., ... Krantic, S. (2016). Expression of phenotypic astrocyte marker is increased in a transgenic mouse model of Alzheimer's disease versus age-matched controls: a presymptomatic stage study. *International Journal of Alzheimer's Disease*, 2016, 5696241.
- Doorduyn, J., de Vries, E.F., Dierck, R.A., & Klein, H.C. (2008). PET imaging of the peripheral benzodiazepine receptor: monitoring disease progression and therapy response in neurodegenerative disorders. *Current Pharmaceutical Design*, 14, 3297–3315.
- Dubois, A., Benavides, J., Peny, B., Duverger, D., Fage, D., Gotti, B., ... Scatton, B. (1988). Imaging of primary and remote ischaemic and excitotoxic brain lesions. An autoradiographic study of peripheral type benzodiazepine binding sites in the rat and cat. *Brain Research*, 445, 77–90.
- Dupont, A.C., Largeau, B., Santiago, R., Guilloteau, D., & Arlicot, N. (2017). Translocator protein-18 kDa (TSPO) positron emission tomography (PET) imaging and its clinical impact in neurodegenerative diseases. *International Journal of Molecular Science*, 18, E785.
- Fairweather, D., Coronado, M.J., Garton, A.E., Dziedzic, J.L., Bucek, A., Cooper, L.T., ... Guilarte, T.R. (2014). Sex differences in translocator protein 18 kDa (TSPO) in the heart: implications for imaging myocardial inflammation. *Journal of Cardiovascular Translational Research*, 7, 192–202.
- Frick, L.R., Williams, K., & Pittenger, C. (2013). Microglial dysregulation in psychiatric disease. *Clinical & Developmental Immunology*, 2013, 608654.
- Garrido, M.J., Valle, M., Calvo, R., & Troconiz, I.F. (1999). Altered plasma and brain disposition and pharmacodynamics of methadone in abstinent rats. *Journal of Pharmacology and Experimental Therapeutics*, 288, 179–187.
- Gavish, M., Weizman, A., Karo, L., Tyano, S., & Tanne, Z. (1986). Decreased peripheral benzodiazepine binding sites in platelets of neuroleptic-treated schizophrenics. *European Journal of Pharmacology*, 121, 275–279.
- Gehlert, D.R., Stephenson, D.T., Schober, D.A., Rash, K., & Clemens, J.A. (1997). Increased expression of peripheral benzodiazepine receptors in the facial nucleus following motor neuron axotomy. *Neurochemistry International*, 31, 705–713.
- Gehlert, D.R., Yamamura, H.I., & Wamsley, J.K. (1985). Autoradiographic localization of "peripheral-type" benzodiazepine binding sites in the rat brain, heart, and kidney. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 328, 454–460.
- Gerhard, A., Banati, R.B., Goerres, G.B., Cagnin, A., Myers, R., Gunn, R.N., ... Brooks, D.J. (2003). [¹¹C]-PK11195 PET imaging of microglial activation in multiple system atrophy. *Neurology*, 61, 686–689.
- Gerhard, A., Pavese, N., Hottot, G., Turkheimer, F., Es, M., Hammers, A., ... Brooks, D.J. (2006). *In vivo* imaging of microglial activation with [¹¹C]-(R)-PK11195 PET in idiopathic Parkinson's disease. *Neurobiology of Disease*, 21, 404–412.
- Gerhard, A., Schwarz, J., Myers, R., Ise, R., & Banati, R.B. (2005). Evolution of microglial activation in patients after ischemic stroke: a [¹¹C]-(R)-PK11195 PET study. *NeuroImage*, 24, 591–595.
- Gerhard, A., Trender-Gerhard, I., Turkheimer, F., Quinn, N.P., Bhatia, K.P., & Brooks, D.J. (2006). *In vivo* imaging of microglia activation with [¹¹C]-(R)-PK11195 PET in Progressive Supranuclear Palsy. *Movement Disorder*, 21, 89–93.
- Gerhard, A., Watts, J., Trender-Gerhard, I., Turkheimer, F., Banati, R.B., Bhatia, K., & Brooks, D.J. (2004). *In vivo* imaging of microglial activation with [¹¹C]-PK11195 PET in corticobasal degeneration. *Movement Disorder*, 19, 1221–1226.
- Ghadery, C., Koshimori, Y., Coakeley, S., Harris, M., Rusjan, P., Kim, J., ... Strafella, A.P. (2017). Microglia activation in Parkinson's disease using [¹⁸F]-FEPPA. *Journal of Neuroinflammation*, 14, 8.
- Goodman, M., Luby, E.D., Froham, C.E., & Gottlieb, J.S. (1961). Orosomucoid in schizophrenia. *Nature*, 192, 370–371.
- Gorina, R., Font-Nieves, M., Marquez-Kisinosky, L., Santalucia, T., & Planas, A.M. (2011). Astrocyte TLR4 activation induces a proinflammatory environment through the interleukin 59, MyD88-dependent NF κ B signaling, MAPK, and Jak1/Stat1 pathways. *Glia*, 59, 242–255.
- Groom, G.N., Junck, L., Foster, N.L., Frey, K.A., & Kuhl, D.E. (1995). PET of peripheral benzodiazepine binding sites in the microgliosis of Alzheimer's disease. *Journal of Nuclear Medicine*, 36, 2207–2210.
- Guilarte, T.R., Kuhlmann, A.C., O'Callaghan, J.P., & Micelli, R.C. (1995). Enhanced expression of peripheral benzodiazepine receptors in the trimethyltin-exposed rat brain: a biomarker of neurotoxicity. *Neurotoxicology*, 16, 441–450.
- Guilarte, T.R., Nihei, M.K., McGlothlan, J.L., & Howard, A.S. (2003). Methamphetamine-induced deficits of brain monoaminergic markers: distal axotomy or neuronal plasticity. *Neuroscience*, 122, 499–513.
- Gut, P., Zweckstetter, M., & Banati, R.B. (2015). Lost in translocation: the functions of the 18-kDa translocator protein. *Trends in Endocrinology Metabolism*, 26, 349–356.
- Hammoud, D.A., Endres, C.J., Chander, A.R., Guilarte, T.R., Wong, D.F., Sacktor, N.C., ... Pomper, M.G. (2005). Imaging glial cell activation with [¹¹C]-R-PK11195 in patients with AIDS. *Journal of Neurovirology*, 11, 346–355.
- Hannestad, J., Dellagioia, N., Gallezot, J., Lim, K., Nabulsi, N., Esterlis, I., ... Carson, R.E. (2013). The neuroinflammation marker translocator protein is not elevated in individuals with mild-to-moderate depression. *Brain Behavior and Immunity*, 33, 131–138.
- Hannestad, J., Gallezot, J.D., Schaffbauer, T., Lim, K., Kloczynski, T., Morris, E.D., ... Cosgrove, K.P. (2012). Endotoxin-induced systemic inflammation activates microglia: [¹¹C]-PBR28 positron emission tomography in nonhuman primates. *NeuroImage*, 63, 232–239.

- Hardwick, M.J., Chen, M.-K., Baidoo, K., Pomper, M.G., & Guilarte, T.R. (2005). *In vivo* imaging of peripheral benzodiazepine receptors in mouse lungs: a biomarker of inflammation. *Molecular Imaging* 4, 432–438.
- Harhausen, D., Sudmann, V., Khojasteh, U., Muller, J., Zille, M., Graham, K., ... Wunder, A. (2013). Specific imaging of inflammation with the 18 kDa translocator protein ligand DPA-714 in animal models of epilepsy and stroke. *PLoS One* 8, e69529.
- Henkel, K., Karitzky, J., Schmid, M., Mader, I., Glatting, G., Unger, J.W., ... Landwehrmeyer, G.B. (2004). Imaging activated microglia with PET and [¹¹C]-PK11195 in corticobasal degeneration. *Movement Disorder* 19, 817–821.
- Hillmer, A.T., Holden, D., Fowles, K., Nabulsi, N., West, B.L., Carson, R.E., & Cosgrove, K.P. (2017). Microglia depletion and activation: a [¹¹C]-PBR28 PET study in nonhuman primates. *EJNMMI Research* 7, 59.
- Hillmer, A.T., Sandiego, C.M., Hannestad, J., Angarita, G.A., Kumar, A., McGovern, E.M., ... Cosgrove, K.P. (2017). *In vivo* imaging of translocator protein, a marker of activated microglia, in alcohol dependence. *Molecular Psychiatry* 22, 1759–1766.
- Holladay, J.W., Dewey, M.J., & Yoo, S.D. (1998). Pharmacokinetics and antidepressant activity of fluoxetine in transgenic mice with elevated serum alpha1-acid glycoproteins levels. *Drug Metabolism and Disposition* 26, 20–24.
- Holmes, S.E., Hinz, R., Conen, S., Gregory, C.J., Matthews, J.C., Anton-Rodriguez, J.M., ... Talbot, P.S. (2017). Elevated translocator protein in anterior cingulate in major depression and a role for inflammation in suicidal thinking: a positron emission tomography study. *Biological Psychiatry* 83, 61–69.
- Iaccarino, L., Moresco, R.M., Presotto, L., Bugiani, O., Iannaccone, G., Tagliavini, F., & Perani, D. (2017). An *in vivo* [¹¹C](R)-PK11195 PET and *in vitro* pathology study of microglia activation in Creutzfeldt-Jakob disease. *Molecular Neurobiology* 55, 2856–2868.
- Iaccarino, L., Presotto, L., Bettinardi, V., Gianolli, L., Roiter, I., Capellari, S., ... Perani, D. (2018). An *in vivo* [¹¹C]-PK PET study of microglia activation in Fatal Familial Insomnia. *Annals of Clinical Translational Neurology* 5, 11–18.
- Israel, I., Ohsiek, A., Al-Momani, E., Albert-Weissenberger, C., Stetter, C., Menci, S., ... Siren, A.-L. (2016). Combined [¹⁸F]-DPA-713 micro-positron emission tomography and autoradiography imaging of microglia activation after closed head injury in mice. *Journal of Neuroinflammation* 13, 140.
- Iwamoto, K., Bundo, M., & Kato, T. (2005). Altered expression of mitochondria-related genes in postmortem brains of patients with bipolar disorder of schizophrenia, as revealed by large-size DNA microarray analysis. *Human Molecular Genetics* 14, 241–253.
- Janssen, B., Vugts, D.J., Funke, U., Molenaar, G.T., Kruijer, P.S., van Berckel, B.N.M., ... Windhorst, A.D. (2017). Imaging of neuroinflammation in Alzheimer's disease, multiple sclerosis, and stroke: recent developments in positron emission tomography. *Biochimica et Biophysica Acta* 1862, 425–441.
- Javadi, J.I., Notorangelo, M.P., Pandey, S.C., Reddy, P.L., Pandey, G.N., & Davis, J.M. (1994). Peripheral benzodiazepine receptors are decreased during cocaine withdrawal in humans. *Biological Psychology* 36, 44–50.
- Jeyakumar, M., Butters, T.D., Dwek, R.A., & Platt, F.M. (2002). Glycosphingolipid lysosomal storage disease: therapy and pathogenesis. *Neuropath and Applied Neurobiol* 28, 343–357.
- Ji, B., Maeda, J., Sawada, M., Ono, M., Okauchi, T., Inaji, M., ... Suhara, T. (2008). Imaging the peripheral benzodiazepine receptor expression as biomarkers of detrimental versus beneficial glial responses in mouse models of Alzheimer's disease and other CNS pathologies. *The Journal of Neuroscience* 28, 12255–12267.
- Joliet-Riant, A., Boukef, M.F., Duche, J.C., Simon, N., & Tillement, J.P. (1998). The genetic variant A of human alpha1-acid glycoprotein limits the blood to brain transfer of drugs it binds. *Pharmacology Letters* 62, 219–226.
- Junck, L., Olson, J.M.M., Ciliax, B.J., Koeppe, R.A., Watkins, G.L., Jewett, D.M., ... Young, A.B. (1989). PET imaging of human gliomas with ligands for the peripheral benzodiazepine binding site. *Annals of Neurology* 26, 752–758.
- Kanaan, N.M., Kordower, J.H., & Collier, T.J. (2010). Age-related changes in glial cells of dopamine midbrain subregions in rhesus monkeys. *Neurobiology of Aging* 31, 937–952.
- Kannan, S., Saadani-Makki, F., Muzik, O., Chakraborty, P., Mangner, T.J., Janisse, J., ... Chugani, D.C. (2007). Microglial activation in perinatal rabbit brain induced by intra-uterine inflammation: detection with [¹¹C]-R-PK11195 and small-animal PET. *Journal of Nuclear Medicine* 48, 946–954.
- Kim, R., Healey, K.L., Sepulveda-Orengo, M.T., & Reissner, K.J. (2017). Astroglial correlates of neuropsychiatric disease: from astrocytopathy to astrogliosis. *Progress in Neuro-psychopharmacology and Biological Psychiatry* 50278-5846(17), 30485–...
- Knezevic, D., Verhoeff, N.P.L., Hafizi, S., Strafella, A.P., Graff-Guerrero, A., Rajji, T., ... Mizhari, R. (2017). Imaging microglia activation and amyloid burden in amnesic mild cognitive impairment. *Journal of Cerebral Blood Flow and Metabolism* 1, 271678X17741395.
- Krueger, K.E., & Papadopoulos, V. (1990). Peripheral-type benzodiazepine receptors mediate translocation of cholesterol from outer to inner mitochondrial membranes in adrenocortical cells. *The Journal of Biological Chemistry* 265, 15015–15022.
- Kuhlmann, A.C., & Guilarte, T.R. (1997). The peripheral benzodiazepine receptor is a sensitive indicator of domoic acid neurotoxicity. *Brain Research* 751, 281–288.
- Kuhlmann, A.C., & Guilarte, T.R. (1999). Regional and temporal expression of the peripheral benzodiazepine receptor in MPTP neurotoxicity. *Toxicological Sciences* 48, 107–116.
- Kuhlmann, A.C., & Guilarte, T.R. (2000). Cellular and subcellular localization of peripheral benzodiazepine receptors after trimethyltin neurotoxicity. *Journal of Neurochemistry* 74, 1694–1704.
- Kurumaji, A., Wakai, T., & Toru, M. (1997). Decreases in peripheral-type benzodiazepine receptors in postmortem brains of chronic schizophrenics. *Journal of Neural Transmission* 104, 1361–1370.
- Lavisse, S., Guillemier, M., Herard, A.-S., Petit, F., Delahaye, M., Van Camp, N., ... Escartin, C. (2012). Reactive astrocytes overexpress TSPO and are detected by TSPO positron emission tomography imaging. *Neurobiology of Disease* 32, 10809–10818.
- Lavisse, S., Inoue, K., Jan, C., Peyronneau, M.A., Petit, F., Goutal, S., ... Hantraye, P. (2015). [¹⁸F]DPA-714 PET imaging of translocator protein TSPO (18 kDa) in the normal and excitotoxically-lesioned nonhuman primate brain. *European Journal of Nuclear Medicine and Molecular Imaging* 42, 478–494.
- Lemus, M.B., Bayliss, J.A., Lockie, S.H., Santos, V.V., Reichenbach, A., Stark, R., & Andrews, Z. B. (2015). A stereological analysis of NPY, POMC, Orexin, GFAP astrocyte, and Iba1 microglial cell number and volume in diet-induced obese male mice. *Endocrinology* 156, 1701–1713.
- Levinson, I., & Levine, S. (1995). Negative correlation between Alpha-1-Acid-Glycoprotein plasma level and response to haloperidol in the acute treatment of schizophrenia. *Biological Psychiatry* 38, 198–200.
- Li, H., Sagar, A.P., & Keri, S. (2018). Translocator protein (18 kDa TSPO) binding, a marker of microglia, is reduced in major depression during cognitive-behavioral therapy. *Progress in Neuro-psychopharmacology Biology Psychiatry* 83, 1–7.
- Liu, B., Le, K.X., Park, M.A., Wang, S., Belanger, A.P., Dubey, S., ... Lemere, C.A. (2015). *In vivo* detection of age- and disease-related increases in neuroinflammation by ¹⁸F-GE180 TSPO microPET imaging in wildtype and Alzheimer's transgenic mice. *The Journal of Neuroscience* 35, 15716–15730.
- Lockhart, A., Davis, B., Matthews, J.C., Rahmoune, H., Hong, G., Gee, A., ... Brown, J. (2003). The peripheral benzodiazepine receptor ligand PK11195 binds with high affinity to the acute phase reactant alpha1-acid glycoprotein: implications for the use of the ligand as a CNS inflammatory marker. *Nuclear Medicine and Biology* 30, 199–206.
- Loth, M.K., Choi, J., McClothar, J.L., Pletnikov, M.V., Pomper, M.G., & Guilarte, T.R. (2016). TSPO in a murine model of Sandhoff disease: presymptomatic marker of neurodegeneration and disease pathophysiology. *Neurobiology of Disease* 85, 174–186.
- Maeda, J., Higuchi, M., Inaji, M., Ji, B., Haneda, E., Okauchi, T., ... Suhara, T. (2007). Phase-dependent roles of reactive microglia and astrocytes in nervous system injury as delineated by imaging of peripheral benzodiazepine receptor. *Brain Research* 1157, 100–111.
- Maeda, J., Zhang, M.-R., Okauchi, T., Ji, B., Ono, M., Hattori, S., ... Higuchi, M. (2011). *In vivo* positron emission tomography imaging of glial response to amyloid-beta and Tau pathologies in mouse models of Alzheimer's disease and related disorders. *The Journal of Neuroscience* 31, 4720–4730.
- Mahuran, D.F. (1999). Biochemical consequences of mutations causing the GM2 gangliosidosis. *Biochemical Biophysical Acta* 1455, 105–138.
- Mankowski, J.L., Queen, S.E., Tarwater, P.J., Adams, R.J., & Guilarte, T.R. (2003). Elevated peripheral benzodiazepine receptor expression in simian immunodeficiency virus encephalitis. *Journal of Neurovirology* 9, 94–100.
- Maragos, P.J., Patel, J., Boulenger, J.P., & Clark-Rosenberg, R. (1982). Characterization of peripheral-type benzodiazepine binding sites in brain using [³H]Ro5-4864. *Molecular Pharmacology* 22, 26–32.
- Masgrau, R., Guaza, C., Ransohoff, R.M., & Galea, E. (2017). Should we stop saying “glial” and “neuroinflammation”. *Trends in Molecular Medicine* 23, 485–500.
- Mattner, F., Linares Bandin, D., Staykova, M., Berghofer, P., Claude Gregoire, M., Ballantyne, P., ... Katsifis, A. (2011). Evaluation of [¹²³I]-CLINDE as a potent SPECT radiotracer to assess the degree of astroglia activation in cuprizone-induced neuroinflammation. *European Journal of Nuclear Medicine and Molecular Imaging* 38, 1516–1528.
- McNeela, A.M., Bernick, C., Hines, R.M., & Hines, D.J. (2017). TSPO regulation in reactive gliotic diseases. *Journal of Neuroscience Research* 96, 978–988.
- Miller, T.R., Wetter, J.B., Jarvis, M.F., & Bitner, R.S. (2013). Spinal microglia activation in rat models of neuropathic osteoarthritic pain: an autoradiographic study using [³H] PK11195. *European Journal of Pain* 17, 692–703.
- Mirzaei, N., Pham Tang, S., Ashworth, S., Coello, C., Plisson, C., Passchier, J., ... Sastre, M. (2016). *In vivo* imaging of microglia activation by positron emission tomography with [¹¹C]-PBR28 in the 5XFAD model of Alzheimer's disease. *GLIA* 64, 993–1006.
- Miyazawa, N., Diksic, M., & Yamamoto, Y. (1995). Chronological study of peripheral benzodiazepine binding sites in the rat stab wounds using [³H]PK11195 as a marker of gliosis. *Acta Neurochirurgica* 137, 207–216.
- Morohaku, K., Pelton, S.H., Daugherty, D.J., Butler, W.R., Deng, W., & Selvaraj, V. (2014). Translocator protein/peripheral benzodiazepine receptor is not required for steroid hormone biosynthesis. *Endocrinology* 155, 89–97.
- Myers, R., Manjil, L.G., Cullen, B.M., Price, G.W., Frackowiak, R.S., & Cremer, J.E. (1991). Macrophage and astrocyte populations in relation to [³H]PK11195 binding in the cerebral cortex following a local ischaemic lesion. *Journal of Cerebral Blood Flow and Metabolism* 11, 314–322.
- Nguyen, D.-L., Wimberley, C., Truillet, C., Jego, B., Caille, F., Pottier, G., ... Boullieret, V. (2018). Longitudinal positron emission tomography imaging of glial cell activation in a mouse model of mesial temporal lobe epilepsy: toward identification of optimal treatment windows. *Epilepsia* 59, 1234–1244.
- Notter, T., Coughlin, J.M., Gschwind, T., Weber-Stadlbauer, U., Wang, Y., Kassiou, M., ... Meyer, U. (2018). Translational evaluation of translocator protein as a marker of neuroinflammation in schizophrenia. *Molecular Psychiatry* 23, 323–334.
- Notter, T., Coughlin, J.M., Sawa, A., & Meyer, U. (2017). Reconceptualization of translocator protein as a biomarker of neuroinflammation in psychiatry. *Molecular Psychiatry* 23, 36–47.
- Ory, D., Planas, A., Dresselaers, T., Gsell, W., Postnov, A., Celen, S., ... Bormans, G. (2015). PET imaging of TSPO in a rat model of local neuroinflammation induced by intracerebral injection of lipopolysaccharide. *Nuclear Medicine and Biology* 42, 753–761.
- Papadopoulos, V., Baraldi, M., Guilarte, T.R., Knudsen, T.B., Lacapère, J.J., Lindemann, P., ... Gavish, M. (2006). Translocator protein (18kDa): new nomenclature for the peripheral-type benzodiazepine receptor based on its structure and molecular function. *Trends in Pharmacological Sciences* 27, 402–409.
- Pappata, S., Levasseur, M., Gunn, R.N., Myers, R., Crouzel, C., Syrota, A., ... Banati, R.B. (2000). Thalamic microglial activation in ischemic stroke detected *in vivo* by [¹¹C]-R-PK11195. *Neurology* 55, 1052–1054.

- Patel, J., & Marangos, P.J. (1982). Differential effects of GABA on peripheral and central type benzodiazepine binding sites in brain. *Neuroscience Letters* 30, 157–160.
- Plaven-Sigay, P., Matheson, G.J., Collste, K., Ashok, A.H., Coughlin, J.M., Howes, O.D., ... Cervenka, S. (2018). Positron Emission Tomography studies in the glial cell marker Translocator Protein in patients with psychosis: A meta-analysis using individual participant data. *Biological Psychiatry* S0006-3223, 31298–31308.
- Politis, M., Lahiri, N., Niccolini, F., Su, P., Wu, K., Giannetti, P., ... Piccini, P. (2015). Increased central microglial activation associated with peripheral cytokine levels in premanifest Huntington's disease gene carriers. *Neurobiology of Disease* 83, 115–121.
- Praet, J., Guglielmetti, C., Berneman, Z., Van der Linden, A., & Ponsaerts, P. (2014). Cellular and molecular neuropathology of the cuprizone mouse model: clinical relevance for multiple sclerosis. *Neuroscience and Biobehavioral Reviews* 47, 485–505.
- Rao, V.L.R., Dogan, A., Bowen, K.K., & Dempsey, R.J. (2000). Traumatic brain injury leads to increased expression of peripheral benzodiazepine receptors, neuronal death, and activation of astrocytes and microglia in rat thalamus. *Experimental Neurology* 161, 102–114.
- Richards, E.M., Zanotti-Fregonara, P., Fujita, M., Newman, L., Farmer, C., Ballard, E.D., ... Zarate, C.A., Jr. (2018). *EJNMMI Research* 8, 57.
- Ritsner, M., Modai, I., Gibel, A., Leschneir, S., Silver, H., Tsinovoy, G., ... Gavish, M. (2003). Decreased platelet peripheral-type benzodiazepine receptors in persistently violent schizophrenic patients. *Journal of Psychiatric Research* 37, 549–556.
- Roberts, R.C. (2017). Postmortem studies on mitochondria in schizophrenia. *Schizophrenia Research* 187, 17–25.
- Rojas, S., Martin, A., Arranz, M.J., Pareto, D., Purroy, J., Verdager, E., ... Planas, A.M. (2007). Imaging brain inflammation with [¹¹C]PK11195 by PET and induction of the peripheral-benzodiazepine receptor after transient focal ischemia in rats. *Journal of Cerebral Blood Flow and Metabolism* 27, 1975–1986.
- Sandiego, C.M., Gallezot, J.-D., Pittman, B., Nabulsi, N., Lim, K., Lin, S.-F., ... Cosgrove, K.P. (2015). Imaging robust microglia activation after lipopolysaccharide administration in humans with PET. *Proceedings of the National Academy Science* 112, 12468–12473.
- Sango, K., Yamanaka, S., Hoffmann, A., Okuda, Y., Grinberg, A., Westphal, H., ... Proia, R.L. (1995). Mouse models of Tay-Sachs and Sandhoff diseases differ in neurologic and phenotype and ganglioside metabolism. *Nature Genetics* 11, 170–176.
- Schain, M., & Kreisl, W.C. (2017). Neuroinflammation in neurodegenerative disorders—a review. *Current Neurology and Neuroscience Reports* 17, 25.
- Schoemaker, H., Morelli, M., Deshmukh, P., & Yamamura, H.I. (1982). [³H]Ro5-4864 benzodiazepine binding in the kainate lesioned striatum and Huntington's disease basal ganglia. *Brain Research* 248, 396–401.
- Selvaraj, V., & Stocco, D.M. (2015). The changing landscape in translocator protein (TSPO) function. *Trends in Endocrinology Metabolism* 26, 341–348.
- Selvaraj, V., & Tu, L.N. (2016). Current status and future perspectives: TSPO in steroid neuroendocrinology. *The Journal of Endocrinology* 231, R1–R30.
- Sérier, S., Tauber, C., Vercouille, J., Mothes, C., Pruckner, C., Guilloleau, D., ... Chalou, A. (2015). Amyloid load and translocator protein 18 kDa in APPswP51-dE9 mice: a longitudinal study. *Neurobiology of Aging* 36, 1639–1652.
- Setiawan, E., Attwells, S., Wilson, A.A., Mizhari, R., Rusjan, P.M., Miler, L., ... Meyer, J.H. (2018). Association of translocator protein total distribution volume with duration of untreated major depressive disorder: a cross-sectional study. *Lancet Psychiatry* 5, 339–347.
- Setiawan, E., Wilson, A.A., Mizhari, R., Rusjan, P.M., Miler, L., Rajkowska, G., ... Meyer, J.H. (2015). Role of translocator protein density, a marker of neuroinflammation, in the brain during major depressive episodes. *JAMA Psychiatry* 72, 268–275.
- Somerville, S.M., Conley, R.R., & Roberts, R.C. (2012). Striatal mitochondria in subjects with chronic undifferentiated vs. chronic paranoid schizophrenia. *Synapse* 66, 29–41.
- Starosta-Rubenstein, S., Ciliax, B.J., Penney, J.B., McKeever, P., & Young, A.B. (1987). Imaging of a glioma using peripheral benzodiazepine receptor ligands. *Proceedings of the National Academic Science* 84, 891–895.
- Stephenson, D.T., Schober, D.A., Smaistig, E.B., Mincy, R.E., Gehlert, D.R., & Clemens, J.A. (1995). Peripheral benzodiazepine receptors are colocalized with activated microglia following transient global forebrain ischemia in the rat. *The Journal of Neuroscience* 15, 5263–5274.
- Streit, W.J., Mrak, R.E., & Griffin, W.S.T. (2004). Microglia and neuroinflammation: a pathological perspective. *Journal of Neuroinflammation* 1, 1–4.
- Suranyi-Cadotte, B., Lafaille, F., Dongier, M., Dumas, M., & Quirion, R. (1988). Decreased density of peripheral benzodiazepine binding sites on platelets of currently drinking, but not abstinent alcoholics. *Neuropharmacology* 27, 443–445.
- Tai, Y.F., Pavese, N., Gerhardt, A., Tabrizi, S.J., Barker, R.A., Brooks, D.J., & Piccini, P. (2007). Imaging microglial activation in Huntington's disease. *Brain Research Bulletin* 72, 148–151.
- Telford, J.E., Bones, J., McManus, C., Saldova, C., Manning, G., Doherty, M., ... Rudd, P.M. (2012). Antipsychotic treatment of acute paranoid schizophrenia patients with olanzapine results in altered glycosylation of serum glycoproteins. *Journal of Proteome Research* 11, 3743–3752.
- Tomasi, G., Edison, P., Bertoldo, A., Roncaroli, F., Singh, P., Gerhard, A., ... Turkheimer, F.E. (2008). Novel reference region model reveals increased microglial and reduced vascular binding of [¹¹C]-R-PK11195 in patients with Alzheimer's disease. *Journal of Nuclear Medicine* 49, 1249–1256.
- Toyama, H., Hatano, K., Suzuki, H., Ichise, M., Momosaki, S., Kudo, G., ... Ito, K. (2008). In vivo imaging of microglial activation using a peripheral benzodiazepine receptor ligand: [¹¹C]-PK11195 and animal PET following ethanol injury in rat striatum. *Annals of Nuclear Medicine* 22, 417–424.
- Tu, L.N., Morohaku, K., Manna, P.R., Pelton, S.H., Butler, W.R., Stocco, D.M., & Selvaraj, V. (2014). Peripheral benzodiazepine receptor/translocator protein global knock-out mice are viable with no effects on steroid hormone biosynthesis. *The Journal of Biological Chemistry* 289, 27444–27454.
- Tu, L.N., Zhao, A.H., Stocco, D.M., & Selvaraj, V. (2015). PK11195 effect on steroidogenesis is not mediated through the translocator protein (TSPO). *Endocrinology* 156, 1033–1039.
- Turkheimer, F.E., Rizzo, G., Bloomfield, P.S., Howes, O., Zanotti-Fregonara, P., Bertoldo, A., & Veronese, M. (2015). The methodology of TSPO imaging with positron emission tomography. *Biochemical Society Transactions* 43, 586–592.
- Turner, M.R., Cagnin, A., Turkheimer, F.E., Miller, C.C., Shaw, C.E., Brooks, D.J., ... Banati, R.B. (2004). Evidence of widespread cerebral microglial activation in amyotrophic lateral sclerosis: an [¹¹C]-R-PK11195 positron emission tomography study. *Neurobiology of Disease* 15, 601–609.
- Van Kesteren, C.F.M.G., Gemmels, H., Witte, L.D., Hoi, E.M., van Gool, A.R., Flakaj, P.G., ... Sommer, I.E.C. (2017). Immune involvement in the pathogenesis of schizophrenia: a meta-analysis on postmortem brain studies. *Translational Psychiatry* 7, e1075.
- Venneti, S., Lopresti, B.J., Wang, G., Bissel, S.J., Mathis, C.A., Meltzer, C.C., ... Wiley, C.A. (2004). PET imaging of brain macrophages using the peripheral benzodiazepine receptor in a macaque model of neuroAIDS. *Journal of Clinical Investigation* 113, 981–989.
- Venneti, S., Lopresti, B.J., & Wiley, C.A. (2006). The peripheral benzodiazepine receptor (Translocator Protein 18 kDa) in microglia: from pathology to imaging. *Progress in Neurobiology* 80, 308–322.
- Venneti, S., Wagner, A.K., Wang, G., Slagel, S.L., Chen, X., Lopresti, B.J., ... Wiley, C.A. (2007). The high affinity peripheral benzodiazepine receptor ligand DAA1106 binds specifically to microglia in a rat model of traumatic brain injury: implications for PET imaging. *Experimental Neurology* 207, 118–127.
- Venneti, S., Wang, G., Nguyen, J., & Wiley, C.A. (2008). The positron emission tomography ligand DAA1106 binds with high affinity to activated microglia in human neurological disorders. *Journal of Neuropathology and Experimental Neurology* 67, 1001–1010.
- Venneti, S.M., Lopresti, B.J., Wang, G., Hamilton, R.L., Mathis, C.A., Klunk, W.E., ... Wiley, C.A. (2009). PK11195 labels activated microglia in Alzheimer's disease and in vivo in a mouse model using PET. *Neurobiology of Aging* 30, 1217–1226.
- Veronese, M., Marques, T.R., Bloomfield, P.S., Rizzo, G., Singh, N., Jones, D., ... Turkheimer, F.E. (2017). Kinetic modelling of [¹¹C]PBR28 for translocator protein PET data: A validation study of vascular modelling in the brain using XBD173 and tissue analysis. *Journal of Cerebral Blood Flow Metabolism* 38, 1227–1242.
- Vowinckel, E., Reutens, D., Becher, B., Verge, G., Evans, A., Owens, T., & Antel, J.P. (1997). PK11195 binding to the peripheral benzodiazepine receptor as a marker of microglia activation in multiple sclerosis and experimental autoimmune encephalomyelitis. *Journal of Neuroscience Research* 50, 345–353.
- Wang, Y., Yue, X., Kiesewetter, D.O., Niu, G., Teng, G., & Chen, X. (2014). PET imaging of neuroinflammation in a rat traumatic brain injury model with radiolabeled TSPO ligand DPA-714. *European Journal of Nuclear Medicine and Molecular Imaging* 41, 1440–1449.
- Weizman, R., Tanne, Z., Karp, L., & Tyano, M. (1986). Peripheral-type benzodiazepine-binding sites in platelets of schizophrenics with and without tardive dyskinesia. *Life Science* 39, 549–555.
- Wierzbicka-Bobrowicz, T., Lewandowska, E., Kosno-Knusewska, E., Lechowicz, W., Pasennik, E., & Schmidt-Sidor, B. (2004). Degeneration of microglial cells in frontal cortex and temporal lobes of chronic schizophrenics. *Folia Neuropathologica* 42, 157–165.
- Williams, R.M., Hampton, T., Pearce, R.K.B., Hirsch, S.R., Ansorge, O., Thom, M., & Maier, M. (2013). Astrocyte decrease in the subgenual cingulate and callosal genu in schizophrenia. *European Archives of Psychiatry and Clinical Neuroscience* 263, 41–52.
- Wodarz, N., Rothenhofer, C., Fischer, R., Stober, G., Kiehl, B., Jungkunz, G., ... Klein, H.E. (1998). Peripheral-type benzodiazepine receptors in diagnostic subtypes of schizophrenia patients. *Psychiatric Research* 81, 363–369.
- Yankam Njiwa, J., Costes, N., Bouillot, C., Bouvard, S., Fioux, S., Becker, G., ... Hammers, A. (2016). Quantitative longitudinal imaging of activated microglia as a marker of inflammation in the pilocarpine rat model of epilepsy using [¹¹C]-R-PK11195 PET and MRI. *Journal of Cerebral Blood Flow and Metabolism* 37, 1251–1263.
- Young, E.C., Patel, A., Meyers, B.S., Kakuma, T., & Alexopoulos, G.S. (1999). Alpha1-acid glycoprotein, age, and sex in mood disorders. *American Journal of Geriatric Psychiatry* 7, 331–334.