

---

# JOURNAL OF CLINICAL EPIDEMIOLOGY

---

VOLUME 113, SEPTEMBER 2019

## TABLE OF CONTENTS

### *Editorial*

---

- vi Trial recruitment a continuing challenge  
*Peter Tugwell and Andre Knottnerus*

### *Commentary*

---

- 83 Coherence in measurement and programming in maternal and newborn health: experience from the BetterBirth trial  
*Natalie Panariello, Amanda Jurczak, Jonathan Spector, Vishwajeet Kumar, and Katherine Semrau*

### *Reviews*

---

- 1 Characteristics and methods of incorporating randomized and nonrandomized evidence in network meta-analyses: a scoping review  
*Kathryn Zhang, Paul Arora, Neba Sati, Audrey Béliveau, Natalie Troke, Areti Angeliki Veroniki, Myanca Rodrigues, Patricia Rios, Wasifa Zarin, and Andrea C. Tricco*
- 44 Minimal clinically important difference of commonly used hip-, knee-, foot-, and ankle-specific questionnaires: a systematic review  
*Derya Çelik, Özge Çoban, and Önder Kılıçoğlu*
- 64 Prevalence of trial registration varies by study characteristics and risk of bias  
*Aidan Christopher Tan, Ivy Jiang, Lisa Askie, Kylie Hunter, Robert John Simes, and Anna Lene Seidler*
- 75 Limited evidence exists on the effectiveness of education and training interventions on trial recruitment; a systematic review  
*Hannah Delaney, Declan Devane, Andrew Hunter, Marita Hennessy, Advoa Parker, Louise Murphy, Patricia Cronin, and Valerie Smith*
- 168 Consolidated Standards of Reporting Trials (CONSORT) extensions covered most types of randomized controlled trials, but the potential workload for authors was high  
*Lina Ghosn, Isabelle Boutron, and Philippe Ravaud*
- 189 Overall bias and sample sizes were unchanged in ICU trials over time: a meta-epidemiological study  
*Carl Thomas Anthon, Anders Granholm, Anders Perner, Jon Henrik Laake, and Morten Hylander Møller*

*Continued on following page*



0895-4356(201909)113:C;1-N

---

# JOURNAL OF CLINICAL EPIDEMIOLOGY

---

VOLUME 113, SEPTEMBER 2019

## TABLE OF CONTENTS *continued*

### *Original Articles*

---

- 11 Harms are assessed inconsistently and reported inadequately Part 2: nonsystematic adverse events  
*Evan Mayo-Wilson, Nicole Fusco, Tianjing Li, Hwanbee Hong, Joseph K. Canner, and Kay Dickersin, for the MUDS investigators*
- 20 Harms are assessed inconsistently and reported inadequately part 1: systematic adverse events  
*Evan Mayo-Wilson, Nicole Fusco, Tianjing Li, Hwanbee Hong, Joseph K. Canner, and Kay Dickersin, for the MUDS investigators*
- 28 Funders improved the management of learning and clustering effects through design and analysis of randomized trials involving surgery  
*Elizabeth J. Conroy, Anna Rosala-Hallas, Jane M. Blazeby, Girvan Burnside, Jonathan A. Cook, and Carrol Gamble*
- 36 Overview: comprehensive and carefully constructed strategies are required when conducting searches for adverse effects data  
*Su Golder, Guy Peryer, and Yoon K. Loke*
- 58 Engaging knowledge users in a systematic review on the comparative effectiveness of geriatrician-led models of care is possible: A cross-sectional survey using the Patient Engagement Evaluation Tool  
*Charlene Soobiah, Sharon E. Straus, Gayle Manley, Sharon Marr, Elliot Pausjenssen, Sylvia Teare, Jemila Hamid, Andrea C. Tricco, and Ainsley Moore*
- 86 A cross-sectional audit showed that most Cochrane intervention reviews searched trial registers  
*Slavica Berber, Ava Grace Tan-Koay, Newton Opiyo, Kerry Dwan, Julie M. Glanville, Toby J. Lasserson, and Melina L. Willson*
- 92 A benefit-harm analysis of adding basal insulin vs. sulfonylurea to metformin to manage type II diabetes mellitus in people with multiple chronic conditions  
*Wendy L. Bennett, Hélène E. Aschmann, Milo A. Puban, Craig W. Robbins, Elizabeth A. Bayliss, Renee Wilson, Richard A. Mularski, Wiley V. Chan, Bruce Leff, Orla Sheehan, Carol Glover, Katie Maslow, Karen Armacost, Suzanne Mintz, and Cynthia M. Boyd*
- 104 Risk of bias assessments for blinding of participants and personnel in Cochrane reviews were frequently inadequate  
*Ognjen Barcot, Matija Boric, Sjetlana Dosenovic, Tina Poklepovic Pericic, Marija Cavar, and Livia Puljak*
- 114 Applying sequential surveillance methods that use regression adjustment or weighting to control confounding in a multisite, rare-event, distributed setting: Part 2 in-depth example of a reanalysis of the measles-mumps-rubella-varicella combination vaccine and seizure risk  
*Andrea J. Cook, Robert D. Wellman, Tracey Marsh, Azadeh Shoaibi, Ram Tiwari, Michael Nguyen, Denise Boudreau, Eric S. Weintraub, Lisa Jackson, and Jennifer C. Nelson*

*Continued on following page*

---

# JOURNAL OF CLINICAL EPIDEMIOLOGY

---

VOLUME 113, SEPTEMBER 2019

## TABLE OF CONTENTS *continued*

- 123 Effect measure modification conceptualized using selection diagrams as mediation by mechanisms of varying population-level relevance  
*Priscilla M. Lopez, S.V. Subramanian, and C. Mary Schooling*
- 129 The Clinician Guideline Determinants Questionnaire was developed and validated to support tailored implementation planning  
*Anna R. Gagliardi, Melissa J. Armstrong, Susanne Bernbardsson, Margot Fleuren, Hector Pardo-Hernandez, Robin W.M. Vernooij, and Melina Willson, Guidelines International Network Implementation Working Group*
- 137 The Trials within Cohorts design faced methodological advantages and disadvantages in the exercise oncology setting  
*Roxanne Gal, Evelyn M. Monninkhof, Carla H. van Gils, Rolf H.H. Groenwold, Desirée H.J.G. van den Bongard, Petra H.M. Peeters, Helena M. Verkooijen, and Anne M. May*
- 147 Several reasons explained the variation in the results of 22 meta-analyses addressing the same question  
*Assem M. Khamis, Mobamad El Mobebe, Jobny Nicolas, Gbida Iskandarani, Marwan M. Refaat, and Elie A. Akl*
- 159 A systematic survey identified 36 criteria for assessing effect modification claims in randomized trials or meta-analyses  
*Stefan Schandelmaier, Yaping Chang, Niveditha Devasenapathy, Tabira Deoji, Joey S.W. Kwong, Luis E. Colunga Lozano, Yung Lee, Arnav Agarwal, Neera Bhatnagar, Hannab Ewald, Ying Zhang, Xin Sun, Lehana Thabane, Michael Walsh, Matthias Briel, and Gordon H. Guyatt*
- 176 Retrieval of individual patient data depended on study characteristics: a randomized controlled trial  
*Areti Angeliki Veroniki, Huda M. Asboor, Susan P.C. Le, Patricia Rios, Lesley A. Stewart, Mike Clarke, Dimitris Mavridis, Sharon E. Straus, and Andrea C. Tricco*
- 200 Factors associated with trial recruitment, preferences, and treatments received were elucidated in a comprehensive cohort study  
*Jenny L. Donovan, Brent Opmeer, Grace J. Young, Nicola Mills, Richard M. Martin, J. Athene Lane, Chris Metcalfe, Tim J. Peters, Michael Davis, Emma L. Turner, Eleanor Walsh, David E. Neal, and Freddie C. Hamdy, ProtecT Study Group*
- 214 Improvement needed in the network geometry and inconsistency of Cochrane network meta-analyses: a cross-sectional survey  
*Ya Gao, Long Ge, Xueni Ma, Xiping Shen, Meng Liu, and Jinbui Tian*

---

## **Letters to the Editor**

- 101 Premarket safety population size associated with approval of expedited program drugs and orphan drugs  
*Kachi Illob*