

STATE-OF-THE-ART REVIEW

Neuromodulation for the Treatment of Heart Rhythm Disorders



Nathan H. Waldron, MD, MHS,^{a,b,*} Marat Fudim, MD,^{b,c,*} Joseph P. Mathew, MD, MHS, MBA,^{a,b}
Jonathan P. Piccini, MD, MHS^{b,c}

JACC: BASIC TO TRANSLATIONAL SCIENCE CME/MOC/ECME

This article has been selected as this month's *JACC: Basic to Translational Science* CME/MOC/ECME activity, available online at <http://www.acc.org/jacc-journals-cme> by selecting the *JACC* Journals CME/MOC/ECME tab.

Accreditation and Designation Statement

The American College of Cardiology Foundation (ACCF) is accredited by the Accreditation Council for Continuing Medical Education (ACCME) and the European Board for Accreditation in Cardiology (EBAC) to provide continuing medical education for physicians.

The ACCF designates this Journal-based CME/MOC/ECME activity for a maximum of 1 AMAPRA Category 1 Credit or 1 EBAC Credit. Physicians should only claim credit commensurate with the extent of their participation in the activity.

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1 Medical Knowledge MOC point in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. Participants will earn MOC points equivalent to the amount of CME credits claimed for the activity. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

Neuromodulation for the Treatment of Heart Rhythm Disorders will be accredited by the European Board for Accreditation in Cardiology (EBAC) for 1 hour of External CME credits. Each participant should claim only those hours of credit that have actually been spent in the educational activity. The Accreditation Council for Continuing Medical Education (ACCME) and the European Board for Accreditation in Cardiology (EBAC) have recognized each other's accreditation systems as substantially equivalent. Apply for credit through the post-course evaluation.

Method of Participation and Receipt of CME/MOC/ECME Certificate

To obtain credit for JACBTS: Basic to Translational Science CME/MOC/ECME, you must:

1. Be an ACC member or JACBTS subscriber.
2. Carefully read the CME/MOC/ECME-designated article available online and in this issue of the *journal*.
3. Answer the post-test questions. At least 2 questions provided must be answered correctly to obtain credit.
4. Complete a brief evaluation.

5. Claim your CME/MOC/ECME credit and receive your certificate electronically by following the instructions given at the conclusion of the activity.

CME/MOC/ECME Objective for This Article: Upon completion of this activity, the learner should be able to: 1) describe the neurohormonal pathways that affect cardiac arrhythmias; 2) identify therapeutic options for refractory ventricular tachycardia; and 3) discuss the potential mechanisms for neurohormonal modulation.

CME/MOC/ECME Editor Disclosure: CME/MOC/ECME Editor L. Kristin Newby, MD, is supported by research grants from Amylin, Bristol-Myers Squibb Company, GlaxoSmithKline, Sanofi, Verily Life Sciences (formerly Google Life Sciences), the MURDOCK Study, NIH, and PCORI; receives consultant fees/honoraria from BioKier, DemeRx, MedScape/TheHeart.org, Metanomics, Philips Healthcare, Roche Diagnostics, CMAC Health Education & Research Institute; serves as an Officer, Director, Trustee, or other fiduciary role for the AstraZeneca HealthCare Foundation and the Society of Chest Pain Centers (now part of ACC); and serves in another role for the American Heart Association and is the Deputy Editor of *JACC: Basic to Translational Science*.

Author Disclosures: Dr. Waldron has received personal fees and grants from Allergan, and AHA grant 16MCPRP30700010 and the NIGMS T32 postdoctoral training grant T32GM008600. Dr. Fudim has received grants from AHA grant 17MCPRP33460225 and NIH T32 grant 5T32HL007101; and has received personal fees from Galvani, Coridea, and Axon Therapies. Dr. Mathew is on the Board of Directors and owns equity in MedBlue Data; and has received grants from NIH R01 grant number HL130443. Dr. Piccini has received grants for clinical research from Abbott, the American Heart Association, Boston Scientific, Gilead, Janssen Pharmaceuticals, and the NHLBI; and serves as a consultant to Abbott, Allergan, ARCA Biopharma, Biotronik, Boston Scientific, Johnson & Johnson, LivaNova, Medtronic, Milestone, Oliver Wyman Health, Sanofi, Philips, and Up-to-Date.

Medium of Participation: Online (article and quiz).

CME/MOC/ECME Term of Approval

Issue Date: August 2019

Expiration Date: July 31, 2020

From the ^aDepartment of Anesthesia, Duke University Medical Center, Durham, North Carolina; ^bDuke Clinical Research Institute, Durham, North Carolina; and the ^cElectrophysiology Section, Duke University Medical Center, Durham, North Carolina. *Drs. Waldron and Fudim contributed equally to this work and are joint first authors. Dr. Waldron has received personal fees and grants from Allergan, and AHA grant 16MCPRP30700010 and the NIGMS T32 postdoctoral training grant T32GM008600. Dr. Fudim has received grants from AHA grant 17MCPRP33460225 and NIH T32 grant 5T32HL007101; and has received personal fees from

Neuromodulation for the Treatment of Heart Rhythm Disorders

Nathan H. Waldron, MD, MHS,^{a,b,*} Marat Fudim, MD,^{b,c,*} Joseph P. Mathew, MD, MHS, MBA,^{a,b}
Jonathan P. Piccini, MD, MHS^{b,c}

HIGHLIGHTS

- Derangement of autonomic nervous signaling is an important contributor to cardiac arrhythmogenesis.
- Modulation of autonomic nervous signaling holds significant promise for the prevention and treatment of cardiac arrhythmias.
- Further clinical investigation is necessary to establish the efficacy and safety of autonomic modulatory therapies in reducing cardiac arrhythmias.

SUMMARY

There is an increasing recognition of the importance of interactions between the heart and the autonomic nervous system in the pathophysiology of arrhythmias. These interactions play a role in both the initiation and maintenance of arrhythmias and are important in both atrial and ventricular arrhythmia. Given the importance of the autonomic nervous system in the pathophysiology of arrhythmias, there has been notable effort in the field to improve existing therapies and pioneer additional interventions directed at cardiac-autonomic targets. The interventions are targeted to multiple and different anatomic targets across the neurocardiac axis. The purpose of this review is to provide an overview of the rationale for neuromodulation in the treatment of arrhythmias and to review the specific treatments under evaluation and development for the treatment of both atrial fibrillation and ventricular arrhythmias. (J Am Coll Cardiol Basic Trans Science 2019;4:546-62) © 2019 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ANATOMY OF THE NEUROCARDIAC AXIS

The neurocardiac axis is composed of multiple intricate areas that provide an interface between the nervous system and the cardiovascular system (Figure 1). This neurocardiac anatomy has been elegantly detailed in a recent review (1). A number of diverse brain structures contribute to efferent cardiac autonomic signaling, including the amygdala, insula, thalamus, and cerebellum (2). In general, second-to-second control of autonomic tone occurs at the level of the brainstem, whereas forebrain structures including the hypothalamus effect more prolonged or gradual shifts in autonomic tone (3). From these

various brain areas, sympathetic signals are first transmitted via the cervical and thoracic spinal cord to the stellate (cervicothoracic) ganglion before synapsing with nerves of the intrinsic cardiac nervous system (4). In contrast, parasympathetic fibers originate in the brainstem and travel primarily via the vagus nerve into the pericardial sack (1). There, sympathetic and parasympathetic efferent fibers form a branching neural network clustering in autonomic ganglionic plexi (AGP) contained in fat pads on the posterior surface of the atria and the superior aspect of the ventricles (Figure 1) (5). AGP contain large populations of colocalized sympathetic and parasympathetic neurons (1), and may serve as

Galvani, Coridea, and Axon Therapies. Dr. Mathew is on the Board of Directors and owns equity in MedBlue Data; and has received grants from NIH R01 grant number HL130443. Dr. Piccini has received grants for clinical research from Abbott, the American Heart Association, Boston Scientific, Gilead, Janssen Pharmaceuticals, and the NHLBI; and serves as a consultant to Abbott, Allergan, ARCA Biopharma, Biotronik, Boston Scientific, Johnson & Johnson, LivaNova, Medtronic, Milestone, Oliver Wyman Health, Sanofi, Philips, and Up-to-Date.

All authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the JACC: Basic to Translational Science [author instructions page](#).

Manuscript received December 31, 2018; revised manuscript received February 22, 2019, accepted February 22, 2019.

ABBREVIATIONS AND ACRONYMS

AERP	= atrial effective refractory period
AF	= atrial fibrillation
AGP	= autonomic ganglionic plexus
ANS	= autonomic nervous system
CABG	= coronary artery bypass grafting
HRV	= heart rate variability
ICD	= implantable cardioverter-defibrillator
LLVNS	= low-level vagal nerve stimulation
OSA	= obstructive sleep apnea
POAF	= post-operative atrial fibrillation
PVI	= pulmonary vein isolation
RDN	= renal denervation
SCS	= spinal cord stimulation
SGB	= stellate ganglion blockade
SNS	= sympathetic nervous system
VF	= ventricular fibrillation
VNS	= vagal nerve stimulation
VT	= ventricular tachycardia

“integration centers” modulating cardiac responses to autonomic input (6). Emerging evidence indicates that these AGP contain an amalgam of afferent neurons, motor neurons, interconnecting local circuit neurons, and that some proportion of these neurons expresses both cholinergic as well as adrenergic immunoreactivity (1). The extensive interconnections as well as the colocalization of cholinergic and adrenergic neurons set the stage for interplay between the parasympathetic and sympathetic nervous system at the level of the AGPs. Beyond the direct communication between the autonomic nervous system (ANS) and the heart, the neurocardiac axis is under modulatory control from additional autonomic reflexes, including afferent nervous input from the baroreflex and the renal nerves, which may modulate the global and regional (cardiac) sympathetic tone.

ATRIAL FIBRILLATION

RATIONALE FOR AUTONOMIC MODULATION FOR ATRIAL FIBRILLATION. Atrial fibrillation (AF) is a common and significant medical problem that affects approximately 3% of the population (7). As the population ages, the incidence and prevalence of AF are projected


to increase even further (8). In addition to being a leading cause of stroke, AF is associated with increased risks of heart failure (HF), myocardial infarction (MI), cognitive impairment, and death (9). The severity or persistence of AF has been associated with worse outcomes, including a higher risk of stroke (10); however, the role of AF burden (proportion of time spent in AF) in adverse outcomes remains incompletely understood (11). Given both its prevalence and impact, AF is a significant public health concern, and reduction of AF and its attendant complications remains a priority for the cardiovascular community.

An increasing body of evidence suggests that activity of the ANS can contribute to the initiation and perpetuation of cardiac arrhythmias, including AF. In isolated atrial myocytes, parasympathetic input shortens the atrial effective refractory period (AERP) (12) and increases dispersion of refractoriness (13), whereas sympathetic input increases calcium transient currents and afterdepolarizations (4), synergistically promoting the onset of AF (14). Similar responses to sympathetic stimulation are observed in pulmonary veins, a common site of ectopy that

triggers AF (15). Although both parasympathetic and sympathetic stimulation may individually promote AF (16,17), data suggest that combined parasympathetic and sympathetic (sympathovagal) discharges are especially potent initiators of AF (15,18) due to the combined effects of shortened AERP and increased calcium transient currents, respectively. Changes in heart rate variability (HRV) before onset of AF (19,20) are consistent with combined sympathovagal discharges. There are also data to suggest that AGP may influence focal sources or drivers of AF. For example, a study of 97 patients undergoing catheter ablation of AF has shown that the majority of patients had a focal source or rotor that overlapped with the anatomic location of an AGP (21). Because AGPs play a crucial role in AF, several strategies to prevent or reduce AF have centered around modulating AGP activity directly or indirectly through the extrinsic cardiac nervous system.

PAST AND PRESENT. Current strategies in autonomic modulation for atrial arrhythmias. A number of strategies to reduce AF partially exert their effects through modulation of the cardiac ANS (Table 1). Beta blockers antagonize the effects of catecholamines on a variety of target tissues, such as cardiac and vascular adrenergic receptors (22). AF ablation may also exert some benefit through autonomic modulation. In addition to creating electrical isolation of the pulmonary veins from the left atrium (23), pulmonary vein isolation (PVI) also reduces AGP-induced firing of pulmonary vein potentials that initiate AF. A positive response to AGP stimulation after extensive PVI independently predicts recurrence of AF (24), suggesting that AGP modulation may contribute to the efficacy of AF ablation.

Aggressive risk factor reduction also reduces the risk of AF. Part of the benefit of weight loss may be explained by alteration of autonomic tone as the metabolic syndrome has been associated with increased sympathetic tone (25). Similarly, obstructive sleep apnea (OSA) results in sympathetic activation (26) and increased focal sources of AF (27) both contributing to an increased risk of AF. Management of these risk factors reduces ANS hyperactivity and may impact the success of AF ablation (28). Treatment of sleep apnea with continuous positive airway pressure decreases risk of AF progression (29), and reduces risk of recurrent AF after catheter ablation (30). Adaptive servo-ventilation, which may be used to treat obstructive, central, or mixed sleep apnea, was recently shown to decrease AF burden in patients with HF and sleep apnea (31). Risk factor reduction such as weight loss and treatment of sleep apnea has the




potential to improve autonomic function and reduce AF through several different mechanisms.

DEVELOPING NEUROMODULATORY APPROACHES.

Ablation-based approaches. AGP ablation. Ablation of AGP by either catheter-based or surgical approaches has shown some promise in improving the success of AF ablation (Central Illustration). Percutaneous anatomically guided AGP ablation as a stand-alone technique has limited efficacy (32) and is inferior to circumferential PVI (33). However, randomized data have shown that the addition of AGP ablation to PVI may increase freedom from AF compared to either technique in isolation (34). Additionally, AGP ablation combined with PVI may result in increased long-term freedom from AF and lower rates of atrial flutter than PVI + linear ablation (35). Not all studies have supported the efficacy of AGP ablation. A recent comparison of video-assisted thoracoscopic surgery for AF with or without AGP ablation showed no difference in freedom from AF; additionally, patients undergoing epicardial AGP ablation had

higher rates of procedural complications and pacemaker implantation (36). Further studies are needed to clarify optimal methods of AGP localization and ablation, and the potential benefits of AGP ablation.

TABLE 1 Current Therapeutic Strategies for Rhythm Control of AF	
Current Therapeutic Strategies for AF	Examples
Antiarrhythmic drug therapy	Class I: flecainide, propafenone Class III: amiodarone, dronedarone, sotalolol, dofetilide
Catheter ablation	Pulmonary vein isolation Substrate modification Ablation of nonpulmonary vein triggers
Surgical ablation	Cox-Maze procedure Thoracoscopic ablation ("mini-maze")
Risk-factor modification	Positive airway pressure for obstructive sleep apnea Blood pressure reduction Weight Loss
AF = atrial fibrillation.	



Renal denervation. Renal denervation (RDN) is a method of modulating central afferent input. Renal afferent and efferent nerves function in a reflex loop where afferent input from the kidney to the central nervous system (CNS) regulates the efferent sympathetic nerve output to the heart and back to the kidney. Renal afferent nerves are diverse and represent a heterogeneous population of fibers including myelinated and nonmyelinated fibers (37,38). Two main types of renal afferents are responsible for

relaying information to the CNS, mechanosensitive (hydrostatic pressure from renal vasculature), and chemosensitive receptors (sensitive to ischemia, osmolar changes, and ionic composition) (39). Functionally, the renal afferent fibers can be divided into: 1) pressor; 2) reno-renal; and 3) depressor types (40,41). In other words, some fibers when activated contribute to the elevation of the sympathetic tone (sympatho-stimulant), whereas others reduce the sympathetic tone (sympatholytic). This is supported

TABLE 2 Current Status of Developing Neuromodulatory Strategies for Arrhythmias

Intervention	Anatomical Target	Current Status	Diseases of Interest	Number of Patients	Ongoing Clinical Trials
AGP ablation	Autonomic Ganglionic Plexi	Clinical trials	AF	60	NCT03535818 : Adjunctive Ganglionated Plexus Ablation in Redo-Pulmonary Vein Isolation (ADD-GP); Randomized trial testing the efficacy of AGP ablation in patients with ongoing paroxysmal arrhythmias after PVI
			AF	14	NCT03636100 : Release of Acetylcholine From the Ganglionated Plexus During the Thaw Phase of Cryoballoon Pulmonary Vein Ablation (GP RESPONSE Study); Observational study designed to study the role of systemic acetylcholine release in response to cryoballoon ablation.
			POAF	62	NCT02035163 : Atrial Fibrillation Prevention in Post Coronary Artery Bypass Graft Surgery with Cryoablation for Ganglionic Plexi; A randomized study to test the ability of AGP cryoablation during cardiac surgery to prevent POAF.
Epicardial botulinum toxin injection	Autonomic ganglionic plexi	Clinical trials	POAF	170	NCT02617069 : Botulinum Toxin Injection Into Epicardial Fat Pads to Prevent Atrial Fibrillation in Patients Undergoing Cardiac Surgery; Randomized trial of epicardial botulinum toxin for the prevention of POAF after cardiac surgery in patients with paroxysmal AF.
			POAF	330	NCT03779841 : A Phase II, Multi-Center, Randomized, Double-Blind, Placebo-Controlled, Dose-Ranging Study to Evaluate the Efficacy and Safety of Botulinum Toxin Type A (AGN-151607) Injections Into the Epicardial Fat Pads to Prevent Post-Operative Atrial Fibrillation in Patients Undergoing Open-Chest Cardiac Surgery; Randomized double-blind, placebo-controlled, dose-ranging trial evaluating the efficacy and safety of epicardial botulinum toxin type injections to prevent POAF in patients undergoing cardiac surgery.
BRS therapy	Carotid and aortic baroreceptors	Preclinical	POAF	95	NCT03243279 : BRS and Outcomes in Cardiothoracic Surgery; Observational study to determine if perioperative baroreceptor sensitivity (BRS) correlates with POAF and outcomes after cardiothoracic surgery
TVNS	Auricular branch of the Vagus Nerve (Afferent)	Clinical trials	POAF	42	NCT03533140 : Postoperative Atrial Fibrillation Suppression By Nerve Stimulation; Randomized trial testing TVNS to prevent POAF and reduce inflammation
			POAF	200	NCT02783157 : Transcutaneous Autonomic Modulation In Thoracic Surgery (TON-POINTS); Trial evaluating TVNS to reduce POAF after cardiothoracic surgery.
			AF	52	NCT02548754 : Transcutaneous Electrical Vagus Nerve Stimulation to Suppress Atrial Fibrillation (TREAT-AF); A small study to evaluate the ability of TVNS to reduce atrial fibrillation burden and inflammation in patients with paroxysmal AF.
RDN	Sympathetic nerves around the renal arteries (afferent)	Clinical trials	AF	300	NCT01635998 : Adjunctive Renal Denervation in the Treatment of Atrial Fibrillation (H-FIB); Phase II trial to evaluate the ability of combined RDN and PVI versus PVI only to reduce recurrent AF.
			AF	100	NCT01990911 : Renal Sympathetic Denervation Prevents Atrial Fibrillation in Patients with Hypertensive Heart Disease: a Pilot Study (RDPAF); Phase II trial testing efficacy of RDN to reduce subclinical AF and restore autonomic imbalance.
			AF	138	NCT02115100 : Treatment of Atrial Fibrillation in Patients by Pulmonary Vein Isolation in Combination With Renal Denervation or Pulmonary Vein Isolation Only (ASAF); Open-label trial testing the ability of combined RDN and PVI to reduce recurrent AF after PVI.
			AF	61	NCT01907828 : A Feasibility Study to Evaluate the Effect of Concomitant Renal Denervation and Cardiac Ablation on AF Recurrence (RDN+AF); Phase II trial evaluating the feasibility of RDN combined with AF ablation in reducing recurrent AF in patients with AF and uncontrolled hypertension.
			AF	40	NCT03246568 : Renal Nerve Denervation After Pulmonary Vein Isolation for Persistent Atrial Fibrillation; Phase II study to evaluate the efficacy of RDN added to PVI in reducing recurrent AF in patients with persistent AF.
			AF	100	NCT01686542 : CPVI Plus Renal Sympathetic Modification Versus CPVI Alone for AF (Atrial Fibrillation) Ablation; Open-label trial evaluating the ability of RDN combined with PVI to reduce recurrent AF compared with PVI alone.
			AF	245	NCT02064764 : Renal Nerve Denervation in Patients with Hypertension and Paroxysmal and Persistent Atrial Fibrillation (Symplicity AF); Phase II trial to assess the feasibility of concurrent RDN and PVI, and the rate of chronic treatment success in patients receiving RDN and PVI vs. PVI only
			AF	40	NCT01952925 : Combined Afib Ablation and RA Denervation for the Maintenance of Sinus Rhythm and Management of Resistant Hypertension; Non-randomized prospective study to evaluate the ability of concurrent AF ablation and RDN to reduce ambulatory blood pressure.
			VT/VF	60	NCT02856373 : Renal Nerve Stimulation and Renal Denervation in Patients With Sympathetic Ventricular Arrhythmias (Redress VT); Will assess impact of renal nerve stimulation before and after percutaneous transluminal RDN on cardiac excitable properties including induction of ventricular tachyarrhythmias before and after RDN
			VT/VF	38	NCT01858194 : REal Sympathetic dEneRvaTion as an Adjunct to Catheter-based VT Ablation (RESET-VT); Prospective, multicenter, randomized control trial testing the impact of catheter-based renal sympathetic denervation (RSDN) as an adjunctive treatment for patients with cardiomyopathy undergoing catheter ablation of ventricular tachycardia (VT).

Continued on the next page

TABLE 2 Continued

Intervention	Anatomical Target	Current Status	Diseases of Interest	Number of Patients	Ongoing Clinical Trials
			VT/VF	462	NCT01747837 : Renal Sympathetic Denervation to Suppress Tachyarrhythmias in ICD Recipients (RESCUE); Randomized, blinded trial to assess the efficacy of RDN in preventing ventricular arrhythmias and ICD shocks.
SCS	Efferent sympathetic fibers in the spinal cord	Clinical trials	POAF	60	NCT03539354 : Temporary Spinal Cord Stimulation to Prevent Atrial Fibrillation After Cardiac Surgery (TerminationAF); Phase II trial to assess the ability of temporary spinal cord stimulation in preventing POAF
TEA	Efferent sympathetic fibers in the spinal cord	Preclinical	AF or VT/VF		None, though TEA infrequently used in some centers to reduce pain and prevent POAF after cardiac surgery
SGB	Stellate ganglion (efferent fibers)	Clinical trials	POAF	707	NCT03269383 : Study to Evaluate the Effectiveness of SGB in Preventing Post-Op Atrial Fibrillation; Phase II trial to evaluate the efficacy of safety of temporary SGB with bupivacaine to prevent POAF.
			VT/VF	20	NCT02646501 : Prospective Randomized Clinical Trial for Effect of Stellate Ganglion Block in Medically Refractory Ventricular Tachycardia; Study evaluates the effect of SGB in patients with recurrent VT/VF despite optimal medical management
			VT/VF or AF	40	NCT03450226 : Effects of Stellate Ganglion Block on Hemodynamics Instability During Beating Heart Surgery; A randomized controlled, blinded trial evaluating the effect of pre-operative SGB on perioperative arrhythmias and myocardial infarction after off-pump CABG.
Invasive VNS	Vagus nerve (efferent fibers)	Preclinical	VT/VF or AF		None, although animal studies have demonstrated the ability of low-level invasive VNS to reduce sympathetic outflow as well as AF inducibility
Surgical sympathectomy	Thoracic sympathetic ganglia (efferent fibers)	Clinical trials	Heart failure - VT/VF prevention	30	NCT03071653 : Left Cardiac Sympathetic Denervation for Cardiomyopathy Feasibility Pilot Study (LCSD); A randomized controlled trial to test the potential safety and efficacy of LCSD in patients with heart failure due to non-ischemic and ischemic cardiomyopathy
Transtacheal cardiac plexus blockade	Cardiac plexus (efferent fibers)	Preclinical	VT/VF	None	

AF = atrial fibrillation; AGP = autonomic ganglionic plexus; BRS = baroreceptor stimulation; ICD = implantable cardioverter-defibrillator; POAF = postoperative atrial fibrillation; PVI = pulmonary vein isolation; RDN = renal denervation; SCS = spinal cord stimulation; SGB = stellate ganglion block; TEA = thoracic epidural anesthesia; TVNS = transcutaneous vagal nerve stimulation; VF = ventricular fibrillation; VNS = vagal nerve stimulation; VT = ventricular tachycardia.


by increasing preclinical and clinical evidence suggesting that endovascular electrical mapping with stimulation of the renal arteries can result in a differential response including an elevation in blood pressure (42-46).

RDN has received a lot of interest for the treatment of hypertension but may also have utility in reducing arrhythmias through autonomic mechanisms. As mentioned above, the renal vasculature is richly innervated by both afferent and efferent sympathetic nerves which contribute to systemic autonomic signaling. Ablating these nerves may reduce both systemic and cardiac catecholamine levels, as well as reduce atrial fibrosis through modulation of the renin-angiotensin-aldosterone system (47). In a rabbit model of ischemic cardiomyopathy, RDN reduced deleterious atrial structural and electrical remodeling and AF inducibility (48). Additionally, percutaneous RDN reduces atrial nerve sprouting, catecholamine levels, atrial fibrosis, and complexity of AF in a goat model of pacing-induced AF (49). Finally, RDN has been shown to attenuate the electrical and structural remodeling of the left atrium induced by long-term intermittent atrial pacing in a canine model (50).

There are few clinical trials evaluating RDN as an adjunct to PVI for the treatment of recurrent AF. A pilot study of 69 patients with chronic kidney disease and paroxysmal AF showed improved freedom from AF and reduction of AF burden at 1 year in patients randomized to PVI and RDN compared with PVI and spironolactone (51). A study of 86 patients with moderate or severe treatment-resistant hypertension randomized to either PVI alone or PVI with RDN showed improved freedom from AF with the addition of RDN; however, the benefit was most apparent in patients with severe hypertension (52). A follow-up analysis from this cohort has shown a reduction in AF recurrences and burden of AF in patients receiving RDN and PVI (53).

As a stand-alone therapy, RDN may also promote reverse-remodeling and improve atrial substrate. A pilot trial of 20 patients with paroxysmal or persistent AF and hypertension showed that RDN reduced AF burden and improved quality of life (54). Moreover, RDN reduced atrial conduction time, complex fractionated atrial activity, and ventricular mass in 14 patients with treatment-resistant hypertension (55). Multiple upcoming trials will help elucidate the role of

FIGURE 2 Stellate Ganglion Block




(A) Illustration of an ultrasound-guided percutaneous stellate ganglion block. (B, C) Ultrasound images of the percutaneous approach on the right side. (B) Image obtained before needle insertion. Dashed arrow indicates the sympathetic ganglion. (C) Image obtained during needle insertion (arrow) and anesthetic infusion. A marks the carotid artery. V marks the jugular vein. Adopted with permission from Armour et al. (104).

RDN as a stand-alone technique or in addition to PVI to reduce sympathetically mediated AF (Table 2).

Pharmacologic approaches. Epicardial botulinum toxin injection. Given the role of the AGP in the initiation and perpetuation of AF, additional nonablative strategies to modulate AGP signaling are currently under active investigation (Table 2). Injection of botulinum toxin into AGP reduces vagal influences on the AERP as well as vulnerability to AF in both canine (12,56) and ovine models (57). Given the temporary nature of this strategy, autonomic modulation using botulinum toxin injected into AGP has recently

gained attention as a potential strategy to reduce post-operative AF (POAF), a common and impactful complication after cardiac surgery (58,59).

This approach has been studied in 2 small clinical trials. In the first study, 60 patients with paroxysmal AF undergoing coronary artery bypass grafting (CABG) received botulinum toxin or placebo injected into the 4 major left atrial fat pads. With this approach, patients receiving epicardial botulinum toxin had reduced incidence of POAF in the first month after surgery (60), as well as reduced recurrent AF and AF burden at 1 and 3 years after surgery,



respectively (61,62). In the second study, 130 patients undergoing CABG and/or valve surgery were randomized to receive epicardial botulinum toxin or placebo injected into the 4 left atrial fat pads as well as the anterior fat pad. Using this technique, there

was no difference between groups in the occurrence of in-hospital POAF; however, initial POAF episodes were shorter and there were no discernible adverse effects of botulinum toxin injections (63). Further large-scale clinical trials are necessary to determine the utility of this strategy in preventing POAF after cardiac surgery. Additionally, future work is necessary to determine whether botulinum toxin could be delivered to AGP in a less invasive fashion.

Stellate ganglion block. Percutaneous stellate ganglion block (SGB) is a minimally invasive interruption of autonomic innervation from the cervical sympathetic ganglia to the heart (Figure 2). As a key relay station for sympathetic nerve signaling, the stellate ganglion is a prime target for autonomic modulation to reduce AF. Ablation of bilateral stellate ganglia eliminates atrial tachycardia episodes in a canine model of pacing-induced HF (64). A promising minimally invasive strategy to temporarily modulate autonomic signaling is SGB with percutaneous injection of local anesthetic—a procedure frequently used to treat pain syndromes in ambulatory clinics. A pilot study of 36 patients undergoing PVI as well as short duration SGB (unilateral, left or right) has shown that

TABLE 3 Current Therapeutic and Adjunctive Strategies for VT/VF	
Current Therapeutic Strategies for VT/VF	Examples
Pharmacotherapy	Beta-adrenergic blockade Nonselective (β-1, β-2): carvedilol, nadolol, propranolol Selective (β-1): metoprolol, esmolol Membrane active antiarrhythmic drugs Class I: lidocaine, mexiletine, procainamide Class III: sotalol, amiodarone, dofetilide
Ablation	Catheter ablation of focal VT, triggers, and substrate modification (endocardial and epicardial)
Mechanical circulatory support	Temporary: Intra-aortic balloon pump, Impella (ABIOMED, Danvers, Massachusetts) extracorporeal membrane oxygenation, et cetera Permanent: Left ventricular assist device, et cetera
Surgery	Surgical ablation Ventricular aneurysm resection Transplantation

Abbreviations as in Table 2.

SGB lengthened the AERP while reducing inducibility and duration of AF (65). Similarly, a pilot study of 25 patients undergoing CABG and/or aortic valve surgery showed the feasibility and potential efficacy of left-sided SGB for the prevention of POAF (66). A larger study of SGB to prevent POAF is forthcoming and will provide important insight regarding the potential effectiveness of this technique (NCT03269383).

Bioelectronic approaches. Low-level vagal nerve stimulation. High-frequency vagal nerve stimulation (VNS) shortens the AERP (67) and action potential duration in the pulmonary veins (68), facilitating the initiation of AF. However, the intensity of VNS appears to modulate its arrhythmogenicity, with low-level VNS (LLVNS) (below the bradycardia threshold) generally showing a protective or antiarrhythmic effect. Left-sided LLVNS causes upregulation of inhibitory small-conductance calcium-activated potassium channels in the stellate ganglia (69). Similarly, intermittent left cervical VNS is able to slow the ventricular rate in a canine model of pacing-induced AF by damaging the ganglion (70) and increasing activity of the AGP (71). Together, these studies suggest that the antiarrhythmic effect of LLVNS may be mediated through decreased stellate ganglia activity as well as alteration of AGP signaling.

The vagus nerve can be stimulated directly with electrodes; however, given the risks of cervical VNS implantation, recent attention has focused on alternative and less-invasive methods of delivering VNS. Stavrakis et al. (72) recently explored the role of LLVNS in preventing POAF. In their study, 54 patients undergoing cardiac surgery who had a temporary bipolar pacing wire placed in vagal preganglionic fibers near the superior vena cava were randomized to LLVNS or sham stimulation for the duration of post-operative intensive care unit stay. They found that LLVNS reduced the occurrence of POAF. They also found that LLVNS ameliorated the post-operative cytokine elevation associated with cardiac surgery, raising the hypothesis that LLVNS has an anti-inflammatory effect (72).

Because vagal nerve projections exist in the ear, LLVNS may be delivered through stimulation of the tragus. In canine models of pacing-induced AF, LLVNS delivered by tragus stimulation reduced the electrical and structural remodeling induced by rapid atrial pacing (73). Transcutaneous LLVNS increases HRV and decreases muscle sympathetic nerve activity in healthy volunteers, reflecting an anti-adrenergic effect (74). In a pilot study of 40 patients undergoing AF ablation, 1 hour of transcutaneous LLVNS via tragus stimulation decreased pacing-induced AF duration as well as systemic cytokine levels (75). Two

trials are currently underway to evaluate the effectiveness of transcutaneous LLVNS in reducing ambulatory (NCT02548754) as well as postoperative AF (NCT02783157).

Baroreceptor stimulation. Mechanoreceptors in the carotid sinus and aortic arch (baroreceptors) generate dynamic feedback to brainstem centers responsible for autonomic tone, thereby exerting beat-to-beat control of blood pressure and heart rate (76). Electric stimulation of carotid baroreceptors reduces sympathetic tone (77) while augmenting vagal tone (78). Thus, baroreceptor stimulation (BRS) has the capacity to generate both pro-arrhythmic and antiarrhythmic influences relative to AF. In a canine model of low-level BRS at a voltage below the threshold to lower systemic blood pressure, 2 h of BRS was associated with increased AERP, increased AF threshold, and decreased cardiac AGP activity (79). This effect was further confirmed in porcine model of high- versus low-level BRS in simulated OSA, where low-level BRS prolonged AERP and reduced AF inducibility, whereas high-level baroreceptor stimulation shortened AERP and did not meaningfully impact AF inducibility. Whether low-level BRS can reduce the occurrence of AF remains to be seen. However, investigation into this potential avenue of neuromodulation for treatment of AF continues.

FUTURE OUTLOOK. Given the increasing prevalence as well as the morbidity associated with AF, new strategies to prevent or reduce AF remain an important unmet need. There has been great progress with current therapeutic strategies to reduce the morbidity of AF, including ablation-based as well as pharmacologic techniques. However, current therapies incompletely use the developing mechanistic understanding of the pathogenesis of AF. Because the ANS is an important contributor in the development of AF, strategies to modulate autonomic signaling hold significant promise in reducing both the incidence and consequences of AF. As outlined above, a number of early-stage clinical trials are ongoing to directly translate promising basic science findings into safe, efficacious strategies to reduce AF (Table 2). As the evidence base surrounding autonomic modulation to reduce AF continues to develop, one can anticipate a growing number of effective strategies using autonomic modulation to reduce AF.

VENTRICULAR TACHYCARDIA

RATIONALE FOR AUTONOMIC MODULATION FOR VENTRICULAR TACHYCARDIA. Despite improvements in our therapies for cardiovascular disease,

ventricular tachycardia (VT) and ventricular fibrillation (VF) remain significant challenges in cardiovascular practice. These life-threatening dysrhythmias are particularly common in cardiac and surgical intensive care units, and are responsible for more than 450,000 deaths every year in the United States alone (80). The sympathetic nervous system (SNS) plays a significant role in the genesis of ventricular arrhythmias, and thus represents an important therapeutic target (**Central Illustration**).

The interaction between the SNS and the heart can be reduced to efferent signals to the heart and afferent signaling from the heart. Sympathetic efferent signaling induces maladaptive changes in ventricular electrophysiology. Sympathetic activation has been shown to be able to either activate or maintain ventricular arrhythmias via the following mechanisms: 1) shortened refractory periods (81,82); 2) abbreviated action potential duration (83,84); 3) increased dispersion of refractoriness (85); 4) increased heterogeneity of repolarization (86), and; 5) induced early afterdepolarizations (87). In sum, sympathetically mediated changes in electrophysiology decrease the ventricular threshold and can induce ventricular arrhythmias. Furthermore, the underlying cardiomyopathy often results in autonomic imbalance and dysregulation as a result of underlying structural changes (scar/fibrosis) or persistent/recurrent states of HF (**Figure 3**).

It is now increasingly well understood that the afferent inputs from the heart and peripheral organs (such as the kidney to the ANS) contribute to maladaptive remodeling of the ANS in the presence of myocardial injury. For example, investigations in a canine model of MI have shown that myocardial injury induces morphologic and neurochemical changes in the bilateral stellate ganglia approximately 5 weeks after the initial insult (88,89). Beyond an increase in neuronal size, MI results in a persistent increase in the synaptic density of bilateral stellate ganglia and is associated with increased stellate ganglion nerve activity for up to 8 weeks.

Furthermore, there is evidence of circadian variation in the stellate ganglion nerve activity. Changes in gene expression and neuronal structure are not restricted to the stellate ganglia but can also be found in the dorsal root ganglia of the spinal cord (90,91). Finally, even without an apparent neuroanatomical link between renal nerves and the cardiac sympathetic tone, direct or indirect (myocardial ischemia) renal nerve activation leads to central nervous remodeling (stellate ganglia) with direct impact on cardiac sympathetic tone (92). These results suggest a

potential mechanistic link between neural remodeling and ventricular arrhythmias.

PAST AND PRESENT. Current strategies in autonomic modulation for ventricular arrhythmias. Clinical management. At present, clinical management of VT/VF is often restricted to pharmacological therapy and catheter ablation (**Table 3**). In the setting of VT/VF storm, the underlying sympathetic hyperactivation is further exacerbated by ventricular arrhythmias, internal or external defibrillations, pain, anxiety, and progressive HF (93). Initial medical management that targets the ANS includes beta blockade.

Use of beta blockade is commonly the first and most important pharmacologic intervention for preventing VT/VF given its effectiveness in reducing sympathetic tone. However, beta blockade in the acute setting of electrical storm (≥ 3 VT/VF episodes in 24 h) is often underused due to concerns for potential beta blockade-induced cardiac decompensation. The recently presented clinical study evaluating a non-selective beta blocker propranolol versus a selective beta blocker metoprolol in the setting of electrical storm has shown the safety and effectiveness of nonselective beta blockers in patients with high morbidity and mortality (94). In these patients, propranolol led to less ventricular arrhythmia compared with metoprolol. However, VT and VF are often refractory to medical management including beta blockade, antiarrhythmic drug therapy, and even mechanical hemodynamic support. Although monomorphic VT and premature ventricular contraction-triggered polymorphic VT may be amenable to successful radiofrequency ablation, this approach is not always immediately feasible, especially in critically ill or hemodynamically unstable patients.

Surgical approach. The concept of physically interrupting cardiac sympathetic innervation to treat ventricular arrhythmias was first introduced by Schwartz et al. (95), who studied the interaction of the ANS with ventricular arrhythmias and showed that surgical cardiac denervation increases the ventricular arrhythmia threshold in healthy dogs ($n = 11$), preventing induction of VT/VF with external stimulation. In the 1990s, unilateral (predominantly left-sided) surgical sympathectomy was applied in patients ($n = 40$) at high risk for ventricular arrhythmias after an MI and was found to reduce risk of sudden death significantly compared to controls without pharmacologic or surgical sympathetic inhibition (96). Given the success with SGB in patients with MI, surgical sympathectomy has been used to

treat patients with long-QT syndrome and catecholaminergic polymorphic VT (97).

Because unilateral sympathetic denervation may be insufficient in some cases, bilateral surgical sympathetic denervation has been proposed as an alternative and more effective therapy (98). Bilateral sympathetic denervation may be more effective due to bilateral stellate ganglion remodeling with cardiac injury (89) and anatomical variation in cardiac innervation (99). Ajijola et al. (98) have shown that bilateral surgical cardiac sympathetic denervation can be useful for the treatment of medically refractory electrical storm due to a number of causes and may result in improved outcomes compared to unilateral surgical denervation (100). With the exception of a single prospective study (96) that compared surgical sympathectomy to placebo, contemporary studies have been limited to retrospective observational studies. The largest published experience now encompasses 121 cases, and shows an association with decreased sustained VT and implantable cardioverter-defibrillator (ICD) shocks in patients with refractory VT (101).

Interventional approaches. Today, the modulation of the autonomic tone via operative sympathetic denervation (i.e., sympathectomy) is considered a useful intervention only when all other therapeutic options have been exhausted. However, not infrequently, patients presenting with electrical storm (≥ 3 VT/VF episodes in 24 h) have either failed traditional therapeutic approaches or are deemed too unstable to undergo surgical sympathectomy. These situations have led some investigators to use minimally invasive approaches to achieve temporary blockade cardiac sympathetic innervation. As a result, minimally invasive approaches to cardiac sympathetic blockade have emerged as therapeutic options and diagnostic tools to help gauge whether a given patient would benefit from the permanent surgical sympathectomy (Figure 2). Notably, interventional approaches to modulate the ANS in patients with VT are not widely adopted and often limited to select institutions across the world.

SGB. Similar to the limited evidence base for the use of SGB in the treatment of AF, the utility of SGB for the treatment of ventricular arrhythmias has been chronicled in individual case reports (102,103). Moreover, SGB has never been investigated in a prospective, controlled clinical study for the treatment of VT/VF. A systematic review of the literature yielded 35 published cases using mostly unilateral SGB (86% of cases) (104). SGB resulted in a significant reduction of ventricular arrhythmia episodes from 24-h pre- to

24-h post-SGB (mean, 16.5; 95% confidence interval [CI]: 9.7 to 23.1 events vs. mean, 1.4; 95% CI: 0.85 to 2.01 events; $p = 0.0002$) (104). The need for defibrillation decreased comparably (pre-SGB mean, 14.2; 95% CI: 6.8 to 21.6 events vs. post-SGB mean, 0.6; 95% CI: 0.3 to 0.9 events; $p = 0.0026$). Importantly, SGB was associated with a reduction of ventricular arrhythmia burden regardless of the etiology of cardiomyopathy (nonischemic vs. ischemic), type of ventricular arrhythmia (monomorphic vs. polymorphic), and degree of contractile dysfunction (preserved vs. reduced ejection fraction). In this series of uncontrolled observations, SGB was followed by surgical sympathectomy in 21% of cases.

An observational study of several different methods of autonomic blockade for the treatment of electrical storm has shown a substantial difference in mortality ($\sim 60\%$ absolute risk reduction compared with standard care) in 49 patients (only 6 received unilateral percutaneous SGB) (105). Finally, the repeated observation that SGB is associated with a prolonged duration of arrhythmia suppression (3 to 4 days) well beyond what is expected from the duration of a pharmacological block (~ 1 day) (102,103,106) suggests the presence of an afferent mechanism with potential effects on neural remodeling. The safety profile of this intervention appears to be favorable given the ability to deliver the nerve block under ultrasound guidance.

Thoracic epidural anesthesia. An alternative pathway to block the central sympathetic output and input to/from the heart is thoracic epidural anesthesia (TEA). Using an established anesthetic approach, TEA blocks afferent and efferent innervation between the heart and CNS. Anti-arrhythmic properties of TEA have been shown in several pre-clinical models (107,108). The safety and efficacy of thoracic epidural anesthesia was explored as part of several small case series (109,110). Successful reduction of ventricular arrhythmia burden was achieved in the majority of patients with refractory arrhythmias (6 of 11 patients, 55%). Notably, the use of thoracic epidural anesthesia is limited in the setting of antiplatelet or oral anticoagulation therapy due to concerns for high-risk bleeding.

Developing neuromodulatory approaches. Alternative approaches to cardiac sympathetic modulation. Beyond the direct surgical or minimally invasive approaches to modulate the autonomic innervation of the heart, there are series of novel interventional approaches to alter the sympathetic nervous tone. Spinal cord stimulation (SCS) at the level of T1 to T5 can modulate autonomic output, likely via inhibition of the stellate

ganglia, and possibly increased vagal activity (111,112). In animal models, SCS reduces HRV, decreases the incidence of ventricular arrhythmias, and reduces left stellate ganglion activity in acute MI (111). There are limited preliminary clinical data regarding the use of SCS. In 2 patients with ischemic and non-ischemic cardiomyopathy, SCS reduced the VT and VF burden up to 75% to 100% (113). A prospective, multicenter randomized clinical study (N = 81 patients) that investigated the effects of SCS therapy for the treatment of systolic HF failed to show improved outcomes (114). Nevertheless, the exact mechanism of SCS on ventricular arrhythmias remains to be determined, and the clinical safety and efficacy of SCS for the prevention of recurrent ventricular arrhythmia remains unknown.

Transtacheal cardiac plexus blockade is a most recent novel attempt to interrupt cardiac sympathetic innervation. Sympathetic nerves converge from both stellate ganglia to form the cardiac plexus. Given the anatomical location of the cardiac plexus between pulmonary artery, aortic arch, and anterior wall of the trachea (115), the cardiac plexus is amenable to a transtacheal access. In a recent study, investigators provided initial evidence that temporary interruption of nerve traffic via lidocaine injection can counteract the effects of stellate ganglion stimulation and thus serve as a new avenue for temporary or permanent minimally invasive cardiac denervation (116).

RDN. As previously mentioned, RDN was initially developed for the treatment of hypertension and has been shown to reduce global and regional sympathetic nerve activity (117). In a porcine ischemia model, RDN reduced the number of spontaneous ventricular extrasystoles and VF in a similar manner to beta blocker therapy (118). The effects of RDN on ventricular electrophysiology appear to be independent of its effects on blood pressure as shown in initial preclinical and clinical studies. The early clinical experience with RDN and ventricular arrhythmias has also been encouraging (119). The largest multicenter case series has shown that bilateral RDN in 13 patients with refractory VT was associated with an 85% freedom from VT at 3 months and no periprocedural complications (120). In a second study including 10 patients, RDN led to a significant decrease in VT/VF burden and ICD shocks at 6 months (121). Despite recent advances for RDN as an antihypertensive strategy (122) and our improved understanding of the best procedural approach (46), definitive evidence to support its use as supplementary strategy for ventricular arrhythmias is needed.

VNS. VNS to augment cardiac parasympathetic tone and oppose sympathetic hyperactivation is also being evaluated for the prevention of ventricular arrhythmias. Preclinical studies support a favorable sympatholytic effect of direct efferent via the cervical vagus nerve or indirect afferent VNS via its auricular branches (123). Post-MI models provide additional evidence for the protective myocardial effects of VNS that are heart rate independent. In canine model of MI, high-intensity vagal stimulation led to a 71% VF-free survival versus 40% with low intensity and 10% in the control group (124). In parallel, prophylactic VNS in the setting of MI minimized the risk of VF onset (125).

FUTURE OUTLOOK. The growing burden of ventricular arrhythmias and their significant morbidity and mortality demands new effective therapeutic approaches. While there are several effective pharmacologic and catheter-based therapies for ventricular arrhythmias, these therapies frequently meet their limits in an increasingly complex medical environment (e.g., use of mechanical circulatory devices). The ANS is a new frontier in the management of ventricular arrhythmias. Before novel therapeutic strategies can be widely adopted in clinical practice, randomized clinical trials are needed to establish evidence of safety and efficacy (Table 2).

Given the myriad of new strategies to modulate the autonomic tone, there is a need to develop methods for individualizing treatment based on a given patient's triggers, comorbidities, and underlying pathology. Future therapies do not have to restrict themselves to secondary prevention nor a device-based approach to neuromodulation.

Recently published work has pioneered the use of optogenetic modulation of cardiac sympathetic nerve activity to prevent ventricular arrhythmias (126). Optogenetics is a relatively novel technology applied in neuroscience to silence or enhance neural activity via targeted genetic modification. Viral vectors are used to introduce an inhibitory light sensitive opsin into stellate ganglion neurons to silence them (127). When ArchT (inhibitory light-sensitive opsin) is genetically expressed in targeted cells, it can be activated via illumination using characteristic wavelength. Activation leads to cell hyperpolarization via activation of ionic channels and as a result silences target cells (128,129).

In a canine model, optogenetic (and reversible) stellate ganglia inhibition was protective against myocardial ischemia-induced ventricular arrhythmias (126). This work builds on prior attempts to modulate norepinephrine release from cardiac neurons via optogenetic modification (130). Targeted

genetic modulation resulted in a desired increase in norepinephrine release with consequent stimulation in cardiac function. Arguably, if this technology can be applied in man (cardiac plexus, stellate ganglia, thoracic sympathetic ganglia, and vagus nerve) it could be used as a preventative tool for high-risk patients with ventricular arrhythmias and as a therapeutic tool for patients with recurrent ventricular arrhythmias. The unique on/off property of the optogenetic modification is the central strength of this technology. A major limitation is the current need for direct application to the target nerve structure via injection for example.

The use of autonomic modulation for primary prevention is currently under investigation and if successful could have potentially far-reaching implications. An ongoing prospective randomized clinical trial is exploring the use of surgical sympathetic denervation for primary prevention of sudden cardiac death in patients with heart failure (131).

CONCLUSIONS

Over the past several decades, there has been increasing recognition of the importance of interactions between the heart and the ANS in the pathophysiology of arrhythmias. There are now a multiplicity of potential therapies under development to target several different neurologic targets and structures for both atrial and ventricular arrhythmias. Moving forward, randomized clinical trials will be essential to establish the safety, efficacy, and comparative effectiveness of these new and promising interventions.

ADDRESS FOR CORRESPONDENCE: Dr. Jonathan P. Piccini, Electrophysiology Section, Duke University Medical Center, Duke Clinical Research Institute, PO Box 17969, Durham, North Carolina 27710. E-mail: jonathan.piccini@duke.edu.

REFERENCES

- Hanna P, Rajendran PS, Ajjola OA, et al. Cardiac neuroanatomy – imaging nerves to define functional control. *Auton Neurosci* 2017;207:48-58.
- Beissner F, Meissner K, Bar KJ, Napadow V. The autonomic brain: an activation likelihood estimation meta-analysis for central processing of autonomic function. *J Neurosci* 2013;33:10503-11.
- McDougall SJ, Munzberg H, Derbenev AV, Zsombok A. Central control of autonomic functions in health and disease. *Front Neurosci* 2014;8:440.
- Shen MJ, Zipes DP. Role of the autonomic nervous system in modulating cardiac arrhythmias. *Circ Res* 2014;114:1004-21.
- Armour JA, Murphy DA, Yuan BX, Macdonald S, Hopkins DA. Gross and microscopic anatomy of the human intrinsic cardiac nervous system. *Anat Rec* 1997;247:289-98.
- Hou Y, Scherlag BJ, Lin J, et al. Ganglionated plexi modulate extrinsic cardiac autonomic nerve input: effects on sinus rate, atrioventricular conduction, refractoriness, and inducibility of atrial fibrillation. *J Am Coll Cardiol* 2007;50:61-8.
- Pistoia F, Sacco S, Tiseo C, Degan D, Ornello R, Carolei A. The epidemiology of atrial fibrillation and stroke. *Cardiol Clin* 2016;34:255-68.
- Krijthe BP, Kunst A, Benjamin EJ, et al. Projections on the number of individuals with atrial fibrillation in the European Union, from 2000 to 2060. *Eur Heart J* 2013;34:2746-51.
- European Heart Rhythm A, European Association for Cardio-Thoracic S, Camm AJ, et al. Guidelines for the management of atrial fibrillation: the Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology (ESC). *Eur Heart J* 2010;31:2369-429.
- Ganesan AN, Chew DP, Hartshorne T, et al. The impact of atrial fibrillation type on the risk of thromboembolism, mortality, and bleeding: a systematic review and meta-analysis. *Eur Heart J* 2016;37:1591-602.
- Chen LY, Chung MK, Allen LA, et al. Atrial fibrillation burden: moving beyond atrial fibrillation as a binary entity: a scientific statement from the American Heart Association. *Circulation* 2018;137:e623-44.
- Oh S, Choi EK, Zhang Y, Mazgalev TN. Botulinum toxin injection in epicardial autonomic ganglia temporarily suppresses vagally mediated atrial fibrillation. *Circ Arrhythm Electrophysiol* 2011;4:560-5.
- Chen PS, Tan AY. Autonomic nerve activity and atrial fibrillation. *Heart Rhythm* 2007;4:S61-4.
- Oliveira M, da Silva MN, Timoteo AT, et al. Inducibility of atrial fibrillation during electrophysiologic evaluation is associated with increased dispersion of atrial refractoriness. *Int J Cardiol* 2009;136:130-5.
- Patterson E, Lazzara R, Szabo B, et al. Sodium-calcium exchange initiated by the Ca²⁺ transient: an arrhythmia trigger within pulmonary veins. *J Am Coll Cardiol* 2006;47:1196-206.
- Zipes DP, Mihalick MJ, Robbins GT. Effects of selective vagal and stellate ganglion stimulation of atrial refractoriness. *Cardiovasc Res* 1974;8:647-55.
- Sharifov OF, Fedorov VV, Beloshapko GG, Glukhov AV, Yushmanova AV, Rosenshtaukh LV. Roles of adrenergic and cholinergic stimulation in spontaneous atrial fibrillation in dogs. *J Am Coll Cardiol* 2004;43:483-90.
- Tan AY, Zhou S, Ogawa M, et al. Neural mechanisms of paroxysmal atrial fibrillation and paroxysmal atrial tachycardia in ambulatory canines. *Circulation* 2008;118:916-25.
- Bettoni M, Zimmermann M. Autonomic tone variations before the onset of paroxysmal atrial fibrillation. *Circulation* 2002;105:2753-9.
- Agarwal SK, Norby FL, Whitsel EA, et al. Cardiac autonomic dysfunction and incidence of atrial fibrillation: results from 20 years follow-up. *J Am Coll Cardiol* 2017;69:291-9.
- Baykaner T, Zografos TA, Zaman JAB, et al. Spatial relationship of organized rotational and focal sources in human atrial fibrillation to autonomic ganglionated plexi. *Int J Cardiol* 2017;240:234-9.
- Mann SJ. Redefining beta-blocker use in hypertension: selecting the right beta-blocker and the right patient. *J Am Soc Hypertens* 2017;11:54-65.
- Pison L, Tiltz R, Jalife J, Haissaguerre M. Pulmonary vein triggers, focal sources, rotors and atrial cardiomyopathy: implications for the choice of the most effective ablation therapy. *J Intern Med* 2016;279:449-56.
- Kurotobi T, Shimada Y, Kino N, et al. Features of intrinsic ganglionated plexi in both atria after extensive pulmonary isolation and their clinical significance after catheter ablation in patients with atrial fibrillation. *Heart Rhythm* 2015;12:470-6.
- Straznicki NE, Grima MT, Sari CI, et al. Neuroadrenergic dysfunction along the diabetes continuum: a comparative study in obese metabolic syndrome subjects. *Diabetes* 2012;61:2506-16.
- Somers VK, Dyken ME, Clary MP, Abboud FM. Sympathetic neural mechanisms in obstructive sleep apnea. *J Clin Invest* 1995;96:1897-904.
- Friedman DJ, Liu P, Barnett AS, et al. Obstructive sleep apnea is associated with

- increased rotor burden in patients undergoing focal impulse and rotor modification guided atrial fibrillation ablation. *Europace* 2018;20:f337-42.
28. Linz D, Elliott AD, Hohl M, et al. Role of autonomic nervous system in atrial fibrillation. *Int J Cardiol* 2019;287:181-8.
 29. Holmqvist F, Guan N, Zhu Z, et al. Impact of obstructive sleep apnea and continuous positive airway pressure therapy on outcomes in patients with atrial fibrillation-Results from the Outcomes Registry for Better Informed Treatment of Atrial Fibrillation (ORBIT-AF). *Am Heart J* 2015;169:647-54.e2.
 30. Deng F, Raza A, Guo J. Treating obstructive sleep apnea with continuous positive airway pressure reduces risk of recurrent atrial fibrillation after catheter ablation: a meta-analysis. *Sleep Med* 2018;46:5-11.
 31. Piccini JP, Pokorney SD, Anstrom KJ, et al. Adaptive servo-ventilation reduces atrial fibrillation burden in patients with heart failure and sleep apnea. *Heart Rhythm* 2019;16:91-7.
 32. Danik S, Neuzil P, d'Avila A, et al. Evaluation of catheter ablation of periatrial ganglionic plexi in patients with atrial fibrillation. *Am J Cardiol* 2008;102:578-83.
 33. Katritsis D, Giazitzoglou E, Sougiannis D, Goumas N, Paxinos G, Camm AJ. Anatomic approach for ganglionic plexi ablation in patients with paroxysmal atrial fibrillation. *Am J Cardiol* 2008;102:330-4.
 34. Katritsis DG, Pokushalov E, Romanov A, et al. Autonomic denervation added to pulmonary vein isolation for paroxysmal atrial fibrillation: a randomized clinical trial. *J Am Coll Cardiol* 2013;62:2318-25.
 35. Pokushalov E, Romanov A, Katritsis DG, et al. Ganglionated plexus ablation vs linear ablation in patients undergoing pulmonary vein isolation for persistent/long-standing persistent atrial fibrillation: a randomized comparison. *Heart Rhythm* 2013;10:1280-6.
 36. Driessen AHG, Berger WR, Krul SPJ, et al. Ganglion plexus ablation in advanced atrial fibrillation: the AFACT study. *J Am Coll Cardiol* 2016;68:1155-65.
 37. Knuepfer MM, Schramm LP. The conduction velocities and spinal projections of single renal afferent fibers in the rat. *Brain Res* 1987;435:167-73.
 38. Barajas L, Wang P. Myelinated nerves of the rat kidney. A light and electron microscopic autoradiographic study. *J Ultrastruct Res* 1978;65:148-62.
 39. DiBona GF, Kopp UC. Neural control of renal function. *Physiol Rev* 1997;77:75-197.
 40. Kopp UC. Role of renal sensory nerves in physiological and pathophysiological conditions. *Am J Physiol Regul Integr Comp Physiol* 2015;308:R79-95.
 41. Kopp UC, Cicha MZ, Smith LA, Mulder J, Hokfelt T. Renal sympathetic nerve activity modulates afferent renal nerve activity by PGE2-dependent activation of alpha1- and alpha2-adrenoceptors on renal sensory nerve fibers. *Am J Physiol Regul Integr Comp Physiol* 2007;293:R1561-72.
 42. Pokushalov E, Romanov A, Corbucci G, et al. A randomized comparison of pulmonary vein isolation with versus without concomitant renal artery denervation in patients with refractory symptomatic atrial fibrillation and resistant hypertension. *J Am Coll Cardiol* 2012;60:1163-70.
 43. Gal P, de Jong MR, Smit JJ, Adiyaman A, Staessen JA, Elvan A. Blood pressure response to renal nerve stimulation in patients undergoing renal denervation: a feasibility study. *J Human Hypertens* 2015;29:292-5.
 44. Chinushi M, Izumi D, Iijima K, et al. Blood pressure and autonomic responses to electrical stimulation of the renal arterial nerves before and after ablation of the renal artery. *Hypertension* 2013;61:450-6.
 45. Lu J, Wang Z, Zhou T, et al. Selective proximal renal denervation guided by autonomic responses evoked via high-frequency stimulation in a pre-clinical canine model. *Circ Cardiovasc Interv* 2015;8. pii:e001847.
 46. Fudim M, Sobotka AA, Yin YH, et al. Selective vs. global renal denervation: a case for less is more. *Curr Hypertens Rep* 2018;20:37.
 47. Linz D, Hohl M, Elliott AD, et al. Modulation of renal sympathetic innervation: recent insights beyond blood pressure control. *Clin Auton Res* 2018;28:375-84.
 48. Yamada S, Fong MC, Hsiao YW, et al. Impact of renal denervation on atrial arrhythmogenic substrate in ischemic model of heart failure. *J Am Heart Assoc* 2018;7. pii:e007312.
 49. Linz D, van Hunnik A, Hohl M, et al. Catheter-based renal denervation reduces atrial nerve sprouting and complexity of atrial fibrillation in goats. *Circ Arrhythm Electrophysiol* 2015;8:466-74.
 50. Wang X, Huang C, Zhao Q, et al. Effect of renal sympathetic denervation on the progression of paroxysmal atrial fibrillation in canines with long-term intermittent atrial pacing. *Europace* 2015;17:647-54.
 51. Kiuchi MG, Chen S, Hoye NA, Purerfellner H. Pulmonary vein isolation combined with spironeolactone or renal sympathetic denervation in patients with chronic kidney disease, uncontrolled hypertension, paroxysmal atrial fibrillation, and a pacemaker. *J Interv Card Electrophysiol* 2018;51:51-9.
 52. Pokushalov E, Romanov A, Katritsis DG, et al. Renal denervation for improving outcomes of catheter ablation in patients with atrial fibrillation and hypertension: early experience. *Heart Rhythm* 2014;11:1131-8.
 53. Romanov A, Pokushalov E, Ponomarev D, et al. Pulmonary vein isolation with concomitant renal artery denervation is associated with reduction in both arterial blood pressure and atrial fibrillation burden: Data from implantable cardiac monitor. *Cardiovasc Ther* 2017;35.
 54. Feyz L, Theuns DA, Bhagwandien R, et al. Atrial fibrillation reduction by renal sympathetic denervation: 12 months' results of the AFFORD study. *Clin Res Cardiol* 2018.
 55. McLellan AJ, Schlaich MP, Taylor AJ, et al. Reverse cardiac remodeling after renal denervation: atrial electrophysiologic and structural changes associated with blood pressure lowering. *Heart Rhythm* 2015;12:982-90.
 56. Lo LW, Chang HY, Scherlag BJ, et al. Temporary suppression of cardiac ganglionated plexi leads to long-term suppression of atrial fibrillation: evidence of early autonomic intervention to break the vicious cycle of "AF begets AF." *J Am Heart Assoc* 2016;5:e003309.
 57. Nazeri A, Ganapathy AV, Massumi A, et al. Effect of botulinum toxin on inducibility and maintenance of atrial fibrillation in ovine myocardial tissue. *Pacing Clin Electrophysiol* 2017;40:693-702.
 58. Mathew JP, Fontes ML, Tudor IC, et al. A multicenter risk index for atrial fibrillation after cardiac surgery. *JAMA* 2004;291:1720-9.
 59. LaPar DJ, Speir AM, Crosby IK, et al. Post-operative atrial fibrillation significantly increases mortality, hospital readmission, and hospital costs. *Ann Thorac Surg* 2014;98:527-33.
 60. Pokushalov E, Kozlov B, Romanov A, et al. Botulinum toxin injection in epicardial fat pads can prevent recurrences of atrial fibrillation after cardiac surgery: results of a randomized pilot study. *J Am Coll Cardiol* 2014;64:628-9.
 61. Pokushalov E, Kozlov B, Romanov A, et al. Long-term suppression of atrial fibrillation by botulinum toxin injection into epicardial fat pads in patients undergoing cardiac surgery: one-year follow-up of a randomized pilot study. *Circ Arrhythm Electrophysiol* 2015;8:1334-41.
 62. Romanov A, Pokushalov E, Ponomarev D, et al. Long-term suppression of atrial fibrillation by botulinum toxin injection into epicardial fat pads in patients undergoing cardiac surgery: three-year follow-up of a randomized study. *Heart Rhythm* 2019;16:172-7.
 63. Waldron NH, Cooter M, Haney JC, et al. Temporary autonomic modulation with botulinum toxin type A to reduce atrial fibrillation after cardiac surgery. *Heart Rhythm* 2019;16:178-84.
 64. Ogawa M, Tan AY, Song J, et al. Cryoablation of stellate ganglia and atrial arrhythmia in ambulatory dogs with pacing-induced heart failure. *Heart Rhythm* 2009;6:1772-9.
 65. Leftheriotis D, Flevari P, Kossyvakis C, et al. Acute effects of unilateral temporary stellate ganglion block on human atrial electrophysiological properties and atrial fibrillation inducibility. *Heart Rhythm* 2016;13:2111-7.
 66. Connors CW, Craig WY, Buchanan SA, Poltak JM, Gagnon JB, Curry CS. Efficacy and efficiency of perioperative stellate ganglion blocks in cardiac surgery: a pilot study. *J Cardiothorac Vasc Anesth* 2018;32:e28-30.
 67. Chen PS, Chen LS, Fishbein MC, Lin SF, Nattel S. Role of the autonomic nervous system in atrial fibrillation: pathophysiology and therapy. *Circ Res* 2014;114:1500-15.
 68. Patterson E, Po SS, Scherlag BJ, Lazzara R. Triggered firing in pulmonary veins initiated by

in vitro autonomic nerve stimulation. *Heart Rhythm* 2005;2:624-31.

69. Shen MJ, Hao-Che C, Park HW, et al. Low-level vagus nerve stimulation upregulates small conductance calcium-activated potassium channels in the stellate ganglion. *Heart Rhythm* 2013;10:910-5.

70. Chinda K, Tsai WC, Chan YH, et al. Intermittent left cervical vagal nerve stimulation damages the stellate ganglia and reduces the ventricular rate during sustained atrial fibrillation in ambulatory dogs. *Heart Rhythm* 2016;13:771-80.

71. Jiang Z, Zhao Y, Tsai WC, et al. Effects of vagal nerve stimulation on ganglionated plexi nerve activity and ventricular rate in ambulatory dogs with persistent atrial fibrillation. *J Am Coll Cardiol EP* 2018;4:1106-14.

72. Stavakis S, Humphrey MB, Scherlag B, et al. Low-level vagus nerve stimulation suppresses post-operative atrial fibrillation and inflammation: a randomized study. *J Am Coll Cardiol EP* 2017;3:929-38.

73. Chen M, Yu L, Liu Q, et al. Low level tragus nerve stimulation is a non-invasive approach for anti-atrial fibrillation via preventing the loss of connexins. *Int J Cardiol* 2015;179:144-5.

74. Clancy JA, Mary DA, Witte KK, Greenwood JP, Deuchars SA, Deuchars J. Non-invasive vagus nerve stimulation in healthy humans reduces sympathetic nerve activity. *Brain Stimul* 2014;7:871-7.

75. Stavakis S, Humphrey MB, Scherlag BJ, et al. Low-level transcutaneous electrical vagus nerve stimulation suppresses atrial fibrillation. *J Am Coll Cardiol* 2015;65:867-75.

76. Suarez-Roca H, Klinger RY, Podgoreanu MV, et al. Contribution of baroreceptor function to pain perception and perioperative outcomes. *Anesthesiology* 2019;130:634-50.

77. Heusser K, Tank J, Engeli S, et al. Carotid baroreceptor stimulation, sympathetic activity, baroreflex function, and blood pressure in hypertensive patients. *Hypertension* 2010;55:619-26.

78. Linz D, Mahfoud F, Schotten U, et al. Effects of electrical stimulation of carotid baroreflex and renal denervation on atrial electrophysiology. *J Cardiovasc Electrophysiol* 2013;24:1028-33.

79. Liao K, Yu L, Zhou X, et al. Low-level baroreceptor stimulation suppresses atrial fibrillation by inhibiting ganglionated plexus activity. *Can J Cardiol* 2015;31:767-74.

80. Zheng ZJ, Croft JB, Giles WH, Mensah GA. Sudden cardiac death in the United States, 1989 to 1998. *Circulation* 2001;104:2158-63.

81. Burgess MJ, Haws CW. Effects of sympathetic stimulation on refractory periods of ischemic canine ventricular myocardium. *J Electrocardiol* 1982;15:1-8.

82. Martins JB, Zipes DP. Effects of sympathetic and vagal nerves on recovery properties of the endocardium and epicardium of the canine left ventricle. *Circ Res* 1980;46:100-10.

83. Taggart P, Sutton P, Chalabi Z, et al. Effect of adrenergic stimulation on action potential duration restitution in humans. *Circulation* 2003;107:285-9.

84. Zaza A, Malfatto G, Schwartz PJ. Sympathetic modulation of the relation between ventricular repolarization and cycle length. *Circ Res* 1991;68:1191-203.

85. Opthof T, Coronel R, Vermeulen JT, Verberne HJ, van Capelle FJ, Janse MJ. Dispersion of refractoriness in normal and ischaemic canine ventricle: effects of sympathetic stimulation. *Cardiovasc Res* 1993;27:1954-60.

86. Vaseghi M, Yamakawa K, Sinha A, et al. Modulation of regional dispersion of repolarization and T-peak to T-end interval by the right and left stellate ganglia. *Am J Physiol Heart Circ Physiol* 2013;305:H1020-30.

87. Ben-David J, Zipes DP. Differential response to right and left ansae subclaviae stimulation of early afterdepolarizations and ventricular tachycardia induced by cesium in dogs. *Circulation* 1988;78:1241-50.

88. Ajjola OA, Yagishita D, Reddy NK, et al. Remodeling of stellate ganglion neurons after spatially targeted myocardial infarction: neuro-peptide and morphologic changes. *Heart Rhythm* 2015;12:1027-35.

89. Han S, Kobayashi K, Joung B, et al. Electro-anatomic remodeling of the left stellate ganglion after myocardial infarction. *J Am Coll Cardiol* 2012;59:954-61.

90. Saddic LA, Howard-Quigano K, Kipke J, et al. Progression of myocardial ischemia leads to unique changes in immediate early gene expression in the spinal cord dorsal horn. *Am J Physiol Heart Circ Physiol* 2018;15:H1592-601.

91. Gao C, Howard-Quigano K, Rau C, et al. Inflammatory and apoptotic remodeling in autonomic nervous system following myocardial infarction. *PLoS One* 2017;12:e0177750.

92. Li C, Huang D, Tang J, et al. CIC-3 chloride channel is involved in isoprenaline-induced cardiac hypertrophy. *Gene* 2018;642:335-42.

93. Fukuda K, Kanazawa H, Aizawa Y, Ardell JL, Shivkumar K. Cardiac innervation and sudden cardiac death. *Circ Res* 2015;116:2005-19.

94. Chatzidou S, Kontogiannis C, Tsilimigras DI, et al. Propranolol versus metoprolol for treatment of electrical storm in patients with implantable cardioverter-defibrillator. *J Am Coll Cardiol* 2018;71:1897-906.

95. Schwartz PJ, Snebold NG, Brown AM. Effects of unilateral cardiac sympathetic denervation on the ventricular fibrillation threshold. *Am J Cardiol* 1976;37:1034-40.

96. Schwartz PJMM, Pollavini G, Lotto A, et al. Prevention of sudden cardiac death after a first myocardial infarction by pharmacologic or surgical antiadrenergic interventions. *J Cardiovasc Electrophysiol* 1992;3:1-16.

97. Wilde AA, Bhuiyan ZA, Crotti L, et al. Left cardiac sympathetic denervation for catecholaminergic polymorphic ventricular tachycardia. *N Engl J Med* 2008;358:2024-9.

98. Ajjola OA, Lellouche N, Bourke T, et al. Bilateral cardiac sympathetic denervation for the management of electrical storm. *J Am Coll Cardiol* 2012;59:91-2.

99. Puddu PE, Jouve R, Langlet F, Guillen JC, Lanti M, Reale A. Prevention of postischemic ventricular fibrillation late after right or left stellate ganglionectomy in dogs. *Circulation* 1988;77:935-46.

100. Vaseghi M, Gima J, Kanaan C, et al. Cardiac sympathetic denervation in patients with refractory ventricular arrhythmias or electrical storm: intermediate and long-term follow-up. *Heart Rhythm* 2014;11:360-6.

101. Vaseghi M, Barwad P, Malavassi Corrales FJ, et al. Cardiac sympathetic denervation for refractory ventricular arrhythmias. *J Am Coll Cardiol* 2017;69:3070-80.

102. Hayase J, Patel J, Narayan SM, Krummen DE. Percutaneous stellate ganglion block suppressing VT and VF in a patient refractory to VT ablation. *J Cardiovasc Electrophysiol* 2013;24:926-8.

103. Tan AY, Abdi S, Buxton AE, Anter E. Percutaneous stellate ganglia block for acute control of refractory ventricular tachycardia. *Heart Rhythm* 2012;9:2063-7.

104. Fudim M, Boortz-Marx R, Ganesh A, et al. Stellate ganglion blockade for the treatment of refractory ventricular arrhythmias: A systematic review and meta-analysis. *J Cardiovasc Electrophysiol* 2017;28:1460-7.

105. Nademanee K, Taylor R, Bailey WE, Rieders DE, Kosar EM. Treating electrical storm: sympathetic blockade versus advanced cardiac life support-guided therapy. *Circulation* 2000;102:742-7.

106. Fudim M, Boortz-Marx R, Patel CB, Sun AY, Piccini JP. Autonomic modulation for the treatment of ventricular arrhythmias: therapeutic use of percutaneous stellate ganglion blocks. *J Cardiovasc Electrophysiol* 2017;28:446-9.

107. Meissner A, Eckardt L, Kirchhoff P, et al. Effects of thoracic epidural anesthesia with and without autonomic nervous system blockade on cardiac monophasic action potentials and effective refractoriness in awake dogs. *Anesthesiology* 2001;95:132-8. discussion 6A.

108. Kamibayashi T, Hayashi Y, Mammoto T, Yamatodani A, Taenaka N, Yoshiya I. Thoracic epidural anesthesia attenuates halothane-induced myocardial sensitization to dysrhythmogenic effect of epinephrine in dogs. *Anesthesiology* 1995;82:129-34.

109. Bourke T, Vaseghi M, Michowitz Y, et al. Neuraxial modulation for refractory ventricular arrhythmias: value of thoracic epidural anesthesia and surgical left cardiac sympathetic denervation. *Circulation* 2010;121:2255-62.

110. Do DH, Bradfield J, Ajjola OA, et al. Thoracic epidural anesthesia can be effective for the short-term management of ventricular tachycardia storm. *J Am Heart Assoc* 2017;6. pii:e007080.

111. Wang S, Zhou X, Huang B, et al. Spinal cord stimulation protects against ventricular arrhythmias by suppressing left stellate ganglion neural activity in an acute myocardial infarction canine model. *Heart Rhythm* 2015;12:1628-35.

112. Odenstedt J, Linderöth B, Bergfeldt L, et al. Spinal cord stimulation effects on myocardial ischemia, infarct size, ventricular arrhythmia, and

- noninvasive electrophysiology in a porcine ischemia-reperfusion model. *Heart Rhythm* 2011;8:892-8.
113. Grimaldi R, de Luca A, Kornet L, Castagno D, Gaita F. Can spinal cord stimulation reduce ventricular arrhythmias? *Heart Rhythm* 2012;9:1884-7.
 114. Zipes DP, Neuzil P, Theres H, et al. Determining the feasibility of spinal cord neuromodulation for the treatment of chronic systolic heart failure: the DEFEAT-HF study. *J Am Coll Cardiol HF* 2016;4:129-36.
 115. Janes RD, Brandys JC, Hopkins DA, Johnstone DE, Murphy DA, Armour JA. Anatomy of human extrinsic cardiac nerves and ganglia. *Am J Cardiol* 1986;57:299-309.
 116. Assis FR, Yu DH, Zhou X, et al. Minimally invasive transtracheal cardiac plexus block for sympathetic neuromodulation. *Heart Rhythm* 2019;16:117-24.
 117. Sobotka PA, Harrison DG, Fudim M. The endpoint on measuring the clinical effects of renal denervation: what are the best surrogates. In: Heuser R, Schlaich M, Sievert H, editors. *Renal Denervation*. London: Springer, London, 2015.
 118. Linz D, Wirth K, Ukena C, et al. Renal denervation suppresses ventricular arrhythmias during acute ventricular ischemia in pigs. *Heart Rhythm* 2013;10:1525-30.
 119. Remo BF, Preminger M, Bradfield J, et al. Safety and efficacy of renal denervation as a novel treatment of ventricular tachycardia storm in patients with cardiomyopathy. *Heart Rhythm* 2014;11:541-6.
 120. Ukena C, Mahfoud F, Ewen S, et al. Renal denervation for treatment of ventricular arrhythmias: data from an International Multicenter Registry. *Clin Res Cardiol* 2016;105:873-9.
 121. Armaganijan LV, Staico R, Moreira DA, et al. 6-Month outcomes in patients with implantable cardioverter-defibrillators undergoing renal sympathetic denervation for the treatment of refractory ventricular arrhythmias. *J Am Coll Cardiol Intv* 2015;8:984-90.
 122. Kandzari DE, Bohm M, Mahfoud F, et al. Effect of renal denervation on blood pressure in the presence of antihypertensive drugs: 6-month efficacy and safety results from the SPYRAL HTN-ON MED proof-of-concept randomised trial. *Lancet* 2018;391:2346-55.
 123. Yu L, Wang S, Zhou X, et al. Chronic intermittent low-level stimulation of tragus reduces cardiac autonomic remodeling and ventricular arrhythmia inducibility in a post-infarction canine model. *J Am Coll Cardiol EP* 2016;2:330-9.
 124. Myers RW, Pearlman AS, Hyman RM, et al. Beneficial effects of vagal stimulation and bradycardia during experimental acute myocardial ischemia. *Circulation* 1974;49:943-7.
 125. Vanoli E, De Ferrari GM, Stramba-Badiale M, Hull SS Jr., Foreman RD, Schwartz PJ. Vagal stimulation and prevention of sudden death in conscious dogs with a healed myocardial infarction. *Circ Res* 1991;68:1471-81.
 126. Yu L, Zhou L, Cao G, et al. Optogenetic modulation of cardiac sympathetic nerve activity to prevent ventricular arrhythmias. *J Am Coll Cardiol* 2017;70:2778-90.
 127. Deisseroth K. Optogenetics: 10 years of microbial opsins in neuroscience. *Nat Neurosci* 2015;18:1213-25.
 128. Tsunematsu T, Tabuchi S, Tanaka KF, Boyden ES, Tominaga M, Yamanaka A. Long-lasting silencing of orexin/hypocretin neurons using archaerhodopsin induces slow-wave sleep in mice. *Behav Brain Res* 2013;255:64-74.
 129. Chow BY, Han X, Dobry AS, et al. High-performance genetically targetable optical neural silencing by light-driven proton pumps. *Nature* 2010;463:98-102.
 130. Wengrowski AM, Wang X, Tapa S, Posnack NG, Mendelowitz D, Kay MW. Optogenetic release of norepinephrine from cardiac sympathetic neurons alters mechanical and electrical function. *Cardiovasc Res* 2015;105:143-50.
 131. Chin A, Ntsekhe M, Viljoen C, Rossouw J, Pennel T, Schwartz PJ. Rationale and design of a prospective study to assess the effect of left cardiac sympathetic denervation in chronic heart failure. *Int J Cardiol* 2017;248:227-31.
 132. Bradfield JS, Ajjola OA, Vaseghi M, Shivkumar K. Mechanisms and management of refractory ventricular arrhythmias in the age of autonomic modulation. *Heart Rhythm* 2018;15:1252-60.
 133. Chatterjee NA, Singh JP. Novel interventional therapies modulate the autonomic tone in heart failure. *J Am Coll Cardiol HF* 2015;3:786-802.

KEY WORDS arrhythmia, atrial fibrillation, autonomic nervous system, ganglionated plexi, neuromodulation, ventricular arrhythmias

