



## Full Length Article

## Gut microbiome: An intermediary to neurotoxicity

Joseph L. Dempsey, Mallory Little, Julia Yue Cui\*

Department of Environmental and Occupational Health Sciences, University of Washington, United States



## ARTICLE INFO

**Keywords:**  
Gut microbiome  
Neurotoxicants  
Gut-brain axis

## ABSTRACT

There is growing recognition that the gut microbiome is an important regulator for neurological functions. This review provides a summary on the role of gut microbiota in various neurological disorders including neurotoxicity induced by environmental stressors such as drugs, environmental contaminants, and dietary factors. We propose that the gut microbiome remotely senses and regulates CNS signaling through the following mechanisms: 1) intestinal bacteria-mediated biotransformation of neurotoxicants that alters the neuro-reactivity of the parent compounds; 2) altered production of neuro-reactive microbial metabolites following exposure to certain environmental stressors; 3) bi-directional communication within the gut-brain axis to alter the intestinal barrier integrity; and 4) regulation of mucosal immune function. Distinct microbial metabolites may enter systemic circulation and epigenetically reprogram the expression of host genes in the CNS, regulating neuroinflammation, cell survival, or cell death. We will also review the current tools for the study of the gut-brain axis and provide some suggestions to move this field forward in the future.

## 1. Introduction

The gut-brain axis is increasingly recognized as an important target for the health of the central nervous system (CNS) (Carabotti et al., 2015; Sharon et al., 2016; Skonieczna-Zydecka et al., 2018; Zhu et al., 2017). As a multidirectional communication network, the key components of the gut-brain axis include the CNS, the autonomic nervous system (ANS), the enteric nervous system (ENS), as well as the hypothalamic pituitary adrenal axis (HPA) (Carabotti et al., 2015). The CNS is crucial for cognitive functions including memory, social, and emotional responses, and it communicates with ANS, ENS, and HPA to orchestrate signal transduction. The ENS regulates the production of intestinal hormones and mucus secretion. In addition, the ENS and enteric immune system interact to maintain gut integrity (Yoo and Mazmanian, 2017). The ENS can polarize macrophages and put them in close proximity to extrinsic and mucosal nerve fibers (Gabanyi et al., 2016). The bioavailability of catecholamines—a sympathetic neurotransmitters and immune modulator—can be modified by gut bacteria; catecholamines are glucuronidated for excretion, whereas some gut bacteria can de-glucuronidate catecholamines and possibly affect leukocytes in the gut (Asano et al., 2012; Yoo and Mazmanian, 2017). Gut permeability and integrity can also be regulated and monitored by the ENS through mechanoreceptors responsive to mucosal abrasions, intrinsic primary afferent neurons responsive to molecular and mechanical aberrations, and tension receptors responsive to stretch. Several

compounds, including short-chain fatty acids from bacteria, activate receptors in neurons that regulate gut motility (Cherbut et al., 1998). The HPA modulates the release of cortisol from adrenal glands during stress response. In addition to host signaling, the gut microbiome is important in the bidirectional communications through modulating gastrointestinal tract (GI) functions, is remotely signaling to the brain and other metabolic organs, and is one of many regulatory targets of brain-to-gut signaling (Carabotti et al., 2015; Fu and Cui, 2017; Sharon et al., 2016; Skonieczna-Zydecka et al., 2018; Zhu et al., 2017).

The human microbiome dwarfs the human genome with a 10:1 ratio of nucleated cells, and 100:1 ratio of genes (Sender et al., 2016). The microorganisms in the GI tract represent the majority of the human microbiome, including bacteria, fungi, parasites, and viruses (Zhu et al., 2017). This review focuses primarily on bacteria as they are the best characterized members of the microbiome at this time. At the phylum level, the dominant intestinal bacteria are Firmicutes and Bacteroidetes in healthy human subjects, whereas other minor phyla include Proteobacteria, Actinomycetes, Verrucomicrobia, and Fusobacteria (Eckburg et al., 2005). As illustrated in Fig. 1, following exposure to environmental stressors such as environmental contaminants, drugs, dietary factors, and other xenobiotics, the gut microbiome is thought to modify the toxicological outcomes through several mechanisms. 1) Gut microbiome can directly metabolize neurotoxicants primarily through reduction and hydrolysis/de-conjugation reactions (Claus et al., 2016; Lu et al., 2015); the gut microbiome may communicate with the liver

\* Corresponding author.

E-mail address: [juliacui@uw.edu](mailto:juliacui@uw.edu) (J.Y. Cui).<https://doi.org/10.1016/j.neuro.2019.08.005>

Received 26 March 2019; Received in revised form 4 August 2019; Accepted 16 August 2019

Available online 24 August 2019

0161-813X/ © 2019 Elsevier B.V. All rights reserved.




Fig. 1. An illustration of the gut-brain axis in environmental stressor induced neurotoxicity.

through the enterohepatic circulation of primary and secondary metabolites, altering hepatic xenobiotic biotransformation and nutrient homeostasis, which are the major functions of the liver (Klaassen and Cui, 2015; Spanogiannopoulos et al., 2016; Swanson, 2015; Visschers et al., 2013). 2) Intestinal dysbiosis as a result of chemical exposure may lead to local inflammation and gut leakiness, subsequently increasing levels of pro-inflammatory cytokines in the systemic circulation, which may contribute to neuroinflammation (Fournier et al., 2018; Janakiraman and Krishnamoorthy, 2018; Lin et al., 2018; Rea et al., 2016; Sampson et al., 2016). 3) Gut microbiota may produce neuro-reactive microbial metabolites, including short-chain fatty acids (SCFAs), ursodeoxycholic acid (UDCA), as well as various neurotransmitters such as gamma-aminobutyric acid (GABA), histamine, acetylcholine, serotonin, melatonin, gut lumen-derived bioreactive free catecholamines, nitric oxide, and hydrogen sulfide (Asano et al., 2012; Iyer et al., 2004; Schicho et al., 2006; Sobko et al., 2006). These microbial metabolites may enter the systemic circulation and reach the molecular targets in the brain to modulate various types of cognitive functions (Carabotti et al., 2015).

In this review, we will first discuss the role of the gut microbiome in various types of neurological diseases, including Alzheimer's disease (AD) and Parkinson's disease (PD), autism spectrum disorder, attention-deficit hyperactivity disorder (ADHD), anorexia nervosa, depression, schizophrenia, bipolar disorders, as well as post-traumatic stress disorder (Table 1). We will then provide a comprehensive analysis of the literature regarding the interactions between various neurotoxicants and gut microbiome (Table 2). These neurotoxicants are selected based on research articles as well as the Casarett and Doull's Toxicology Textbook (Klaassen CD 2013), including various drugs and environmental contaminants, and are summarized in Table 3. While a substantial portion of the research body in this area still remains at the

association level, several mechanistic studies have been performed, taking advantage of novel research tools, including germ-free (GF) models, fecal/single microbial strain inoculations, anaerobic cultures, and microbial metabolite supplementations (Table 2). We will also discuss the challenges and opportunities characterizing the gut-brain axis in neurotoxicology.

## 2. The gut-brain axis and neurological diseases

### 2.1. Neurodevelopmental disorders

#### 2.1.1. Attention-deficit hyperactivity disorder

Attention-deficit hyperactivity disorder (ADHD) is a common condition characterized by inattention, hyperactivity, and/or impulsivity that affects 5% of children and 2.5% of adults (Faraone et al., 2015). Factors that increase the risk of ADHD include age, sex (4:1 male to female) (Faraone and Glatt, 2010), socioeconomic status (low family income) (Larsson et al., 2014), genetic variants, as well as environmental factors such as parental behavior (Harold et al., 2013; Stevens et al., 2008), prenatal metrics, and some environmental contaminants including organophosphate pesticides, polychlorinated biphenyls (PCBs), and lead (Banerjee et al., 2007; Scassellati et al., 2012). Recent studies demonstrate that the composition and predicted functions of gut microbiome are altered in ADHD patients. In one study, a decrease in parent-reported ADHD symptoms was associated with an increase in the abundance of *Faecalibacterium* spp. with no alteration in the richness of the gut microbiota (Jiang et al., 2018). A second study of male juveniles found that relative to healthy controls, subjects with ADHD had lower species richness as well as an increase in *Bacteroidaceae* (Prehn-Kristensen et al., 2018). Young adult subjects with ADHD had increased *Bifidobacterium* spp. associated with a predicted increase in

**Table 1**  
Studies investigating neurological diseases in the gut-brain axis.

Disease	Model	Bacteria influence	Source
ADHD	Human adolescents and adults	Association	(Aarts et al., 2017)
	Human children	Association (biomarker)	(Jiang et al., 2018)
	Humans: juvenile males	Association (diversity; biomarker)	(Prehn-Kristensen et al., 2018)
	Humans: cesarean delivery or antibiotic use during first two years of life	No association	(Axelsson et al., 2019a)
Alzheimer's disease	Mice	Association (diversity)	(Hoffman et al., 2017)
	Artificial A $\beta$ aggregation assays	Inverse association with SCFAs	(Ho et al., 2018)
Anorexia nervosa	Human adult females	Association	(Borgo et al., 2017; Kleiman et al., 2015; Mack et al., 2016; Morita et al., 2015; Morkl et al., 2017)
Autism	Mice	Association	(Chen et al., 2016)
	Human children	No association	(Gondalia et al., 2012)
	Human children	Association	(Coretti et al., 2018; De Angelis et al., 2013; Finegold et al., 2010; Hicks et al., 2018; Kang et al., 2013; Liu et al., 2019; Pulikkan et al., 2018; Qiao et al., 2018; Rose et al., 2018; Son et al., 2015; Wang et al., 2019a)
	Mice valproic acid-induced	Association	(de Theije et al., 2014)
	BTBR T <sup>+</sup> Itpr3 <sup>fl</sup> /J mouse model of ASD	Association	(Coretti et al., 2017; Golubeva et al., 2017)
	Juvenile hamsters clindamycin and propionic acid induced	Mechanistic	(El-Ansary et al., 2018)
	Shank3-null mice	Mechanistic	(Tabouy et al., 2018)
	Rat valproic acid-induced	Association (biomarker)	(Liu et al., 2018)
Autism and antibiotics	Human children in Manitoba, Canada or Denmark	No association	(Axelsson et al., 2019b; Hamad et al., 2018)
Autism and Vitamin A Behavior - generic germ-free	Human children with autism	Association	(Liu et al., 2017a)
	Germ-free mice	Association	(Arentsen et al., 2015; Lu et al., 2018; Neufeld et al., 2011; Sudo et al., 2004)
	Germ-free mice	Mechanistic	(Diaz Heijtz et al., 2011)
	Germ-free, stress-sensitive rats	Association	(Crumevolle-Arias et al., 2014)
	Antibiotic-exposed mice	Association	(Desbonnet et al., 2015)
Bipolar	Human adults	Association	(Evans et al., 2017; Painold et al., 2018; Vinberg et al., 2019)
	Human adults	Mechanistic	(Painold et al., 2018)
Depression	Mice	Mechanistic	(Sun et al., 2018a; Zheng et al., 2016) 30521978 (Huang et al., 2019)
	Rats	Association	(Tillmann et al., 2018; Yu et al., 2017)
	Rats	Mechanistic	(Abildgaard et al., 2017; Kelly et al., 2016)
	Human adults	Association	(Aizawa et al., 2016; Chen et al., 2018d, c; Jiang et al., 2015; Lurie et al., 2015; Naseribafrouei et al., 2014)
Parkinson's disease	Human adults	Mechanistic	(Kazemi et al., 2018; Miyaoka et al., 2018; Rudzki et al., 2019)
	Human adults	Association	(Hill-Burns et al., 2017; Unger et al., 2016)
	Mice	Mechanistic	(Sampson et al., 2016)
Postpartum depression	Adult women	Mechanistic	(Slykerman et al., 2017)
	Pregnant women	Association	(Murphy et al., 2018)
PTSD	Humans	Association	(Hemmings et al., 2017)
	Male mice	Association	(Gautam et al., 2015, 2018)
Schizophrenia	Rats	Association	(Dunphy-Doherty et al., 2018; Pyndt Jorgensen et al., 2015)
	Humans	Association	(Castro-Nallar et al., 2015; Flowers et al., 2019; Nagamine et al., 2018; Nguyen et al., 2019; Olde Loohuis et al., 2018; Shen et al., 2018)
	Humans	Mechanistic	(Okubo et al., 2019)
	Mice	Mechanistic	(Zheng et al., 2019)

cyclohexadienyl dehydratase I, which is important for the generation of the dopamine and noradrenaline precursor phenylalanine (Aarts et al., 2017). Dysregulation of dopamine and noradrenaline is implicated in ADHD etiology; however, it is unclear whether bacteria can independently induce ADHD-like behavior.

### 2.1.2. Autism spectrum disorder

Autism spectrum disorder (ASD) is a severe neurodevelopmental disorder characterized by persistent social and communicative deficits as well as repetitive and restrictive patterns of behaviors, interests, and activities. It is increasingly recognized that both environmental and genetic factors play important roles in the etiology of ASD (Hallmayer et al., 2011), including the gut microbiome, and ASD in children (Coretti et al., 2018; De Angelis et al., 2013; Finegold et al., 2010; Hicks et al., 2018; Kang et al., 2013; Liu et al., 2019; Pulikkan et al., 2018; Qiao et al., 2018; Rose et al., 2018; Son et al., 2015; Wang et al., 2019a). Interestingly, there is a high prevalence of gastrointestinal disorders in children with ASD. Gastrointestinal-related symptoms in ASD were associated with a less diverse gut microbiome and had decreased abundances of the genera *Prevotella*, *Coprococcus*, and an

unclassified taxon in *Veillonellaceae*, which are carbohydrate-degrading and fermenting bacteria, suggesting that carbohydrate metabolism or SCFAs may be involved in ASD (Kang et al., 2013). In ASD patients, a positive correlation was identified between high levels of the SCFA butyrate and *Faecalibacterium prausnitzii* (Coretti et al., 2018). Male mice perinatally exposed to valproic acid to induce ASD-like behaviors also had increased caecum butyrate (de Theije et al., 2014). ASD children were found to have higher fecal abundances of *Caloramator*, *Sarcina* and *Clostridium* genera along with a decrease in *Bifidobacterium*, which is known to produce SCFAs (De Angelis et al., 2013). A similar study comparing children with ASD and neurotypical healthy controls found no differences in fecal microbiome diversity, but did identify a significant interaction between ASD and the *Cyanobacteria/Chloroplast* genus (Son et al., 2015). A comparison of Indian children with ASD relative to healthy controls identified ASD as a significant factor to explain the differences in gut phylotypes; *Lactobacillus* spp. was significantly associated with ASD using a meta-analysis of the Indian children cohort with a US cohort (Pulikkan et al., 2018). Behavioral tests and hazard models from studies investigating antibiotics or vitamin A and ASD do not support a causal relationship (Axelsson et al.,

**Table 2**  
Studies investigating neurotoxicants and microbiome.

Category	Toxicant	Model	Dose or exposure	End point and bacteria result	Source
Air pollution	PM <sub>2.5</sub>	Mouse (8-12 weeks old)	135.4 ± 6.4 mg/m <sup>3</sup> 8 hours for 5 days/week for 3 weeks	Increased genera, as well as alpha and beta diversity	(Mutlu et al., 2018)
	PM <sub>10</sub>	Mouse (6-8 weeks old)	18 µg/g/day (oral gavage)	Altered short chain fatty acid concentrations and microbial composition	(Kish et al., 2013)
Dietary and herbal medicine	Fructooligosaccharides	Rat (adults given 100 mg/kg/d d-galactose i.p. for inducing AD)	100 mg/kg/d (oral gavage)	Maintains gut microbiota diversity while improving neurological endpoints	(Chen et al., 2017)
	Ketogenic diet	Mouse (adult 6-Hz induced seizure model of refractory epilepsy)	6:1 fat:protein ketogenic diet ( <i>ad libitum</i> )	Increased hippocampal GABA/glutamate levels mechanistically by <i>Akkermansia</i> and <i>Parabacteroides</i>	(Olson et al., 2018)
Drugs	Amiodarone	Rat (adult)	Probiotic: 1.5 × 10 <sup>9</sup> CFU/dose once daily for 7 days Amiodarone: 50 mg/kg (single oral dose)	<i>E. coli</i> Nissle 1917 increased the bioavailability whereas <i>L. casei</i> DN-114 001 delayed max plasma concentration	(Matuskova et al., 2017, 2014)
	Chemotherapeutics	Bacteria ( <i>in vitro</i> )	Variable—dose-response relationship (screens to identify bacteria susceptible to chemotherapeutics)	Lactic acid bacteria and bifidobacteria may be susceptible whereas other bacteria may affect efficacy	(Florez et al., 2016; Lehouritis et al., 2015)
	Cisplatin	Mouse (8-12 weeks)	Cisplatin: 10 mg/kg Probiotic: healthy donor fecal pellet or 2 × 10 <sup>8</sup> <i>Ruminococcus gnavus</i>	Probiotic ameliorated intestinal toxicity and systemic inflammation of cisplatin	(Perales-Puchalt et al., 2018)
	Disulfiram	Bacteria ( <i>in vitro</i> )	Variable (disulfiram and metabolites)	Antimicrobial activity against gram-positive bacteria	(Frazier et al., 2019; Long, 2017; Sheppard et al., 2018)
	Doxorubicin	Mouse (8-10-week-old females)	20 mg/kg by i.p.	Enteric injury (crypt depth, crypt number, and proliferative cell number) is dependent on presence of bacteria	(Rigby et al., 2016)
		Bacteria ( <i>in vitro</i> )	150 µg/mL	Deglycosylation of doxorubicin reduces toxicity	(Yan et al., 2018)
		Mouse (7-10-week-old females)	10 mg/kg by i.p.	Pre-treatment with pectin (1.5 mg per day orally) protected against ileitis independent of SCFAs	(Sahasrabudhe et al., 2018)
	Lithium, valproate and aripiprazole	Rat (adult)	Variable	Increased microbial species richness and diversity; some antimicrobial activity	(Cussotto et al., 2018)
	Metronidazole	Rat (adult males)	1 mg/ml in drinking water for 1 week	Increased in <i>Bifidobacterium spp.</i> and <i>Enterobacteriaceae</i> and increased mucosal thickness	(Pelissier et al., 2010)
	MPTP	Mouse (8-week-old males)	30 mg/kg by i.p.	Decreased Firmicutes and order Clostridiales; increased Proteobacteria, Turicibacterales and Enterobacteriales; PD-like effects abrogated by FMT	(Sun et al., 2018b)
Mouse (male and/or female adults, WT and metabotropic glutamate receptors knock-out)		18 mg/kg 2 times per week for 5 weeks 10 mg/kg for 5 days, rest 2 days, 20 mg/kg for 5 days by i.p.	Altered microbiome composition; Gender- and genotype-specific results; associations between microbiome diversity and sensorimotor performance	(Lai et al., 2018; Torres et al., 2018)	
Nano-encapsulated doxorubicin and paclitaxel	Mouse (8-week-old males)	12.93 µmol/kg	Altered microbial activity indicated by levels of hippurate and indoxyl sulfate	(Song et al., 2015)	
Nitrofurantoin	Human (adults)	100 mg twice per day	Decreased <i>Clostridium sp.</i> and increased <i>Faecalibacterium sp.</i>	(Stewardson et al., 2015; Vervoort et al., 2015)	
Metals	Arsenic	Mouse (8-week-old males and females)	100 ppb sodium arsenite for 13 weeks Up to 250 ppb Antibiotic-treated for 3 days then exposed up to 1 ppm arsenic in drinking water for 2 weeks (females only)	Altered gene abundances for genes involved in carbohydrate metabolism, pyruvate fermentation, short-chain fatty acid synthesis, and starch utilization Arsenic eroded bacterial biofilms adjacent to the mucosa	(Chi et al., 2016, 2017a; Chi et al., 2019; Dheer et al., 2015; Lu et al., 2014)
		Mouse (7-13-week-old males and females)	25 and 100 ppm sodium arsenate	1 ppm arsenic did not alter gut microbiome Antibiotic-treated and GF mice accumulate more arsenic than controls; human fecal transplants protect mice lacking the arsenic detoxification enzyme from arsenic-induced mortality, but may depend on <i>Faecalibacterium spp.</i>	(Coryell et al., 2018)
			Devon Great Consols (DGC) mine site		(Pass et al., 2015)

(continued on next page)

Table 2 (continued)

Category	Toxicant	Model	Dose or exposure	End point and bacteria result	Source
		Earthworm ( <i>Lumbricus rubellus</i> )		Slight trend for an association between worm microbiome diversity and arsenic contamination	
		Human ( <i>in vitro</i> ; SHIME)	Soil arsenic concentrations (varies) 100 µg/L As, 600 µg/L As, 600 µg/L As and 0.1 mg/L Fe, 600 µg/L As and 0.3 mg/L Fe, and 600 µg/L and As + 3 mg/L Fe	Bioaccessibility in the Arsenic was 1.8–2.8 times more bioaccessible in the colon than in the small intestinal phase Iron decreased bioaccessibility of arsenic with increased arsenic methylation;	(Yin et al., 2015; Yu et al., 2016)
		Human ( <i>in vitro</i> ; Caco-2 cells)	Soil arsenic samples ranging from 15.5 to 3225.6 mg/kg	Gut microbiota can directly release soil arsenic; arsenic in colon is digested more quickly than in soil	(Yin et al., 2017)
		Human	Children: high group (218.8 µg/L in drinking water) and low group (1.7 µg/L in drinking water) Infants at 6 weeks of age Adult men and women	Children: high arsenic group had increased Gammaproteobacteria class, Enterobacteriales order, and <i>Enterobacteriaceae</i> family Infants: 8 genera were enriched with higher arsenic exposure, whereas 15 genera were decreased; changes were associated with males, but not females Time-weighted urinary arsenic associated with order Campylobacteriales and the genera <i>Anaerostipes</i> and <i>Faecalibacterium</i>	(Dong et al., 2017) (Hoen et al., 2018) (Wu et al., 2019)
		Zebrafish (larva)	10, 50, and 100 ppb for 20 days	Increased the genera <i>Acinetobacter</i> , <i>Sediminibacterium</i> , and <i>Janthinobacterium</i> ; decreased the genera <i>Bdellovibrio</i> and <i>Pseudomonas</i>	(Dahan et al., 2018)
Copper		Broiler chicken (Start from Day 0 as chicks)	8 or 187.5 mg/kg of Cu from Cu Sul or 187.5 mg/kg of Cu from TBCC	No affect on performance by Cu source or concentration	(Pang et al., 2009)
		Holstein-Friesian calves (Dairy cows; 7 month old males)	3 g/100 L of copper supplementation as cupric sulphate in drinking water for 75 days	Increased microbial alpha diversity in rumen; altered bacteria <i>S24-7</i> , <i>Planctomycetaceae p-1088-a5</i> gut group, and <i>Azospira spp.</i>	(Biscarini et al., 2018)
Gold		Mouse (7-8-week-old males)	Up to 25 µg gold/kg bodyweight for 8 7 different types of gold nanoparticles for 8 days during and after 5-day dextran sodium sulfate exposure	5 nm/Citrate and Au-5 nm/polyvinylpyrrolidone attenuated colonic and systemic inflammation; gold nanoparticles decreased alpha diversity;	(Zhu et al., 2018b)
Lead		Mouse	8 week old females: 10 ppm for 13 weeks; 32 ppm to dams in drinking water 2 weeks prior to breeding and up to 40 weeks of age for pups 6-week-old males and females: 0.01, 0.03, or 0.1 mg/L Pb for 15 weeks	Majority of bacteria decreased following lead exposure; Cultivable aeobes decreased and anaerobes increased;	(Gao et al., 2017c; Wu et al., 2016; Xia et al., 2018a)
		Zebrafish (adult males)	10 and 30 µg/L for 7 days	at highest exposure, decreased Firmicutes and increased Bacteroidetes and Proteobacteria 52 microbes altered by 30 µg/L PB group; altered metabolites in pathways for glucose and lipid metabolism, amino acid metabolism, nucleotide metabolism	(Xia et al., 2018b)
		Carp ( <i>Cyprinus carpio</i> ; 105 days old)	1 mg/L	Pb decreased the expressions of pro-inflammatory cytokines; <i>Lactobacillus reuteri</i> P16 decreased mortality, improved the growth performance, and abrogated changes in gene expression	(Giri et al., 2018)
Magnesium		Mouse (8-week-old males)	Standard diet with 500 mg Mg/kg food and magnesium deficient diet with 50 mg Mg/kg food for 6 weeks	Increased depressive-like behavior with magnesium deficiency; the gut microbiome was altered by magnesium deficiency and was positively correlated with hippocampal interleukin-6	(Winther et al., 2015)
Manganese		Mouse (8-week-old males and females)	100 ppm in drinking water for 13 weeks	Altered microbial tryptophan and phenylalanine biosynthesis pathways; bacteria of male mice had altered GABA/putrescine metabolism	(Chi et al., 2017b)
Mercury		Mummichog ( <i>Fundulus heteroclitus</i> )	Total mercury in food: 0.08, 24.4, or 131 µg/g dry weight for 15 days	Microbial Hg resistance gene mercuric reductase 8 times higher in fish at Hg contamination site	(Lloyd et al., 2016)
Methylmercury		Human (36–39 weeks pregnant)	Hair: 57 ng total Hg/g hair; Stool: 150 (2.1–810) ng total mercury /g stool; Cord blood: 0.23 (0.061–0.73) µg MeHg/L cord blood	17 genera were correlated with mercury concentration in stool or hair	(Rothenberg et al., 2016)
		Fathead minnow and mouse	Minnow: 0.02, 0.72, 0.87, or 5.50 µg/g, mercury in food twice daily for 30 days Mice: 0.02 µg/g, 0.43 µg/g, or 4.39 ± 0.57 µg/g mercury in food twice daily for 30 days	Minnow gut microbiome adapted to detoxify MeHg; Mouse midbrain L-glutamine, O-phosphatidylcholine, dopamine, tagatose, hydroquinone, L-ascorbic acid, inosine 5'-monophosphate, and uracil decreased	(Bridges et al., 2018)
		Bacteria ( <i>in vitro</i> ; fresh human fecal samples in anaerobic chamber)	10 ng/g monomethylmercury and 1 ng/g mercury for 0, 12, 24, 36, and 48 hours	Monomethylmercury concentration decreased under a balanced or protein rich diet, but not a carbohydrate rich diet	(Guo et al., 2018a)

(continued on next page)

Table 2 (continued)

Category	Toxicant	Model	Dose or exposure	End point and bacteria result	Source
	Arsenic and zinc	Mouse (4-week-old females)	Zinc-adequate diet with 0, 50, or 500 ppb arsenic or zinc-deficient diet with 0, 50, or 500 ppb arsenic And drinking water with 0, 50, or 500 ppb sodium arsenite for 6 weeks	No interaction between arsenic exposure and zinc restriction; plasma zinc concentration was positively correlated with the genera <i>Shewanella</i> , <i>Rheinheimera</i> , and <i>Bifidobacterium</i>	(Gaulke et al., 2018)
	Arsenic, cadmium, cobalt, chromium, and nickel	Rat (adults)	15, 22, or 31 mg/kg/day sodium arsenite; 35, 54, or 85 mg/kg/day cadmium chloride; 44, 62, or 88 mg/kg/day sodium dichromate, 27, 47, or 82 mg/kg/day cobalt chloride, or 177, 232, or 300 mg/kg/day nickel chloride for 5 days	47 genera were affected by at least one metal exposure; nickel uniquely altered 25 genera; bacteria with higher iron importing genes were increased by arsenic and nickel	(Richardson et al., 2018)
	Lead and cadmium	Mouse (6-week-old females)	For 8 weeks: Cadmium: 20 or 100 ppm Lead: 100 or 500 ppm	Decreased <i>Lachnospiraceae</i> and increased <i>Lactobacillaceae</i> and <i>Erysipelotrichaceae</i>	(Breton et al., 2013)
	Mercury and copper	Mouse (8-week-old females)	5 mg/kg copper, 2 mg/kg mercury, or 2.5 mg/kg copper and 1 mg/kg mercury	Copper decreased <i>Rikenella spp.</i> , <i>Jeotgailcoccus spp.</i> , and <i>Staphylococcus spp.</i> ; mercury decreased <i>Rikenella spp.</i> , <i>Jeotgailcoccus spp.</i> , and <i>Staphylococcus spp.</i>	(Ruan et al., 2018)
	Mercury, lead, arsenic, and cadmium	Humans (children and pregnant women)	Lead (µg/L): 22.6 or 47.1 Mercury (nmol/L): 8.8 and 9.5 Arsenic (nmol/L): 3.0 and 6.5 Cadmium (nmol/L): 1.1 and 1.2	<i>Lactobacillus rhamnosus</i> GR-1 protected against mercury and arsenic blood levels in pregnant women; Increased blood lead levels was associated with increased <i>Succinivibrionaceae</i> and <i>Gammaproteobacteria</i>	(Bisanz et al., 2014)
Noise	Lead and PCBs	Human (males and females age 60-84)	Lead geometric mean of 2.17 µg/dL; many PCBs; geometric mean range of individual PCBs in blood of 7.86-66.99 ng/g lipid	Increased PCB-146 and lead concentrations result in lower Digit Symbol Coding Test of the Weschler Adult Intelligence Scale	(Przybyla et al., 2017)
	Noise	Mouse (3-month-old male senescence-accelerated mouse prone 8)	< 40 (background), 88, or 98 decibels for 30 days	Cognitive impairment and Aβ accumulation; changes similar to aged mice; decreased gut microbiota diversity; fecal transplant of noised-exposed mice induced epithelial integrity impairment and Aβ accumulation to unexposed mice	(Cui et al., 2018)
Environmental chemicals - PBDEs	BDE-47 and BDE-99	Mouse (9-week-old male CV and GF)	100 µmol/kg for 4 days	Lack of gut microbiome altered PBDE metabolite profiles; increased <i>Akkermansia muciniphila</i> and <i>Allobaculum spp.</i> ; BDE-99 increased unconjugated bile acids	(Li et al., 2018, 2017)
	DE-71	Zebrafish (male and female adult)	5.0 ng/L for 7 days	Decreased <i>Mycoplasma spp.</i> , <i>Ruminiclostridium spp.</i> , <i>unclassified Firmicutes sensu stricto spp.</i> , and <i>Fusobacterium spp.</i> ; disrupted intestinal neural signaling, epithelial barrier integrity in males	(Chen et al., 2018a)
Environmental chemicals - PCBs	PCB-126	Mouse (8-week-old male <i>Ldlr</i> <sup>-/-</sup> )	1 µmol/kg at weeks 2 and 4 of a 12 week study	Decreased <i>S24.7</i> , <i>Clostridiales</i> , <i>Bifidobacterium spp.</i> , <i>Ruminococcus spp.</i> , <i>Oscillospira spp.</i> , and <i>Lactobacillus spp.</i> and increased <i>Akkermansia spp.</i> ; positive associations between <i>Bifidobacterium spp.</i> and GLP-1, as well as <i>Akkermansia spp.</i> and fasting blood glucose	(Petriello et al., 2018)
	Multiple	Mouse	Adult females: Varied up to 50 mg/kg 11-13 month old males with voluntary exercise 5 weeks prior: 150 µmol/kg	<i>Parasutterella</i> , <i>Ruminococcus</i> , <i>Prevotellaceae</i> UCG-001, <i>Alloprevotella</i> and <i>Parabacteroides</i> were decreased by PCBs; exercise attenuated PCB-induced changes in gut microbiome	(Chi et al., 2018; Choi et al., 2013)
Pesticide - Carbamates	Aldicarb	Mouse (8-week-old males)	2 ppm for 13 weeks in drinking water	Increased genes involved in virulence, adhesion, bacteriocins, antioxidant defense, protein degradation, DNA repair	(Gao et al., 2018b)
	Many	Bacteria from rumen of Holstein dairy cows ( <i>in vitro</i> )	Concentration varied; screened for bacteria that degrade carbamates through esterase activity	26 isolates had esterase activity and degraded at least 1 polyurethane and pesticide carbamate	(Ufarte et al., 2017)
Pesticide - Neonicotinoid	Imidacloprid	Fruit fly ( <i>Drosophila melanogaster</i> ; adults and larva)	10, 50, and 100 µM in food	Increased <i>Acetobacter spp.</i> and <i>Lactobacillus spp.</i> ; survival decreased with co-exposure to bacterial infection or heat stress	(Daisley et al., 2017)
		Honey bee ( <i>Apis mellifera</i> ; adults)	500 µg/liter imidacloprid suspended in sterilized sucrose syrup for 3 days	Little or no impact on the gut microbiome of adult worker bees	(Raymann et al., 2018)
Pesticide - Organochlorine	p, p'-DDE and β-HCH	Mouse (adult males)	p, p'-DDE: 1 mg/kg body weight/day β-HCH: 10 mg/kg body weight/day	Both chemicals decreased the genera <i>Parabacteroides</i> , <i>Prevotella</i> , <i>Bacteroides</i> , <i>Clostridium XIVa</i> and <i>Clostridium IV</i> and increased <i>Barnesiella</i> , <i>Alloprevotella</i> , <i>Oscillibacter</i> , <i>Lactobacillus</i> , <i>Parasutterella</i> and <i>Akkermansia</i>	(Liu et al., 2017b)
Pesticide - methyl	Azinphos-methyl	Human	0.021 to 6.192 ng azinphos-methyl/g blood serum	Decreased <i>Streptococcus</i> , <i>Micrococineae</i> , <i>Gemella</i> , <i>Haemophilus</i> , <i>Halomonas</i> , <i>Actinomycineae</i> , and <i>Granulicatella</i>	(Stanaway et al., 2017)

(continued on next page)

Table 2 (continued)

Category	Toxicant	Model	Dose or exposure	End point and bacteria result	Source
Organo-phosphate	Chlorpyrifos	Rat and human (in vitro; SHIME)	SHIME: 1 mg/day for 30 days Rat dams and pups: 1 mg/kg/day from gestation till 60 days of age	SHIME: increased <i>Enterococcus</i> spp. and <i>Bacteroides</i> spp.; decreased <i>Lactobacillus</i> spp. and <i>Bifidobacterium</i> spp. Rats: Decreased <i>Lactobacillus</i> spp. and <i>Bifidobacterium</i> spp.	(Joly et al., 2013)
		Rat	Adult males exposed to 0.3 or 3 mg/kg/day for 9 weeks Dams exposed orally from gestation to weaning and pups exposed thereafter at 1 or 3.5 mg/kg/day Dams exposed orally from gestation to weaning and pups exposed thereafter at 1 or 5 mg/kg/day	<i>Sutterella</i> spp. consistently enriched regardless of diet (normal or high-fat) In ileum, increased <i>Enterococcus</i> spp., <i>Clostridium</i> spp., <i>Staphylococcus</i> spp., and <i>Bacteroides</i> spp.; Decreased <i>Bifidobacterium</i> spp. in ileum and colon Inulin supplementation in drinking water abrogated chlorpyrifos-induced metabolic disorders in adults exposed in utero	(Fang et al., 2018; Joly Condette et al., 2015; Reygner et al., 2016b)
		Mouse (10-week-old male mice)	1 mg/kg/day for 30 days	Increase lipopolysaccharide and diamine oxidase in the serum; increased <i>Lactobacillaceae</i> and decreased <i>Bacteroidaceae</i>	(Zhao et al., 2016)
		Human (in vitro; SHIME)	0.35 and 1 mg/mL working solution	inulin co-treatment partially reversed dysbiosis and inhibited pro-inflammatory signaling when effluent was applied to Caco-2/TC7 intestinal cells; Increased in <i>Enterobacteriaceae</i> , <i>Bacteroides</i> spp. and <i>Clostridia</i>	(Requile et al., 2018; Reygner et al., 2016a)
	Diamondback moth larva ( <i>Plutella xylostella</i> )	Cabbage leaves dipped in 50 g/L solution for 10 minutes	Larva resistant to insecticides had increased levels of Lactobacillales order; Isolated from <i>P. xylostella</i> , <i>Enterococcus</i> sp. increased, <i>Serratia</i> sp. decreased, and <i>Enterobacter</i> sp. had not effect on insecticide resistance	(Xia et al., 2018c, 2013)	
	Fruit fly ( <i>Drosophila melanogaster</i> )	10 µM in food with or without <i>Lactobacillus rhamnosus</i> GG	<i>L. rhamnosus</i> GG prevented chlorpyrifos toxicity; abutbiotic-treated and GF flies live longer than CV flies; gut-derived <i>Lactobacillus plantarum</i> metabolizes chlorpyrifos to the oxon	(Daisley et al., 2018; Trinder et al., 2016)	
	Zebrafish (adult males)	30, 100 and 300 µg/L for 21 days	Altered 25 microbial genera and increased malondialdehyde and decreased glutathione in the gut	(Wang et al., 2019b)	
Pesticide - Pyrethrin	Diazinon	Mouse (8-week-old males and females)	4 mg/L in drinking water for 13 weeks	Altered expression of 677 microbial genes; disrupted quorum sensing, and enriched motility, sporulation, and stress response genes; sex-specific changes in altered bacteria	(Gao et al., 2017a, b)
	Fenitrothion	Mosquito ( <i>Anopheles albimanus</i> )	Captured mosquitos were exposed to bottles containing 50 µg fenitrothion and categorized as susceptible or resistant	Resistant mosquitos had a lower bacterial diversity but an enrichment for OP-degrading bacteria and enzymes	(Dada et al., 2018)
	Malathion	Mouse (8-week-old males)	2 mg/L in drinking water (~0.6 mg/kg/day) for 13 weeks	Enrichment of genes encoding virulence, mobility, and cell wall; altered quorum sensing	(Gao et al., 2018a)
	Monocrotophos and other OPs	Mouse (8-week-old females)	28 µg/kg/day in drinking water for 30 days	Mice given fecal microbiota of monocrotophos-exposed mice had significant blood glucose intolerance	(Velmurugan et al., 2017)
Pesticide - rotenoids	Rotenone	Rat (8-week-old)	2.75 mg/kg 5 days a week for 4 weeks	Decreased <i>Provatella</i> family, increased <i>Bacteroides-Prevotella-Porphyrmonas</i> spp. and <i>Bifidobacterium</i> spp.	(Nasuti et al., 2016)
		Mouse	15-17 week old males (after training and restrain stress): 10 mg/kg/day for 6 weeks 7-week-old males: 10 mg/kg/day for 4 weeks 8-9 week-old males: 30 mg/kg/day for 4 weeks	Butyrate exposure before treatment improves proinflammatory phenotype mediated by TLR4 Increased <i>Bifidobacterium</i> spp. in colon; changes in microbiota were consistent with PD patients Increased relative abundance of fecal <i>Akkermansia</i> Decreased <i>Bifidobacterium</i> spp. Gastrointestinal dysfunction and microbiome dysbiosis occurred before motor dysfunction; increased <i>Lactobacillus</i> spp. and decreased <i>Desulfovibrio</i> spp. associated with gastrointestinal dysfunction and motor dysfunction	(Seth et al., 2018) (Johnson et al., 2018) (Dodiya et al., 2018; Pereze-Pardo et al., 2018; Yang et al., 2017)
Pesticide-Herbicides	Atrazine	Zebrafish (4 months)	0.42 ± 0.02 µg/L for 7 days	Altered gut microbiota composition and function, but did not differ overall from control; decreased body weight and gonadosomatic index of females; induced intestinal inflammation in males	(Chen et al., 2018b)
			178.2 ± 7.8 µg/L for 6 days	No effect on gut bacteria of tadpoles or adults	

(continued on next page)

Table 2 (continued)

Category	Toxicant	Model	Dose or exposure	End point and bacteria result	Source
		Tree frog tadpoles and adults ( <i>Osteopilus spetentionalis</i> )			(Knutie et al., 2018)
	Glyphosate	Bacteria ( <i>in vitro</i> isolated from poultry)	5.0, 2.40, 1.20, 0.60, 0.30, 0.15 and 0.075 mg/ml	Pathogenic bacteria are resistant to glyphosate whereas beneficial bacteria are susceptible	(Shehata et al., 2013)
		<i>Enterococcus spp.</i> ( <i>in vitro</i> from cattle and horses)	Serial dilutions from 0.001 to 10 mg/mL	All tested <i>Enterococcus spp.</i> inhibit <i>Clostridium botulinum</i> ; higher concentrations of glyphosate inhibited <i>E. faecalis</i> growth but not <i>C. botulinum</i>	(Kruger et al., 2013)
		Rat	1.75 mg/kg bw/day from gestational day 6 to postnatal day 125 0.1 ppb, 400 ppm and 5000 ppm in drinking water in adults for 673 days 2.5 mg/kg/day in adults for 2 weeks	Increased <i>Prevotella spp.</i> and decreased <i>Lactobacillus spp.</i> Increased Bacteroidetes family S24-7 and a decreased Lactobacillaceae across all doses No observable short term effects; aromatic amino acids alleviate the antimicrobial effect of glyphosate	(Lozano et al., 2018; Mao et al., 2018; Nielsen et al., 2018)
		Cultural bacteria from green turtles ( <i>Chelonia mydas</i> )	0.00022, 0.00044, 0.056, 0.1125, 1.8, and 3.6 g/L of glyphosate from Rodeo® for 24 hours	Reduced growth and decreased survival at concentrations greater than 0.00022	(Kittle et al., 2018)
		Mouse (4-week-old males)	250 or 500 mg/kg/day for 1 day, 6 weeks, or 12 weeks	Increase of anxiety and depression-like behaviors associated with decreased Firmicutes, Bacteroidetes <i>Corynebacterium spp.</i> and <i>Lactobacillus spp.</i>	(Aitbali et al., 2018)
		Honey bee ( <i>Apis mellifera</i> ; larva and adults)	Larva: 0.8, 4, and 20 mg/L Adults: 5 and 10 mg/L	Larva: no effect on development rate, but higher doses decreased survival; decrease in beta diversity of 20 mg/L group; increased <i>Acidobacteria</i> and <i>Gemmatimonadaceae</i> Adult: sensitivity was dependent on microbiome containing an insensitive 5-enolpyruvylshikimate-3-phosphate synthase gene; increased for opportunistic pathogen <i>Serratia marcescens</i>	(Dai et al., 2018; Motta et al., 2018)
Pesticide - Fungicide	Copper sulfate	Piglets (28-days-old)	Up to 175 mg/kg food for 2 weeks	May increase villi and crypt depth in duodenum; decreased <i>Enterobacteriaceae</i> and <i>Streptococci spp.</i>	(Hojberg et al., 2005) (Di Giancamillo et al., 2018)
Pesticide - Mixture	Boscalid, captan, chlorpyrifos, thiofanate, thiacloprid, and ziram	Mouse (16-week-old males and females)	Ziram < 0.01 mg/kg food, chlorpyrifos 47 µg/kg food, thiacloprid 56 µg/kg food, boscalid 240 µg/kg food, thiofanate 205 µg/kg food, captan 165 µg/kg food for 52 weeks	In females, increased microbial associated metabolites 3-indoxyl sulphate and phenyl derivatives phenylacetyl glycine and p-cresol glucuronide	(Lukowicz et al., 2018)
	Coumaphos, tau-fluvalinate, and chlorothalonil	Honey bee ( <i>Apis mellifera</i> )	Colonies treated with tau-fluvalinate and coumaphos given strips with ~10% of the active ingredient; Colonies treated with chlorothalonil at 10 µg/L in 30% sucrose; all treatments for 6 weeks	Pesticide-dependent changes in microbiome and fungal communities; chlorothalonil increased genes for oxidative phosphorylation and decreased sugar and peptidase metabolism	(Kakumanu et al., 2016)
	Lambda-cyhalothrin, deltamethrin, chlorpyrifos ethyl, spinosad and lufenuron	Bacteria from insecticide-resistant fall armyworm ( <i>Spodoptera frugiperda</i> )	<i>In vitro</i> culturing with insecticide doses of 10, 20, 40, 80, and 160 µg/mL	16 microbial strains were isolated and shown to be resistant against at least 1 insecticide	(Almeida et al., 2017)
Plant and Animal Toxins	Bamboo (cyanide)	Giant panda ( <i>Ailuropoda melanoleuca</i> ) and red panda ( <i>Ailurus fulgens</i> )	N/A	Metagenome enriched with cyanide degrading enzymes; high abundance of <i>Pseudomonas</i>	(Zhu et al., 2018a)
	Domoic acid	Mollusks	N/A	Blue mussels ( <i>Mytilus edulis</i> ) and soft-shell clams ( <i>Mya arenaria</i> ) carry bacteria that can degrade domoic acid	(Stewart et al., 1998)
	Nicotine		60 mg/L for 13 weeks in drinking water		(Chi et al., 2017c)

(continued on next page)

Table 2 (continued)

Category	Toxicant	Model	Dose or exposure	End point and bacteria result	Source
SCFA	pro-pionic acid (PPA)	Mouse (8-week-old males and females)		Microbiome of male mice enriched for oxidative stress response, DNA repair genes, and acetate synthesis	(Frye et al., 2016)
		Human ( <i>in vitro</i> ; lymphoblastoid cell lines from male children with autism)	0.1, 0.5 and 1 mM	Mitochondrial function increased with PPA; however high PPA and long exposure duration increased proton leaking	
Solvents	Ethanol	Mouse (6-8-week-old females)	5% for 10 days and 5 g/kg via oral gavage 9 hours prior to euthanizing	Neuroinflammation and increased SI cytokines were abrogated in antibiotic-treated mice	(Lowe et al., 2018)
		Mouse (8-10-week-old males)	5% for 6 weeks and/or <i>Lactobacillus rhamnosus</i> GG $1 \times 10^9$ cfu daily	Increased <i>Alcaligenes</i> sp. and <i>Corynebacterium</i> sp.; <i>Lactobacillus rhamnosus</i> GG prevented hepatic injury	(Bull-Otterson et al., 2013)
	Human (adults)	Group mean 177-188 g/day	<i>Dorea</i> spp. and <i>Blautia</i> spp. were increased in alcohol dependent subjects and correlated with intestinal permeability	(Leclercq et al., 2014)	
	Mouse (7-week-old females)	Up to 20% for 8 weeks	Altered genera in <i>Lachnospiraceae</i> family and decreased <i>Alistipes</i> spp.; decreased <i>Clostridium</i> spp. with saccharin co-consumption; decreased <i>Adlercreutzia</i> spp. was positively correlated with alcohol preference	(Labrecque et al., 2015; Wang et al., 2018; Xu et al., 2018)	
	Rat (adults)	20% for 13 weeks (voluntary consumption)	Decreased microbiome diversity; decreased <i>Lactobacillus</i> spp.	(Kosnicki et al., 2018)	
Formaldehyde	Trichloroethylene	Mouse (6 weeks old)	1 or 3 ng/mL intragastrically	Increased abundance of 13 genera and decreased 4 genera	(Guo et al., 2018b)
		Mouse (gestational day 0 to postnatal day 154 or 259; females)	0.05 or 500 µg/ml	High dose, 259 day exposure, decreased <i>Bacteroides</i> spp. and <i>Lactobacillus</i> spp. and increased <i>Bifidobacterium</i> spp. and <i>Enterobacteriaceae</i>	(Khare et al., 2019)

2019b; Hamad et al., 2018; Liu et al., 2017a). Other studies have found decreased alpha diversity and altered composition of the oral microbiome (Hicks et al., 2018; Qiao et al., 2018).

In 12 month old BTBR T + tf/J (BTBR) inbred mice (an established model of ASD for social interaction and behavior) 16 taxa were altered in a sex-specific manner relative to sex-matched C57BL/6 mice (Coretti et al., 2017). For female BTBR mice, this included an increase in *Akkermansia* spp. and a decrease in *Oscillospira* spp., whereas in male BTBR mice, there was an increase in *Lactobacillus* spp. and a decrease in *Desulfovibrio* spp. (Coretti et al., 2017). It was later shown that compared to C57BL/6 J mice, adult male BTBR mice have decreased *Bifidobacterium* spp. and *Blautia* spp. associated with dysregulated bile acid and tryptophan metabolism related to ASD behavior (Golubeva et al., 2017). Male Syrian hamsters exposed to a neurotoxic dose of propionic acid exhibit ASD-like behaviors and glutamate excitotoxicity in the brain. A probiotic mixture of *Bifidobacterium breve*, *Bifidobacterium infantis*, *Lactobacillus acidophilus*, *Lactobacillus bulgaricus*, *Lactobacillus casei*, *Lactobacillus rhamnosus*, and *Streptococcus thermophiles* abrogated the ASD effects of high-dose propionic acid (El-Ansary et al., 2018). Deletion of the gene SHANK3 is associated with dysregulated neurodevelopment and autistic like behaviors (Durand et al., 2007). In Shank3 knockout mice, there was a decrease in the levels of *Lactobacillus reuteri* that correlated with decreased expression of gamma-aminobutyric acid (GABA) receptor subunits in the brain; supplementation of *L. reuteri* attenuated unsocial behavior in male Shank3 knockout mice and increased GABA receptor expression (Tabouy et al., 2018).

In summary, both ADHD and ASD patients have higher levels of *Faecalibacterium*, whereas *Bifidobacterium* is higher in ADHD but lower in ASD. Conflicting observations in the richness of gut microbiome were made in both diseases.

## 2.2. Neurodegenerative diseases

### 2.2.1. Alzheimer's disease

AD is a progressive degenerative neurological disorder that is among the most common causes of dementia worldwide. An estimated 5.5 million Americans are living with AD (Hebert et al., 2013). These

include 5.3 million patients above 65 years of age (10% of all Americans older than 65) and 200,000 individuals below 65 (early-onset AD). More than 90% of AD cases are idiosyncratic and late-onset (after age 65).

Growing evidence supports a critical role of environmental factors interacting with genes of susceptibility to influence the initiation and progression of AD. There is an immense impetus to understand the interactions between host genes and the functional microbiome that may lead to the pathogenesis of AD (Bhattacharjee and Lukiw, 2013). In humans, gut microbiome from AD patients has decreased microbial diversity and is compositionally distinct from age- and gender-matched healthy controls, evidenced by a decrease in Firmicutes and *Bifidobacterium* spp., but an increase in Bacteroidetes, and such dysbiosis correlated with cerebrospinal fluid biomarkers of AD (Vogt et al., 2017). In the APPS1 transgenic AD mouse model (human transgenes for both amyloid beta precursor protein [APP] and presenilin 1 [PSEN1]), GF mice had a drastic reduction of cerebral amyloid  $\beta$  (A $\beta$ ) pathology, a hallmark for clinical AD, whereas colonizing these GF mice with microbiota from conventional mice increased cerebral A $\beta$  pathology (Harach et al., 2017). Similarly, life-long combinatorial antibiotics treatment of AD mice also reduced A $\beta$  pathology (Minter et al., 2016). Interestingly, probiotic supplementation reduced cognitive decline, accumulation of A $\beta$  aggregates, and plasma inflammatory cytokines (Bonfili et al., 2017). Acute perinatal antibiotic treatment resulted in long-term alterations of gut microbiota, reduction in brain A $\beta$  deposition, and inflammatory signaling in serum and brain of aged APPS1 mice, indicating that there is a critical time window early in life to target the microbiome and modulate late-onset of AD (Minter et al., 2017).

Microbial metabolites may modulate the gut-brain axis during the pathogenesis of AD. Fecal SCFAs were decreased in AD mice (Zhang et al., 2017). In an AD mouse model, administration of butyrate, a major microbial-derived SCFA, led to epigenetic reprogramming (by inhibiting histone deacetylation [HDAC]) and improved memory, even at an advanced stage of disease progression (Govindarajan et al., 2011). Inflammation is a critical contributor to the pathogenesis of AD (Akiyama, 1994; Ardura-Fabregat et al., 2017; Eikelenboom et al.,

**Table 3**  
List of neurotoxicants and type of toxicity.

Toxicant	Neurotoxicity	Sources
Amiodarone	Myelinopathy	(Graham Lantos, 1997)
Antibiotics (unspecified)	Peripheral neuropathy	(Fraser et al., 1985)
Arsenic	Postpartum depression	(Murphy et al., 2018)
Atrazine	Wallerian degeneration of axons	(Klaassen, 2013)
Carbamates	Decreased serotonin	(Chen et al., 2018b)
Cisplatin	Inhibition of hippocampal neurogenesis, memory dysfunctions, impaired myelination	(Seth et al., 2019)
Copper imbalance	Peripheral neuropathy	(Graham Lantos, 1997)
	Aceruloplasminemia, Alzheimer's disease, amyotrophic lateral sclerosis, Huntington's disease, Menkes disease, occipital horn syndrome, Parkinson's disease, prion disease, and Wilson disease	(Desai and Kaler, 2008)
Disulfiram	Peripheral neuropathy	(Graham Lantos, 1997)
Doxorubicin	Progressive ataxia	(Graham Lantos, 1997; Spencer and Schaumburg, 2000)
Ethanol	Worsened short-term memory, impaired neuronal signaling	(Huf et al., 2019)
	Glial dysfunction, neuroinflammation	(Gomez et al., 2018)
Formaldehyde	Neurobehavioral impairment and seizures	(Tang et al., 2011)
Gold sodium thiomalate	Decreased unmyelinated axons	(Levine et al., 1986)
Lead	ADHD	(Banerjee et al., 2007; Scassellati et al., 2012)
	Cerebral edema, lead encephalopathy, peripheral neuropathy	(Johnston and Goldstein, 1998)
Magnesium deficiency	Depression	(Winther et al., 2015)
Manganese	Parkinson's disease	(Calderon-Garciduenas et al., 2016; Dobson et al., 2004)
Methylmercury	Neuronal degeneration, ataxia, paresthesia, psychomotor retardation, developmental disabilities, and cognitive deficits	(Klaassen, 2013)
Metronidazole	Peripheral neuropathy	(Goolsby et al., 2018)
MPTP (1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine)	Parkinson's disease	(Kopin, 1987)
Nitrofurantoin	Peripheral neuropathy	(Spencer and Schaumburg, 2000)
Organochlorines	Blocks chloride channels of the GABA-A receptor, seizures, weak association with Parkinson's disease	(Costa, 2015)
Organophosphates	ADHD	(Banerjee et al., 2007; Scassellati et al., 2012)
	Over-stimulation of neurons	(Ruark et al., 2013)
	Altered serotonin signaling	(Slotkin et al., 2008)
Ozone	Alzheimer's disease	(Calderon-Garciduenas et al., 2016)
PM2.5	Alzheimer's disease	(Calderon-Garciduenas et al., 2016)
	Parkinson's disease	(Shin et al., 2018)
Polybrominated diphenyl ethers (PBDEs)	Developmental neurotoxicity	(Dorman et al., 2018)
	Inhibited adult neurogenesis	(Li et al., 2013)
Polychlorinated biphenyls (PCBs)	ADHD	(Banerjee et al., 2007; Scassellati et al., 2012)
	Neurological deficits (differences in neuromotor development, decrements in cognition and behavioral deficits)	(Korrick and Sagiv, 2008)
	Lower cognitive functioning	(Przybyla et al., 2017)
Pyrethroids	Gulf War Illness (GWI), hippocampal volume, neural stem cell activity, and neurogenesis	(Parihar et al., 2013)
Rotenoids	Apoptosis in serotonergic and dopaminergic neurons	(Bisbal and Sanchez, 2019)
	Parkinson's disease	(Betarbet et al., 2000)
Trichloroethylene (TCE)	CNS depressant by inhibiting GABA <sub>A</sub> and glycine receptors	(Beckstead et al., 2000; Krasowski and Harrison, 2000)

1994), and SCFAs also have anti-inflammatory effects through HDAC inhibition (Vinolo et al., 2011). These studies suggest that SCFAs can be neuroprotective. In addition to SCFAs, the microbial-derived ursodeoxycholic acid (UDCA) has been shown to be neuroprotective in various *in vitro* and *in vivo* AD models by reducing apoptosis and promoting cell survival (Ramalho et al., 2008). UDCA treatment can reduce inflammation and increase intestinal *Akkermansia muciniphila*, which is a normal gut bacterial species responsible for SCFA production (Van den Bossche et al., 2017).

In summary, gut microbiome may contribute to the pathogenesis of AD through dysbiosis-induced pro-inflammatory signaling, whereas beneficial microbial metabolites and/or bacteria that produce these metabolites may serve as novel therapeutic modalities for AD.

### 2.2.2. Parkinson's disease

Parkinson's disease (PD) affects an estimated 1 million people in the US and is the second most common neurodegenerative disease; however, less than 10% of cases are considered hereditary (Nalls et al., 2014). Symptoms of PD include motor deficits (tremors), muscle

rigidity, bradykinesia, and impaired gait. PD is one of a group of neurodegenerative diseases called synucleinopathies characterized by an aggregation of  $\alpha$ -synuclein ( $\alpha$ Syn). Interestingly, there is a rostrocaudal gradient of  $\alpha$ Syn in the ENS with higher expression in the upper gastrointestinal tract, and gastrointestinal symptoms often occur before motor symptoms in PD (Cersosimo et al., 2013). Mice overexpressing  $\alpha$ Syn were protected from aggregation by antibiotic treatment, and exposing  $\alpha$ Syn-overexpressing mice to microbiota of PD patients increased physical impairments compared to colonization by healthy donors (Sampson et al., 2016). In an effort to identify microbial metabolites that may antagonize PD development, Sampson et al. (2016) demonstrated that oral exposure to high concentrations of SCFAs (propionate, acetate, and butyrate) was sufficient to produce the motor deficits similar to PD in mice (Sampson et al., 2016). A human study, however, found decreased SCFAs in PD patients compared to healthy controls (Unger et al., 2016). Gut dysbiosis in PD patients included increased *Proteus* spp., *Bilophila* spp., and *Roseburia* spp. along with decreased *Lachnospiraceae*, *Rikenellaceae*, *Peptostreptococcaceae*, and *Butyrivococcus* spp. (Sampson et al., 2016). A cross-sectional study

comparing PD patients to healthy controls found increased *Ruminococcaceae* to be associated with disease duration (Hill-Burns et al., 2017).

In summary, for AD and PD, fecal SFCAs were lower, suggesting that this class of microbial metabolites may be beneficial to prevent the disease onset. In mouse models, SCFA supplementation improved memory likely due to the HDAC inhibitor and anti-inflammation properties; however, conflicting results were observed in PD mouse models, because oral exposure to high concentrations of SCFAs actually produced motor deficit similar to PD.

## 2.3. Psychiatric disorders

### 2.3.1. Depression

Major depressive disorder (MDD) is a common, life-disrupting mental health disorder that is a leading cause of disability worldwide (Moussavi et al., 2007). Diagnosed by at least two weeks of low mood, MDD is accompanied by low self-esteem, loss of interest, and low energy. Environmental factors are associated with MDD, and several human studies have examined the association between gut microbiome and patients with MDD (Aizawa et al., 2016; Chen et al., 2018c; Jiang et al., 2015; Lurie et al., 2015; Naseribafrouei et al., 2014). Of note, *Oscillibacter* spp. and *Alistipes* spp. showed a high association with MDD (Naseribafrouei et al., 2014). A second study also found increased *Alistipes* spp. and *Enterobacteriaceae*, as well as decreased *Faecalibacterium* spp. (Jiang et al., 2015). *Bifidobacterium* spp. and *Lactobacillus* spp. were decreased in another MDD cohort (Aizawa et al., 2016), and increased Actinobacteria and decreased Bacteroidetes were identified in female MDD patients (Chen et al., 2018c). In the chronic variable stress rat model for depression, the genera *Candidatus Arthromitus* and *Oscillibacter* were increased and *Marvinbryantia*, *Corynebacterium*, *Psychrobacter*, *Christensenella*, *Lactobacillus*, *Peptostreptococcaceae incertae sedis*, *Anaerovorax*, *Clostridiales incertae sedis* and *Coprococcus* were decreased in depressed rats compared to control rats (Yu et al., 2017). The Flinders sensitive rat line—a strain used for depression studies—had decreased sample richness (alpha diversity) compared to controls with increased *Proteobacteria* and decreased *Elusimicrobia* and *Saccharibacteria* (Tillmann et al., 2018). Interestingly, transplantation of fecal microbiota from human MDD patients to microbiota-depleted rats or mice induced depression-like behaviors, such as anhedonia and anxiety (Kelly et al., 2016; Zheng et al., 2016). The transplantation of fecal microbiota from depressed patients to GF mice suggested that the gut microbiota could disrupt cAMP Responsive Element Binding Protein 1 (Creb1) signaling by down-regulating calcium voltage-gated channel subunit alpha 1 E (CACNA1E) and disrupting axonogenesis of the olfactory bulb (Huang et al., 2019). Clinical trials of patients with MDD and one of three probiotics improved cognitive performance. Treatment with *Lactobacillus plantarum* 299v decreased the concentration of kynurenine, suggesting that microbial tryptophan metabolism may be important for MDD's etiology (Kazemi et al., 2018; Rudzki et al., 2019). In a male C57BL/6 mouse model of chronic unpredictable mild stress-induced (CUMS) depression, *Clostridium butyricum* treatment improved depressive behaviors, possibly because *C. butyricum* upregulated glucagon-like peptide-1 (GLP-1) in the intestine, activated GLP-1R in the brain, and increased cerebral serotonin (Sun et al., 2018a). Overall, gut dysbiosis may contribute to MDD and supplementation of probiotics may alleviate the disorder.

### 2.3.2. Bipolar affective disorder and schizophrenia

Bipolar affective disorder (BD) is a spectrum of psychiatric syndromes through a cyclical pattern of excitement and depressive behavior (Akiskal et al., 2000; Dacquino et al., 2015). BD is a major cause of global disability and premature mortality (Painold et al., 2018). Several human studies have investigated the association between the gut microbiome and BD. A relatively small cross-sectional comparison of fecal microbiome revealed a negative correlation between duration of BD

and sample richness (Painold et al., 2018), and decreased microbiome richness was also observed in a monozygotic twin study comparing the risk of developing or having BD (Vinberg et al., 2019). The abundance of *Faecalibacterium* spp. was decreased in a small cohort of BD patients (Painold et al., 2018), and this was replicated in a larger cohort of BD patients that also showed a positive correlation between *Faecalibacterium* spp. abundance and better self-reported outcomes (Flowers et al., 2017). The use of atypical antipsychotics as a treatment for BD increased *Lachnospiraceae* and decreased *Akkermansia* spp. and *Sutterella* spp. (Flowers et al., 2017). Of clinical interest, patients diagnosed with BD and given the probiotic OMNi-BiOTiC® Stress Repair (*Bifidobacterium bifidum* W23, *B. lactis* W51, *B. lactis* W52, *Lactobacillus acidophilus* W22, *L. casei* W56, *L. paracasei* W20, *L. plantarum* W62, *L. salivarius* W24, *Lactococcus lactis* W19) had significant improvement in attention and psychomotor processing speed after taking the probiotic for 1–3 months (Painold et al., 2018).

Schizophrenia is a debilitating chronic mental health disorder characterized by delusional thoughts, disorganized behavior, and decreased participation in daily activities. Schizophrenia has increased mortality (Saha et al., 2007) and creates a heavy financial burden (Knapp et al., 2004). Although the etiology of schizophrenia is not well understood, risk factors include genetic inheritance (Schizophrenia Psychiatric Genome-Wide Association Study, 2011) and environmental interactions (Severance et al., 2015; van Os et al., 2010). Both microbial richness index (Chao) and diversity index (Shannon) were lower in patients with schizophrenia compared to healthy controls (Zheng et al., 2019). GF mice given human microbiome fecal transplants of people with schizophrenia exhibited schizophrenic-like behaviors and displayed similar glutamatergic hypofunction as other mouse schizophrenic models, which have decreased glutamate and increased glutamine and GABA (Zheng et al., 2019). A metagenomics analysis of the oropharyngeal microbiome comparing 16 schizophrenia patients with 16 controls demonstrated increased *Lactobacillus* spp. and *Bifidobacterium* spp., as well as an increase in the fungal phylum Ascomycota (Castro-Nallar et al., 2015). Of note, a single-arm study treating schizophrenia patients with the probiotic *Bifidobacterium breve* A-1 had improved anxiety and depression scores, and they reported fewer negative symptoms (Okubo et al., 2019). *Bifidobacterium* spp. was increased in the fecal microbiota of older schizophrenic subjects given 4<sup>α</sup>-β-D-galactosylsucrose as a prebiotic treatment for underweight patients (Nagamine et al., 2018).

A 16S rDNA gene sequencing study of the fecal microbiome found increased relative abundance of the genera *Succinivibrio*, *Megasphaera*, *Collinsella*, *Clostridium*, *Klebsiella* and *Methanobrevibacter* in schizophrenia patients compared to healthy controls, whereas the abundance of *Blautia*, *Coprococcus*, *Roseburia* was decreased (Shen et al., 2018). A second group found increased *Anaerococcus* whereas *Haemophilus*, *Sutterella*, and *Clostridium* were decreased; there was also an association between worsening depressive symptoms and the abundance of *Bacteroides* (Nguyen et al., 2019). Interestingly, metatranscriptomics of whole blood from healthy individuals compared to schizophrenia patients found increased microbial sample richness; increased diversity was inversely associated with CD8+ CD28− CD45RA−, indicating an association between the microbiome, immunity, and schizophrenia (Olde Loohuis et al., 2018). A schizophrenia model in rats using social isolation had increased Actinobacteria and decreased Clostridia compared to controls and also noted associations between gut microbiota and hippocampal interleukin (IL)-6 and IL-10 (Dunphy-Doherty et al., 2018).

In summary, among the three types of psychiatric disorders, decreased gut microbiome richness appeared to be a common feature. *Faecalibacterium* spp. was decreased in both depression and BD; whereas *Lactobacillus* spp. supplementation was beneficial to improve both depression and BD. Depression and schizophrenia appeared to have opposite microbial patterns: for example, depression has lower *Bifidobacterium* spp. and *Lactobacillus* spp., whereas schizophrenia has

higher levels of these bacteria.

## 2.4. Other neurological disorders associated with stress

### 2.4.1. Postpartum depression

Postpartum depression is a clinically diagnosed form of MDD with onset during the peripartum period; several risk factors for postpartum depression include history of depression, pregnancy and birth complications, and psychoneuroimmune dysregulation (A.P. Association, 2013; Corwin et al., 2015; Dunn et al., 2015; Osborne and Monk, 2013). As many as 19.2% of women have a major depressive episode within the first three months postpartum (Gavin et al., 2005). Remarkably, antibiotic exposure was identified as an independent risk factor up to 2 months postpartum (Murphy et al., 2018). A cohort of 380 women in New Zealand were given the probiotic *Lactobacillus rhamnosus* HN001 or a placebo from 14 to 16 weeks gestation through 6 months postpartum; women given the probiotic reported lower depression and anxiety scores during the postpartum period (Slykerman et al., 2017). Dysbiosis may increase the risk of postpartum depression, whereas probiotic supplementation may decrease the risk of depressive episodes.

### 2.4.2. Post-traumatic stress disorder

Post-traumatic stress disorder (PTSD) is a commonly occurring condition manifesting after exposure to trauma, such as war, sexual assault, and other distressing experiences, and has a high rate of psychiatric comorbidity (Kessler, 2000). PTSD can often occur for many years and is frequently associated with exposure to multiple traumas (Hemmings et al., 2017). In a mouse model of PTSD, in which adult male C57BL/6 J mice were exposed to extreme aggression by SJL albino male mice, the microbial-influenced metabolites 3-phenylpropionate, hippurate, and phenylpropionylglycine were increased in PTSD mice (Gautam et al., 2015), indicating an altered gut microbiome. A follow-up study using the same PTSD aggressive mouse model showed an immediate and inconsistent dysregulation of microbiome composition, specifically for *Akkermansia* spp., *Anaeroplasm* spp., *Lactobacillus* spp., and *Oscillospira* spp. (Gautam et al., 2018). In a preliminary human study investigating the fecal microbiome in South African PTSD-affected individuals, there was no difference in the diversity of the microbiota, but there was a decrease in the total abundance of the phyla Actinobacteria, Lentisphaerae, and Verrucomicrobia associated with PTSD (Hemmings et al., 2017).

### 2.4.3. Anorexia nervosa

Anorexia nervosa is an eating disorder characterized by severe weight loss (lack of appropriate weight gain or maintenance) often caused by the limitation of calories, fear of gaining weight, and denial of being underweight (Miller and Golden, 2010). Anorexia nervosa is more likely to affect females than males (Herpertz-Dahlmann, 2009). The fecal microbiota of 16 female patients showed increased sample richness at the time of admission compared to discharge (Kleiman et al., 2015). Relative to controls, samples from anorexic patients during admission were higher in the genera *Anaerostipes* and *Faecalibacterium*, but an undefined genus in the family *Coriobacteriales* was lower (Kleiman et al., 2015). Another study identified decreased *Clostridium coccoides*, *Clostridium leptum*, and *Bacteroides fragilis*, as well as decreased SCFAs acetic acid and propionic acid, in the stool of female patients with anorexia nervosa compared to healthy controls (Morita et al., 2015). In another cohort of anorexia nervosa patients, propionate and butyrate were decreased, corresponding to increased *Enterobacteriaceae* and *Methanobrevibacter smithii* and decreased *Roseburia* spp., *Ruminococcus* spp., and *Clostridium* spp. (Borgo et al., 2017). Anorexia nervosa patients who gained weight had increased sample richness; the genera in *Clostridium XI* and *Bacteroidetes* were unique to anorexia nervosa patients before weight gain (Mack et al., 2016). In BALB/c mice, a chronic caloric restriction study was conducted to determine the effect on gut microbiota (Chen et al., 2016). Briefly, caloric intake was limited to

prevent weight gain starting at 28 days of age and restored to *ad libitum* feeding after day 97 until day 120 when the tissues were collected. Interestingly, the changes in gut microbiome persisted even after lifting the caloric restriction. Collectively, it is unclear if the compositional changes mechanistically contribute to anorexia nervosa.

## 3. Gut microbiome mechanistically contributes to behavior changes – lessons learned from germ-free (GF) research models

Studies have established the association between intestinal dysbiosis and neurological disorders of various etiologies (Alam et al., 2017; Bourassa et al., 2016; Galland, 2014; Heiss and Olofsson, 2019; Kelly et al., 2017; McKay et al., 2017; Moos et al., 2016; Tremlett et al., 2017; Zhu et al., 2017), but the role of the gut microbiome in modulating various behavior changes requires mechanistic investigations using laboratory models such as GF rats and mice. GF rats had exacerbated neuroendocrine and behavior responses to acute stress, coinciding with alterations of the dopaminergic turnover rate in the upper structures of the brain known to regulate stress response and anxiety-like behavior (Crumevolle-Arias et al., 2014). The GF rats also had higher serum corticosterone levels and elevated mRNA of corticotropin releasing factor but reduced mRNA of glucocorticoid receptor in hippocampus (Crumevolle-Arias et al., 2014). Similarly, GF mice had impaired cognitive behaviors in response to novel objects, corresponding to decreased expression of genes involved in brain-derived neurotrophic factor (BDNF) signaling in amygdala, which is a key region for the social brain network (Arentsen et al., 2015). In response to restraint stress, plasma levels of adrenocorticotrophic hormone (ATCH) and corticosterone were also substantially higher in GF mice (Sudo et al., 2004). Conversely, under basal conditions (i.e. without an environmental stimulus), GF mice exhibited anxiolytic behavior accompanied by decreased mRNA expression of the N-methyl-D-aspartate receptor subunit NR2B in the central amygdala, as well as increased mRNA of BDNF but decreased mRNA of serotonin receptor 1A in the dentate granule layer of the hippocampus (Neufeld et al., 2011). There are also many morphological differences between specific pathogen-free (SPF) and GF mice. For example, GF mice have delayed brain maturation and organization, evidenced by lower volumes and fractional anisotropy in major gray and white matter areas, as well as lower levels of myelination in total brain and major white and grey matter structures at either 4 or 12-weeks of age (Lu et al., 2018). This coincided with lower mobility and higher anxiety of GF mice in an open field test, which is a photo beam tracking method to monitor the movement of animals with two sets of infrared beams, at 4-weeks of age (Lu et al., 2018). At 12-weeks of age, GF mice also had reduced spatial and learning memory in the Morris water maze test, which measures spatial learning for subjects that use distal cues to navigate from start locations around the perimeter of an open swimming field to locate a submerged escape platform. At this age, GF mice also exhibited reduced contextual memory in contextual fear conditioning test, which measures persistent freezing behavior using a foot shock context arena and a conditioned stimulus, in order to quantify hippocampus-dependent learning and memory test (Lu et al., 2018). In a three-chamber social test, which assesses cognition in the form of interest in a never-before-met intruder (i.e. social novelty), GF mice had reduced social novelty at 12-week of age. Similar behavioral changes were also observed in antibiotic-treated SPF mice (Desbonnet et al., 2015).

While the absence of gut microbiota leads to marked behavioral changes, microbial colonization of GF mice reprogrammed the post-natal development of the HPA system. For example, the heightened HPA stress response in GF mice was attenuated by inoculation with *Bifidobacterium infantis*, but was enhanced by the enteropathogen *Escherichia coli* (Sudo et al., 2004). There appears to be a critical time window for microbiome inoculation to affect behavior in GF mice as fecal transplants from SPF mouse donor to 6 week old but not 8 week old GF mice reduced elevated HPA response in GF mice (Sudo et al.,

2004). Early-life inoculation using SPF feces reduced the expression of postsynaptic density protein 95 (PSD-95) and synaptophysin in the striatum, which are involved in the neuronal circuits for motor control and anxiety behavior (Diaz Hejtz et al., 2011). Together these studies demonstrate that the presence of the commensal gut microbiota has dual functions in modulating both cognitive functions under basal conditions as well as stress response.

#### 4. Microbiome and neurotoxics

##### 4.1. Air pollution

Recent epidemiological studies have established a positive correlation between exposure to ubiquitous traffic-related air pollution and the exacerbations of various neurological disorders such as Alzheimer's disease and Parkinson's disease (Babadjouni et al., 2017; Calderon-Garciduenas et al., 2016). Large populations living in highly polluted metropolitan regions are at higher risk for robust central nervous system pathology (Babadjouni et al., 2017). The primary cause of urban air pollution is through vehicular emissions. Exposure to fine particulate matter (PM<sub>2.5</sub>) and ozone above US EPA standards, have been linked to both Alzheimer's and Parkinson's diseases (Calderon-Garciduenas et al., 2016; Palacios, 2017; Shin et al., 2018). Air pollution positively associates with the hallmark clinical characteristics of neuroinflammation and CNS diseases, including increased expression of pro-inflammatory markers in brain, diffused amyloid plaques, neuronal cell loss, and impaired cognition (Block and Calderon-Garciduenas, 2009). It has been suggested that inhaled PM can directly translocate to the brain and cause neuroinflammation, which in turn contributes to neurodegeneration (Calderon-Garciduenas et al., 2016; Oberdorster et al., 2005). In addition, the novel contribution of the gut-brain axis in several CNS disorders suggests that gut-derived pro-inflammatory signaling accompanied with dys-regulated microbial or host-derived lipid profiles may contribute to neurotoxicity (Russo et al., 2018; Valles and Francino, 2018). In human subjects, exposure to freeway air pollution correlated with decreased Bacteroidaceae and increased Coriobacteriaceae as well as increased fasting glucose levels (Alderete et al., 2018). In mice, inhalation of PM altered the composition of the gut microbiome and was suggested to play a role in PM-induced GI inflammation (Beamish et al., 2011; Kish et al., 2013; Mutlu et al., 2018). Because intestinal bacteria contribute to both local and systemic inflammation, the latter may lead to neuroinflammation (Grigg and Sonnenberg, 2017; Reinoso Webb et al., 2016). Investigation of the interplay among air pollution, gut microbiome, and inflammatory signaling is an intriguing direction in research on mechanisms of neurotoxicity.

##### 4.2. Antibiotics and drugs

The important role of gut microbiome in the biotransformation of various therapeutic drugs has been extensively reviewed (Bisanz et al., 2018; Carmody and Turnbaugh, 2014; Haiser and Turnbaugh, 2013; Spanogiannopoulos et al., 2016). Briefly, gut microbiome can either utilize their own enzymes to metabolize drugs into reactive or inactive metabolites or can secrete endogenous microbial metabolites into circulation to interact with host receptors in various organs. We will focus on studies on gut microbiome and the drug categories that have neurotoxic side effects as summarized in Table 2. These drugs are selected based on Chapter 16 of the Casarett & Doull's Toxicology Textbook (Klaassen, 2013) as well as literature search regarding the involvement of the gut microbiome in the metabolism and/or toxicities of these chemicals.

**Microbiome and compounds associated with neuropathies.** The anticancer drug doxorubicin is known to produce progressive ataxia in laboratory animal models through degeneration of dorsal root ganglion cells and axonal degeneration (Graham Lantos, 1997; Spencer and Schaumburg, 2000). In the human gut microbiome, doxorubicin can be

inactivated by *Raoultella planticola* via reductive deglycosylation (Yan et al., 2018), thus targeting this bacteria is promising to reduce its side effects including neurotoxicity. Intestinal bacteria contribute to doxorubicin-induced intestinal damage (Rigby et al., 2016), whereas the dietary fiber pectin, which is a substrate of microbial fermentation, can reduce doxorubicin-induced intestinal inflammation through direct interaction with host Toll-like receptors 1 and 2 (Sahasrabudhe et al., 2018). In addition, gut microbiota can affect the efficacy of doxorubicin and many other anticancer drugs (Florez et al., 2016; Lehouritis et al., 2015), whereas doxorubicin can alter the gut microbial activity (Song et al., 2015). It may be interesting to investigate whether the interactions between doxorubicin and gut microbiota may modify the CNS toxicity.

MPTP (1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine) is an industrial chemical and contaminant of illicit narcotics that selectively destroys dopaminergic neurons of the nigrostriatal pathway, leading to Parkinsonian syndromes (Calne et al., 1985; Kopin, 1987). In mice, sub-chronic MPTP exposure affected the gut microbiome diversity, which was associated with sensorimotor performance and fear learning (Torres et al., 2018). Importantly, dysbiosis and GI dysfunction precedes motor dysfunction (Lai et al., 2018), whereas fecal microbiota transplantation from MPTP-exposed mice alone can induce motor impairment and striatal neurotransmitter decreased in normal mice (Sun et al., 2018b). Therefore, microbiota may mechanistically contribute to the pathogenesis of Parkinson's disease.

**Microbiome and compounds associated with axonopathies.** The antibiotic and antiprotozoal drug metronidazole is a drug that can produce peripheral neuropathy (Goolsby et al., 2018), leading to axonal degeneration, lesions of cerebellar nuclei, mostly affecting myelinated fibers (Graham Lantos, 1997; Spencer and Schaumburg, 2000). Studies showed that metronidazole also altered gut microbiota in the colon of healthy rats, evidenced by increased *Bifidobacteria* spp. (especially *Bifidobacterium pseudolongum*) and *Enterobacteria*, and this associated with increased mucus layer thickness (Pelissier et al., 2010). While this effect has been suggested to benefit the treatment of inflammatory bowel diseases, potential systemic effects following changes in the microbiota, especially on the CNS, need further investigation.

The antibiotic nitrofurantoin produces peripheral neuropathy via axonal degeneration (Spencer and Schaumburg, 2000). In human subjects with urinary tract infections, nitrofurantoin treatment correlated with a reduced relative abundance of the genus *Clostridium* and an increased relative abundance of the genus *Facalibacterium* (Stewardson et al., 2015), whereas another clinical study showed that nitrofurantoin treatment resulted in a temporary increase in the genus *Bifidobacterium* (Vervoort et al., 2015). Overall, it is generally considered that this drug only has a mild effect on the gut microbiota community, although the potential changes in microbial functions locally and in distal organs need additional investigations.

The anticancer drug cisplatin produces peripheral neuropathy evidenced by axonal degeneration and microtubule accumulation in early stages (Graham Lantos, 1997; Spencer and Schaumburg, 2000). In mice, cisplatin treatment disrupts the gut microbiome as well as the mucosal integrity, and this can be partially corrected by *Ruminococcus gnavus*, which is one of the bacteria that is depleted by cisplatin, or by fecal gavage (Perales-Puchalt et al., 2018).

**Microbiome and compounds associated with myelinopathies.** Amiodarone is an anti-arrhythmic drug that is known to produce myelinopathy (Graham Lantos, 1997; Spencer and Schaumburg, 2000) as well as peripheral neuropathy following long-term high-dose therapy (Fraser et al., 1985). In rats, oral administration of *Lactobacillus casei* DN-114 001 slowed the absorption of amiodarone without altering the pharmacokinetics of its main metabolite (Matuskova et al., 2017); however, oral administration of *Escherichia coli* Nissel 1917 (EcN) increased the oral bioavailability of the parent compound as well as the P450-mediated metabolism (Matuskova et al., 2014). These studies at the single-strain resolution have suggested that targeting distinct

microbiota can alter the pharmacokinetics and potentially the toxicity of amiodarone.

Disulfiram, which treats chronic alcoholism by inhibiting aldehyde dehydrogenase, produces peripheral neuropathy and swelling in distal axons (Graham Lantos, 1997). Interestingly, disulfiram was repurposed as an antibiotic for multi-drug resistant *Staphylococcus aureus* infections, and it can increase the vancomycin susceptibility of three clinical vancomycin-resistant *S. aureus* strains (Long, 2017). The disulfiram metabolite diethyldithiocarbamate also has antibacterial activity towards *Bacillus anthracis* (Frazier et al., 2019). In addition, disulfiram-based disulfide derivatives have been shown to exhibit antibacterial activity against gram-positive *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacillus*, and *Listeria* spp. (Sheppard et al., 2018). It remains unknown whether the antibacterial effect of disulfiram can affect the CNS.

**Psychotropic drugs.** Several psychotropic drugs, including fluoxetine and escitalopram, have antimicrobial effects *in vitro* (Cussotto et al., 2018). Alternatively, lithium, valproate, and aripiprazole increased the richness and diversity of gut microbiome in humans. In addition, escitalopram, venlafaxine, fluoxetine, and aripiprazole have also been shown to increase the gut permeability in the ileum. Because it is known that microbiota plays an important role in regulating the gut-brain interactions, it has been suggested that the intestinal effect of the psychotropics may contribute to the mechanism of action and side effects of these medications (Cussotto et al., 2018).

#### 4.3. Metals and metalloids

**Arsenic.** Arsenic is a carcinogenic metalloid that is widely distributed in nature in its trivalent (arsenic trioxide and sodium arsenite) and pentoxide forms (sodium arsenate, arsenic pentoxide, and arsenic acid). Other organoarsenicals are also common. Occupational exposure to arsenic compounds comes in the form of pesticides, herbicides, and other agricultural products. Environmental arsenic is in drinking water with sources in the United States less than 5 µg/L (Environmental Protection Agency maximum contaminant limit is 10 µg/L), however it is estimated that many people in Bangladesh drink water in excess of 50 µg/L (Klaassen, 2013). Acute exposure to arsenic has many symptoms, including fever, anorexia, hepatomegaly, melanosis, cardiac arrhythmia, and terminal cardiac failure, as well as a delay in sensory loss in the peripheral nervous system (Wallerian degeneration of axons) (Klaassen, 2013). Chronic arsenic exposure can cause diffuse or spotted hyperpigmentation or hypopigmentation, as well as liver injury. Interestingly, peripheral neuropathy also occurs with the development of numbness or “pins and needles” in the hands and feet, caused by dying-back axonopathy with demyelination.

From environmental observations, there was a non-significant trend in microbiome composition dependent on arsenic contamination observed in soil and earthworm microbiome samples around an arsenic mine, with about 47 taxa (about 7% of the abundance) driving the differences (Pass et al., 2015). Zebrafish exposed to varying low concentrations of arsenic (10, 50, and 100 ppb for 20 days) found 43 amplicon sequence variants that increased in abundance and 43 that decreased at the genus level (Dahan et al., 2018).

The Simulator of the Human Microbial Intestinal Ecosystem (SHIME) system is a dynamic series of six compartments and pumps that are used to mimic the intestines by controlling the pH, relative volume of fluid, and residence time, as well as the inoculation of the human microbiome (Joly et al., 2013; Requile et al., 2018). A study using the human SHIME model showed that arsenite (As[III]) is more readily metabolized than arsenate (As[V]), 66.5–92% to 22.1–38.2% respectively (Yin et al., 2015), and gut microbiota can also release arsenic from the soil in solid phase for absorption (Yin et al., 2017). Another SHIME model demonstrated that co-administration of drinking water with ferric iron and arsenic increased arsenic methylation and decreased absorption in the colon. Effluents from iron-exposed SHIME

colons decreased toxicity in the human hepatoma cell line HepG2 cells (Yu et al., 2016). *Bacteroides* spp., *Clostridium* spp., *Alistipes* spp., and *Bilophila* spp. had resistance to and ability to methylate arsenic (Yu et al., 2016).

Several mouse studies have looked at the effect of arsenic exposure on the microbiome using 16S rDNA gene sequencing data. One of the first arsenic microbiome studies exposed C57BL/6 mice to 10 ppm arsenic in drinking water for 4 weeks and showed clear microbiome compositional differences between control and exposed mice with 2 microbial classes increased and 7 classes decreased (Lu et al., 2014). Another study showed that microbiome compositional and functional changes could be sex-specific (Chi et al., 2016), and a follow-up study exposing mice to 100 ppb arsenic for 13 weeks showed altered abundance of microbial genes involved in carbohydrate metabolism, especially pyruvate fermentation, short-chain fatty acid synthesis, and starch utilization, as well as stress response genes (Chi et al., 2017a). However, mice exposed to arsenite (10 or 250 ppb) did not have altered expression of arsenate reductase (arsA) or the arsenite exporter (arsB), thus bacteria with genes protective against arsenic may not be the bacteria that were increased during arsenic exposure (Dheer et al., 2015).

Importantly, Coryell et al. (2018) showed that GF mice and antibiotic-treated conventional mice exposed to arsenate (25 and 100 ppm) excreted less arsenic in stool and accumulated more arsenic in tissues relative to control mice. GF arsenite methyltransferase (As3mt)-knockout mice are hypersensitive to arsenic-induced mortality but are protected following human fecal microbiome transplants, likely in part due to protective properties of *Faecalibacterium prausnitzii* (Coryell et al., 2018). Similarly, antibiotic-treated mice exposed to arsenite (III; 250 ppb and 1 ppm) had increased total urinary arsenic, but decreased fecal arsenic than conventionally raised mice, indicating that a depleted microbiome could absorb more arsenic (Chi et al., 2019). This evidence suggests that gut microbiota may limit the arsenic absorption, but no conclusions are definitive between arsenic-induced microbiome neurotoxicity. Additionally, many of the studies used exposures greater than the described human exposures (50 ppb or 50 µg/L in the population from Bangladesh); therefore, translational conclusions regarding arsenic exposure and microbiome in humans should remain cautious.

**Copper.** Copper is a trace element found in all tissues and is required for a plethora of molecular events including cellular respiration, peptide amidation, neurotransmitter biosynthesis, pigment formation, and connective tissue strength. Copper is present throughout the brain with many enzymes in the CNS being copper-dependent for proper function. Both a deficiency and an excess of copper due to genetics or environmental factors can lead to several neurological conditions, including aceruloplasminemia, Alzheimer’s disease, amyotrophic lateral sclerosis, Huntington’s disease, Menkes disease, occipital horn syndrome, Parkinson’s disease, prion disease, and Wilson disease (Desai and Kaler, 2008). Copper ions are naturally antimicrobial, but few papers have investigated the antimicrobial properties of copper on the gut microbiome. Culturing of chicken ileal microbiome exposed to copper increased *Lactobacillus* spp. while decreasing *Escherichia coli*, but the effect of copper on the *in vivo* microbiome was negligible (Pang et al., 2009). Supplementation of copper to dairy calves for 75 days increased the richness of the rumen microbiome and decreased the lipopolysaccharide biosynthetic pathway (Biscarini et al., 2018). It appears that copper supplementation, not necessarily excess copper, may alter the microbiome composition, but it is unclear if this may play a role in copper neurotoxicity.

**Gold.** Elemental gold is considered nontoxic and is used as an aesthetic enhancement in food and drinks. However, gold in its ionic form can have neurotoxic effects. For example, used as an anti-inflammatory treatment for rheumatoid arthritis, gold sodium thiomalate decreases the number of unmyelinated axons, suggesting gold could slow signal response to the peripheral nervous system (Levine et al., 1986; Salzer and Zalc, 2016). A study investigating gold nanoparticle

protection against colitis also identified a gut dysbiosis in mice following 8 days of exposure with decreased species richness as well as decreased *Lactobacillus* spp. and other SCFA-producing bacteria (Kase et al., 1987). Further research is needed to understand the effect of ionic gold on the gut microbiome and neurotoxicity.

**Lead.** Lead is a non-essential, heavy metal toxicant that can cause severe neurotoxicity with the EPA dictating the action level at 15 ppb. Adults can be exposed to lead through contamination of drinking water via lead pipes or by its natural occurrence in the environment, as well as in occupational settings such as smelting and soldering. Children can also be exposed through mouthing or consuming objects other than food, such as dust and lead contaminated paint. Children more readily absorb lead compared to adults with limited protection by the developing blood-brain barrier (Perlstein and Attala, 1966). Shown in Fig. 2A, lead concentrations in mussels on the coastal regions of the US are highest on the East coast between Washington and Boston with large areas of exposure around the Lake Erie region in northern Ohio, Chicago, and along the West coast near San Francisco and Seattle (see Methods for GIS data analysis). Concentrations of lead in bivalves ranged from 0.172 to 73.6  $\mu\text{g}/\text{dry g}$ . Large acute exposures result in severe cerebral edema, likely due to damaged endothelial cells, with children more likely to develop lead encephalopathy than adults (Johnston and Goldstein, 1998). Although not well understood, chronic lead exposure results in peripheral neuropathy and can also include gastritis, abdominal pain, anemia, and deposits of lead in gums and long bones.

Animal studies showed that lead exposure (acute, chronic, and perinatal) alters the composition of the gut microbiome, which may contribute to metabolic disorders. For instance, perinatal lead exposure in drinking water (32 ppm; 2 weeks before mating to postnatal day 21) increased the adult body weight of male mouse offspring, but not female offspring, and this was highly associated with microbiome compositional change, suggesting that lead-exposed gut microbiome may play a role in sex-specific differences (Wu et al., 2016). Adult female C57BL/6J mice exposed to lead (10 ppm) for 13 weeks decreased the abundance of the genera *Blautia*, *Coprococcus*, two species of *Ruminococcus*, and 1 taxon in each of the families *Clostridiales* and *Lachnospiraceae*, as well as increased abundance of a genus in the *Clostridiaceae* family (Gao et al., 2017c). Lead exposure reduced microbial metabolites in the vitamin E and bile acid pathways as well as altered the gene abundance for nitrogen metabolism and energy metabolism (Gao et al., 2017c). Dysregulated energy metabolism and altered microbiome composition were also found in 5-week-old mice exposed up to 0.1 mg/L (0.1 ppm) of lead for 15 weeks; a dose-dependent decrease in the phylum Bacteroides and an increase in Firmicutes was observed (Xia et al., 2018a). Adult male zebrafish exposed to 0.03 mg/L (0.03 ppm) lead for 7 days had an increase in Firmicutes, but a decrease in Proteobacteria; in total 30 genera were altered by acute lead exposure (Lai et al., 2018). Many of the microbiome-lead exposure studies identified changes in gut microbiome, however the exposures are greater than the EPA action limit of 15 ppb and no studies related the changes in microbiome to neurotoxic effects in the host. More studies are needed to identify if gut microbiome is a contributing factor to lead-induced neurotoxicity.


**Magnesium.** Magnesium is an important cofactor that helps regulate diverse chemical reactions in the body, including cardiovascular, alimentary, endocrine, and osteoarticular systems, as well as brain biochemistry by influencing neurotransmission pathways associated with depression (Rajizadeh et al., 2017; Serefko et al., 2016). Mice exposed to a magnesium-deficient diet for 6 weeks had increased depressive-like behaviors and a gut microbiome composition correlated with magnesium deficiency (Winther et al., 2015). However, it is unclear if the compositional changes in gut microbiome contribute to the observed depressive-like behaviors.

**Manganese.** Manganese is an essential trace element needed for many enzymatic reactions, such as arginase and glutamine synthetase.

Excessive consumption of manganese can lead to Parkinson's disease-like symptoms, including dysfunction of the basal ganglia (Dobson et al., 2004). In the United States, the highest concentration of manganese in bivalves is in northern Ohio and northwestern New York around the Great Lakes region (Lake Erie and Lake Ontario; Fig. 2A). Adult mice exposed to neurotoxic concentrations of manganese for 13 weeks induced a gut dysbiosis for 12 genera in female mice and 16 genera in male mice. Furthermore, there was an increase in microbial tryptophan synthesis pathways in females, but in males, there was a general decrease in GABA/putrescine metabolism (Chi et al., 2017b). Again, it is not known if the changes in gut microbiome contribute to manganese toxicity.

**Mercury and methylmercury.** Elemental mercury (Hg) is a heavy, silvery liquid metal that is used in products such as thermometers, barometers, float valves, and fluorescent lamps. Exposure to elemental mercury may cause emotional disturbances, tremors, and fatigue, but there is limited evidence of toxicity in humans (Klaassen, 2013). The National Institute for Occupational Safety and Health (NIOSH) limits occupational exposures to 0.05 mg/m<sup>3</sup> (about 6 ppb) over a 10-h workday and the EPA limits mercury to 2 ppb in drinking water. In the context of the microbiome, one study examined the development of Hg and antibiotic resistance in bacteria of the estuarine fish mummichog (*Fundulus heteroclitus*) upon exposure to Hg in the diet for 15 days. The microbiome of fish exposed to mercury developed resistance to at least three antibiotics compared to control fish; furthermore, there was an 8-fold increase in the mercury resistance gene mercuric reductase at a mercury contamination site (Lloyd et al., 2016). This indicates that mercury exposure induced the co-selection of mercury and antibiotic resistance. An epidemiological study of pregnant women and children found that consumption of probiotic yogurt was protective against increased blood levels of mercury and arsenic in pregnant women but not children (Bisanz et al., 2014). Interestingly, a combinational exposure of copper and mercury in female Kunming mice decreased the abundance of the genera *Sporosarcina*, *Jeotgailcoccus*, and *Staphylococcus*, but increased the genus *Anaeroplasm* (Ruan et al., 2018). It is unclear how the shift would affect neurotoxicity, but this demonstrates the need to consider mixtures when assessing risk.

Methylmercury (organic mercury) is highly toxic, causing neuronal degeneration, ataxia, paresthesia, and psychomotor retardation in adults (Klaassen, 2013). In children, exposure to methylmercury can cause developmental disabilities, retardation, and cognitive deficits, which is attributed to an immature blood-brain barrier and higher disposition of mercury in the developing brain. In an epidemiological study, methylmercury concentration in the stool or total mercury in hair of pregnant women (36–39 weeks gestation) was correlated with 17 bacterial genera, but there was no detection of the mercury methylation gene or the methylmercury detoxification genes (Rothenberg et al., 2016). The inability to identify known mercury biotransformation genes in the gut microbiome of pregnant women could be attributed to small sample size ( $n = 6$ ) or there could be new alternative pathways for microbial mercury metabolism. One group cultured the microbiome of two human subjects to investigate methylmercury metabolism rate in relation to nutrients; there was a dependency of diet on demethylation of methylmercury by an increase in protein content, but the responsible genes and mechanism could not be identified (Guo et al., 2018a). Changes in gut microbiome following dietary exposure to methylmercury is further supported by a study in fathead minnows which showed significant compositional differences between methylmercury exposed and unexposed groups with 18 genera altered by exposure (Bridges et al., 2018). They found an increase in bacteria that generally have genes for xenobiotic metabolism and metal removal, including Deltaproteobacteria FAC87, *Xanthomonadaceae*, *Comamonadaceae*, *Cloacibacterium* spp., and *Pirellula* spp. (Bridges et al., 2018). Overall, more studies at human relative exposures are needed to conclude the effect of mercury and microbiome on host neurotoxicity.



**Fig. 2.** Collection sites and concentrations of BDE-47, BDE-99, PCB-95, PCB-153, lead, and manganese in bivalves collected from the coastal United States. (A) Green circles indicate BDE-47 sampling sites while purple circles correspond to BDE-99 sampling sites in bivalves with size corresponding to concentration (ng/dry g). (B) Green circles indicate PCB-95 sampling sites while purple circles correspond to PCB-153 sampling sites in bivalves with size corresponding to concentration (ng/dry g). (C) Green circles indicate lead sampling sites while purple circles correspond to manganese sampling sites in bivalves with size corresponding to concentration (ug/dry g) (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).

**4.4. Persistent environmental toxicants**

**PCBs.** Polychlorinated biphenyls (PCBs) were formerly used in industrial and consumer projects before they were banned in the United States in 1979. Due to their bio-accumulative nature, these persistent environmental contaminants are still of significant public health concern and are listed on the 2015 Agency for Toxic Substances and Disease Registry Substance Priority List (<http://www.atsdr.cdc.gov/SPL/index.html>). PCBs are frequently detected in fatty food (Gorchev and Jelinek, 1985; Newsome et al., 1998), air in public schools (Herrick, 2010; Herrick et al., 2016), industrial paint pigments (Anezaki and Nakano, 2014; Hu and Hornbuckle, 2010), as well as human samples including blood, adipose tissue, milk, and placenta (Nakagawa et al., 1987; Suzuki et al., 2005; Wang et al., 2010). In humans, epidemiological studies showed a positive link between developmental PCB exposure and neurological deficits in infants and children (Korrick and Sagiv, 2008; Schantz, 1996; Winneke, 2011). Symptoms associated with PCB neurotoxicity include decreased motor and cognitive skills in newborns, greater incidence of behavioral problems and lower IQ scores in children, and lower cognitive functioning in older adults (Przybyla et al., 2017; Ribas-Fito et al., 2001; Schantz, 1996).

Developmental neurotoxicity following PCB exposure has also been observed in laboratory animals including mouse and rat pups, as well as chicken embryos (Kania-Korwel et al., 2017; Roelens et al., 2005; Yang et al., 2009) and in neuronal cell cultures (Lyng and Seegal, 2008; Lyng et al., 2007). Suggested molecular mechanisms of PCB neurotoxicity include binding to aryl hydrocarbon receptor (for dioxin-like PCB congeners), decreasing dopamine content, GABAergic neuronal dysfunction, interference with calcium signaling, as well as altering the dendritic growth and plasticity by promoting the activity of ryanodine receptors (Lyng et al., 2007; Tilson and Kodavanti, 1998; Yang et al., 2009).

As shown in Fig. 2B, Concentrations of PCB-95 in bivalves ranged from 0.12 to 153.88 ng/dry g while concentrations of PCB-153 ranged from 0.04 to 472.5 ng/dry g. Similar to PBDEs, concentrations of both PCB congeners tended to increase with increasing proximity to coastal cities with large populations, with overall higher concentrations of PCB-153 measured. One exception to this trend appears south of Miami, Florida where PCB-95 is present in higher concentrations than PCB-153. Coinciding with PCB-associated neurodevelopmental disorders, there is strong evidence that intestinal dysbiosis also plays a role in the etiology of these diseases (Borre et al., 2014; Warner, 2019). In adult

mice, oral exposure to an environmentally relevant PCB mixture (PCB-153, PCB-138, and PCB-180) was shown to decrease the levels of *Proteobacteria* correlating with the activity level of the mice, whereas exercise attenuated PCB-induced changes in gut microbiome (Choi et al., 2013). Dioxin-like PCB-126 increased both intestinal and systemic inflammation correlating with gut dysbiosis (Petriello et al., 2018). Gut dysbiosis may lead to altered microbial metabolites as well as increased pro-inflammatory cytokines in circulation. Oral exposure to PCBs increased pro-inflammatory mediators in brain and other metabolic organs (Chi et al., 2018; Sipka et al., 2008). Because systemic inflammation has been linked to neuroinflammation (Bendorius et al., 2018), it is possible that PCB-induced gut dysbiosis contributes to PCB-induced neurotoxicity partially through inflammation. In addition, there is a positive association between circulating glucagon-like peptide-1 (GLP-1) and *Bifidobacterium* spp. following PCB-126 exposure (Petriello et al., 2018) with GLP-1 protecting against neurotoxicity of various etiologies (Athauda and Foltynie, 2016; Chang et al., 2018; Khalilnezhad and Taskiran, 2018).

**PBDEs.** Polybrominated diphenyl ethers (PBDEs) are a class of persistent environmental contaminants that were previously used as flame retardants in various consumer products. Although the commercial use of PBDEs has been banned in the United States recently, PBDEs are still ubiquitously found in the environment due to their resistance to degradation, bio-accumulation along the food chain, as well as the recycling of PBDE-containing products worldwide (Schechter et al., 2010, 2003). PBDE levels in human specimens, such as blood, breast milk, and adipose tissue, have increased exponentially over the last 30 years, and are especially high in North America as compared to European and Asian countries (Costa et al., 2014; Costa and Giordano, 2007). Infants and toddlers are particularly vulnerable to PBDE-induced adverse effects due to exposure to PBDE-contaminated breast milk and dust (Costa et al., 2014). In contrast to many studies, a 2017 study in middle-aged and older Californian women showed a modest average annual percent increase in serum concentrations of 3 major PBDE congeners (BDE-47, BDE-100, and BDE-153) from 2011 to 2015 (Hurley et al., 2017). The Geographic Information System (GIS)-based distribution of 2 breast milk-enriched PBDE congeners (BDE-47 and BDE-99) in coastal areas of the United States is shown in Fig. 2C. Concentrations of BDE-47 in bivalves ranged from 0.3 to 68.4 ng/dry g and concentrations of BDE-99 ranged from 0.2 to 38.4 ng/dry g. The concentrations of both congeners tended to increase with proximity to a major coastal city, and overall higher concentrations of BDE-47 measured. Higher brominated congeners are thought to debrominate into lower forms through photolytic degradation, which possibly explains this overall trend (Lagalante et al., 2011). BDE-47 was present in notably higher concentrations than BDE-99 near Houston, Texas, Biloxi, Mississippi, Mobile, Alabama, and Tampa, Florida.

PBDEs are present in 97% of adults in the United States as measured by the National Health and Nutrition Examination Survey (NHANES), with BDE-47 and BDE-99 present in the highest serum concentrations (Sjodin et al., 2008). Pregnant Mexican immigrant women living in California were found to have increasing levels of total PBDEs with each year of residence as measured by the Center for the Health Assessment of Mothers and Children of Salinas (CHAMACOS) cohort, with BDE-47 the most common congener found (Castorina et al., 2011). In fact, BDE-47 can reach higher levels in Americans than total PBDE levels in humans in other countries (Fromme et al., 2009). For 223 Mexican women living in California as measured through the CHAMACOS cohort, increasing total PBDE (including BDE-47 and BDE-99) serum concentrations were associated with lower chance of pregnancy (Harley et al., 2010).

PBDEs have been linked to developmental neurotoxicity. A 2018 review and meta-analysis of animal evidence showed that exposure to BDE-47, BDE-99, and BDE-209 affects learning (Dorman et al., 2018). In humans, large prospective cohorts showed that *in utero* exposure to PBDEs is associated with impaired executive function and poor control

of attention in children and that prenatal and postnatal PBDE exposure adversely impact externalizing behavior (Vuong et al., 2018). However, the exact mechanisms of PBDE-induced neurotoxicity in humans is unknown. In mice and rats, perinatal PBDE exposure at environmentally relevant concentrations produced persistent changes in spontaneous motor activity, including hyperactivity and decreased habituation, as well as disruptions in learning and memory (Costa and Giordano, 2007). Proposed mechanisms of PBDE developmental neurotoxicity include reduction of circulating thyroid hormones, induction of apoptotic cell death of neurons and astrocytes, oxidative stress-induced damage, interference with calcium signaling and neurotransmitters, as well as biotransformation of PBDEs (Costa et al., 2014; Costa and Giordano, 2007; Giordano et al., 2008). In zebrafish, developmental neurotoxicity of DE-71 (a commercial PBDE mixture) decreased expression of genes involved in central nervous system development and decreased locomotion activity (Chen et al., 2012).

Investigation of the involvement of gut microbiome in PBDE-induced neurotoxicity is still at an early stage. We hypothesize that PBDE neurotoxicity may be regulated through the gut-liver axis. This hypothesis is based on 1) the evidence that PBDEs are primarily oxidized by hepatic cytochrome P450s (CYPs) into hydroxylated metabolites, which are considered more toxic than the parent compound (Dingemans et al., 2008), and 2) the observations that the hepatic P450 levels are profoundly modified by lack of gut microbiome in mice (Kuno et al., 2016; Selwyn et al., 2015; Toda et al., 2009). In primary adult neural stem/progenitor cells isolated from subventricular zone of mice, the hydroxylated BDE-47 metabolite 6-OH-BDE-47, but not its parent compound, inhibited adult neurogenesis demonstrated by decreased survival, proliferation, and neuronal differentiation associated with inhibition of ERK5 signaling (Li et al., 2013). We have reported that in BDE-47 orally exposed GF mice, there was an increase in 5-OH-BDE-47 but lower levels of 4 other BDE-47 hydroxylated metabolites in liver; whereas in BDE-99 orally exposed GF mice, there was a decrease in 4 minor BDE-99 hydroxylated metabolites in liver (Liu et al., 2017b). Interestingly, the lack of gut microbiome potentiated PBDE-mediated up-regulation of Cyp1a2 and Cyp3a11 at both mRNA and protein levels in mouse liver (Liu et al., 2017b). We also showed that in large intestinal content of adult male mice, PBDEs induced a dysbiosis, including an increase in *Akkermansia muciniphila* and *Erysipelotrichaceae* *Allobaculum* spp., positively associated with increased unconjugated bile acids in serum as well as increased DNA abundance in microbial 7 $\alpha$ -dehydroxylation enzymes involved in secondary bile acid synthesis (Li et al., 2018). Although evidence in humans is limited, it has been shown in mice that deoxycholic acid (a microbial-derived secondary bile acid) may contribute to a neurological decline (McMillin and DeMorrow, 2016; McMillin et al., 2016). Microbial metabolic functions in the intestine have also been shown to be altered by DE-71 in zebrafish, associated with disruption of neural signaling, epithelial barrier integrity, inflammatory response, oxidative stress, and compromised detoxification potential in males (Vuong et al., 2018). Similar to the studies on heavy metal exposure, a mechanism linking microbiome and PBDE-induced neurotoxicity remains elusive.

**Organochlorines.** Organochlorines are a class of pesticides that include dichlorodiphenyltrichloroethane (DDT) and hexachlorocyclohexane (HCH), which were banned in the 1970s and 1980s due to their reproductive (Salihovic et al., 2016), neurobehavioral (Saeedi Saravi and Dehpour, 2016), and immunological (Mrema et al., 2013) toxicities, as well as carcinogenicity (Freund et al., 2014). In addition, organochlorines are environmentally persistent (Jayaraj et al., 2016) and are still detected in Atlantic bluefin tuna (*Thunnus thynnus*), an indicator organism for the oceanic environment and therefore a potential risk to human health if consumed chronically (Maisano et al., 2016). Chronic exposure to the major metabolites of DDT, dichlorodiphenyldichloroethylene (p,p'-DDE) and  $\beta$ -hexachlorocyclohexane ( $\beta$ -HCH), in mice altered the composition of the gut microbiota and increased the presence of bacteria with microbial

bile salt hydrolase, which is required for the production of secondary bile acids (Liu et al., 2017b). However, it is unclear how the metabolites modified the gut microbiota and how this may contribute to organochlorine neurotoxicity.

#### 4.5. Pesticides

**Carbamates.** Carbamates are derived from carbamic acid that kill insects by reversible inhibition of acetylcholinesterase (AChE). The inhibition of AChE prevents the degradation of the neurotransmitter acetylcholine causing over-stimulation of a nerve or muscle, resulting in exhaustion and tetany (Fukuto, 1990). Carbamate pesticides have increased along with organophosphorus compounds and some compounds can have toxicities for mammals and aquatic organisms.

In the rumen microbiome of cows, Ufarté et al. (2017) identified a bacterium named clone 44112 that could metabolize the carbamate fenobucarb to a less toxic compound, but failed to metabolize two other carbamates, fenoxycarb and prosulfocarb (Ufarte et al., 2017). The carbamate-specific metabolism by microbes is similar to a study showing that some isolated bacteria could metabolize fenobucarb from rice paddy soils, but not other carbamates (Kim et al., 2014). However, neither of these studies associated the microbiome to neurological health. Consumption of aldicarb in drinking water of C57BL/6 mice for 13 weeks resulted in a differential abundance of 17 genera and also significantly disrupted the brain metabolism marked by a reduction in glucose and malic acid, whereas 3-hydroxybutyric acid was increased (Gao et al., 2018b). Although toxicity was not shown to be dependent on a gut dysbiosis, this study demonstrated a potential association between aldicarb neurotoxicity and gut microbiome.

**Neonicotinoids.** Neonicotinoids are a popular and growing class of insecticide thought to selectively and irreversibly target the nicotinic acetylcholine receptors in the CNS of insects, making them generally safe for humans and animals (Han et al., 2018). Because of this selectivity, neonicotinoids also increase the mortality of pollinators, such as honey bees (*Apis mellifera*) (Ihara and Matsuda, 2018). For example, imidacloprid, the first neonicotinoid, suppresses immune function and increases the risk of pathogen-infection in honey bees (Di Prisco et al., 2013). Using *Drosophila melanogaster* (fruit fly) as a model for honey bees, it was shown that sublethal exposure to imidacloprid induced a gut dysbiosis and decreased survival following *Serratia marcescens* infection (Daisley et al., 2017). However, a study examining this effect in honey bees showed that decreased survival due to imidacloprid exposure and metabolism of the compound is not gut microbiome-dependent (Raymann et al., 2018).

**Organophosphates.** Organophosphates (OPs) are esters of phosphoric acid that covalently bind to AChE, causing over-stimulation of neurons (Ruark et al., 2013). More recently, some OPs were recognized to have off-target toxicities at concentrations below the inhibition of AChE. For example, some OPs can inhibit or activate serine hydrolases and inhibit cannabinoid receptor 1 (CNR1) (Casida and Quistad, 2004). Diazinon exposure can also alter signaling of the neurotransmitter serotonin, production of which is influenced by the microbial metabolites short-chain fatty acids (SCFAs) and tryptophan (Slotkin et al., 2008; Timofeeva et al., 2008; Waclawikova and El Aidy, 2018). Gao et al. (2017b) demonstrated that C57BL/6 mice exposed to diazinon in drinking water for 13 weeks altered gut microbiome composition (19 genera), function, and metabolic profiles in a sex-specific manner (16S rDNA sequencing and whole genome shotgun sequencing) (Gao et al., 2017b), including decreased tryptophan synthase, which is required for the biosynthesis of tryptophan in bacteria. In a follow-up study, meta-transcriptomic analysis revealed an increase in the RNA of quorum sensing genes as well as bacterial motility and sporulation, which could explain the detection of pathogenic bacteria (Gao et al., 2017a). Similarly, C57BL/6 male mice exposed to malathion for 13 weeks exhibited increased DNA expression of quorum sensing, motility, and pathogenicity genes, indicating that exposure to some OPs may increase the

gut microbiota susceptibility to colonization by pathogenic bacteria (Gao et al., 2018a). One study examined the gut microbiome of the malaria vector-containing mosquito *Anopheles albimanus* and demonstrated that mosquitos resistant to the OP fenitrothion had a relatively lower diversity than mosquitos who were not resistant, possibly due to fenitrothion-exposure selecting for a few bacteria which utilize fenitrothion as an energy source, leading to their overabundance (Dada et al., 2018). Interestingly, BALB/c mice exposed to monocrotophos for 180 days had increased production of the microbial SCFA acetate, which induced gluconeogenesis and glucose intolerance (Velmurugan et al., 2017). Fecal microbiome transplants of monocrotophos-exposed mice to unexposed mice showed that glucose intolerance was microbiome-dependent. In humans, an examination of the oral microbiome of farmworkers by season demonstrated that microbiome perturbations may act persistently as the dysbiosis persisted in winter after the potential for exposure during the spring and summer (Stanaway et al., 2017). Therefore, while the majority of these chemicals have not been investigated for gut microbiome-dependent neurotoxicity, it has been noted that gut dysbiosis causes neurotoxicity via multiple mechanisms (Galland, 2014), thus exposure to chemicals such as carbamates could potentially further these effects.

Chlorpyrifos is the most well studied OP insecticide in relation to changes in the microbiomes of humans and animal models, with some evidence for neurotoxic effects. Most notably, adult male Wistar rats exposed to chlorpyrifos for 9 weeks increased bacteria associated with neurotoxicity such as *Candidatus Arthromitus* spp., in addition to inducing metabolic disorders (obesity and diabetes) (Fang et al., 2018). Rats exposed perinatally to chlorpyrifos had altered gut microbiome profiles as adults and were associated with impaired epithelium protection (Joly Condetta et al., 2015) and induced metabolic disorders, but these conditions were partially abrogated by inulin supplementation (Reygner et al., 2016b). The microbial metabolites SCFAs and bile acids were altered in the urine of mice exposed to chlorpyrifos for 30 days (Zhao et al., 2016). The metabolome profiles correlated with changes in gut microbiome. Increased lipopolysaccharide, intestinal inflammation, and intestinal permeability indicates that the perturbations of the mouse microbiome by chlorpyrifos significantly affect gut health and the microbial metabolites that can act as signaling to the host (Zhao et al., 2016). Zebrafish exposed to chlorpyrifos also had altered gut microbiome composition (25 taxa) as well as metabolic changes in glucose and lipid metabolism, TCA cycle, and amino acid metabolism (Wang et al., 2019b). In the SHIME model, chlorpyrifos decreased the colony forming units (cfu) of *Bifidobacterium* spp. and *Lactobacillus* spp. after 15–30 days of exposure (Requile et al., 2018). Decreased *Bifidobacterium* spp. and *Lactobacillus* spp. was reflected in two other SHIME studies, as well as increased *Enterococcus* spp. (Joly et al., 2013; Reygner et al., 2016a).

Two other environmentally relevant animal models, which have been used to explore gut microbiome-chlorpyrifos interactions, are the diamondback moth (*Plutella xylostera*) and fruit fly (*Drosophila melanogaster*). The diamondback moth is an economically deleterious pest of cruciferous crops, and it is suspected that the development of pesticide-resistance is partly due to the microbiome of this insect. *In vitro* experiments from bacteria isolated from the gut microbiome of the diamondback moth revealed that *Enterococcus* spp., vitamin C, and acetylsalicylic acid enhanced resistance to chlorpyrifos; however, the bacteria do not detoxify the insecticide (Xia et al., 2018c). An earlier sequencing study found that an insecticide resistant line of diamondback moth larva had increased *Lactobacillales*, *Pseudomonadales*, and *Xanthomonadales* (Xia et al., 2013). Contradictorily, GF and antibiotic treated fruit flies exposed to chlorpyrifos lived significantly longer than conventionally raised flies, and this was due to metabolism of chlorpyrifos to the stronger AChE inhibitor chlorpyrifos oxon by *Lactobacillus plantarum* (Daisley et al., 2018). This effect was mitigated by supplementation of the human probiotic *Lactobacillus rhamnosus* GG, which binds, but does not metabolize chlorpyrifos as demonstrated in a

previous study along with *L. rhamnosus* GR-1 (Trinder et al., 2016). Studies in the diamondback moth and fruit fly demonstrate that certain bacteria species can increase or decrease the toxicity of chlorpyrifos.

Overall, microbiome may contribute to neurotoxicity and other toxicological endpoints due to organophosphate exposure, but more evidence (behavior and biochemical data) is needed to ascertain a direct mechanism of OP exposure to gut dysbiosis to neurotoxicity.

**Pyrethroids.** Pyrethroids are derived from the chrysanthemum flower (originally known as pyrethrum) and are found in many household insecticides and repellents. Pyrethroids prevent the closure of voltage-gated sodium channels, making them axonic excitotoxins that prevent repolarization that leads to paralysis of an affected organism. Pyrethroids are broad spectrum pesticides, affecting insects indiscriminately, including bees, but are poorly absorbed by humans and often are used to treat clothing and prevent mosquito-borne diseases.

Permethrin has been linked to the depressive- and anxiety-like behavior of Gulf War Illness (GWI), as well as reduced hippocampal volume, neural stem cell activity, and neurogenesis (Parihar et al., 2013). Co-exposure of permethrin and another GWI compound pyridostigmine bromide in a mouse model decreased *Lactobacillus* spp. and *Bifidobacterium* spp. and supplementation of the microbial SCFA butyrate restored gut homeostasis (Seth et al., 2018). Chronic exposure to permethrin in rats decreased abundance of the beneficial bacteria *Bifidobacterium* spp. and *Lactobacillus paracasei* in the feces (Nasuti et al., 2016). Overall, pyrethrins, and specifically permethrin, have an effect on the microbiome, but more research is needed to identify if changes in the gut microbiome are associated with the neuroinflammation and behavioral changes associated with GWI.

**Rotenoids.** Rotenoids are naturally occurring substances, particularly in the flowering plant subfamily *Faboideae* as well as *Nyctaginaceae*. Many rotenoids such as rotenone have broad spectrum insecticidal properties. A hydrophobic compound, rotenone is an inhibitor of the mitochondrial complex I that easily crosses the blood-brain barrier causing toxicity to the central nervous system. Specifically, rotenone selectively induces apoptosis in serotonergic and dopaminergic neurons (Bisbal and Sanchez, 2019; Ren and Feng, 2007; Ren et al., 2005). Interestingly, rotenone exposure in Lewis rats for 7 days induces the pathology of Parkinson's disease and is therefore used in animal models for Parkinson's disease (Betarbet et al., 2000; Bisbal and Sanchez, 2019). Mice orally exposed to rotenone for 28 days were shown to have 14 bacteria taxa at the family level dysregulated with a marked decrease in *Bifidobacterium* spp. (Pereze-Pardo et al., 2018). Furthermore, dopaminergic cell loss was inversely correlated with increased *Rikenellaceae*, *Erysipelotrichaceae*, *Ruminococcaceae*, and *S24-7*, as well as a positive correlation with *Bifidobacteriaceae*. A second study in mice that used unpredictable restraint stress for 12 weeks and 6 weeks of rotenone exposure found that rotenone exposure furthered the restraint stress-associated increase in the relative abundance of the mucin-degrading bacteria *Akkermansia muciniphila* (Dodiya et al., 2018). This result is interesting because an increase in *A. muciniphila* has been noted in patients with Parkinson's disease, possibly due to the bacterium's ability to activate TLR2 as explored in a review by Radisavljevic et al (Radisavljevic et al., 2018). In a 4-week longitudinal rotenone exposure study in mice, gastrointestinal dysfunction started at 3 weeks along with significant inflammation; rotenone also decreased the microbial richness at 3 weeks compared to baseline with decreased in Bacteroidetes and increased Firmicutes (Yang et al., 2017). However, in adult male rats exposed to rotenone for 4 weeks, there was a tendency for *Lactobacillus* spp. and *Bifidobacterium* spp. to increase despite matching intestinal inflammatory properties (Johnson et al., 2018). It should be noted that some species of *Lactobacillus* produce D-lactic acid, which can cause neurological effects (Yilmaz et al., 2018). Overall, it is unclear if gut microbiome altered by rotenone may contribute to Parkinson's disease or if rotenone neuronal toxicity induces a gut dysbiosis; further studies are necessary to identify mechanisms of toxicity and

disease pathology.

**Herbicides and fungicides.** In general, herbicides and fungicides are designed to prevent and kill unwanted plants, molds, and fungi; however, their chemical properties can have unintended neurotoxicity and hormonal disruption in humans and animals. For example, atrazine, a triazine herbicide used in corn and sugarcane fields as well as grass turfs (lawns and golf courses), demasculinizes male gonads during developmental exposure (Hayes et al., 2011). This occurs across amphibians, reptiles, and mammals and causes an increase in the size of testicular tubules, loss of Sertoli cells, and a marked loss of germ cells. One experiment in Cuban tree frogs (*Osteopilus septentrionalis*) found that atrazine did not affect the amphibian chytrid fungus *Batrachochytrium dendrobatidis* in a gut microbiome-dependent manner, however the intensity of the fungus was negatively associated with the microbial phylum Fusobacteria (Knutie et al., 2018). In zebrafish, atrazine decreased serotonin and increased inflammation in males and affected the abundance of many genera of bacteria (Chen et al., 2018b).

Glyphosate is a broad-spectrum herbicide that is the leading product for weed management by acting as a competitive inhibitor of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), preventing the biosynthesis of aromatic amino acids and other metabolites in plants and some microorganisms that have the shikimate pathway (Motta et al., 2018). However, recent evidence suggests that glyphosate has off target effects on non-shikimate organisms. For example, chronic exposure of rat hippocampal cells to 1% Roundup® (commercial product with 0.38% glyphosate) showed decreased <sup>3</sup>H-glutamate uptake with increased glutamine uptake, indicating glutamatergic excitotoxicity (Cattani et al., 2014). Glyphosate alters the behavior of honey bees, and, interestingly, the genomes of the bee gut microbiome contain EPSPS (Motta et al., 2018). Young worker bees fed glyphosate for 5 days significantly decreased the abundance of the dominant gut microbiota species and increased their susceptibility to the opportunistic pathogen *Serratia marcescens* (Motta et al., 2018). The gut microbiome of honey bee larva fed a high concentration of glyphosate (20 mg/L) had 9 taxa that were decreased relative to controls or low dose glyphosate exposures (Dai et al., 2018). Cultured gut bacteria from the gastrointestinal tracts of green turtles (*Chelonia mydas*) exposed to six different glyphosate concentrations, as well as a deionized water control, showed a decreased in four bacteria genera: *Pantoea*, *Proteus*, *Shigella*, and *Staphylococcus* (Kittle et al., 2018). Interestingly, exposing the gut bacteria of poultry *in vitro* to glyphosate demonstrated that pathogenic bacteria such as pathogenic bacteria as *Salmonella Enteritidis*, *Salmonella Gallinarum*, *Salmonella Typhimurium*, *Clostridium perfringens*, and *Clostridium botulinum* are highly resistant to glyphosate, whereas beneficial bacteria *Enterococcus faecalis*, *Enterococcus faecium*, *Bacillus badius*, *Bifidobacterium adolescentis*, and *Lactobacillus* spp. were susceptible to glyphosate (Shehata et al., 2013), indicating that glyphosate could increase the risk for pathogenic diseases. Indeed, glyphosate exposure reduces the most prevalent *Enterococcus* spp. in German cattle, which can produce bacteriocins against *Clostridium botulinum* (Kruger et al., 2013).

Several studies have used rodent models to evaluate glyphosate-induced perturbations to the gut microbiome as well as possible neurotoxic associations. Mice exposed chronically to 250 or 500 mg/kg/day of glyphosate had increased anxiety and depression like behaviors as well as decreased *Corynebacterium* spp. and *Lactobacillus* spp., indicating that key microbes could increase neurobehavioral alterations (Aitbali et al., 2018). Exposing female Sprague-Dawley rats to three doses of Roundup® increased the family *S24-7* and decreased *Lactobacillaceae*; culturable bacteria in the rat microbiome showed varying sensitivities to glyphosate, with one highly resistant *Escherichia coli* strain lacking EPSPS (Lozano et al., 2018). Another study examining the rat pups from *in utero* and developmental exposure to postnatal day 125 of glyphosate in drinking water showed an increase in *Prevotella* spp., but a decrease in *Lactobacillus* spp. at postnatal day 31 (Mao et al., 2018). However, another study found that a diet sufficient in aromatic

amino acids would prevent the antimicrobial effects of glyphosate on the rat microbiome (Nielsen et al., 2018). Overall, some bacteria that decrease from glyphosate exposure, such as *Lactobacillus* spp., may increase the neurotoxicity of glyphosate, but further studies are warranted.

#### 4.6. Plant and animal toxins

In addition to the manmade neurotoxicants, microbiome may contribute to the detoxification/toxication of plant and animal derived toxins. For example, the neurotoxin domoic acid is produced by the diatom *Pseudo-nitzschia* multiseriis and is responsible for amnesic shellfish poisoning. Domoic acid exerts its neurotoxicity through activating the excitatory glutamate receptors as a structural analog to glutamate (Bates et al., 2018; Chegini and Metcalfe, 2005; Mills et al., 2016; Smith and Swoboda, 2019). Autochthonous bacteria have been suggested to contribute to the biodegradation and disposal of domoic acid, and blue mussels and soft-shell clams have been demonstrated as the unique sources of these domoic acid-utilizing bacteria (Stewart et al., 1998).

Nicotine, which is the primary active substance in tobacco, interacts with the nicotinic receptors in CNS for excitatory signaling. Studies using mouse models have shown that nicotine exposure affects the gut microbiome in a sex-dependent manner, showing that there is a male gut microbiome-specific modulation of microbial pathways involved in carbohydrate metabolism, oxidative stress, and DNA repair. In addition, fecal metabolomics studies showed that multiple neurotransmitters, including glutamate, GABA, and glycine, as well as the neuro-active metabolites leucine and uric acid, were altered by nicotine in both sexes (Chi et al., 2017c). This indicates that gut microbiome may at least partly contribute to nicotine's CNS effect.

Cyanide can induce tremors through neuronal calcium signaling (Johnson et al., 1986) and is enriched in secondary metabolites in bamboo. Interestingly, bamboo-eating pandas are deficient in rhodanese, which is one of the essential cyanide-detoxifying enzymes as compared to other herbivores. Correspondingly, the gut microbiota of pandas have high proportions of *Pseudomonas* spp. and enriched putative genes coding for rhodanese-like enzymes involved in cyanide degradation. This diet-driven symbiotic relationship is advantageous for the host survival during evolution (Zhu et al., 2018a).

#### 4.7. Solvents

Solvents refer to a class of organic chemicals of varying lipophilicity and volatility classified by molecular structure and functional groups that generally lack a charge and are of small molecular size. Toxicity of solvents is determined by the number of carbon atoms, saturation (number of bonds between carbon atoms), configuration of the carbon atoms, halogenation, and presence of functional groups (Klaassen, 2013).

One of the greatest solvent exposures for most humans is ethanol through intoxicating beverages as well as from its use in household products, pharmaceuticals, industry, and gasoline additive. The metabolism of ethanol by the liver is well characterized by the formation of acetaldehyde by CYP2E1 or alcohol dehydrogenase followed by acetate by aldehyde dehydrogenase, and it is well known that chronic alcohol ingestion can induce alcoholic liver disease. Mice fed the liquid Liber-DeCarli diet with alcohol (5% v/v) for 6 weeks to induce alcoholic liver disease increased the fecal pH and increased fecal *Corynebacterium* spp. as well as the alkaline tolerant *Alcaligenes* spp. compared to mice not exposed to alcohol (Bull-Otterson et al., 2013). Supplementation with the probiotic bacterium *Lactobacillus rhamnosus* GG prevented the ethanol induced pathogenic changes. A combination of ethanol and the artificial sweetener saccharin increased Eubacteria in pregnant mice, whereas it decreased in non-pregnant mice; in pregnant mice, ethanol and saccharin increased the abundance of *Clostridium* spp. compared to

ethanol and weight (Labrecque et al., 2015). Another study found that alcohol dependent mice had increased *Bifidobacterium* spp., and metabolomics analysis revealed changes in bacteria relevant metabolism, including secondary bile acids and serotonin (Wang et al., 2018). Ethanol exposure for 3 weeks in mice induced anxiety and depression-like behaviors also increased *Adlercreutzia* spp., *Allobaculum* spp. and *Turicibacter* spp. and decreased *Helicobacter* spp. (Xu et al., 2018). Interestingly, decreased *Adlercreutzia* spp. was positively correlated with alcohol preference and negatively correlated with anxiety-like behavior and the decreased expression of brain-derived neurotrophic factor (BDNF) and  $\alpha 1$  subunit of  $\gamma$ -aminobutyric acid A receptor (Gabra1) in prefrontal cortex.

Alcohol dependent human subjects admitted to a gastroenterology ward for a 3-week detoxification and rehabilitation program had increased gut permeability by the second day of alcohol withdrawal. Alcohol dependent subjects with high gut permeability had decreased abundance of the genera *Ruminococcus*, *Faecalibacterium*, *Subdoligranulum*, *Oscillibacter*, and *Anaerofilum* as well as increased *Dorea* (Leclercq et al., 2014). Interestingly, increased gut permeability was negatively correlated with the reduced total number of bacteria and microbial metabolites from tryptophan metabolism, which is used for generating neurotransmission molecules such as serotonin, were low or not present in high gut permeable subjects. Mice fed a liquid diet with alcohol were protected from alcohol-induced neuroinflammation, increased small intestine and brain cytokine expression by antibiotic treatment, and antibiotics abrogated microglia activation and morphological changes in the cortex and hippocampus, demonstrating a direct connection between the gut microbiome and the CNS effects of alcohol (Lowe et al., 2018).

Associations between gut microbiome and solvents have also been investigated for formaldehyde and trichloroethylene (TCE). Formaldehyde is an irritating, gaseous solvent that is an environmental toxic hazard found in paint, cloth, cigarette smoke and exhaust gas. Formaldehyde has many detrimental effects on various tissues including skin, eye, gonads, the gastrointestinal system and the respiratory tract, as well as the nervous system, which can cause neuro-behavioral impairment and seizures (Kilburn, 1994; Tang et al., 2011). Balb/c mice exposed to formaldehyde had increased abundance of the genera *Prevotella*, *Dorea*, *Desulfovibrio*, *Adlercreutzia*, *Anaeroplasm*, *Coproccoccus*, *Candidatus Arthromitus*, *Delftia*, *Lactococcus*, and *Serratia* and decreased abundance of *Bacteroides* (Guo et al., 2018b). TCE is a clear sweet-smelling industrial solvent with a variety of uses, including dry cleaning, film cleaning, and degreaser. TCE is a CNS depressant by inhibiting GABA<sub>A</sub> and glycine receptors (Beckstead et al., 2000; Krasowski and Harrison, 2000). Mice exposed to TCE perinatally and up to postnatal day 259 decreased the abundance of the genera *Bacteroides* and *Lactobacillus* and increased the abundance of *Bifidobacterium* (Khare et al., 2019). Overall, more research is needed to understand the contribution of the microbiome to neurotoxicity by solvents.

### 5. Closing remarks: microbiome as a contributor to neurotoxicity

Taken together, our review provides a comprehensive literature update regarding the role of gut microbiome in various neurological disorders, especially during chemical-induced neurotoxicity in both human subjects and animal models. Although the research in neurotoxicants and gut-brain axis is still in its early phase, we hypothesize that gut microbiome may contribute to the pathogenesis and/or resolution of neurotoxicity by 1) direct biotransformation of the xenobiotics into neuro-reactive or inactive metabolites; 2) alteration of endogenous microbial neuro-reactive metabolites, some of which may have epigenetic reprogramming potential in regulating the transcription of host genes involved in cognitive functions in brain; 3) modulation of neuroinflammation by affecting the intestinal barrier integrity and the systemic availability of gut-derived pro-inflammatory cytokines; and 4) regulation of mucosal immune function (Fig. 1).

Among various neurological disorders and chemical-induced neurotoxicity, distinct neuroactive microbial metabolites appear to be common targets, this includes decreased SCFAs and/or decreased abundance of SCFA-producing bacteria in AD mouse models (Zhang et al., 2017), anorexia nervosa patients (Borgo et al., 2017; Morita et al., 2015), PD patients (Unger et al., 2016), ASD patients (De Angelis et al., 2013; Kang et al., 2013), as well as ionic gold (Kase et al., 1987) and permethrin/pyridostigmine bromide co-exposed mice (Seth et al., 2018). Conversely, SCFA supplementation has been shown to improve AD (Govindarajan et al., 2011; Vinolo et al., 2011), restore permethrin-induced gut dysbiosis, and stall microbiome-induced GI inflammation in Gulf War illness mouse model (Seth et al., 2018). This is likely contributed by the HDAC inhibitor and anti-inflammation properties of SCFAs (Bourassa et al., 2016; Matt et al., 2018; Vinolo et al., 2011). Tryptophan metabolism is another important common target in neurotoxicity, as it is dysregulated in ASD (Golubeva et al., 2017), MDD (Kazemi et al., 2018; Rudzki et al., 2019), alcohol addiction (Leclercq et al., 2014), as well as manganese- and organophosphate-induced neurotoxicity (Chi et al., 2017b; Slotkin et al., 2008; Timofeeva et al., 2008; Waclawikova and El Aidy, 2018). These common targets should be especially focused on as they may represent common signaling pathways that can be harnessed therapeutically to mitigate neurotoxicity.

In the endeavor to move forward the research on gut-brain axis and neurotoxicity, we propose that it will be important to consider the following aspects:

1) Beyond associations: more mechanistic investigations are needed *in vitro* and at the single microbial species resolution to establish the causality. A combination of research tools including anaerobic culture of bacteria, GF mice, fecal transplant/single strain inoculations, metabolomics, as well as biochemical and behavioral assessment of CNS functions, will likely lead to a more thorough understanding of the exact molecular mechanisms of gut microbiota-mediated regulation of CNS functions. For example, Dr. Elaine Hsiao's group demonstrated how the ketogenic diet exerts its anti-seizure effects in mice through modulating the gut microbiota by initially using a multi-omics approach of 16S rDNA sequencing and metabolomics followed by identification of key bacteria that mitigate seizures (Olson et al., 2018). The ketogenic diet promoted the enrichment of the genera *Akkermansia* and *Parabacteroides* in the intestine and correlated with reduction in systemic gamma-glutamylated amino acids and increased hippocampal GABA/glutamate levels. The dependency of gut microbiota in ketogenic diet mediated anti-seizure effect was confirmed using antibiotic-exposed mice and GF mice, inoculation of specific bacteria (*Akkermansia* and *Parabacteroides*), and fecal microbiome transfer (Olson et al., 2018). Overall, this study demonstrated which bacteria modified by ketogenic diet mechanistically contributed to seizure protection in mice.

The SHIME system, or a similar microbial culturing method, is another tool that could be used to study the specific effects of chemical exposures on bacteria and understand how the chemicals may alter gut microbiome composition in humans. Because the system uses a dynamic series of six compartments that can mimic the intestines (Joly et al., 2013; Requile et al., 2018), some bacteria that have not been cultured by traditional methods may colonize in the system. Furthermore, the bacteria cultured before and after exposure could be used to colonize GF mice to elucidate a microbiome dependent effect such that exposure causes a gut dysbiosis which leads to neurotoxicity. The SHIME system was used to characterize the effect of exposures on the microbiome, including arsenic (Yin et al., 2017, 2015; Yu et al., 2016) and chlorpyrifos (Joly et al., 2013; Requile et al., 2018; Reygnier et al., 2016a).

In general, an obstacle to study the role of intestinal bacteria using single strain anaerobic cultures is that a substantial number of the intestinal bacteria are not culturable *in vitro* (Lagier et al., 2015; Lagkouvardos et al., 2017). This is partly because the artificial media

may lack key growth factors that are provided by other symbiotic bacteria in the gut (D'Onofrio et al., 2010; Fenn et al., 2017). Co-cultures of multiple microbial species may be the solution to this problem, and this method has been shown to be successful in characterizing GABA-modulating bacteria in the human gut (Strandwitz et al., 2019). Because microbes live in a symbiotic environment in the gut, metabolites produced by one microbe may serve as important substrates for the function of another microbe. Conversely, antagonistic microbes can compete with each other for the same nutrient niche and may inhibit a particular signaling pathway under physiological or pathophysiological conditions (Das et al., 2018). Therefore, in the effort to determine the mechanistic roles of microbes at single species resolution, it is important to perform co-culture experiments to investigate the microbe-microbe interactions and identify key microbial metabolites in the context of microbial networks that contribute to the pathogenesis of neurological diseases. In addition, it is important to include host enterocytes in the co-culturing system, so as to determine the interactions between distinct microbes and host receptors (Sadaghian Sadabad et al., 2015).

In addition, steady-state pre-rRNA analysis, which quantifies the ribosomal RNA precursors instead of mature 16S rDNA, has been shown to be a more promising culture-independent tool to assess the active growth status of the bacteria. Pre-rRNA is rapidly replenished when growth-limited bacteria encounter a more favorable growth environment and such changes can occur only in viable bacteria but not in dead cells or with free nucleic acids. Pre-rRNAs also appear to be a more sensitive biomarker to environmental stress as compared to the mature rDNA signals (Cangelosi and Brabant, 1997; Cangelosi and Meschke, 2014; Mackow and Chang, 1985; Oerther et al., 2000; Srivastava and Schlessinger, 1990). Lastly, taking the advantage of the CRISPR-Cas9 gene-editing system, one could engineer individual intestinal bacteria to investigate the necessity of the microbial genes at single species resolution (Mimee et al., 2015).

2) Beyond 16S rDNA survey: investigating microbial genes and neuroactive microbial metabolites to further decipher the bacterial functions. As reviewed in Tables 1 and 2, a substantial amount of research in gut microbiome and neurological disorders has utilized 16S rDNA gene sequencing, which has provided important and valuable information regarding what taxa are differentially regulated during the progression of diseases. However, 16S rDNA gene sequencing often does not resolve taxonomy past the genus level and provides only moderately accurate predictions of the functional changes. Deep whole-metagenome shotgun sequencing, as well as a more cost-effective approach, namely shallow shotgun metagenomic sequencing, are highly recommended alternatives for high resolution taxonomic and functional microbiome analysis (Hillmann et al., 2018). In addition, we believe that more research using metatranscriptomics (Bashiardes et al., 2016) and metaproteomics (Mills et al., 2019) are needed to further characterize the gut microbiome functions beyond DNA level. Computational tools for functional predictions of the microbiome, such as FishTaco and AGORA (Magnusdottir et al., 2017; Manor and Borenstein, 2017), followed by wet lab validations using metabolomics approach (Daliri et al., 2017), will also provide further mechanistic insights into further understanding the gut-brain axis in neurotoxicity.

Specifically, building on the known neuroactive microbial metabolites that contribute to the pathogenesis and/or mitigation of neurotoxicity, it is important to determine the regulation of these microbial genes that are responsible for the production of these metabolites. As summarized in this review, SCFAs are commonly reduced in AD mouse models (Zhang et al., 2017), anorexia nervosa patients (Borgo et al., 2017; Morita et al., 2015), PD patients (Unger et al., 2016); whereas decreased SCFA-producing bacteria were also observed in intestines of ASD patients (De Angelis et al., 2013; Kang et al., 2013), ionic gold exposed mice (Kase et al., 1987), and permethrin and pyridostigmine bromide co-exposed mice (Seth et al., 2018). Dysregulation in microbial and/or host tryptophan metabolism was found associated with ASD

behavior (Golubeva et al., 2017), MDD (Kazemi et al., 2018; Rudzki et al., 2019), alcohol addition (Leclercq et al., 2014), as well as manganese- and organophosphate-induced neurotoxicities (Chi et al., 2017b; Slotkin et al., 2008; Timofeeva et al., 2008; Waclawikova and El Aidy, 2018). It is especially important to focus on the neuroactive metabolites that are commonly regulated in neurological disorders of various etiology, such as SCFAs and tryptophan microbial metabolites.

3) Beyond intestinal bacteria: characterizing the involvement of other microorganisms in neurotoxicity. Although most of the current endeavors in exploring the gut microbiome have focused on intestinal bacteria, other microorganisms are important for neurodegenerative diseases and may also provide alternative therapeutic options. These other microorganisms that could be essential to the normal function of the host or contribute to pathogenicity of diseases include fungi, protozoa, and viruses (Barko et al., 2018). Fungi have been detected in different brain regions in Alzheimer's disease patients (Alonso et al., 2018; Pisa et al., 2015), as well as in patients with amyotrophic lateral sclerosis (Alonso et al., 2017). One hypothesis is that these fungi are of gut origin and may be a result of gut leakage. Regarding viruses, it is not known what viruses in the gut microbiome may be beneficial to host health, however bacteriophage therapy may be method to alter the gut microbiome. The gut microbiome of gnotobiotic mice colonized with known human gut bacteria was modulated by lytic phages that shifted the gut microbiome composition and the gut metabolome, suggesting that bacteriophage could be used to modulate the microbiome (Hsu et al., 2019). Indeed, intravenous treatment with engineered phages was used to treat a cystic fibrosis patient with a drug-resistant strain of *Mycobacterium abscessus* (Dedrick et al., 2019). More research is needed in this field to further characterize the sources, types, and mechanistic involvement of these microorganisms during various types of neurological diseases. The bioavailability of gut-derived microorganisms and microbial products should also be assessed when considering the presence of the blood-brain barrier.

4) Beyond early-life exposure: epigenetic reprogramming via microbial metabolites. Early-life exposure to environmental chemicals has been shown to be an important contributing factor for developmental origins of human diseases. Along those lines, early-life induced dysbiosis may also predispose the host organism to delayed onset of adverse health outcomes including neurodegenerative diseases. There is a critical time window early in life to target the microbiome and modulate late-onset of Alzheimer's disease, because acute antibiotic-treatment in perinatal period resulted in long-term alterations of gut microbiota (especially Lachnospiraceae and S24-7) and reduction in brain A $\beta$  deposition, as well as reduced inflammatory signaling in serum and brain of aged AD mice (Minter et al., 2017). Neuroactive microbial metabolites, such as SCFAs which have anti-inflammatory HDAC inhibitor properties, may be beneficial against the pathogenesis of AD (Govindarajan et al., 2011; Vinolo et al., 2011). Therefore, linking gut microbiome and metabolome to host epigenome and inflammasome is an exciting approach to investigate the remote-sensing mechanisms between the gut and brain.

#### Transparency document

The Transparency document associated with this article can be found in the online version.

#### Declaration of Competing Interest

None.

#### Acknowledgements

This work is supported by the United States National Institute of Health grants T32 ES007032-39, R01GM111381, R01ES019487, and R01ES025708, the University of Washington Interdisciplinary Center of

for Exposures, Diseases, and Environment (EDGE) (P30ES007033), and the University of Washington Department of Exposure and Occupational Health Sciences Sheldon Murphy Endowment.

#### References

- Aarts, E., Ederveen, T.H.A., Naaijen, J., Zwieters, M.P., Boekhorst, J., Timmerman, H.M., Smeekens, S.P., Netea, M.G., Buitelaar, J.K., Franke, B., van Hijum, S., Arias Vasquez, A., 2017. Gut microbiome in ADHD and its relation to neural reward anticipation. *PLoS One* 12 (9), e0183509.
- Abildgaard, A., Elfving, B., Hokland, M., Wegener, G., Lund, S., 2017. Probiotic treatment reduces depressive-like behaviour in rats independently of diet. *Psychoneuroendocrinology* 79, 40–48.
- Aitbali, Y., Ba-M'hamed, S., Elhidar, N., Nafis, A., Soraa, N., Bennis, M., 2018. Glyphosate based- herbicide exposure affects gut microbiota, anxiety and depression-like behaviors in mice. *Neurotoxicol. Teratol.* 67, 44–49.
- Aizawa, E., Tsuji, H., Asahara, T., Takahashi, T., Teraishi, T., Yoshida, S., Ota, M., Koga, N., Hattori, K., Kunugi, H., 2016. Possible association of Bifidobacterium and Lactobacillus in the gut microbiota of patients with major depressive disorder. *J. Affect. Disord.* 202, 254–257.
- Akiskal, H.S., Bourgeois, M.L., Angst, J., Post, R., Moller, H., Hirschfeld, R., 2000. Re-evaluating the prevalence of and diagnostic composition within the broad clinical spectrum of bipolar disorders. *J. Affect. Disord.* 59 (Suppl 1), S5–S30.
- Akiyama, H., 1994. Inflammatory response in Alzheimer's disease. *Tohoku J. Exp. Med.* 174 (3), 295–303.
- Alam, R., Abdolmaleky, H.M., Zhou, J.R., 2017. Microbiome, inflammation, epigenetic alterations, and mental diseases. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 174 (6), 651–660.
- Alderete, T.L., Jones, R.B., Chen, Z., Kim, J.S., Habre, R., Lurmann, F., Gilliland, F.D., Goran, M.I., 2018. Exposure to traffic-related air pollution and the composition of the gut microbiota in overweight and obese adolescents. *Environ. Res.* 161, 472–478.
- Almeida, L.G., Moraes, L.A., Trigo, J.R., Omoto, C., Consoli, F.L., 2017. The gut microbiota of insecticide-resistant insects houses insecticide-degrading bacteria: a potential source for biotechnological exploitation. *PLoS One* 12 (3), e0174754.
- Alonso, R., Pisa, D., Fernandez-Fernandez, A.M., Carrasco, L., 2018. Infection of Fungi and Bacteria in brain tissue from elderly persons and patients with alzheimer's disease. *Front. Aging Neurosci.* 10, 159.
- Alonso, R., Pisa, D., Fernandez-Fernandez, A.M., Rabano, A., Carrasco, L., 2017. Fungal infection in neural tissue of patients with amyotrophic lateral sclerosis. *Neurobiol. Dis.* 108, 249–260.
- Anezaki, K., Nakano, T., 2014. Concentration levels and congener profiles of polychlorinated biphenyls, pentachlorobenzene, and hexachlorobenzene in commercial pigments. *Environ. Sci. Pollut. Res. Int.* 21 (2), 998–1009.
- Ardura-Fabregat, A., Boddeke, E., Boza-Serrano, A., Brioschi, S., Castro-Gomez, S., Ceyzeriat, K., Dansokho, C., Dierkes, T., Gelders, G., Heneka, M.T., Hoeijmakers, L., Hoffmann, A., Iaccarino, L., Jahnert, S., Kuhbandner, K., Landreth, G., Lonnemann, N., Loschmann, P.A., McManus, R.M., Paulus, A., Reemst, K., Sanchez-Caro, J.M., Tiberi, A., Van der Perren, A., Vautheny, A., Venegas, C., Webers, A., Weydt, P., Wijasa, T.S., Xiang, X., Yang, Y., 2017. Targeting neuroinflammation to treat alzheimer's disease. *CNS Drugs* 31 (12), 1057–1082.
- Arentsen, T., Raith, H., Qian, Y., Forssberg, H., Diaz Heijt, R., 2015. Host microbiota modulates development of social preference in mice. *Microb. Ecol. Health Dis.* 26, 29719.
- Asano, Y., Hiramoto, T., Nishino, R., Aiba, Y., Kimura, T., Yoshihara, K., Koga, Y., Sudo, N., 2012. Critical role of gut microbiota in the production of biologically active, free catecholamines in the gut lumen of mice. *Am. J. Physiol. Gastrointest. Liver Physiol.* 303 (11), G1288–1295.
- A.P. Association, 2013. Diagnostic and Statistical Manual of Mental Disorders, 5th ed. American Psychiatric Publishing, Arlington.
- Athauda, D., Foltynie, T., 2016. The glucagon-like peptide 1 (GLP) receptor as a therapeutic target in Parkinson's disease: mechanisms of action. *Drug Discov. Today* 21 (5), 802–818.
- Axelsson, P.B., Clausen, T.D., Petersen, A.H., Hageman, I., Pinborg, A., Kessing, L.V., Bergholt, T., Rasmussen, S.C., Keiding, N., Lokkegaard, E.C.L., 2019a. Investigating the effects of cesarean delivery and antibiotic use in early childhood on risk of later attention deficit hyperactivity disorder. *J. Child Psychol. Psychiatry* 60 (2), 151–159.
- Axelsson, P.B., Clausen, T.D., Petersen, A.H., Hageman, I., Pinborg, A., Kessing, L.V., Bergholt, T., Rasmussen, S.C., Keiding, N., Lokkegaard, E.C.L., 2019b. Relation between infant microbiota and autism?: results from a national cohort sibling design study. *Epidemiology* 30 (1), 52–60.
- Babadjouni, R.M., Hodis, D.M., Radwanski, R., Durazo, R., Patel, A., Liu, Q., Mack, W.J., 2017. Clinical effects of air pollution on the central nervous system; a review. *J. Clin. Neurosci.* 43, 16–24.
- Banerjee, T.D., Middleton, F., Faraone, S.V., 2007. Environmental risk factors for attention-deficit hyperactivity disorder. *Acta Paediatr.* 96 (9), 1269–1274.
- Barko, P.C., McMichael, M.A., Swanson, K.S., Williams, D.A., 2018. The gastrointestinal microbiome: a review. *J. Vet. Intern. Med.* 32 (1), 9–25.
- Bashiardes, S., Zilberman-Schapira, G., Elinav, E., 2016. Use of metatranscriptomics in microbiome research. *Bioinform. Biol. Insights* 10, 19–25.
- Bates, S.S., Hubbard, K.A., Lundholm, N., Montresor, M., Leaw, C.P., 2018. Pseudo-nitzschia, Nitzschia, and domoic acid: new research since 2011. *Harmful Algae* 79, 3–43.
- Beamish, L.A., Osornio-Vargas, A.R., Wine, E., 2011. Air pollution: an environmental factor contributing to intestinal disease. *J. Crohns Colitis* 5 (4), 279–286.
- Beckstead, M.J., Weiner, J.L., Eger 2nd, E.I., Gong, D.H., Mihic, S.J., 2000. Glycine and gamma-aminobutyric acid(A) receptor function is enhanced by inhaled drugs of abuse. *Mol. Pharmacol.* 57 (6), 1199–1205.
- Bendorius, M., Po, C., Muller, S., Jeltsch-David, H., 2018. From systemic inflammation to

- neuroinflammation: the case of neuropilus. *Int. J. Mol. Sci.* 19 (11).
- Betarbet, R., Sherer, T.B., MacKenzie, G., Garcia-Osuna, M., Panov, A.V., Greenamyre, J.T., 2000. Chronic systemic pesticide exposure reproduces features of Parkinson's disease. *Nat. Neurosci.* 3 (12), 1301–1306.
- Bhattacharjee, S., Lukiw, W.J., 2013. Alzheimer's disease and the microbiome. *Front. Cell. Neurosci.* 7, 153.
- Bisanz, J.E., Enos, M.K., Mwanga, J.R., Chungalucha, J., Burton, J.P., Gloor, G.B., Reid, G., 2014. Randomized open-label pilot study of the influence of probiotics and the gut microbiome on toxic metal levels in Tanzanian pregnant women and school children. *MBio* 5 (5), e01580–01514.
- Bisanz, J.E., Spanogiannopoulos, P., Pieper, L.M., Bustion, A.E., Turnbaugh, P.J., 2018. How to determine the role of the microbiome in drug disposition. *Drug Metab. Dispos.* 46 (11), 1588–1595.
- Bisbal, M., Sanchez, M., 2019. Neurotoxicity of the pesticide rotenone on neuronal polarization: a mechanistic approach. *Neural Regen. Res.* 14 (5), 762–766.
- Biscarini, F., Palazzo, F., Castellani, F., Masetti, G., Grotta, L., Cichelli, A., Martino, G., 2018. Rumen microbiome in dairy calves fed copper and grape-pomace dietary supplementations: composition and predicted functional profile. *PLoS One* 13 (11), e0205670.
- Block, M.L., Calderon-Garciduenas, L., 2009. Air pollution: mechanisms of neuroinflammation and CNS disease. *Trends Neurosci.* 32 (9), 506–516.
- Bonfili, L., Cecarini, V., Berardi, S., Scarpona, S., Suchodolski, J.S., Nasuti, C., Fiorini, D., Boarelli, M.C., Rossi, G., Eleuteri, A.M., 2017. Microbiota modulation counteracts Alzheimer's disease progression influencing neuronal proteolysis and gut hormones plasma levels. *Sci. Rep.* 7 (1), 2426.
- Borgo, F., Riva, A., Benetti, A., Casiraghi, M.C., Bertelli, S., Garbossa, S., Anselmetti, S., Scarone, S., Pontiroli, A.E., Morace, G., Borghi, E., 2017. Microbiota in anorexia nervosa: the triangle between bacterial species, metabolites and psychological tests. *PLoS One* 12 (6), e0179739.
- Borre, Y.E., O'Keefe, G.W., Clarke, G., Stanton, C., Dinan, T.G., Cryan, J.F., 2014. Microbiota and neurodevelopmental windows: implications for brain disorders. *Trends Mol. Med.* 20 (9), 509–518.
- Bourassa, M.W., Alim, I., Bultman, S.J., Ratan, R.R., 2016. Butyrate, neuroepigenetics and the gut microbiome: Can a high fiber diet improve brain health? *Neurosci. Lett.* 625, 56–63.
- Breton, J., Massart, S., Vandamme, P., De Brandt, E., Pot, B., Foligne, B., 2013. Ecotoxicology inside the gut: impact of heavy metals on the mouse microbiome. *BMC Pharmacol. Toxicol.* 14, 62.
- Bridges, K.N., Zhang, Y., Curran, T.E., Magnuson, J.T., Venables, B.J., Durrer, K.E., Allen, M.S., Roberts, A.P., 2018. Alterations to the intestinal microbiome and metabolism of *Pimephales promelas* and *Mus musculus* following exposure to dietary methylmercury. *Environ. Sci. Technol.* 52 (15), 8774–8784.
- Bull-Otterson, L., Feng, W., Kirpich, I., Wang, Y., Qin, X., Liu, Y., Gobejishvili, L., Joshi-Barve, S., Ayvaz, T., Petrosino, J., Kong, M., Barker, D., McClain, C., Barve, S., 2013. Metagenomic analyses of alcohol induced pathogenic alterations in the intestinal microbiome and the effect of *Lactobacillus rhamnosus* GG treatment. *PLoS One* 8 (1), e53028.
- Calderon-Garciduenas, L., Avila-Ramirez, J., Calderon-Garciduenas, A., Gonzalez-Heredia, T., Acuna-Ayala, H., Chao, C.K., Thompson, C., Ruiz-Ramos, R., Cortes-Gonzalez, V., Martinez-Martinez, L., Garcia-Perez, M.A., Reis, J., Mukherjee, P.S., Torres-Jardon, R., Lachmann, I., 2016. Cerebrospinal fluid biomarkers in highly exposed PM2.5 urbanites: the risk of Alzheimer's and Parkinson's diseases in Young Mexico city residents. *J. Alzheimers Dis.* 54 (2), 597–613.
- Calne, D.B., Langston, J.W., Martin, W.R., Stoessl, A.J., Ruth, T.J., Adam, M.J., Pate, B.D., Schulzer, M., 1985. Positron emission tomography after MPTP: observations relating to the cause of Parkinson's disease. *Nature* 317 (6034), 246–248.
- Cangelosi, G.A., Brabant, W.H., 1997. Depletion of pre-16S rRNA in starved *Escherichia coli* cells. *J. Bacteriol.* 179 (14), 4457–4463.
- Cangelosi, G.A., Meschke, J.S., 2014. Dead or alive: molecular assessment of microbial viability. *Appl. Environ. Microbiol.* 80 (19), 5884–5891.
- Carabotti, M., Scirocco, A., Maselli, M.A., Severi, C., 2015. The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems. *Ann. Gastroenterol.* 28 (2), 203–209.
- Carmody, R.N., Turnbaugh, P.J., 2014. Host-microbial interactions in the metabolism of therapeutic and diet-derived xenobiotics. *J. Clin. Invest.* 124 (10), 4173–4181.
- Casida, J.E., Quistad, G.B., 2004. Organophosphate toxicology: safety aspects of non-acetylcholinesterase secondary targets. *Chem. Res. Toxicol.* 17 (8), 983–998.
- Castorina, R., Bradman, A., Sjodin, A., Fenster, L., Jones, R.S., Harley, K.G., Eisen, E.A., Eskenazi, B., 2011. Determinants of serum polybrominated diphenyl ether (PBDE) levels among pregnant women in the CHAMACOS cohort. *Environ. Sci. Technol.* 45 (15), 6553–6560.
- Castro-Nallar, E., Bendall, M.L., Perez-Losada, M., Sabuncyan, S., Severance, E.G., Dickerson, F.B., Schroeder, J.R., Yolken, R.H., Crandall, K.A., 2015. Composition, taxonomy and functional diversity of the oropharynx microbiome in individuals with schizophrenia and controls. *PeerJ* 3, e1140.
- Cattani, D., de Liz, Oliveira, Cavalli, V.L., Heinz Rieg, C.E., Domingues, J.T., Dal-Cim, T., Tasca, C.I., Mena Barreto Silva, F.R., Zamoner, A., 2014. Mechanisms underlying the neurotoxicity induced by glyphosate-based herbicide in immature rat hippocampus: involvement of glutamate excitotoxicity. *Toxicology* 320, 34–45.
- Cersosimo, M.G., Raina, G.B., Pecci, C., Pellene, A., Calandra, C.R., Gutierrez, C., Micheli, F.E., Benarroch, E.E., 2013. Gastrointestinal manifestations in Parkinson's disease: prevalence and occurrence before motor symptoms. *J. Neurol.* 260 (5), 1332–1338.
- Chang, C.C., Lin, T.C., Ho, H.L., Kuo, C.Y., Li, H.H., Korolenko, T.A., Chen, W.J., Lai, T.J., Ho, Y.J., Lin, C.L., 2018. GLP-1 analogue liraglutide attenuates mutant huntingtin-induced neurotoxicity by restoration of neuronal insulin signaling. *Int. J. Mol. Sci.* 19 (9).
- Chegin, S., Metcalfe, D.D., 2005. Contemporary issues in food allergy: seafood toxin-induced disease in the differential diagnosis of allergic reactions. *Allergy Asthma Proc.* 26 (3), 183–190.
- Chen, D., Yang, X., Yang, J., Lai, G., Yong, T., Tang, X., Shuai, O., Zhou, G., Xie, Y., Wu, Q., 2017. Prebiotic effect of fructooligosaccharides from *Morinda officinalis* on Alzheimer's disease in rodent models by targeting the microbiota-gut-Brain Axis. *Front. Aging Neurosci.* 9, 403.
- Chen, J., Toyomasu, Y., Hayashi, Y., Linden, D.R., Szurszewski, J.H., Nelson, H., Farrugia, G., Kashyap, P.C., Chia, N., Ordog, T., 2016. Altered gut microbiota in female mice with persistent low body weights following removal of post-weaning chronic dietary restriction. *Genome Med.* 8 (1), 103.
- Chen, J.J., Zheng, P., Liu, Y.Y., Zhong, X.G., Wang, H.Y., Guo, Y.J., Xie, P., 2018d. Sex differences in gut microbiota in patients with major depressive disorder. *Neuropsychiatr. Dis. Treat.* 14, 647–655.
- Chen, L., Hu, C., Lok-Shun Lai, N., Zhang, W., Hua, J., Lam, P.K.S., Lam, J.C.W., Zhou, B., 2018a. Acute exposure to PBDEs at an environmentally realistic concentration causes abrupt changes in the gut microbiota and host health of zebrafish. *Environ Pollut* 240, 17–26.
- Chen, L., Yu, K., Huang, C., Yu, L., Zhu, B., Lam, P.K., Lam, J.C., Zhou, B., 2012. Prenatal transfer of polybrominated diphenyl ethers (PBDEs) results in developmental neurotoxicity in zebrafish larvae. *Environ. Sci. Technol.* 46 (17), 9727–9734.
- Chen, L., Zhang, W., Hua, J., Hu, C., Lok-Shun Lai, N., Qian, P.Y., Lam, P.K.S., Lam, J.C.W., Zhou, B., 2018b. Dysregulation of intestinal health by environmental pollutants: involvement of the estrogen receptor and aryl hydrocarbon receptor. *Environ. Sci. Technol.* 52 (4), 2323–2330.
- Chen, Z., Li, J., Gui, S., Zhou, C., Chen, J., Yang, C., Hu, Z., Wang, H., Zhong, X., Zeng, L., Chen, K., Li, P., Xie, P., 2018c. Comparative metaproteomics analysis shows altered fecal microbiota signatures in patients with major depressive disorder. *Neuroreport* 29 (5), 417–425.
- Cherbut, C., Ferrier, L., Roze, C., Anini, Y., Blottiere, H., Lecannu, G., Galmiche, J.P., 1998. Short-chain fatty acids modify colonic motility through nerves and polypeptide YY release in the rat. *Am. J. Physiol.* 275 (6), G1415–G1422.
- Chi, L., Bian, X., Gao, B., Ru, H., Tu, P., Lu, K., 2016. Sex-specific effects of arsenic exposure on the trajectory and function of the gut microbiome. *Chem. Res. Toxicol.* 29 (6), 949–951.
- Chi, L., Bian, X., Gao, B., Tu, P., Ru, H., Lu, K., 2017a. The effects of an environmentally relevant level of arsenic on the gut microbiome and its functional metagenome. *Toxicol. Sci.* 160 (2), 193–204.
- Chi, L., Gao, B., Bian, X., Tu, P., Ru, H., Lu, K., 2017b. Manganese-induced sex-specific gut microbiome perturbations in C57BL/6 mice. *Toxicol. Appl. Pharmacol.* 331, 142–153.
- Chi, L., Mahbub, R., Gao, B., Bian, X., Tu, P., Ru, H., Lu, K., 2017c. Nicotine alters the gut microbiome and metabolites of gut-brain interactions in a sex-specific manner. *Chem. Res. Toxicol.* 30 (12), 2110–2119.
- Chi, L., Xue, J., Tu, P., Lai, Y., Ru, H., Lu, K., 2019. Gut microbiome disruption altered the biotransformation and liver toxicity of arsenic in mice. *Arch. Toxicol.* 93 (1), 25–35.
- Chi, Y., Lin, Y., Zhu, H., Huang, Q., Ye, G., Dong, S., 2018. PCBs-high-fat diet interactions as mediators of gut microbiota dysbiosis and abdominal fat accumulation in female mice. *Environ Pollut* 239, 332–341.
- Choi, J.J., Eum, S.Y., Rampersaud, E., Daunert, S., Abreu, M.T., Toborek, M., 2013. Exercise attenuates PCB-induced changes in the mouse gut microbiome. *Environ. Health Perspect.* 121 (6), 725–730.
- Claus, S.P., Guillou, H., Ellero-Simatos, S., 2016. The gut microbiota: a major player in the toxicity of environmental pollutants? *NPJ Biofilms Microbiomes* 2, 16003.
- Coretti, L., Cristiano, C., Florio, E., Scala, G., Lama, A., Keller, S., Cuomo, M., Russo, R., Pero, R., Paciello, O., Mattace Raso, G., Meli, R., Coccoza, S., Calignano, A., Chiariotti, L., Lembo, F., 2017. Sex-related alterations of gut microbiota composition in the BTBR mouse model of autism spectrum disorder. *Sci. Rep.* 7, 45356.
- Coretti, L., Paparo, L., Riccio, M.P., Amato, F., Cuomo, M., Natale, A., Borrelli, L., Corrado, G., Comegna, M., Buommino, E., Castaldo, G., Bravaccio, C., Chiariotti, L., Berni Canani, R., Lembo, F., 2018. Gut microbiota features in young children with autism spectrum disorders. *Front. Microbiol.* 9, 3146.
- Corwin, E.J., Pajer, K., Paul, S., Lowe, N., Weber, M., McCarthy, D.O., 2015. Bidirectional psychoneuroimmune interactions in the early postpartum period influence risk of postpartum depression. *Brain Behav. Immun.* 49, 86–93.
- Coryell, M., McAlpine, M., Pinkham, N.V., McDermott, T.R., Walk, S.T., 2018. The gut microbiome is required for full protection against acute arsenic toxicity in mouse models. *Nat. Commun.* 9 (1), 5424.
- Costa, L.G., 2015. The neurotoxicity of organochlorine and pyrethroid pesticides. *Handb. Clin. Neurol.* 131, 135–148.
- Costa, L.G., de Laat, R., Tagliaferri, S., Pellacani, C., 2014. A mechanistic view of polybrominated diphenyl ether (PBDE) developmental neurotoxicity. *Toxicol. Lett.* 230 (2), 282–294.
- Costa, L.G., Giordano, G., 2007. Developmental neurotoxicity of polybrominated diphenyl ether (PBDE) flame retardants. *Neurotoxicology* 28 (6), 1047–1067.
- Crumeyrolle-Arias, M., Jaglin, M., Bruneau, A., Vancassel, S., Cardona, A., Dauge, V., Naudon, L., Rabot, S., 2014. Absence of the gut microbiota enhances anxiety-like behavior and neuroendocrine response to acute stress in rats. *Psychoneuroendocrinology* 42, 207–217.
- Cui, B., Su, D., Li, W., She, X., Zhang, M., Wang, R., Zhai, Q., 2018. Effects of chronic noise exposure on the microbiome-gut-brain axis in senescence-accelerated prone mice: implications for Alzheimer's disease. *J. Neuroinflammation* 15 (1), 190.
- Cusotto, S., Strain, C.R., Fouhy, F., Strain, R.G., Peterson, V.L., Clarke, G., Stanton, C., Dinan, T.G., Cryan, J.F., 2018. Differential effects of psychotropic drugs on microbiome composition and gastrointestinal function. *Psychopharmacology (Berl.)*.
- D'Onofrio, A., Crawford, J.M., Stewart, E.J., Witt, K., Gavrish, E., Epstein, S., Clardy, J., Lewis, K., 2010. Siderophores from neighboring organisms promote the growth of uncultured bacteria. *Chem. Biol.* 17 (3), 254–264.
- Dacquino, C., De Rossi, P., Spalletta, G., 2015. Schizophrenia and bipolar disorder: the road from similarities and clinical heterogeneity to neurobiological types. *Clin. Chim. Acta* 449, 49–59.
- Dada, N., Sheth, M., Liebman, K., Pinto, J., Lenhart, A., 2018. Whole metagenome sequencing reveals links between mosquito microbiota and insecticide resistance in malaria vectors. *Sci. Rep.* 8 (1), 2084.

- Dahan, D., Jude, B.A., Lamendella, R., Keesing, F., Perron, G.G., 2018. Exposure to arsenic alters the microbiome of larval zebrafish. *Front. Microbiol.* 9, 1323.
- Dai, P., Yan, Z., Ma, S., Yang, Y., Wang, Q., Hou, C., Wu, Y., Liu, Y., Diao, Q., 2018. The herbicide glyphosate negatively affects midgut bacterial communities and survival of honey bee during larvae reared in vitro. *J. Agric. Food Chem.* 66 (29), 7786–7793.
- Daisley, B.A., Trinder, M., McDowell, T.W., Collins, S.L., Sumarah, M.W., Reid, G., 2018. Microbiota-mediated modulation of organophosphorus insecticide toxicity by species-dependent interactions with lactobacilli in a *Drosophila melanogaster* insect model. *Appl. Environ. Microbiol.* 84 (9).
- Daisley, B.A., Trinder, M., McDowell, T.W., Welle, H., Dube, J.S., Ali, S.N., Leong, H.S., Sumarah, M.W., Reid, G., 2017. Neonicotinoid-induced pathogen susceptibility is mitigated by *Lactobacillus plantarum* immune stimulation in a *Drosophila melanogaster* model. *Sci. Rep.* 7 (1), 2703.
- Daliri, E.B., Wei, S., Oh, D.H., Lee, B.H., 2017. The human microbiome and metabolomics: current concepts and applications. *Crit. Rev. Food Sci. Nutr.* 57 (16), 3565–3576.
- Das, P., Ji, B., Kovatcheva-Datchary, P., Backhed, F., Nielsen, J., 2018. In vitro co-cultures of human gut bacterial species as predicted from co-occurrence network analysis. *PLoS One* 13 (3), e0195161.
- De Angelis, M., Piccolo, M., Vannini, L., Siragusa, S., De Giacomo, A., Serrazanetti, D.I., Cristofori, F., Guerzoni, M.E., Gobbetti, M., Francavilla, R., 2013. Fecal microbiota and metabolome of children with autism and pervasive developmental disorder not otherwise specified. *PLoS One* 8 (10), e76993.
- de Theije, C.G., Wopereis, H., Ramadan, M., van Eijndthoven, T., Lambert, J., Knol, J., Garssen, J., Kraneveld, A.D., Oozeer, R., 2014. Altered gut microbiota and activity in a murine model of autism spectrum disorders. *Brain Behav. Immun.* 37, 197–206.
- Dedrick, R.M., Guerrero-Bustamante, C.A., Garland, R.A., Russell, D.A., Ford, K., Harris, K., Gilmour, K.C., Soothill, J., Jacobs-Sera, D., Schooley, R.T., Hatfull, G.F., Spencer, H., 2019. Engineered bacteriophages for treatment of a patient with a disseminated drug-resistant *Mycobacterium abscessus*. *Nat. Med.* 25 (5), 730–733.
- Desai, V., Kaler, S.G., 2008. Role of copper in human neurological disorders. *Am. J. Clin. Nutr.* 88 (3), 855S–858S.
- Desbonnet, L., Clarke, G., Traplin, A., O'Sullivan, O., Crispie, F., Moloney, R.D., Cotter, P.D., Dinan, T.G., Cryan, J.F., 2015. Gut microbiota depletion from early adolescence in mice: implications for brain and behaviour. *Brain Behav. Immun.* 48, 165–173.
- Dheer, R., Patterson, J., Dudash, M., Stachler, E.N., Bibby, K.J., Stolz, D.B., Shiva, S., Wang, Z., Hazen, S.L., Barchowsky, A., Stolz, J.F., 2015. Arsenic induces structural and compositional colonic microbiome change and promotes host nitrogen and amino acid metabolism. *Toxicol. Appl. Pharmacol.* 289 (3), 397–408.
- Di Giancillo, A., Rossi, R., Martino, P.A., Aidos, L., Maghin, F., Domeneghini, C., Corino, C., 2018. Copper sulphate forms in piglet diets: microbiota, intestinal morphology and enteric nervous system glial cells. *Anim. Sci. J.* 89 (3), 616–624.
- Di Prisco, G., Cavaliere, V., Annosia, D., Varricchio, P., Caprio, E., Nazzi, F., Gargiulo, G., Pennacchio, F., 2013. Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. *Proc Natl Acad Sci U S A* 110 (46), 18466–18471.
- Diaz Hejtz, R., Wang, S., Anuar, F., Qian, Y., Bjorkholm, B., Samuelsson, A., Hibberd, M.L., Forsberg, H., Pettersson, S., 2011. Normal gut microbiota modulates brain development and behavior. *Proc Natl Acad Sci U S A* 108 (7), 3047–3052.
- Dingemans, M.M., de Groot, A., van Kleef, R.G., Bergman, A., van den Berg, M., Vijverberg, H.P., Westerink, R.H., 2008. Hydroxylation increases the neurotoxic potential of BDE-47 to affect exocytosis and calcium homeostasis in PC12 cells. *Environ. Health Perspect.* 116 (5), 637–643.
- Dobson, A.W., Erikson, K.M., Aschner, M., 2004. Manganese neurotoxicity. *Ann. N. Y. Acad. Sci.* 1012, 115–128.
- Dodiya, H.B., Forsyth, C.B., Voigt, R.M., Engen, P.A., Patel, J., Shaikh, M., Green, S.J., Naqib, A., Roy, A., Kordower, J.H., Pahan, K., Shannon, K.M., Keshavarzian, A., 2018. Chronic stress-induced gut dysfunction exacerbates Parkinson's disease phenotype and pathology in a rotenone-induced mouse model of Parkinson's disease. *Neurobiol. Dis.*
- Dong, X., Shulzhenko, N., Lemaitre, J., Greer, R.L., Peremyslova, K., Quamruzzaman, Q., Rahman, M., Hasan, O.S., Joya, S.A., Golam, M., Christiani, D.C., Morgun, A., Kile, M.L., 2017. Arsenic exposure and intestinal microbiota in children from Sirajdikhan, Bangladesh. *PLoS One* 12 (12), e0188487.
- Dorman, D.C., Chiu, W., Hales, B.F., Hauser, R., Johnson, K.J., Mantus, E., Martel, S., Robinson, K.A., Rooney, A.A., Rudel, R., Sathyanarayana, S., Schantz, S.L., Waters, K.M., 2018. Polybrominated diphenyl ether (PBDE) neurotoxicity: a systematic review and meta-analysis of animal evidence. *J. Toxicol. Environ. Health B Crit. Rev.* 21 (4), 269–289.
- Dunn, A.B., Paul, S., Ware, L.Z., Corwin, E.J., 2015. Perineal injury during childbirth increases risk of postpartum depressive symptoms and inflammatory markers. *J. Midwifery Womens Health* 60 (4), 428–436.
- Dunphy-Doherty, F., O'Mahony, S.M., Peterson, V.L., O'Sullivan, O., Crispie, F., Cotter, P.D., Wigmore, P., King, M.V., Cryan, J.F., Fone, K.C.F., 2018. Post-weaning social isolation of rats leads to long-term disruption of the gut microbiota-immune-brain axis. *Brain Behav. Immun.* 68, 261–273.
- Durand, C.M., Betancur, C., Boeckers, T.M., Bockmann, J., Chaste, P., Fauchereau, F., Nygren, G., Rastam, M., Gillberg, I.C., Anckarsater, H., Sponheim, E., Goubran-Botros, H., Delorme, R., Chabane, N., Mouren-Simeoni, M.C., de Mas, P., Bieth, E., Roge, B., Heron, D., Burglen, L., Gillberg, C., Leboyer, M., Bourgeron, T., 2007. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat. Genet.* 39 (1), 25–27.
- Eckburg, P.B., Bik, E.M., Bernstein, C.N., Purdom, E., Dethlefsen, L., Sargent, M., Gill, S.R., Nelson, K.E., Relman, D.A., 2005. Diversity of the human intestinal microbial flora. *Science* 308 (5728), 1635–1638.
- Eikelenboom, P., Zhan, S.S., van Gool, W.A., Allsop, D., 1994. Inflammatory mechanisms in Alzheimer's disease. *Trends Pharmacol. Sci.* 15 (12), 447–450.
- El-Ansary, A., Bacha, A.B., Bjorklund, G., Al-Orfi, N., Bhat, R.S., Moubayed, N., Abed, K., 2018. Probiotic treatment reduces the autistic-like excitation/inhibition imbalance in juvenile hamsters induced by orally administered propionic acid and clindamycin. *Metab. Brain Dis.* 33 (4), 1155–1164.
- Evans, S.J., Bassis, C.M., Hein, R., Assari, S., Flowers, S.A., Kelly, M.B., Young, V.B., Ellingrod, V.E., McInnis, M.G., 2017. The gut microbiome composition associates with bipolar disorder and illness severity. *J. Psychiatr. Res.* 87, 23–29.
- Fang, B., Li, J.W., Zhang, M., Ren, F.Z., Pang, G.F., 2018. Chronic chlorpyrifos exposure elicits diet-specific effects on metabolism and the gut microbiome in rats. *Food Chem. Toxicol.* 111, 144–152.
- Faraone, S.V., Asherson, P., Banaschewski, T., Biederman, J., Buitelaar, J.K., Ramos-Quiroga, J.A., Rohde, L.A., Sonuga-Barke, E.J., Tannock, R., Franke, B., 2015. Attention-deficit/hyperactivity disorder. *Nat. Rev. Dis. Primers* 1, 15020.
- Faraone, S.V., Glatt, S.J., 2010. A comparison of the efficacy of medications for adult attention-deficit/hyperactivity disorder using meta-analysis of effect sizes. *J. Clin. Psychiatry* 71 (6), 754–763.
- Fenn, K., Strandwitz, P., Stewart, E.J., Dimise, E., Rubin, S., Gurubacharya, S., Clardy, J., Lewis, K., 2017. Quinones are growth factors for the human gut microbiota. *Microbiome* 5 (1), 161.
- Finegold, S.M., Dowd, S.E., Gontcharova, V., Liu, C., Henley, K.E., Wolcott, R.D., Youn, E., Summan, P.H., Granpeesheh, D., Dixon, D., Liu, M., Molitoris, D.R., Green 3rd, J.A., 2010. Pyrosequencing study of fecal microflora of autistic and control children. *Anaerobe* 16 (4), 444–453.
- Florez, A.B., Sierra, M., Ruas-Madiedo, P., Mayo, B., 2016. Susceptibility of lactic acid bacteria, bifidobacteria and other bacteria of intestinal origin to chemotherapeutic agents. *Int. J. Antimicrob. Agents* 48 (5), 547–550.
- Flowers, S.A., Baxter, N.T., Ward, K.M., Kraal, A.Z., McInnis, M.G., Schmidt, T.M., Ellingrod, V.L., 2019. Effects of atypical antipsychotic treatment and resistant starch supplementation on gut microbiome composition in a cohort of patients with bipolar disorder or schizophrenia. *Pharmacotherapy* 39 (2), 161–170.
- Flowers, S.A., Evans, S.J., Ward, K.M., McInnis, M.G., Ellingrod, V.L., 2017. Interaction between atypical antipsychotics and the gut microbiome in a bipolar disease cohort. *Pharmacotherapy* 37 (3), 261–267.
- Fournier, C.N., Houser, M., Tansey, M.G., Glass, J.D., Hertzberg, V.S., 2018. The gut microbiome and neuroinflammation in amyotrophic lateral sclerosis? Emerging clinical evidence. *Neurobiol. Dis.*
- Fraser, A.G., McQueen, I.N., Watt, A.H., Stephens, M.R., 1985. Peripheral neuropathy during long-term high-dose amiodarone therapy. *J. Neurol Neurosurg Psychiatry* 48 (6), 576–578.
- Frazier, K.R., Moore, J.A., Long, T.E., 2019. Antibacterial activity of disulfiram and its metabolites. *J. Appl. Microbiol.* 126 (1), 79–86.
- Freund, R., Kelsberg, G., Safranek, S., 2014. Clinical Inquiry: do oral contraceptives put women with a family history of breast cancer at increased risk? *J. Fam. Pract.* 63 (9), 540–549.
- Fromme, H., Korner, W., Shahin, N., Wanner, A., Albrecht, M., Boehmer, S., Parlar, H., Mayer, R., Liebl, B., Bolte, G., 2009. Human exposure to polybrominated diphenyl ethers (PBDE), as evidenced by data from a duplicate diet study, indoor air, house dust, and biomonitoring in Germany. *Environ. Int.* 35 (8), 1125–1135.
- Frye, R.E., Rose, S., Chacko, J., Wynne, R., Bennuri, S.C., Slatery, J.C., Tippett, M., Delhey, L., Melnyk, S., Kahler, S.G., MacFabe, D.F., 2016. Modulation of mitochondrial function by the microbiome metabolite propionic acid in autism and control cell lines. *Transl. Psychiatry* 6 (10), e927.
- Fu, Z.D., Cui, J.Y., 2017. Remote sensing between liver and intestine: importance of microbial metabolites. *Curr. Pharmacol. Rep.* 3 (3), 101–113.
- Fukuto, T.R., 1990. Mechanism of action of organophosphorus and carbamate insecticides. *Environ. Health Perspect.* 87, 245–254.
- Gabanyi, I., Muller, P.A., Feighery, L., Oliveira, T.Y., Costa-Pinto, F.A., Mucida, D., 2016. Neuro-immune interactions drive tissue programming in intestinal macrophages. *Cell* 164 (3), 378–391.
- Galland, L., 2014. The gut microbiome and the brain. *J. Med. Food* 17 (12), 1261–1272.
- Gao, B., Bian, X., Chi, L., Tu, P., Ru, H., Lu, K., 2017a. Editor's highlight: Organophosphate Diazinon altered quorum sensing, cell motility, stress response, and carbohydrate metabolism of gut microbiome. *Toxicol. Sci.* 157 (2), 354–364.
- Gao, B., Bian, X., Mahub, R., Lu, K., 2017b. Sex-specific effects of organophosphate diazinon on the gut microbiome and its metabolic functions. *Environ. Health Perspect.* 125 (2), 198–206.
- Gao, B., Chi, L., Mahub, R., Bian, X., Tu, P., Ru, H., Lu, K., 2017c. Multi-omics reveals that lead exposure disturbs gut microbiome development, key metabolites, and metabolic pathways. *Chem. Res. Toxicol.* 30 (4), 996–1005.
- Gao, B., Chi, L., Tu, P., Bian, X., Thomas, J., Ru, H., Lu, K., 2018a. The organophosphate malathion disturbs gut microbiome development and the quorum-sensing system. *Toxicol. Lett.* 283, 52–57.
- Gao, B., Chi, L., Tu, P., Gao, N., Lu, K., 2018b. The carbamate aldicarb altered the gut microbiome, metabolome and lipidome of C57BL/6J mice. *Chem. Res. Toxicol.*
- Gaulke, C.A., Rolshoven, J., Wong, C.P., Hudson, L.G., Ho, E., Sharpton, T.J., 2018. Marginal zinc deficiency and environmentally relevant concentrations of arsenic elicit combined effects on the gut microbiome. *mSphere* 3 (6).
- Gautam, A., D'Arpa, P., Donohue, D.E., Muhie, S., Chakraborty, N., Luke, B.T., Grapov, D., Carroll, E.E., Meyerhoff, J.L., Hammamieh, R., Jett, M., 2015. Acute and chronic plasma metabolomic and liver transcriptomic stress effects in a mouse model with features of post-traumatic stress disorder. *PLoS One* 10 (1), e0117092.
- Gautam, A., Kumar, R., Chakraborty, N., Muhie, S., Hoke, A., Hammamieh, R., Jett, M., 2018. Altered fecal microbiota composition in all male aggressor-exposed rodent model simulating features of post-traumatic stress disorder. *J. Neurosci. Res.* 96 (7), 1311–1323.
- Gavin, N.I., Gaynes, B.N., Lohr, K.N., Meltzer-Brody, S., Gartlehner, G., Swinson, T., 2005. Perinatal depression: a systematic review of prevalence and incidence. *Obstet. Gynecol.* 106 (5 Pt 1), 1071–1083.
- Giordano, G., Kavanagh, T.J., Costa, L.G., 2008. Neurotoxicity of a polybrominated diphenyl ether mixture (DE-71) in mouse neurons and astrocytes is modulated by intracellular glutathione levels. *Toxicol. Appl. Pharmacol.* 232 (2), 161–168.
- Giri, S.S., Yun, S., Jun, J.W., Kim, H.J., Kim, S.G., Kang, J.W., Kim, S.W., Han, S.J., Sukumaran, V., Park, S.C., 2018. Therapeutic effect of intestinal autochthonous

- Lactobacillus reuteri P16 against waterborne lead toxicity in *Cyprinus carpio*. *Front. Immunol.* 9, 1824.
- Golubeva, A.V., Joyce, S.A., Moloney, G., Burokas, A., Sherwin, E., Arbolea, S., Flynn, I., Khochanskiy, D., Moya-Perez, A., Peterson, V., Rea, K., Murphy, K., Makarova, O., Buravkov, S., Hyland, N.P., Stanton, C., Clarke, G., Gahan, C.G.M., Dinan, T.G., Cryan, J.F., 2017. Microbiota-related changes in bile acid & tryptophan metabolism are associated with gastrointestinal dysfunction in a mouse model of autism. *EBioMedicine* 24, 166–178.
- Gomez, G.I., Falcon, R.V., Maturana, C.J., Labra, V.C., Salgado, N., Rojas, C.A., Oyarzun, J.E., Cerpa, W., Quintanilla, R.A., Orellana, J.A., 2018. Heavy alcohol exposure activates astroglial hemichannels and pannexons in the Hippocampus of adolescent rats: effects on neuroinflammation and astrocyte arborization. *Front. Cell. Neurosci.* 12, 472.
- Gondalia, S.V., Palombo, E.A., Knowles, S.R., Cox, S.B., Meyer, D., Austin, D.W., 2012. Molecular characterisation of gastrointestinal microbiota of children with autism (with and without gastrointestinal dysfunction) and their neurotypical siblings. *Autism Res.* 5 (6), 419–427.
- Goolsby, T.A., Jakeman, B., Gaynes, R.P., 2018. Clinical relevance of metronidazole and peripheral neuropathy: a systematic review of the literature. *Int. J. Antimicrob. Agents* 51 (3), 319–325.
- Gorchev, H.G., Jelinek, C.F., 1985. A review of the dietary intakes of chemical contaminants. *Bull. World Health Organ.* 63 (5), 945–962.
- Govindarajan, N., Agis-Balboa, R.C., Walter, J., Sananbenesi, F., Fischer, A., 2011. Sodium butyrate improves memory function in an Alzheimer's disease mouse model when administered at an advanced stage of disease progression. *J. Alzheimers Dis.* 26 (1), 187–197.
- Graham Lantos, P.L., 1997. *Greenfield's Neuropathology*. Arnold, London.
- Grigg, J.B., Sonnenberg, G.F., 2017. Host-microbiota interactions shape local and systemic inflammatory diseases. *J. Immunol.* 198 (2), 564–571.
- Guo, G., Yumvihoze, E., Poulain, A.J., Man Chan, H., 2018a. Monomethylmercury degradation by the human gut microbiota is stimulated by protein amendments. *J. Toxicol. Sci.* 43 (12), 717–725.
- Guo, J., Zhao, Y., Jiang, X., Li, R., Xie, H., Ge, L., Xie, B., Yang, X., Zhang, L., 2018b. Exposure to formaldehyde perturbs the mouse gut microbiome. *Genes (Basel)* 9 (4).
- Haiser, H.J., Turnbaugh, P.J., 2013. Developing a metagenomic view of xenobiotic metabolism. *Pharmacol. Res.* 69 (1), 21–31.
- Hallmayer, J., Cleveland, S., Torres, A., Phillips, J., Cohen, B., Torigoe, T., Miller, J., Fedele, A., Collins, J., Smith, K., Lotspeich, L., Croen, L.A., Ozonoff, S., Lajonchere, C., Grether, J.K., Risch, N., 2011. Genetic heritability and shared environmental factors among twin pairs with autism. *Arch. Gen. Psychiatry* 68 (11), 1095–1102.
- Hamad, A.F., Alessi-Severini, S., Mahmud, S.M., Brownell, M., Kuo, I.F., 2018. Early childhood antibiotics use and autism spectrum disorders: a population-based cohort study. *Int. J. Epidemiol.* 47 (5), 1497–1506.
- Han, W., Tian, Y., Shen, X., 2018. Human exposure to neonicotinoid insecticides and the evaluation of their potential toxicity: an overview. *Chemosphere* 192, 59–65.
- Harach, T., Marungu, N., Duthilleul, N., Cheatham, V., Mc Coy, K.D., Frisoni, G., Neher, J.J., Fak, F., Jucker, M., Lasser, T., Bolmont, T., 2017. Reduction of Abeta amyloid pathology in APPPS1 transgenic mice in the absence of gut microbiota. *Sci. Rep.* 7, 41802.
- Harley, K.G., Marks, A.R., Chevrier, J., Bradman, A., Sjodin, A., Eskenazi, B., 2010. PBDE concentrations in women's serum and fecundability. *Environ. Health Perspect.* 118 (5), 699–704.
- Harold, G.T., Leve, L.D., Barrett, D., Elam, K., Neiderhiser, J.M., Natsuaki, M.N., Shaw, D.S., Reiss, D., Thapar, A., 2013. Biological and rearing mother influences on child ADHD symptoms: revisiting the developmental interface between nature and nurture. *J. Child Psychol. Psychiatry* 54 (10), 1038–1046.
- Hayes, T.B., Anderson, L.L., Beasley, V.R., de Solla, S.R., Iguchi, T., Ingraham, H., Kestemont, P., Kniwald, J., Kniwald, Z., Langlois, V.S., Luque, E.H., McCoy, K.A., Munoz-de-Toro, M., Oka, T., Oliveira, C.A., Orton, F., Ruby, S., Suzawa, M., Tavera-Mendoza, L.E., Trudeau, V.L., Victor-Costa, A.B., Willingham, E., 2011. Demasculinization and feminization of male gonads by atrazine: consistent effects across vertebrate classes. *J. Steroid Biochem. Mol. Biol.* 127 (1–2), 64–73.
- Hebert, L.E., Weuve, J., Scherr, P.A., Evans, D.A., 2013. Alzheimer disease in the United States (2010–2050) estimated using the 2010 census. *Neurology* 80 (19), 1778–1783.
- Heiss, C.N., Olofsson, L.E., 2019. The role of the gut microbiota in development, function and disorders of the central nervous system and the enteric nervous system. *J. Neuroendocrinol.* e12684.
- Hemmings, S.M.J., Malan-Muller, S., van den Heuvel, L.L., Demmitt, B.A., Stanislawski, M.A., Smith, D.G., Bohr, A.D., Stamper, C.E., Hyde, E.R., Morton, J.T., Marotz, C.A., Siebler, P.H., Braspenning, M., Van Crieling, W., Hoisington, A.J., Brenner, L.A., Postolache, T.T., McQueen, M.B., Krauter, K.S., Knight, R., Seedat, S., Lowry, C.A., 2017. The microbiome in posttraumatic stress disorder and trauma-exposed controls: an exploratory study. *Psychosom. Med.* 79 (8), 936–946.
- Herpertz-Dahlmann, B., 2009. Adolescent eating disorders: definitions, symptomatology, epidemiology and comorbidity. *Child Adolesc. Psychiatr. Clin. N. Am.* 18 (1), 31–47.
- Herrick, R.F., 2010. PCBs in school-persistent chemicals, persistent problems. *New Solut.* 20 (1), 115–126.
- Herrick, R.F., Stewart, J.H., Allen, J.G., 2016. Review of PCBs in US schools: a brief history, an estimate of the number of impacted schools, and an approach for evaluating indoor air samples. *Environ. Sci. Pollut. Res. Int.* 23 (3), 1975–1985.
- Hicks, S.D., Uhlig, R., Afshari, P., Williams, J., Chronos, M., Tierney-Aves, C., Wagner, K., Middleton, F.A., 2018. Oral microbiome activity in children with autism spectrum disorder. *Autism Res.* 11 (9), 1286–1299.
- Hill-Burns, E.M., Debelius, J.W., Morton, J.T., Wissemann, W.T., Lewis, M.R., Wallen, Z.D., Peddada, S.D., Factor, S.A., Molho, E., Zabetian, C.P., Knight, R., Payami, H., 2017. Parkinson's disease and Parkinson's disease medications have distinct signatures of the gut microbiome. *Mov. Disord.* 32 (5), 739–749.
- Hillmann, B., Al-Ghalith, G.A., Shields-Cutler, R.R., Zhu, Q., Gohl, D.M., Beckman, K.B., Knight, R., Knights, D., 2018. Evaluating the information content of shallow shotgun metagenomics. *mSystems* 3 (6).
- Ho, L., Ono, K., Tsuji, M., Mazzola, P., Singh, R., Pasinetti, G.M., 2018. Protective roles of intestinal microbiota derived short chain fatty acids in Alzheimer's disease-type beta-amyloid neuropathological mechanisms. *Expert Rev. Neurother.* 18 (1), 83–90.
- Hoen, A.G., Madan, J.C., Li, Z., Coker, M., Lundgren, S.N., Morrison, H.G., Palys, T., Jackson, B.P., Sogin, M.L., Cottingham, K.L., Karagas, M.R., 2018. Sex-specific associations of infants' gut microbiome with arsenic exposure in a US population. *Sci. Rep.* 8 (1), 12627.
- Hoffman, J.D., Parikh, I., Green, S.J., Chlipala, G., Mohnhey, R.P., Keaton, M., Bauer, B., Hartz, A.M.S., Lin, A.L., 2017. Age drives distortion of brain metabolic, vascular and cognitive functions, and the gut microbiome. *Front. Aging Neurosci.* 9, 298.
- Hojberg, O., Canibe, N., Poulsen, H.D., Hedemann, M.S., Jensen, B.B., 2005. Influence of dietary zinc oxide and copper sulfate on the gastrointestinal ecosystem in newly weaned piglets. *Appl. Environ. Microbiol.* 71 (5), 2267–2277.
- Hsu, B.B., Gibson, T.E., Yeliseyev, V., Liu, Q., Lyon, L., Bry, L., Silver, P.A., Gerber, G.K., 2019. Dynamic modulation of the gut microbiota and metabolome by bacteriophages in a mouse model. *Cell Host Microbe* 25 (6), 803–814 e805.
- Hu, D., Hornbuckle, K.C., 2010. Inadvertent polychlorinated biphenyls in commercial paint pigments. *Environ. Sci. Technol.* 44 (8), 2822–2827.
- Huang, C., Yang, X., Zeng, B., Zeng, L., Gong, X., Zhou, C., Xia, J., Lian, B., Qin, Y., Yang, L., Liu, L., Xie, P., 2019. Proteomic analysis of olfactory bulb suggests CACNA1E as a promoter of CREB signaling in microbiota-induced depression. *J. Proteomics* 194, 132–147.
- Huf, F., Bandiera, S., Muller, C.B., Gea, L., Carvalho, F.B., Rahmeier, F.L., Reiter, K.C., Tortorelli, L.S., Gomez, R., da Cruz Fernandes, M., 2019. Comparative study on the effects of cigarette smoke exposure, ethanol consumption and association: behavioral parameters, apoptosis, glial fibrillary acid protein and S100beta immunoreactivity in different regions of the rat hippocampus. *Alcohol* 77, 101–112.
- Hurley, S., Goldberg, D., Nelson, D.O., Guo, W., Wang, Y., Baek, H.G., Park, J.S., Petreas, M., Bernstein, L., Anton-Culver, H., Reynolds, P., 2017. Temporal evaluation of polybrominated diphenyl ether (PBDE) serum levels in middle-aged and older California women, 2011–2015. *Environ. Sci. Technol.* 51 (8), 4697–4704.
- Ihara, M., Matsuda, K., 2018. Neonicotinoids: molecular mechanisms of action, insights into resistance and impact on pollinators. *Curr. Opin. Insect Sci.* 30, 86–92.
- Iyer, L.M., Aravind, L., Coon, S.L., Klein, D.C., Koonin, E.V., 2004. Evolution of cell-cell signaling in animals: did late horizontal gene transfer from bacteria have a role? *Trends Genet.* 20 (7), 292–299.
- Janakiraman, M., Krishnamoorthy, G., 2018. Emerging role of diet and microbiota interactions in neuroinflammation. *Front. Immunol.* 9, 2067.
- Jayaraj, R., Megha, P., Sreedev, P., 2016. Organochlorine pesticides, their toxic effects on living organisms and their fate in the environment. *Interdiscip. Toxicol.* 9 (3–4), 90–100.
- Jiang, H., Ling, Z., Zhang, Y., Mao, H., Ma, Z., Yin, Y., Wang, W., Tang, W., Tan, Z., Shi, J., Li, L., Ruan, B., 2015. Altered fecal microbiota composition in patients with major depressive disorder. *Brain Behav. Immun.* 48, 186–194.
- Jiang, H.Y., Zhou, Y.Y., Zhou, G.L., Li, Y.C., Yuan, J., Li, X.H., Ruan, B., 2018. Gut microbiota profiles in treatment-naïve children with attention deficit hyperactivity disorder. *Behav. Brain Res.* 347, 408–413.
- Johnson, J.D., Meisenheimer, T.L., Isom, G.E., 1986. Cyanide-induced neurotoxicity: role of neuronal calcium. *Toxicol. Appl. Pharmacol.* 84 (3), 464–469.
- Johnson, M.E., Stringer, A., Bobrovskaya, L., 2018. Rotenone induces gastrointestinal pathology and microbiota alterations in a rat model of Parkinson's disease. *Neurotoxicology* 65, 174–185.
- Johnston, M.V., Goldstein, G.W., 1998. Selective vulnerability of the developing brain to lead. *Curr. Opin. Neurol.* 11 (6), 689–693.
- Joly, C., Gay-Queheillard, J., Leke, A., Chardon, K., Delanaud, S., Bach, V., Khorsi-Cauet, H., 2013. Impact of chronic exposure to low doses of chlorpyrifos on the intestinal microbiota in the Simulator of the Human Intestinal Microbial Ecosystem (SHIME) and in the rat. *Environ. Sci. Pollut. Res. Int.* 20 (5), 2726–2734.
- Joly Condet, C., Bach, V., Mayeur, C., Gay-Queheillard, J., Khorsi-Cauet, H., 2015. Chlorpyrifos exposure during perinatal period affects intestinal microbiota associated with delay of maturation of digestive tract in rats. *J. Pediatr. Gastroenterol. Nutr.* 61 (1), 30–40.
- Kakumani, M.L., Reeves, A.M., Anderson, T.D., Rodrigues, R.R., Williams, M.A., 2016. Honey bee gut microbiome is altered by in-hive pesticide exposures. *Front. Microbiol.* 7, 1255.
- Kang, D.W., Park, J.G., Ilhan, Z.E., Wallstrom, G., Labaer, J., Adams, J.B., Krajmalnik-Brown, R., 2013. Reduced incidence of *Prevotella* and other fermenters in intestinal microflora of autistic children. *PLoS One* 8 (7), e68322.
- Kania-Korwel, I., Lukaszewicz, T., Barnhart, C.D., Stamou, M., Chung, H., Kelly, K.M., Bandiera, S., Lein, P.J., Lehmler, H.J., 2017. Editor's highlight: congener-specific disposition of chiral polychlorinated biphenyls in lactating mice and their offspring: implications for PCB developmental neurotoxicity. *Toxicol. Sci.* 158 (1), 101–115.
- Kase, H., Kaneko, M., Yamada, K., 1987. K-13, a novel inhibitor of angiotensin I converting enzyme produced by *Micromonospora halophytica* subsp. *xilixia*. I. Fermentation, isolation and biological properties. *J. Antibiot (Tokyo)* 40 (4), 450–454.
- Kazemi, A., Noorbala, A.A., Azam, K., Eskandari, M.H., Djafarian, K., 2018. Effect of probiotic and prebiotic vs placebo on psychological outcomes in patients with major depressive disorder: a randomized clinical trial. *Clin. Nutr.*
- Kelly, J.R., Borre, Y.C.O.B., Patterson, E., El Aïdy, S., Deane, J., Kennedy, P.J., Beers, S., Scott, K., Moloney, G., Hoban, A.E., Scott, L., Fitzgerald, P., Ross, P., Stanton, C., Clarke, G., Cryan, J.F., Dinan, T.G., 2016. Transferring the blues: depression-associated gut microbiota induces neurobehavioural changes in the rat. *J. Psychiatr. Res.* 82, 109–118.
- Kelly, J.R., Minuto, C., Cryan, J.F., Clarke, G., Dinan, T.G., 2017. Cross talk: the microbiota and neurodevelopmental disorders. *Front. Neurosci.* 11, 490.
- Kessler, R.C., 2000. Posttraumatic stress disorder: the burden to the individual and to society. *J. Clin. Psychiatry* 61 (Suppl 5), 4–12 discussion 13–14.
- Khalilnezhad, A., Taskiran, D., 2018. Protective effects of glucagon-like peptide-1 (GLP-1) analogue exenatide against glucose and fructose-induced neurotoxicity. *Int. J.*

- Neurosci. 1–24.
- Khare, S., Gokulan, K., Williams, K., Bai, S., Gilbert, K.M., Blossom, S.J., 2019. Irreversible effects of trichloroethylene on the gut microbial community and gut-associated immune responses in autoimmune-prone mice. *J. Appl. Toxicol.* 39 (2), 209–220.
- Kilburn, K.H., 1994. Neurobehavioral impairment and seizures from formaldehyde. *Arch. Environ. Health* 49 (1), 37–44.
- Kim, I., Kim, D.U., Kim, N.H., Ka, J.O., 2014. Isolation and characterization of fenobucarb-degrading bacteria from rice paddy soils. *Biodegradation* 25 (3), 383–394.
- Kish, L., Hotte, N., Kaplan, G.G., Vincent, R., Tso, R., Ganzle, M., Rioux, K.P., Thiesen, A., Barkema, H.W., Wine, E., Madsen, K.L., 2013. Environmental particulate matter induces murine intestinal inflammatory responses and alters the gut microbiome. *PLoS One* 8 (4), e62220.
- Kittle, R.P., McDermid, K.J., Muehlstein, L., Balazs, G.H., 2018. Effects of glyphosate herbicide on the gastrointestinal microflora of Hawaiian green turtles (*Chelonia mydas*) Linnaeus. *Mar. Pollut. Bull.* 127, 170–174.
- Klaassen, C., 2013. In: Klaassen, C. (Ed.), *Casarett and Doull's Toxicology: The Basic Science of Poisons*, 8th ed. McGraw-Hill Education.
- Klaassen, C.D., Cui, J.Y., 2015. Review: mechanisms of how the intestinal microbiota alters the effects of drugs and bile acids. *Drug Metab. Dispos.* 43 (10), 1505–1521.
- Kleiman, S.C., Watson, H.J., Bulik-Sullivan, E.C., Huh, E.Y., Tarantino, L.M., Bulik, C.M., Carroll, I.M., 2015. The intestinal microbiota in acute anorexia nervosa and during renourishment: relationship to depression, anxiety, and eating disorder psychopathology. *Psychosom. Med.* 77 (9), 969–981.
- Knapp, M., Mangalore, R., Simon, J., 2004. The global costs of schizophrenia. *Schizophr. Bull.* 30 (2), 279–293.
- Knutie, S.A., Gabor, C.R., Kohl, K.D., Rohr, J.R., 2018. Do host-associated gut microbiota mediate the effect of an herbicide on disease risk in frogs? *J. Anim. Ecol.* 87 (2), 489–499.
- Kopin, I.J., 1987. MPTP: an industrial chemical and contaminant of illicit narcotics stimulates a new era in research on Parkinson's disease. *Environ. Health Perspect.* 75, 45–51.
- Korrick, S.A., Sagiv, S.K., 2008. Polychlorinated biphenyls, organochlorine pesticides and neurodevelopment. *Curr. Opin. Pediatr.* 20 (2), 198–204.
- Kosnicki, K.L., Penprase, J.C., Cintora, P., Torres, P.J., Harris, G.L., Brasser, S.M., Kelley, S.T., 2018. Effects of moderate, voluntary ethanol consumption on the rat and human gut microbiome. *Addict. Biol.*
- Krasowski, M.D., Harrison, N.L., 2000. The actions of ether, alcohol and alkane general anaesthetics on GABA and glycine receptors and the effects of TM2 and TM3 mutations. *Br. J. Pharmacol.* 129 (4), 731–743.
- Kruger, M., Shehata, A.A., Schrod, W., Rodloff, A., 2013. Glyphosate suppresses the antagonistic effect of *Enterococcus* spp. on *Clostridium botulinum*. *Anaerobe* 20, 74–78.
- Kuno, T., Hirayama-Kurogi, M., Ito, S., Ohtsuki, S., 2016. Effect of intestinal flora on protein expression of drug-metabolizing enzymes and transporters in the liver and kidney of germ-free and antibiotics-treated mice. *Mol. Pharm.* 13 (8), 2691–2701.
- Labrecque, M.T., Malone, D., Caldwell, K.E., Allan, A.M., 2015. Impact of ethanol and saccharin on fecal microbiome in pregnant and non-pregnant mice. *J. Pregnancy Child Health* 2 (5).
- Lagalante, A.F., Shedden, C.S., Greenbacker, P.W., 2011. Levels of polybrominated diphenyl ethers (PBDEs) in dust from personal automobiles in conjunction with studies on the photochemical degradation of decabromodiphenyl ether (BDE-209). *Environ. Int.* 37 (5), 899–906.
- Lagier, J.C., Hugon, P., Khelaifa, S., Fournier, P.E., La Scola, B., Raouf, D., 2015. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin. Microbiol. Rev.* 28 (1), 237–264.
- Lagkouvardos, I., Overmann, J., Clavel, T., 2017. Cultured microbes represent a substantial fraction of the human and mouse gut microbiota. *Gut Microbes* 8 (5), 493–503.
- Lai, F., Jiang, R., Xie, W., Liu, X., Tang, Y., Xiao, H., Gao, J., Jia, Y., Bai, Q., 2018. Intestinal pathology and gut microbiota alterations in a Methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) mouse model of parkinson's disease. *Neurochem. Res.* 43 (10), 1986–1999.
- Larsson, H., Sariassan, A., Langstrom, N., D'Onofrio, B., Lichtenstein, P., 2014. Family income in early childhood and subsequent attention deficit/hyperactivity disorder: a quasi-experimental study. *J. Child Psychol. Psychiatry* 55 (5), 428–435.
- Leclercq, S., Matamoros, S., Cani, P.D., Neyrinck, A.M., Jamar, F., Starkel, P., Windy, K., Tremaroli, V., Backhed, F., Verbeke, K., de Timary, P., Delzenne, N.M., 2014. Intestinal permeability, gut-bacterial dysbiosis, and behavioral markers of alcohol-dependence severity. *Proc Natl Acad Sci U S A* 111 (42), E4485–4493.
- Lehouritis, P., Cummins, J., Stanton, M., Murphy, C.T., McCarthy, F.O., Reid, G., Urbaniak, C., Byrne, W.L., Tangney, M., 2015. Local bacteria affect the efficacy of chemotherapeutic drugs. *Sci. Rep.* 5, 14554.
- Levine, J.D., Goldstine, J., Mayes, M., Moskowitz, M.A., Basbaum, A.I., 1986. The neurotoxic effect of gold sodium thiomalate on the peripheral nerves of the rat. Insights into the antiinflammatory actions of gold therapy. *Arthritis Rheum.* 29 (7), 897–901.
- Li, C.Y., Dempsey, J.L., Wang, D., Lee, S., Weigel, K.M., Fei, Q., Bhatt, D.K., Prasad, B., Rafferty, D., Gu, H., Cui, J.Y., 2018. PBDEs altered gut microbiome and bile acid homeostasis in male C57BL/6 mice. *Drug Metab. Dispos.* 46 (8), 1226–1240.
- Li, C.Y., Lee, S., Cade, S., Kuo, L.J., Schultz, I.R., Bhatt, D.K., Prasad, B., Bammler, T.K., Cui, J.Y., 2017. Novel interactions between gut microbiome and host drug-processing genes modify the hepatic metabolism of the environmental chemicals polybrominated diphenyl ethers. *Drug Metab. Dispos.* 45 (11), 1197–1214.
- Li, T., Wang, W., Pan, Y.W., Xu, L., Xia, Z., 2013. A hydroxylated metabolite of flame-retardant PBDE-47 decreases the survival, proliferation, and neuronal differentiation of primary cultured adult neural stem cells and interferes with signaling of ERK5 MAP kinase and neurotrophin 3. *Toxicol. Sci.* 134 (1), 111–124.
- Lin, L., Zheng, L.J., Zhang, L.J., 2018. Neuroinflammation, gut microbiome, and Alzheimer's disease. *Mol. Neurobiol.* 55 (11), 8243–8250.
- Liu, F., Horton-Sparks, K., Hull, V., Li, R.W., Martinez-Cerdeno, V., 2018. The valproic acid rat model of autism presents with gut bacterial dysbiosis similar to that in human autism. *Mol. Autism* 9, 61.
- Liu, J., Liu, X., Xiong, X.Q., Yang, T., Cui, T., Hou, N.L., Lai, X., Liu, S., Guo, M., Liang, X.H., Cheng, Q., Chen, J., Li, T.Y., 2017a. Effect of vitamin A supplementation on gut microbiota in children with autism spectrum disorders - a pilot study. *BMC Microbiol.* 17 (1), 204.
- Liu, Q., Shao, W., Zhang, C., Xu, C., Wang, Q., Liu, H., Sun, H., Jiang, Z., Gu, A., 2017b. Organochlorine pesticides modulated gut microbiota and influenced bile acid metabolism in mice. *Environ. Pollut.* 226, 268–276.
- Liu, S., Li, E., Sun, Z., Fu, D., Duan, G., Jiang, M., Yu, Y., Mei, L., Yang, P., Tang, Y., Zheng, P., 2019. Altered gut microbiota and short chain fatty acids in Chinese children with autism spectrum disorder. *Sci. Rep.* 9 (1), 287.
- Lloyd, N.A., Janssen, S.E., Reinfelder, J.R., Barkay, T., 2016. Co-selection of mercury and multiple antibiotic resistances in Bacteria Exposed to mercury in the *Fundulus heteroclitus* gut microbiome. *Curr. Microbiol.* 73 (6), 834–842.
- Long, T.E., 2017. Repurposing thiram and disulfiram as antibacterial agents for multi-drug-resistant *Staphylococcus aureus* infections. *Antimicrob. Agents Chemother.* 61 (9).
- Lowe, P.P., Gyongyosi, B., Satishchandran, A., Iracheta-Vellve, A., Cho, Y., Ambade, A., Szabo, G., 2018. Reduced gut microbiome protects from alcohol-induced neuroinflammation and alters intestinal and brain inflammasome expression. *J. Neuroinflammation* 15 (1), 298.
- Lozano, V.L., Defarge, N., Rocque, L.M., Mesnage, R., Hennequin, D., Cassier, R., de Vendomois, J.S., Panoff, J.M., Seralini, G.E., Amiel, C., 2018. Sex-dependent impact of Roundup on the rat gut microbiome. *Toxicol. Rep.* 5, 96–107.
- Lu, J., Synowiec, S., Lu, L., Yu, Y., Bretherick, T., Takada, S., Yarnykh, V., Caplan, J., Caplan, M., Claud, E.C., Drobyshesky, A., 2018. Microbiota influence the development of the brain and behaviors in C57BL/6J mice. *PLoS One* 13 (8), e021829.
- Lu, K., Abo, R.P., Schlieper, K.A., Graffam, M.E., Levine, S., Wishnok, J.S., Swenberg, J.A., Tannenbaum, S.R., Fox, J.G., 2014. Arsenic exposure perturbs the gut microbiome and its metabolic profile in mice: an integrated metagenomics and metabolomics analysis. *Environ. Health Perspect.* 122 (3), 284–291.
- Lu, K., Mahubb, R., Fox, J.G., 2015. Xenobiotics: interaction with the intestinal microflora. *ILAR J.* 56 (2), 218–227.
- Lukowicz, C., Eller-Simatos, S., Regnier, M., Polizzi, A., Lasserre, F., Montagner, A., Lippi, Y., Jamin, E.L., Martin, J.F., Naylies, C., Canlet, C., Debrauwer, L., Bertrand-Michel, J., Al Saati, T., Theodorou, V., Loiseau, N., Mselli-Lakhal, L., Guillou, H., Gamet-Payrastré, L., 2018. Metabolic effects of a chronic dietary exposure to a low-dose pesticide cocktail in mice: sexual dimorphism and role of the constitutive androstane receptor. *Environ. Health Perspect.* 126 (6), 067007.
- Lurie, I., Yang, Y.X., Haynes, K., Mamtani, R., Boursi, B., 2015. Antibiotic exposure and the risk for depression, anxiety, or psychosis: a nested case-control study. *J. Clin. Psychiatry* 76 (11), 1522–1528.
- Lyng, G.D., Seegal, R.F., 2008. Polychlorinated biphenyl-induced oxidative stress in organotypic co-cultures: experimental dopamine depletion prevents reductions in GABA. *Neurotoxicology* 29 (2), 301–308.
- Lyng, G.D., Snyder-Keller, A., Seegal, R.F., 2007. Polychlorinated biphenyl-induced neurotoxicity in organotypic cocultures of developing rat ventral mesencephalon and striatum. *Toxicol. Sci.* 97 (1), 128–139.
- Mack, I., Cuntz, U., Gramer, C., Niedermaier, S., Pohl, C., Schwierz, A., Zimmermann, K., Zipfel, S., Enck, P., Penders, J., 2016. Weight gain in anorexia nervosa does not ameliorate the faecal microbiota, branched chain fatty acid profiles, and gastrointestinal complaints. *Sci. Rep.* 6, 26752.
- Mackow, E.R., Chang, F.N., 1985. Processing of precursor ribosomal RNA and the presence of a modified ribosome assembly scheme in *Escherichia coli* relaxed strain. *FEBS Lett.* 182 (2), 407–412.
- Magnusdottir, S., Heinken, A., Kutt, L., Ravcheev, D.A., Bauer, E., Noronha, A., Greenhalgh, K., Jager, C., Baginska, J., Wilmes, P., Fleming, R.M., Thiele, I., 2017. Generation of genome-scale metabolic reconstructions for 773 members of the human gut microbiota. *Nat. Biotechnol.* 35 (1), 81–89.
- Maisano, M., Cappello, T., Oliva, S., Natalotto, A., Giannetto, A., Parrino, V., Battaglia, P., Romeo, T., Salvo, A., Spano, N., Maueri, A., 2016. PCB and OCP accumulation and evidence of hepatic alteration in the Atlantic bluefin tuna, *T. thynnus*, from the Mediterranean Sea. *Mar. Environ. Res.* 121, 40–48.
- Manor, O., Borenstein, E., 2017. Systematic characterization and analysis of the taxonomic drivers of functional shifts in the human microbiome. *Cell Host Microbe* 21 (2), 254–267.
- Mao, Q., Manservigi, F., Panzacchi, S., Mandrioli, D., Menghetti, I., Vornoli, A., Bua, L., Falcioni, L., Lesseur, C., Chen, J., Belpoggi, F., Hu, J., 2018. The Ramazzini Institute 13-week pilot study on glyphosate and Roundup administered at human-equivalent dose to Sprague Dawley rats: effects on the microbiome. *Environ. Health* 17 (1), 50.
- Matt, S.M., Allen, J.M., Lawson, M.A., Mailing, L.J., Woods, J.A., Johnson, R.W., 2018. Butyrate and dietary soluble Fiber improve neuroinflammation associated with aging in mice. *Front. Immunol.* 9, 1832.
- Matuskova, Z., Anzenbacher, P., Vecera, R., Siller, M., Tlaskalova-Hogenova, H., Strojil, J., Anzenbacherova, E., 2017. Effect of *Lactobacillus casei* on the pharmacokinetics of amidarone in male wistar rats. *Eur. J. Drug Metab. Pharmacokinet.* 42 (1), 29–36.
- Matuskova, Z., Anzenbacherova, E., Vecera, R., Tlaskalova-Hogenova, H., Kolar, M., Anzenbacher, P., 2014. Administration of a probiotic can change drug pharmacokinetics: effect of *E. Coli* Nissle 1917 on amidarone absorption in rats. *PLoS One* 9 (2), e87150.
- McKay, K.A., Kowalec, K., Brinkman, F., Finlay, B.B., Horwitz, M., Manges, A.R., Osborne, L., Tremlett, H., 2017. From bugs to brains: the microbiome in neurological health. *Mult. Scler. Relat. Disord.* 12, 1–3.
- McMillin, M., DeMorrow, S., 2016. Effects of bile acids on neurological function and disease. *FASEB J.* 30 (11), 3658–3668.
- McMillin, M., Frampton, G., Quinn, M., Ashfaq, S., de los Santos 3rd, M., Grant, S., DeMorrow, S., 2016. Bile acid signaling is involved in the neurological decline in a murine model of acute liver failure. *Am. J. Pathol.* 186 (2), 312–323.
- Miller, C.A., Golden, N.H., 2010. An introduction to eating disorders: clinical

- presentation, epidemiology, and prognosis. *Nutr. Clin. Pract.* 25 (2), 110–115.
- Mills, B.D., Pearce, H.L., Khan, O., Jarrett, B.R., Fair, D.A., Lahvis, G.P., 2016. Prenatal domoic acid exposure disrupts mouse pro-social behavior and functional connectivity MRI. *Behav. Brain Res.* 308, 14–23.
- Mills, R.H., Vazquez-Baeza, Y., Zhu, Q., Jiang, L., Gaffney, J., Humphrey, G., Smarr, L., Knight, R., Gonzalez, D.J., 2019. Evaluating metagenomic prediction of the metaproteome in a 4.5-Year study of a patient with Crohn's disease. *mSystems* 4 (1).
- Mimee, M., Tucker, A.C., Voigt, C.A., Lu, T.K., 2015. Programming a human commensal bacterium, *Bacteroides thetaiotaomicron*, to sense and respond to stimuli in the murine gut microbiota. *Cell Syst.* 1 (1), 62–71.
- Minter, M.R., Hinterleitner, R., Meisel, M., Zhang, C., Leone, V., Zhang, X., Oyler-Castrillo, P., Zhang, X., Musch, M.W., Shen, X., Jabri, B., Chang, E.B., Tanzi, R.E., Sisodia, S.S., 2017. Antibiotic-induced perturbations in microbial diversity during post-natal development alters amyloid pathology in an aged APPSWE/PS1DeltaE9 murine model of Alzheimer's disease. *Sci. Rep.* 7 (1), 10411.
- Minter, M.R., Zhang, C., Leone, V., Ringus, D.L., Zhang, X., Oyler-Castrillo, P., Musch, M.W., Liao, F., Ward, J.F., Holtzman, D.M., Chang, E.B., Tanzi, R.E., Sisodia, S.S., 2016. Antibiotic-induced perturbations in gut microbial diversity influences neuroinflammation and amyloidosis in a murine model of Alzheimer's disease. *Sci. Rep.* 6, 30028.
- Miyaoka, T., Kanayama, M., Wake, R., Hashioka, S., Hayashida, M., Nagahama, M., Okazaki, S., Yamashita, S., Miura, S., Miki, H., Matsuda, H., Koike, M., Izuhara, M., Araki, T., Tsuchie, K., Azis, I.A., Arauchi, R., Abdullah, R.A., Oh-Nishi, A., Horiguchi, J., 2018. Clostridium butyricum MIYAIRI 588 as adjunctive therapy for treatment-resistant major depressive disorder: a prospective open-label trial. *Clin. Neuropharmacol.* 41 (5), 151–155.
- Moos, W.H., Fallor, D.V., Harpp, D.N., Kanara, I., Pernokas, J., Powers, W.R., Steliou, K., 2016. Microbiota and neurological disorders: a gut feeling. *Biores. Open Access* 5 (1), 137–145.
- Morita, C., Tsuji, H., Hata, T., Gondo, M., Takakura, S., Kawai, K., Yoshihara, K., Ogata, K., Nomoto, K., Miyazaki, K., Sudo, N., 2015. Gut dysbiosis in patients with anorexia nervosa. *PLoS One* 10 (12), e0145274.
- Morkl, S., Lackner, S., Muller, W., Gorkiewicz, G., Kashofer, K., Oberascher, A., Painold, A., Holl, A., Holzer, P., Meinitzer, A., Mangge, H., Holasek, S., 2017. Gut microbiota and body composition in anorexia nervosa inpatients in comparison to athletes, overweight, obese, and normal weight controls. *Int. J. Eat. Disord.* 50 (12), 1421–1431.
- Motta, E.V.S., Raymann, K., Moran, N.A., 2018. Glyphosate perturbs the gut microbiota of honey bees. *Proc Natl Acad Sci U S A* 115 (41), 10305–10310.
- Moussavi, S., Chatterji, S., Verdes, E., Tandon, A., Patel, V., Ustun, B., 2007. Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *Lancet* 370 (9590), 851–858.
- Mrema, E.J., Rubino, F.M., Brambilla, G., Moretto, A., Tsatsakis, A.M., Colosio, C., 2013. Persistent organochlorinated pesticides and mechanisms of their toxicity. *Toxicology* 307, 74–88.
- Murphy, J.R., Paul, S., Dunlop, A.L., Corwin, E.J., 2018. Maternal peripartum antibiotic exposure and the risk of postpartum depression. *Res. Nurs. Health.*
- Mutlu, E.A., Comba, I.Y., Cho, T., Engen, P.A., Yazici, C., Soberanes, S., Hamanaka, R.B., Nigdelioglu, R., Meliton, A.Y., Ghio, A.J., Budinger, G.R.S., Mutlu, G.M., 2018. Inhalational exposure to particulate matter air pollution alters the composition of the gut microbiome. *Environ Pollut* 240, 817–830.
- Nagamine, T., Ido, Y., Nakamura, M., Okamura, T., 2018. 4(G)-beta-D-galactosylsucrose as a prebiotic may improve underweight in inpatients with schizophrenia. *Biosci. Microbiota Food Health* 37 (2), 45–47.
- Nakagawa, R., Iida, T., Takenaka, S., Fukamachi, K., Mori, A., Takahashi, K., Asahi, M., 1987. PCBs in human blood and adipose tissue: levels and similarities on gas chromatograms. *Fukuoka Igaku Zasshi* 78 (5), 309–313.
- Nalls, M.A., Pankratz, N., Lill, C.M., Do, C.B., Hernandez, D.G., Saad, M., et al., 2014. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat. Genet.* 46 (9), 989–993.
- Naseribafrouei, A., Hestad, K., Avershina, E., Sekelja, M., Linlokken, A., Wilson, R., Rudi, K., 2014. Correlation between the human fecal microbiota and depression. *Neurogastroenterol. Motil.* 26 (8), 1155–1162.
- Nasuti, C., Coman, M.M., Olek, R.A., Fiorini, D., Verdenelli, M.C., Cecchini, C., Silvi, S., Fedeli, D., Gabbianelli, R., 2016. Changes on fecal microbiota in rats exposed to permethrin during postnatal development. *Environ. Sci. Pollut. Res. Int.* 23 (11), 10930–10937.
- Neufeld, K.M., Kang, N., Bienenstock, J., Foster, J.A., 2011. Reduced anxiety-like behavior and central neurochemical change in germ-free mice. *Neurogastroenterol. Motil.* 23 (3), 255–264 e119.
- Newsome, W.H., Davies, D.J., Sun, W.F., 1998. Residues of polychlorinated biphenyls (PCB) in fatty foods of the Canadian diet. *Food Addit. Contam.* 15 (1), 19–29.
- Nguyen, T.T., Kosciolk, T., Maldonado, Y., Daly, R.E., Martin, A.S., McDonald, D., Knight, R., Jeste, D.V., 2019. Differences in gut microbiome composition between persons with chronic schizophrenia and healthy comparison subjects. *Schizophr. Res.* 204, 23–29.
- Nielsen, L.N., Roager, H.M., Casas, M.E., Frandsen, H.L., Gosewinkle, U., Bester, K., Licht, T.R., Hendriksen, N.B., Bahl, M.L., 2018. Glyphosate has limited short-term effects on commensal bacterial community composition in the gut environment due to sufficient aromatic amino acid levels. *Environ Pollut* 233, 364–376.
- Oberdorster, G., Oberdorster, E., Oberdorster, J., 2005. Nanotoxicology: an emerging discipline evolving from studies of ultrafine particles. *Environ. Health Perspect.* 113 (7), 823–839.
- Oerther, D.B., Pernthaler, J., Schramm, A., Amann, R., Raskin, L., 2000. Monitoring precursor 16S rRNAs of *Acinetobacter* spp. in activated sludge wastewater treatment systems. *Appl. Environ. Microbiol.* 66 (5), 2154–2165.
- Okubo, R., Koga, M., Katsumata, N., Odamak, T., Matsuyama, S., Oka, M., Narita, H., Hashimoto, N., Kusumi, I., Xiao, J., Matsuoka, Y.J., 2019. Effect of bifidobacterium breve A-1 on anxiety and depressive symptoms in schizophrenia: a proof-of-concept study. *J. Affect. Disord.* 245, 377–385.
- Olde Loohuis, L.M., Mangul, S., Ori, A.P.S., Jospin, G., Koslicki, D., Yang, H.T., Wu, T., Boks, M.P., Lomen-Hoerth, C., Wiedau-Pazos, M., Cantor, R.M., de Vos, W.M., Kahn, R.S., Eskin, E., Ophoff, R.A., 2018. Transcriptome analysis in whole blood reveals increased microbial diversity in schizophrenia. *Transl. Psychiatry* 8 (1), 96.
- Olson, C.A., Vuong, H.E., Yano, J.M., Liang, Q.Y., Nusbaum, D.J., Hsiao, E.Y., 2018. The gut microbiota mediates the anti-seizure effects of the ketogenic diet. *Cell* 173 (7), 1728–1741 e1713.
- Osborne, L.M., Monk, C., 2013. Perinatal depression—the fourth inflammatory morbidity of pregnancy?: theory and literature review. *Psychoneuroendocrinology* 38 (10), 1929–1952.
- Painold, A., Morkl, S., Kashofer, K., Halwachs, B., Dalkner, N., Bengesser, S., Birner, A., Fellendorf, F., Platzer, M., Queissner, R., Schutze, G., Schwarz, M.J., Moll, N.M., Holzer, P., Holl, A.K., Kapfhammer, H.P., Gorkiewicz, G., Reininghaus, E.Z., 2018. A step ahead: exploring the gut microbiota in inpatients with bipolar disorder during a depressive episode. *Bipolar Disord.*
- Palacios, N., 2017. Air pollution and Parkinson's disease - evidence and future directions. *Rev. Environ. Health* 32 (4), 303–313.
- Pang, Y., Patterson, J.A., Applegate, T.J., 2009. The influence of copper concentration and source on ileal microbiota. *Poult. Sci.* 88 (3), 586–592.
- Parihar, V.K., Hattiangady, B., Shuai, B., Shetty, A.K., 2013. Mood and memory deficits in a model of Gulf War illness are linked with reduced neurogenesis, partial neuron loss, and mild inflammation in the hippocampus. *Neuropsychopharmacology* 38 (12), 2348–2362.
- Pass, D.A., Morgan, A.J., Read, D.S., Field, D., Weightman, A.J., Kille, P., 2015. The effect of anthropogenic arsenic contamination on the earthworm microbiome. *Environ. Microbiol.* 17 (6), 1884–1896.
- Pelissier, M.A., Vasquez, N., Balamurugan, R., Pereira, E., Dossou-Yovo, F., Suau, A., Pochart, P., Magne, F., 2010. Metronidazole effects on microbiota and mucus layer thickness in the rat gut. *FEMS Microbiol. Ecol.* 73 (3), 601–610.
- Perales-Puchalt, A., Perez-Sanz, J., Payne, K.K., Svoronos, N., Allegrezza, M.J., Chaurio, R.A., Anadon, C., Calmette, J., Biswas, S., Mine, J.A., Costich, T.L., Nickels, L., Wickramasinghe, J., Rutkowski, M.R., Conejo-Garcia, J.R., 2018. Frontline Science: microbiota reconstitution restores intestinal integrity after cisplatin therapy. *J. Leukoc. Biol.* 103 (5), 799–805.
- Pereze-Pardo, P., Dodiya, H.B., Engen, P.A., Naqib, A., Forsyth, C.B., Green, S.J., Garssen, J., Keshavarzian, A., Kraneveld, A.D., 2018. Gut bacterial composition in a mouse model of Parkinson's disease. *Benef. Microbes* 9 (5), 799–814.
- Perlstein, M., Attala, R., 1966. Neurologic sequelae of plumbism in children. *Clin. Pediatr. (Phila)* 5 (5), 292–298.
- Petriello, M.C., Hoffman, J.B., Vsevolozhskaya, O., Morris, A.J., Hennig, B., 2018. Dioxin-like PCB 126 increases intestinal inflammation and disrupts gut microbiota and metabolic homeostasis. *Environ Pollut* 242 (Pt A), 1022–1032.
- Pisa, D., Alonso, R., Rabano, A., Rodal, I., Carrasco, L., 2015. Different brain regions are infected with Fungi in Alzheimer's disease. *Sci. Rep.* 5, 15015.
- Prehn-Kristensen, A., Zimmermann, A., Tittmann, L., Lieb, W., Schreiber, S., Baving, L., Fischer, A., 2018. Reduced microbiome alpha diversity in young patients with ADHD. *PLoS One* 13 (7), e0200728.
- Przybyla, J., Houseman, E.A., Smit, E., Kile, M.L., 2017. A path analysis of multiple neurotoxic chemicals and cognitive functioning in older US adults (NHANES 1999–2002). *Environ. Health* 16 (1), 19.
- Pulikkan, J., Maji, A., Dhakan, D.B., Saxena, R., Mohan, B., Anto, M.M., Agarwal, N., Grace, T., Sharma, V.K., 2018. Gut microbial dysbiosis in Indian children with autism spectrum disorders. *Microb. Ecol.* 76 (4), 1102–1114.
- Pyndt Jorgensen, B., Krych, L., Pedersen, T.B., Plath, N., Redrobe, J.P., Hansen, A.K., Nielsen, D.S., Pedersen, C.S., Larsen, C., Sorensen, D.B., 2015. Investigating the long-term effect of subchronic phencyclidine-treatment on novel object recognition and the association between the gut microbiota and behavior in the animal model of schizophrenia. *Physiol. Behav.* 141, 32–39.
- Qiao, Y., Wu, M., Feng, Y., Zhou, Z., Chen, L., Chen, F., 2018. Alterations of oral microbiota distinguish children with autism spectrum disorders from healthy controls. *Sci. Rep.* 8 (1), 1597.
- Radisavljevic, N., Cirstea, M., Brett Finlay, B., 2018. Bottoms up: the role of gut microbiota in brain health. *Environ. Microbiol.*
- Rajzadeh, A., Mozaffari-Khosravi, H., Yassini-Ardakani, M., Dehghani, A., 2017. Effect of magnesium supplementation on depression status in depressed patients with magnesium deficiency: a randomized, double-blind, placebo-controlled trial. *Nutrition* 35, 56–60.
- Ramallo, R.M., Viana, R.J., Low, W.C., Steer, C.J., Rodrigues, C.M., 2008. Bile acids and apoptosis modulation: an emerging role in experimental Alzheimer's disease. *Trends Mol. Med.* 14 (2), 54–62.
- Raymann, K., Motta, E.V.S., Girard, C., Riddington, I.M., Dinsler, J.A., Moran, N.A., 2018. Imidacloprid decreases honey bee survival rates but does not affect the gut microbiome. *Appl. Environ. Microbiol.* 84 (13).
- Rea, K., Dinan, T.G., Cryan, J.F., 2016. The microbiome: a key regulator of stress and neuroinflammation. *Neurobiol. Stress* 4, 23–33.
- Reinoso Webb, C., Koboziev, I., Furr, K.L., Grisham, M.B., 2016. Protective and pro-inflammatory roles of intestinal bacteria. *Pathophysiology* 23 (2), 67–80.
- Ren, Y., Feng, J., 2007. Rotenone selectively kills serotonergic neurons through a microtubule-dependent mechanism. *J. Neurochem.* 103 (1), 303–311.
- Ren, Y., Liu, W., Jiang, H., Jiang, Q., Feng, J., 2005. Selective vulnerability of dopaminergic neurons to microtubule depolymerization. *J. Biol. Chem.* 280 (40), 34105–34112.
- Requile, M., Gonzalez Alvarez, D.O., Delanaud, S., Rhazi, L., Bach, V., Depeint, F., Khorsi-Cauet, H., 2018. Use of a combination of in vitro models to investigate the impact of chlorophylls and inulin on the intestinal microbiota and the permeability of the intestinal mucosa. *Environ. Sci. Pollut. Res. Int.* 25 (23), 22529–22540.
- Reygnier, J., Joly Condet, C., Bruneau, A., Delanaud, S., Rhazi, L., Depeint, F., Abdennebi-Najar, L., Bach, V., Mayeur, C., Khorsi-Cauet, H., 2016a. Changes in composition and function of human intestinal microbiota exposed to chlorophylls in oil as assessed by the SHIME(R) model. *Int. J. Environ. Res. Public Health* 13 (11).

- Reygner, J., Lichtenberger, L., Elmhiri, G., Dou, S., Bahi-Jaber, N., Rhazi, L., Depeint, F., Bach, V., Khorsi-Cauet, H., Abdennebi-Najar, L., 2016b. Inulin supplementation lowered the metabolic defects of prolonged exposure to chlorpyrifos from gestation to young adult stage in offspring rats. *PLoS One* 11 (10), e0164614.
- Ribas-Fito, N., Sala, M., Kogevinas, M., Sunyer, J., 2001. Polychlorinated biphenyls (PCBs) and neurological development in children: a systematic review. *J. Epidemiol. Community Health* 55 (8), 537–546.
- Richardson, J.B., Dancy, B.C.R., Horton, C.L., Lee, Y.S., Madejczyk, M.S., Xu, Z.Z., Ackermann, G., Humphrey, G., Palacios, G., Knight, R., Lewis, J.A., 2018. Exposure to toxic metals triggers unique responses from the rat gut microbiota. *Sci. Rep.* 8 (1), 6578.
- Rigby, R.J., Carr, J., Orgel, K., King, S.L., Lund, P.K., Dekaney, C.M., 2016. Intestinal bacteria are necessary for doxorubicin-induced intestinal damage but not for doxorubicin-induced apoptosis. *Gut Microbes* 7 (5), 414–423.
- Roelens, S.A., Beck, V., Aerts, G., Clerens, S., Vanden Bergh, G., Arckens, L., Darras, V.M., van der Geysen, S., 2005. Neurotoxicity of polychlorinated biphenyls (PCBs) by disturbance of thyroid hormone-regulated genes. *Ann. N. Y. Acad. Sci.* 1040, 454–456.
- Rose, D.R., Yang, H., Serena, G., Sturgeon, C., Ma, B., Careaga, M., Hughes, H.K., Angkustiri, K., Rose, M., Hertz-Picciotto, I., Van de Water, J., Hansen, R.L., Ravel, J., Fasano, A., Ashwood, P., 2018. Differential immune responses and microbiota profiles in children with autism spectrum disorders and co-morbid gastrointestinal symptoms. *Brain Behav. Immun.* 70, 354–368.
- Rothenberg, S.E., Keiser, S., Ajami, N.J., Wong, M.C., Gesell, J., Petrosino, J.F., Johs, A., 2016. The role of gut microbiota in fetal methylmercury exposure: insights from a pilot study. *Toxicol. Lett.* 242, 60–67.
- Ruan, Y., Wu, C., Guo, X., Xu, Z., Xing, C., Cao, H., Zhang, C., Hu, G., Liu, P., 2018. High doses of copper and mercury changed cecal microbiota in female mice. *Biol. Trace Elem. Res.*
- Ruark, C.D., Hack, C.E., Robinson, P.J., Anderson, P.E., Gearhart, J.M., 2013. Quantitative structure-activity relationships for organophosphates binding to acetylcholinesterase. *Arch. Toxicol.* 87 (2), 281–289.
- Rudzi, L., Ostrowska, L., Pawlak, D., Malus, A., Pawlak, K., Waszkiewicz, N., Szulc, A., 2019. Probiotic *Lactobacillus Plantarum* 299v decreases kynurenine concentration and improves cognitive functions in patients with major depression: a double-blind, randomized, placebo controlled study. *Psychoneuroendocrinology* 100, 213–222.
- Russo, R., Cristiano, C., Avagliano, C., De Caro, C., La Rana, G., Raso, G.M., Canani, R.B., Meli, R., Calignano, A., 2018. Gut-brain Axis: role of lipids in the regulation of inflammation, pain and CNS diseases. *Curr. Med. Chem.* 25 (32), 3930–3952.
- Sadaghian Sadabad, M., von Martels, J.Z., Khan, M.T., Blokzijl, T., Paglia, G., Dijkstra, G., Harmsen, H.J., Faber, K.N., 2015. A simple coculture system shows mutualism between anaerobic faecalibacteria and epithelial Caco-2 cells. *Sci. Rep.* 5, 17906.
- Saeedi Saravi, S.S., Dehpour, A.R., 2016. Potential role of organochlorine pesticides in the pathogenesis of neurodevelopmental, neurodegenerative, and neurobehavioral disorders: a review. *Life Sci.* 145, 255–264.
- Saha, S., Chant, D., McGrath, J., 2007. A systematic review of mortality in schizophrenia: is the differential mortality gap worsening over time? *Arch. Gen. Psychiatry* 64 (10), 1123–1131.
- Sahasrabudhe, N.M., Beukema, M., Tian, L., Troost, B., Scholte, J., Bruininx, E., Bruggeman, G., van den Berg, M., Scheurink, A., Schols, H.A., Faas, M.M., de Vos, P., 2018. Dietary Fiber pectin directly blocks toll-like receptor 2-1 and prevents doxorubicin-induced ileitis. *Front. Immunol.* 9, 383.
- Salihovic, S., Ganna, A., Fall, T., Broeckling, C.D., Prenni, J.E., van Bavel, B., Lind, P.M., Ingelsson, E., Lind, L., 2016. The metabolic fingerprint of p,p'-DDE and HCB exposure in humans. *Environ. Int.* 88, 60–66.
- Salzer, J.L., Zalc, B., 2016. Myelination. *Curr. Biol.* 26 (20), R971–R975.
- Sampson, T.R., Debelius, J.W., Thron, T., Janssen, S., Shastri, G.G., Ilhan, Z.E., Challis, C., Schretter, C.E., Rocha, S., Gradinaru, V., Chesselet, M.F., Keshavarzian, A., Shannon, K.M., Krajmalnik-Brown, R., Wittung-Stafshede, P., Knight, R., Mazmanian, S.K., 2016. Gut microbiota regulate motor deficits and neuroinflammation in a model of parkinson's disease. *Cell* 167 (6), 1469–1480 e1412.
- Scassellati, C., Bonvicini, C., Faraone, S.V., Gennarelli, M., 2012. Biomarkers and attention-deficit/hyperactivity disorder: a systematic review and meta-analysis. *J. Am. Acad. Child Adolesc. Psychiatry* 51 (10), 1003–1019 e1020.
- Schantz, S.L., 1996. Developmental neurotoxicity of PCBs in humans: what do we know and where do we go from here? *Neurotoxicol. Teratol.* 18 (3), 217–227 discussion 229–276.
- Schecter, A., Haffner, D., Colacino, J., Patel, K., Papke, O., Opel, M., Birnbaum, L., 2010. Polybrominated diphenyl ethers (PBDEs) and hexabromocyclodecane (HBCD) in composite U.S. Food samples. *Environ. Health Perspect.* 118 (3), 357–362.
- Schecter, A., Pavuk, M., Papke, O., Ryan, J.J., Birnbaum, L., Rosen, R., 2003. Polybrominated diphenyl ethers (PBDEs) in U.S. mothers' milk. *Environ. Health Perspect.* 111 (14), 1723–1729.
- Schicho, R., Krueger, D., Zeller, F., Von Weyhern, C.W., Frieling, T., Kimura, H., Ishii, I., De Giorgio, R., Campi, B., Schemann, M., 2006. Hydrogen sulfide is a novel prosecretory neuromodulator in the Guinea-pig and human colon. *Gastroenterology* 131 (5), 1542–1552.
- Schizophrenia Psychiatric Genome-Wide Association Study, C., 2011. Genome-wide association study identifies five new schizophrenia loci. *Nat. Genet.* 43 (10), 969–976.
- Selwyn, F.P., Cui, J.Y., Klaassen, C.D., 2015. RNA-seq quantification of hepatic drug processing genes in germ-free mice. *Drug Metab. Dispos.* 43 (10), 1572–1580.
- Sender, R., Fuchs, S., Milo, R., 2016. Revised estimates for the number of human and Bacteria cells in the body. *PLoS Biol.* 14 (8), e1002533.
- Serefko, A., Szopa, A., Poleszak, E., 2016. Magnesium and depression. *Magnes. Res.* 29 (3), 112–119.
- Seth, B., Yadav, A., Tandon, A., Shankar, J., Chaturvedi, R.K., 2019. Carbofuran hampers oligodendrocytes development leading to impaired myelination in the hippocampus of rat brain. *Neurotoxicology* 70, 161–179.
- Seth, R.K., Kimono, D., Alhasson, F., Sarkar, S., Albadrani, M., Lasley, S.K., Horner, R., Janulewicz, P., Nagarkatti, M., Nagarkatti, P., Sullivan, K., Chatterjee, S., 2018. Increased butyrate priming in the gut stalls microbiome associated-gastrointestinal inflammation and hepatic metabolic reprogramming in a mouse model of Gulf War Illness. *Toxicol. Appl. Pharmacol.* 350, 64–77.
- Severance, E.G., Prandovszky, E., Castiglione, J., Yolken, R.H., 2015. Gastroenterology issues in schizophrenia: why the gut matters. *Curr. Psychiatry Rep.* 17 (5), 27.
- Sharon, G., Sampson, T.R., Geschwind, D.H., Mazmanian, S.K., 2016. The central nervous system and the gut microbiome. *Cell* 167 (4), 915–932.
- Shehata, A.A., Schrodli, W., Aldin, A.A., Hafez, H.M., Kruger, M., 2013. The effect of glyphosate on potential pathogens and beneficial members of poultry microbiota in vitro. *Curr. Microbiol.* 66 (4), 350–358.
- Shen, Y., Xu, J., Li, Z., Huang, Y., Yuan, Y., Wang, J., Zhang, M., Hu, S., Liang, Y., 2018. Analysis of gut microbiota diversity and auxiliary diagnosis as a biomarker in patients with schizophrenia: a cross-sectional study. *Schizophr. Res.*
- Sheppard, J.G., Frazier, K.R., Saralkar, P., Hossain, M.F., Geldenhuys, W.J., Long, T.E., 2018. Disulfiram-based disulfides as narrow-spectrum antibacterial agents. *Bioorg. Med. Chem. Lett.* 28 (8), 1298–1302.
- Shin, S., Burnett, R.T., Kwong, J.C., Hystad, P., van Donkelaar, A., Brook, J.R., Copes, R., Tu, K., Goldberg, M.S., Villeneuve, P.J., Martin, R.V., Murray, B.J., Wilton, A.S., Kopp, A., Chen, H., 2018. Effects of ambient air pollution on incident Parkinson's disease in Ontario, 2001 to 2013: a population-based cohort study. *Int. J. Epidemiol.* 47 (6), 2038–2048.
- Sipka, S., Eum, S.Y., Son, K.W., Xu, S., Gavalas, V.G., Hennig, B., Toborek, M., 2008. Oral Administration of PCBs INDUCES PROINFLAMMATORY and PROMETASTATIC RESPONSES. *Environ. Toxicol. Pharmacol.* 25 (2), 251–259.
- Sjodin, A., Wong, L.Y., Jones, R.S., Park, A., Zhang, Y., Hodge, C., Dipietro, E., McClure, C., Turner, W., Needham, L.L., Patterson Jr., D.G., 2008. Serum concentrations of polybrominated diphenyl ethers (PBDEs) and polybrominated biphenyl (PBB) in the United States population: 2003–2004. *Environ. Sci. Technol.* 42 (4), 1377–1384.
- Skonieczna-Zydecka, K., Marlicz, W., Misera, A., Koulaouzidis, A., Loniewski, I., 2018. Microbiome-the missing link in the gut-brain Axis: focus on its role in gastrointestinal and mental health. *J. Clin. Med.* 7 (12).
- Slotkin, T.A., Ryde, I.T., Levin, E.D., Seidler, F.J., 2008. Developmental neurotoxicity of low dose diazinon exposure of neonatal rats: effects on serotonin systems in adolescence and adulthood. *Brain Res. Bull.* 75 (5), 640–647.
- Slykerman, R.F., Hood, F., Wickens, K., Thompson, J.M.D., Barthow, C., Murphy, R., Kang, J., Rowden, J., Stone, P., Crane, J., Stanley, T., Abels, P., Purdie, G., Maude, R., Mitchell, E.A., Probiotic in Pregnancy Study, G., 2017. Effect of *Lactobacillus rhamnosus* HN001 in pregnancy on postpartum symptoms of depression and anxiety: a randomised double-blind placebo-controlled trial. *EBioMedicine* 24, 159–165.
- Smith, M.E., Swoboda, H.D., 2019. Shellfish Toxicity. *StatPearls*. Treasure Island, FL.
- Sobko, T., Huang, L., Midtvedt, T., Norin, E., Gustafsson, L.E., Norman, M., Jansson, E.A., Lundberg, J.O., 2006. Generation of NO by probiotic bacteria in the gastrointestinal tract. *Free Radic. Biol. Med.* 41 (6), 985–991.
- Son, J.S., Zheng, L.J., Rowehl, L.M., Tian, X., Zhang, Y., Zhu, W., Litcher-Kelly, L., Gadaw, K.D., Gathungu, G., Robertson, C.E., Ir, D., Frank, D.N., Li, E., 2015. Comparison of fecal microbiota in children with autism Spectrum disorders and neurotypical siblings in the simons simplex collection. *PLoS One* 10 (10), e0137725.
- Song, Y., Zhao, R., Hu, Y., Hao, F., Li, N., Nie, G., Tang, H., Wang, Y., 2015. Assessment of the biological effects of a multifunctional nano-drug-carrier and its encapsulated drugs. *J. Proteome Res.* 14 (12), 5193–5201.
- Spanogiannopoulos, P., Bess, E.N., Carmody, R.N., Turnbaugh, P.J., 2016. The microbial pharmacists within us: a metagenomic view of xenobiotic metabolism. *Nat. Rev. Microbiol.* 14 (5), 273–287.
- Spencer, P., Schaumburg, H., 2000. *Experimental and Clinical Neurotoxicology*, 2nd ed. Oxford University Press, New York.
- Srivastava, A.K., Schlessinger, D., 1990. Mechanism and regulation of bacterial ribosomal RNA processing. *Annu. Rev. Microbiol.* 44, 105–129.
- Stanaway, I.B., Wallace, J.C., Shojia, A., Griffith, W.C., Hong, S., Wilder, C.S., Green, F.H., Tsai, J., Knight, M., Workman, T., Vigoren, E.M., McLean, J.S., Thompson, B., Faustman, E.M., 2017. Human oral buccal microbiomes are associated with farm-worker status and azinphos-methyl agricultural pesticide exposure. *Appl. Environ. Microbiol.* 83 (2).
- Stevens, S.E., Sonuga-Barke, E.J., Kreppner, J.M., Beckett, C., Castle, J., Colvert, E., Groothues, C., Hawkins, A., Rutter, M., 2008. Inattention/overactivity following early severe institutional deprivation: presentation and associations in early adolescence. *J. Abnorm. Child Psychol.* 36 (3), 385–398.
- Stewardson, A.J., Gaia, N., Francois, P., Malhotra-Kumar, S., Delemont, C., Martinez de Tejada, B., Schrenzel, J., Harbarth, S., Lazarevic, V., Saturn, W.P., Groups, W.P.S., 2015. Collateral damage from oral ciprofloxacin versus nitrofurantoin in outpatients with urinary tract infections: a culture-free analysis of gut microbiota. *Clin. Microbiol. Infect.* 21 (4), 344 e341–311.
- Stewart, J.E., Marks, L.J., Gilgan, M.W., Pfeiffer, E., Zwicker, B.M., 1998. Microbial utilization of the neurotoxin domoic acid: blue mussels (*Mytilus edulis*) and soft shell clams (*Mya arenaria*) as sources of the microorganisms. *Can. J. Microbiol.* 44 (5), 456–464.
- Strandwitz, P., Kim, K.H., Terekhova, D., Liu, J.K., Sharma, A., Levering, J., McDonald, D., Dietrich, D., Ramadhar, T.R., Lekbua, A., Mroue, N., Liston, C., Stewart, E.J., Dubin, M.J., Zengler, K., Knight, R., Gilbert, J.A., Clardy, J., Lewis, K., 2019. GABA-modulating bacteria of the human gut microbiota. *Nat. Microbiol.* 4 (3), 396–403.
- Sudo, N., Chida, Y., Aiba, Y., Sonoda, J., Oyama, N., Yu, X.N., Kubo, C., Koga, Y., 2004. Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. *J. Physiol. (Paris)* 558 (Pt 1), 263–275.
- Sun, J., Wang, F., Hu, X., Yang, C., Xu, H., Yao, Y., Liu, J., 2018a. Clostridium butyricum attenuates chronic unpredictable mild stress-induced depressive-like behavior in mice via the gut-brain Axis. *J. Agric. Food Chem.* 66 (31), 8415–8421.
- Sun, M.F., Zhu, Y.L., Zhou, Z.L., Jia, X.B., Xu, Y.D., Yang, Q., Cui, C., Shen, Y.Q., 2018b. Neuroprotective effects of fecal microbiota transplantation on MPTP-induced Parkinson's disease mice: gut microbiota, glial reaction and TLR4/TNF-alpha signaling pathway. *Brain Behav. Immun.* 70, 48–60.
- Suzuki, G., Nakano, M., Nakano, S., 2005. Distribution of PCDDs/PCDFs and Co-PCBs in

- human maternal blood, cord blood, placenta, milk, and adipose tissue: dioxins showing high toxic equivalency factor accumulate in the placenta. *Biosci. Biotechnol. Biochem.* 69 (10), 1836–1847.
- Swanson, H.L., 2015. Drug Metabolism by the Host and Gut Microbiota: A Partnership or Rivalry? *Drug Metab. Dispos.* 43 (10), 1499–1504.
- Tabouy, L., Getselter, D., Ziv, O., Karpuj, M., Tabouy, T., Lukic, I., Maayouf, R., Werbner, N., Ben-Amram, H., Nuriel-Ohayon, M., Koren, O., Elliott, E., 2018. Dysbiosis of microbiome and probiotic treatment in a genetic model of autism spectrum disorders. *Brain Behav. Immun.* 73, 310–319.
- Tang, X.Q., Ren, Y.K., Chen, R.Q., Zhuang, Y.Y., Fang, H.R., Xu, J.H., Wang, C.Y., Hu, B., 2011. Formaldehyde induces neurotoxicity to PC12 cells involving inhibition of paraoxonase-1 expression and activity. *Clin. Exp. Pharmacol. Physiol.* 38 (4), 208–214.
- Tillmann, S., Abildgaard, A., Winther, G., Wegener, G., 2018. Altered fecal microbiota composition in the Flinders sensitive line rat model of depression. *Psychopharmacology (Berl.)*.
- Tilson, H.A., Kodavanti, P.R., 1998. The neurotoxicity of polychlorinated biphenyls. *Neurotoxicology* 19 (4–5), 517–525.
- Timofeeva, O.A., Roegge, C.S., Seidler, F.J., Slotkin, T.A., Levin, E.D., 2008. Persistent cognitive alterations in rats after early postnatal exposure to low doses of the organophosphate pesticide, diazinon. *Neurotoxicol. Teratol.* 30 (1), 38–45.
- Toda, T., Ohi, K., Kudo, T., Yoshida, T., Ikarashi, N., Ito, K., Sugiyama, K., 2009. Ciprofloxacin suppresses Cyp3a in mouse liver by reducing lithocholic acid-producing intestinal flora. *Drug Metab. Pharmacokinet.* 24 (3), 201–208.
- Torres, E.R.S., Akinyeke, T., Stagaman, K., Duvoisin, R.M., Meshul, C.K., Sharpton, T.J., Raber, J., 2018. Effects of sub-chronic MPTP exposure on behavioral and cognitive performance and the microbiome of wild-type and mGlu8 knockout female and male mice. *Front. Behav. Neurosci.* 12, 140.
- Tremlett, H., Bauer, K.C., Appel-Cresswell, S., Finlay, B.B., Waubant, E., 2017. The gut microbiome in human neurological disease: a review. *Ann. Neurol.* 81 (3), 369–382.
- Trinder, M., McDowell, T.W., Daisley, B.A., Ali, S.N., Leong, H.S., Sumarah, M.W., Reid, G., 2016. Probiotic *Lactobacillus rhamnosus* reduces organophosphate pesticide absorption and toxicity to *Drosophila melanogaster*. *Appl. Environ. Microbiol.* 82 (20), 6204–6213.
- Ufarte, L., Laville, E., Duquesne, S., Morgavi, D., Robe, P., Klopp, C., Rizzo, A., Pizzut-Serin, S., Potocki-Veronese, G., 2017. Discovery of carbamate degrading enzymes by functional metagenomics. *PLoS One* 12 (12), e0189201.
- Unger, M.M., Spiegel, J., Dillmann, K.U., Grundmann, D., Philippeit, H., Burmann, J., Fassbender, K., Schwierz, A., Schafer, K.H., 2016. Short chain fatty acids and gut microbiota differ between patients with Parkinson's disease and age-matched controls. *Parkinsonism Relat. Disord.* 32, 66–72.
- Valles, Y., Francino, M.P., 2018. Air pollution, early life microbiome, and development. *Curr. Environ. Health Rep.* 5 (4), 512–521.
- Van den Bossche, L., Hindryckx, P., Devisscher, L., Devriese, S., Van Welden, S., Holvoet, T., Vilchez-Vargas, R., Vital, M., Pieper, D.H., Vandev Bussche, J., Vanhaecke, L., Van de Wiele, T., De Vos, M., Laukens, D., 2017. Ursodeoxycholic acid and its taurine- or glycine-conjugated species reduce colitogenic dysbiosis and equally suppress experimental colitis in mice. *Appl. Environ. Microbiol.* 83 (7).
- van Os, J., Kenis, G., Rutten, B.P., 2010. The environment and schizophrenia. *Nature* 468 (7321), 203–212.
- Velmurugan, G., Ramprasath, T., Swaminathan, K., Mithieux, G., Rajendhran, J., Dhivakar, M., Parthasarathy, A., Babu, D.D., Thumbaraj, L.J., Freddy, A.J., Dinakaran, V., Puhari, S.S., Rekha, B., Christy, Y.J., Anusha, S., Divya, G., Suganya, K., Meganathan, B., Kalyanaram, N., Vasudevan, V., Kamaraj, R., Karthik, M., Jeyakumar, B., Abhishek, A., Paul, E., Pushpanathan, M., Rajmohan, R.K., Velayutham, K., Lyon, A.R., Ramasamy, S., 2017. Gut microbial degradation of organophosphate insecticides induces glucose intolerance via gluconeogenesis. *Genome Biol.* 18 (1), 8.
- Vervooft, J., Xavier, B.B., Stewardson, A., Coenen, S., Godycki-Cwirko, M., Adriaenssens, N., Kowalczyk, A., Lammens, C., Harbarth, S., Goossens, H., Malhotra-Kumar, S., 2015. Metagenomic analysis of the impact of nitrofurantoin treatment on the human faecal microbiota. *J. Antimicrob. Chemother.* 70 (7), 1989–1992.
- Vinberg, M., Ottesen, N.M., Meluken, I., Sorensen, N., Pedersen, O., Kessing, L.V., Miskowiak, K.W., 2019. Remitted affective disorders and high familial risk of affective disorders associate with aberrant intestinal microbiota. *Acta Psychiatr. Scand.* 139 (2), 174–184.
- Vinolo, M.A., Rodrigues, H.G., Nachbar, R.T., Curi, R., 2011. Regulation of inflammation by short chain fatty acids. *Nutrients* 3 (10), 858–876.
- Visschers, R.G., Luyer, M.D., Schaap, F.G., Olde Damink, S.W., Soeters, P.B., 2013. The gut-liver axis. *Curr. Opin. Clin. Nutr. Metab. Care* 16 (5), 576–581.
- Vogt, N.M., Kerby, R.L., Dill-McFarland, K.A., Harding, S.J., Merluzzi, A.P., Johnson, S.C., Carlsson, C.M., Asthana, S., Zetterberg, H., Blennow, K., Bendlin, B.B., Rey, F.E., 2017. Gut microbiome alterations in Alzheimer's disease. *Sci. Rep.* 7 (1), 13537.
- Vuong, A.M., Yoltou, K., Dietrich, K.N., Braun, J.M., Lanphear, B.P., Chen, A., 2018. Exposure to polybrominated diphenyl ethers (PBDEs) and child behavior: current findings and future directions. *Horm. Behav.* 101, 94–104.
- Waclawikova, B., El Aidy, S., 2018. Role of microbiota and tryptophan metabolites in the remote effect of intestinal inflammation on brain and depression. *Pharmaceuticals Basel (Basel)* 11 (3).
- Wang, G., Liu, Q., Guo, L., Zeng, H., Ding, C., Zhang, W., Xu, D., Wang, X., Qiu, J., Dong, Q., Fan, Z., Zhang, Q., Pan, J., 2018. Gut microbiota and relevant metabolites analysis in alcohol dependent mice. *Front. Microbiol.* 9, 1874.
- Wang, M., Zhou, J., He, F., Cai, C., Wang, H., Wang, Y., Lin, Y., Rong, H., Cheng, G., Xu, R., Zhou, W., 2019a. Alteration of gut microbiota-associated epitopes in children with autism spectrum disorders. *Brain Behav. Immun.* 75, 192–199.
- Wang, N., Kong, D., Cai, D., Shi, L., Cao, Y., Pang, G., Yu, R., 2010. Levels of polychlorinated biphenyls in human adipose tissue samples from southeast China. *Environ. Sci. Technol.* 44 (11), 4334–4340.
- Wang, X., Shen, M., Zhou, J., Jin, Y., 2019b. Chlorpyrifos disturbs hepatic metabolism associated with oxidative stress and gut microbiota dysbiosis in adult zebrafish. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 216, 19–28.
- Warner, B.B., 2019. The contribution of the gut microbiome to neurodevelopment and neuropsychiatric disorders. *Pediatr. Res.* 85 (2), 216–224.
- Winneke, G., 2011. Developmental aspects of environmental neurotoxicology: lessons from lead and polychlorinated biphenyls. *J. Neurol. Sci.* 308 (1–2), 9–15.
- Winther, G., Pyndt Jorgensen, B.M., Elfving, B., Nielsen, D.S., Kihl, P., Lund, S., Sorensen, D.B., Wegener, G., 2015. Dietary magnesium deficiency alters gut microbiota and leads to depressive-like behaviour. *Acta Neuropsychiatr.* 27 (3), 168–176.
- Wu, F., Yang, L., Islam, M.T., Jasmine, F., Kibriya, M.G., Nahar, J., Barmon, B., Parvez, F., Sarwar, G., Ahmed, A., Eunos, M., Islam, T., Slavkovich, V., Hu, J., Li, H., Graziano, J.H., Pei, Z., Ahsan, H., Chen, Y., 2019. The role of gut microbiome and its interaction with arsenic exposure in carotid intima-media thickness in a Bangladesh population. *Environ. Int.* 123, 104–113.
- Wu, J., Wen, X.W., Faulk, C., Boehnke, K., Zhang, H., Dolinoy, D.C., Xi, C., 2016. Perinatal lead exposure alters gut microbiota composition and results in sex-specific body-weight increases in adult mice. *Toxicol. Sci.* 151 (2), 324–333.
- Xia, J., Jin, C., Pan, Z., Sun, L., Fu, Z., Jin, Y., 2018a. Chronic exposure to low concentrations of lead induces metabolic disorder and dysbiosis of the gut microbiota in mice. *Sci. Total Environ.* 631–632, 439–448.
- Xia, J., Lu, L., Jin, C., Wang, S., Zhou, J., Ni, Y., Fu, Z., Jin, Y., 2018b. Effects of short term lead exposure on gut microbiota and hepatic metabolism in adult zebrafish. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 209, 1–8.
- Xia, X., Sun, B., Gurr, G.M., Vasseur, L., Xue, M., You, M., 2018c. Gut microbiota mediate insecticide resistance in the diamondback moth, *plutella xylostella* (L.). *Front. Microbiol.* 9, 25.
- Xia, X., Zheng, D., Zhong, H., Qin, B., Gurr, G.M., Vasseur, L., Lin, H., Bai, J., He, W., You, M., 2013. DNA sequencing reveals the midgut microbiota of diamondback moth, *Plutella xylostella* (L.) and a possible relationship with insecticide resistance. *PLoS One* 8 (7), e68852.
- Xu, Z., Wang, C., Dong, X., Hu, T., Wang, L., Zhao, W., Zhu, S., Li, G., Hu, Y., Gao, Q., Wan, J., Liu, Z., Sun, J., 2018. Chronic alcohol exposure induced gut microbiota dysbiosis and its correlations with neuropsychic behaviors and brain BDNF/Gabra1 changes in mice. *Biofactors*.
- Yan, A., Culp, E., Perry, J., Lau, J.T., MacNeil, L.T., Surette, M.G., Wright, G.D., 2018. Transformation of the anticancer drug doxorubicin in the human gut microbiome. *ACS Infect. Dis.* 4 (1), 68–76.
- Yang, D., Kim, K.H., Phimister, A., Bachstetter, A.D., Ward, T.R., Stackman, R.W., Mervis, R.F., Wisniewski, A.B., Klein, S.L., Kodavanti, P.R., Anderson, K.A., Wayman, G., Pessah, I.N., Lein, P.J., 2009. Developmental exposure to polychlorinated biphenyls interferes with experience-dependent dendritic plasticity and ryanodine receptor expression in weanling rats. *Environ. Health Perspect.* 117 (3), 426–435.
- Yang, X., Qian, Y., Xu, S., Song, Y., Xiao, Q., 2017. Longitudinal analysis of fecal microbiome and pathological processes in a rotenone induced mice model of parkinson's disease. *Front. Aging Neurosci.* 9, 441.
- Yilmaz, B., Schibli, S., Macpherson, A.J., Sokollik, C., 2018. D-lactic acidosis: successful suppression of D-lactate-producing *Lactobacillus* by probiotics. *Pediatrics* 142 (3).
- Yin, N., Cai, X., Du, H., Zhang, Z., Li, Z., Chen, X., Sun, G., Cui, Y., 2017. In vitro study of soil arsenic release by human gut microbiota and its intestinal absorption by Caco-2 cells. *Chemosphere* 168, 358–364.
- Yin, N., Zhang, Z., Cai, X., Du, H., Sun, G., Cui, Y., 2015. In vitro method to assess soil arsenic metabolism by human gut microbiota: arsenic speciation and distribution. *Environ. Sci. Technol.* 49 (17), 10675–10681.
- Yoo, B.B., Mazmanian, S.K., 2017. The enteric network: interactions between the immune and nervous systems of the gut. *Immunity* 46 (6), 910–926.
- Yu, H., Wu, B., Zhang, X.X., Liu, S., Yu, J., Cheng, S., Ren, H.Q., Ye, L., 2016. Arsenic metabolism and toxicity influenced by ferric iron in simulated gastrointestinal tract and the roles of gut microbiota. *Environ. Sci. Technol.* 50 (13), 7189–7197.
- Yu, M., Jia, H., Zhou, C., Yang, Y., Zhao, Y., Yang, M., Zou, Z., 2017. Variations in gut microbiota and fecal metabolic phenotype associated with depression by 16S rRNA gene sequencing and LC/MS-based metabolomics. *J. Pharm. Biomed. Anal.* 138, 231–239.
- Zhang, L., Wang, Y., Xiayu, X., Shi, C., Chen, W., Song, N., Fu, X., Zhou, R., Xu, Y.F., Huang, L., Zhu, H., Han, Y., Qin, C., 2017. Altered gut microbiota in a mouse model of alzheimer's disease. *J. Alzheimers Dis.* 60 (4), 1241–1257.
- Zhao, Y., Zhang, Y., Wang, G., Han, R., Xie, X., 2016. Effects of chlorpyrifos on the gut microbiome and urine metabolome in mouse (*Mus musculus*). *Chemosphere* 153, 287–293.
- Zheng, P., Zeng, B., Liu, M., Chen, J., Pan, J., Han, Y., Liu, Y., Cheng, K., Zhou, C., Wang, H., Zhou, X., Gui, S., Perry, S.W., Wong, M.L., Licinio, J., Wei, H., Xie, P., 2019. The gut microbiome from patients with schizophrenia modulates the glutamate-glutamine-GABA cycle and schizophrenia-relevant behaviors in mice. *Sci. Adv.* 5 (2), eaau8317.
- Zheng, P., Zeng, B., Zhou, C., Liu, M., Fang, Z., Xu, X., Zeng, L., Chen, J., Fan, S., Du, X., Zhang, X., Yang, D., Yang, Y., Meng, H., Li, W., Melgiri, N.D., Licinio, J., Wei, H., Xie, P., 2016. Gut microbiome remodeling induces depressive-like behaviors through a pathway mediated by the host's metabolism. *Mol. Psychiatry* 21 (6), 786–796.
- Zhu, L., Yang, Z., Yao, R., Xu, L., Chen, H., Gu, X., Wu, T., Yang, X., 2018a. Potential mechanism of detoxification of cyanide compounds by gut microbiomes of bamboo-eating pandas. *mxSphere* 3 (3).
- Zhu, S., Jiang, X., Boudreau, M.D., Feng, G., Miao, Y., Dong, S., Wu, H., Zeng, M., Yin, J.J., 2018b. Orally administered gold nanoparticles protect against colitis by attenuating Toll-like receptor 4- and reactive oxygen/nitrogen species-mediated inflammatory responses but could induce gut dysbiosis in mice. *J. Nanobiotechnology* 16 (1), 86.
- Zhu, X., Han, Y., Du, J., Liu, R., Jin, K., Yi, W., 2017. Microbiota-gut-brain axis and the central nervous system. *Oncotarget* 8 (32), 53829–53838.