

Molecular characterization and phylogenetic analysis of *Trypanosoma* spp. detected from striped leaf-nosed bats (*Hipposideros vittatus*) in Zambia

Yongjin Qiu^a, Masahiro Kajihara^b, Hayato Harima^a, Bernard Mudenda Hang'ombe^{c,d,e}, Ryo Nakao^f, Kyoko Hayashida^g, Akina Mori-Kajihara (Dr.)^b, Katendi Changula^c, Yoshiki Eto^b, Joseph Ndebe^h, Reiko Yoshida^b, Yoshihiro Takadate (Dr.)^b, Daniel Mwizabiⁱ, Hiroki Kawabata^j, Martin Simuunza^h, Aaron Mweene^{d,e,h}, Hirofumi Sawa^{d,e,h,k,l,m}, Ayato Takada (Professor)^{b,h,k}, Chihiro Sugimoto^{g,h,k,*}

^a Hokudai Center for Zoonosis Control in Zambia, School of Veterinary Medicine, The University of Zambia, PO Box 32379, Lusaka, 10101, Zambia

^b Division of Global Epidemiology, Hokkaido University Research Center for Zoonosis Control, Kita-20 Nishi-10, Kita-ku, Sapporo, Hokkaido, 001-0020, Japan

^c Department of Para-clinical Studies, School of Veterinary Medicine, The University of Zambia, P.O. Box 32379, Lusaka, 10101, Zambia

^d Global Virus Network Affiliate Center of Excellence, The University of Zambia, P.O. Box 32379, Lusaka, 10101, Zambia

^e African Center of Excellence for Infectious Diseases of Humans and Animals, P.O. Box 32379, Lusaka, 10101, Zambia

^f Laboratory of Veterinary Parasitology, Graduate School of Infectious Diseases, Hokkaido University, Kita-18 Nishi-9, Kita-ku, Sapporo, Hokkaido, 060-0818, Japan

^g Division of Collaboration and Education, Hokkaido University Research Center for Zoonosis Control, Kita-20 Nishi-10, Kita-ku, Sapporo, Hokkaido, 001-0020, Japan

^h Department of Disease Control, School of Veterinary Medicine, University of Zambia, P.O. Box 32379, Lusaka, 10101, Zambia

ⁱ Department of National Parks and Wildlife, Ministry of Tourism and Arts, Chilanga, 101010, Zambia

^j Department of Bacteriology-I, National Institute of Infectious Diseases, Toyama 1-23-1, Shinjyuku-ku, Tokyo, 162-8640, Japan

^k Global Station for Zoonosis Control, Global Institution for Collaborative Research and Education (GI-CoRE), Hokkaido University, Kita-20 Nishi-10, Kita-ku, Sapporo, Hokkaido, 001-0020, Japan

^l Global Virus Network, 725 West Lombard St, Baltimore, MD, 21201, USA

^m Division of Molecular Pathology, Hokkaido University Research Center for Zoonosis Control, Kita-20 Nishi-10, Kita-ku, Sapporo, Hokkaido, 001-0020, Japan

ARTICLE INFO

Keywords:

Zambia

Bat

Trypanosoma cruzi clade

Trypanosoma conorhini

Trypanosoma dionisii

ABSTRACT

Bat trypanosomes consist of more than 30 trypanosome species from over 70 species of bats. Recent studies suggest that bats play a role in disseminating trypanosomes from African continent to the terrestrial mammals both in the Afrotropic-Palearctic Ecozones and Nearctic Ecozone. However, the diversity, distribution, and evolution of bat trypanosomes are still unclear. To better understand their evolution, more genetic data of bat trypanosomes from a variety of locations are required. During a survey of *Borrelia* spp. of bats inhabiting a cave in Zambia, we observed flagellate parasites from 5 of 43 hemocultures. Sequence and phylogenetic analyses of the glycosomal glyceraldehyde 3-phosphate dehydrogenase gene (*gGAPDH*; 572 bp) and the 18S ribosomal RNA gene (18S rRNA gene; 1,079–1,091 bp) revealed that all were *Trypanosoma* spp. belonged to the *Trypanosoma cruzi* clade. Three and two of them exhibited the similarity with *T. conorhini* and *T. dionisii*, respectively. The present study provides the first genetic data on *Trypanosoma* spp. of bats inhabiting Zambia.

1. Introduction

Approximately 1,240 bat species are recognized in the world, representing about 20% of all classified mammalian species worldwide. They play a role as a natural reservoir of wide variety of pathogens including virus, bacteria, and protozoa (Calisher et al., 2006). Trypanosomes, which are blood parasites widespread in all continents and commonly transmitted by blood sucking arthropods and leeches

(Hoare, 1972; Hamilton et al., 2007), have been reported in over 70 bat species in the world. These bat trypanosome species belong to three subgenera, namely *Herpetosoa*, *Megatrypanum*, and *Schizotrypanum*. *T. cruzi*, which is a causative agent of human Chagas disease in South America (Bern, 2015), has been detected in a variety of terrestrial animals including bats. Many species in the subgenus *Schizotrypanum* constitute a large monophyletic assemblage that has been designated as *T. cruzi* clade. The members of bat trypanosomes in the *T. cruzi* clade

* Corresponding author. Division of Collaboration and Education, Hokkaido University Research Center for Zoonosis Control, Kita 20, Nishi 10, Kita-Ku, Sapporo, Hokkaido, 001-0020, Japan.

E-mail address: sugimoto@czc.hokudai.ac.jp (C. Sugimoto).

<https://doi.org/10.1016/j.ijppaw.2019.04.009>

Received 17 December 2018; Received in revised form 23 April 2019; Accepted 24 April 2019

2213-2244/ © 2019 The Authors. Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

include *T. vespertilionis* in European bats and *T. vespertilionis*-like *Trypanosoma* spp. in West African bats, *T. cruzi marinkellei* in South American bats, and *T. dionisii* in European, Asian, and South American bats (Hoare, 1972; Mafie et al., 2018). Recent molecular evidence from studies on bat trypanosomes suggest that the *T. cruzi* clade evolved from a broader clade of bat trypanosomes, and that bat trypanosomes had successfully made the host switch to other terrestrial mammalian species in both the Nearctic Ecozone and Afrotropic-Palearctic Ecozones (Hamilton et al., 2012a). Therefore, molecular investigation of the genus *Trypanosoma* in bats all over the world is important for better understanding of evolution and diversity of pathogenic trypanosomes.

In Sub-Saharan Africa, tsetse-transmitted trypanosomes, including *T. brucei* sensu lato, *T. congolense*, and *T. vivax*, threaten human and animal health (Büscher et al., 2017; Morrison et al., 2016). Non-tsetse-transmitted trypanosomes such as *T. lewisi* are also known to be prevalent in rodents in the region (Hoare, 1972; Keymer, 1971). In bats, *T. livingstonei* and *T. erneyi* belonging to the *T. cruzi* clade were detected in Mozambique (Lima et al., 2012, 2013). The ancestral species of the parasites in the *T. cruzi* clade in bats is considered as the origin of trypanosomes in land mammals. The phylogenetic position of *T. livingstonei* is peripheral to *T. cruzi* clade; this fact supports the hypothesis that *T. cruzi* originated from bats (Lima et al., 2013). Thus, genetic investigation of trypanosomes in bats is important to understand whole picture of evolution and distribution of trypanosomes and their genetic diversity.

During the survey on *Borrelia* spp. in bats captured in a cave in Zambia, where a patient suffering from relapsing fever got a tick bite (Qiu et al., in press), we unexpectedly observed flagellate protozoa in the five hemocultures. We found that these parasites were phylogenetically divided into two distinct subclades within the *T. cruzi* clade. This study provides the first genetic characterization of bat trypanosomes in Zambia.

2. Materials and methods

2.1. Culture

In December 2017, a total of 43 bats (32 *Rousettus aegyptiacus* and 11 *Hipposideros vittatus*) were captured at the cave (15.44 S, 28.51 E) in Zambia as part of a surveillance program of filovirus infection and *Borrelia* spp. in accordance with the ethical standards approved by the Department of National Parks and Wildlife (formerly Zambia Wildlife Authority), Ministry of Tourism and Arts of the Republic of Zambia (Act No. 12 of 1998) (Changula et al., in press). Uncoagulated whole blood samples were collected from bats. For the purpose of *Borrelia* spp. isolation, 100 µl of the peripheral blood were added into the Barbour-Stoenner-Kelly (BSK)-M medium and incubated for 4 weeks at 34 °C, with 5.0% CO₂ as previously described (Takano et al., 2014). The culture was observed microscopically every 2–3 days.

2.2. Morphological investigation

Giemsa staining was carried out for visualizing flagellate protozoa from one of the bat samples in the BSK-M medium (ZB17-105). Microscopy (Shimadzu Motic BA210E) was used for observation and image acquisition of the parasites.

2.3. DNA extraction and polymerase chain reaction (PCR)

The culture media that contained the parasites were harvested at 2 weeks after culture initiation. After centrifugation at 1,600 g for 10 min, DNA was extracted from the pellet using DNAzol (Invitrogen, MA, USA) according to the manufacturer's instructions.

PCR amplification of the glycosomal glyceraldehyde 3-phosphate dehydrogenase (*gGAPDH*), and 18S ribosomal RNA genes was carried out using previously described primers (Table 1) (Hamilton et al., 2004;

da Silva et al., 2004; Lemos et al., 2015). All PCR reactions were conducted in a 20 µl-reaction mixture containing 2 µl of 10 × Ex Taq Buffer (TaKaRa Bio Inc., Shiga, Japan), 0.1 µl of Ex Taq Hot Start Version (TaKaRa Bio Inc.), 1.6 µl of 2.5 mM dNTPs mixture, 200 nM of each primer, and 2 µl of template DNA. The reaction conditions were 98 °C for 1 min and 35 cycles of 98 °C for 10 s, 55 °C for 30 s, and 72 °C for 60 s, followed by a final extension at 72 °C for 5 min. The PCR products were subjected to electrophoresis in a 1.2% agarose gel and stained with ethidium bromide.

2.4. Sequencing and phylogenetic analyses

Cycle sequencing was performed using the BigDye Terminator version 3.1 chemistry (Applied Biosystems, MA, USA). Sequencing products were run on a 3130xl Genetic Analyzer (Applied Biosystems). Sanger sequencing data were analyzed using GENETYX version 9.1 (GENETYX Corporation, Tokyo, Japan). In order to obtain a longer sequence of the 18S rRNA gene, sequences of each sample from 2 PCRs targeting 18S rRNA gene were combined. Approximately 1,080-bp fragments of the 18S rRNA gene sequence were obtained.

The obtained *gGAPDH* and 18S rRNA gene sequences of trypanosomes from bats were aligned with closely related parasite sequences deposited in the database (DDBJ/EMBL/GenBank) using ClustalW multiple alignment program. Phylogenetic trees were constructed by using three methods; the maximum likelihood, neighbor joining, and minimum evolution methods embedded in the MEGA version 6.06 (Tamura et al., 2013). Phylograms were constructed with concatenated *gGAPDH* and 18S rRNA gene sequences, as previously described (Cottonatil et al., 2014; Espinosa-Álvarez et al., 2018). The DDBJ/EMBL/GenBank accession numbers obtained in this study are as follows: *gGAPDH*: LC415422 and LC415423, 18S rRNA gene: LC415424 and LC415425.

2.5. Species delimitation

Poisson tree processes (PTP) model for species delimitation was used for inferring the relationship between *Trypanosoma* spp. detected in this study and their closely related species. PTP species delimitation analysis was carried out via the bPTP webserver (Zhang et al., 2013). We employed a maximum likelihood phylogeny of the concatenated sequences of 18S rDNA and *gGAPDH* with default parameters as reported elsewhere (Cottontail et al., 2014).

3. Results

3.1. Detection of flagellate protozoa

Of the 43 bat blood samples that were added to the BSK-M medium, flagellate protozoa were observed at about 1 week after culture initiation in 5 samples from *H. vittatus*. During the 4 weeks of observation period, the number of parasites decreased gradually. Though the parasites were co-cultured with primary cells derived from *H. vittatus*, the number of parasites did not increase (data not shown). Microscopic observation of the slides prepared from one culture (ZB17-105) clearly showed flagellate protozoa with stumpy and slender forms of variable sizes (13.0–36.0 µm from the anterior to the posterior) (Fig. 1). Unfortunately, the morphological data from other cultures were not recorded.

3.2. Sequence analysis of *Trypanosoma* spp.

Sequence analysis of *gGAPDH* revealed the presence of two distinct sequence types. The *gGAPDH* sequence type 1 from the samples ZB17-105, –111, and –115 (GenBank no. LC415422) showed 97.0% (555/572 bp) identity with the *T. conorhini* isolate TCC2156, which was detected from a triatomine (*Triatomia rubrofasciata*) of Hawaii

Table 1
Primers used in this study.

Primer	Sequence 5'-3'	Target gene	Amplicon size	Reference
TRY927F	GAAACAAGAAACACGGGAG	18S rRNA gene	900 bp	Hamilton et al. (2004)
TRY927R	CTACTGGGCAGCTTGGGA			
609F	CACCCGCGGTAATTCCAGC	18S rRNA gene	900 bp	da Silva et al. (2004)
706R	TCTGAGACTGTAACCTCAA			
GAP3F	GTGAAGGCGCAGCGCAAC	<i>gGAPDH</i> gene	600 bp	Lemos et al. (2015)
GAP5R	CCGAGGATGYCCTTCATG			

(GenBank no. MF144731) (Espinosa-Álvarez et al., 2018). The sequence type 2 from the samples ZB17-108 and –109 (GenBank no. LC415423) showed 94.2% (539/572 bp) identity with the *T. dionisii* isolate TryCC495 from a bat (*Carollia perspicillata*) in Brazil (GenBank no. GQ140363) (Cavazzana et al., 2010).

The same grouping was obtained by the sequence analysis of 18S rRNA. The 18S rRNA gene of the sequence type 1 from the samples ZB17-105, –111, and –115 (GenBank no. LC415424) showed 97.7% (1074/1099 bp) identity with *Trypanosoma* sp. isolate NanDoum1 from the African palm civet (*N. binotata*) in Cameroon (GenBank no. FM202492) (Hamilton et al., 2009). The sequence type 2 from the samples ZB17-108 and –109 (GenBank no. LC415425) showed 95.2% (1033/1084 bp) identity with the *T. dionisii* isolate P3 from a bat (*Pipistrellus pipistrellus*) captured in the United Kingdom (GenBank no. AJ009151) (Stevens et al., 1998).

3.3. Phylogenetic analysis of *Trypanosoma* spp.

In the phylogenetic trees based on the concatenated 18S rRNA and *gGAPDH* gene sequences, the sequence type 1 (ZB17-105, –111, and –115) and type 2 (ZB17-108 and –109) belonged to the clades clustering respectively with *T. conorhini* and *T. dionisii* (Fig. 2). The topologies of the trees generated with three different methods are in good agreement, except for the position of parasites in the Australian clade (*T. noyesi* and *Trypanosoma* sp. isolate H25) and the Neobats clade (*T. wauwau*) (Fig. 2). The PTP analysis inferred two types of *Trypanosoma* spp. detected in this study as putative species (Fig. 3). *Trypanosoma* sp. sequence type 1 and *T. conorhini* were clustered together, while *Trypanosoma* sp. sequence type 2 and *T. dionisii* were clustered together, both of which formed a distinct monophyletic group (Fig. 3).




Fig. 1. Giemsa staining of *Trypanosoma* sp. from ZB17-105 in BSK-M medium. Representative images of ZB17-105 in the BSK-M medium are displayed at the same magnification (x1000). (a,b) flagellates resembling promastigote forms. (c) possibly epimastigote forms under division. K: kinetoplast, N: nucleus, F: flagellum.

4. Discussion

We observed two distinct sequence types of *Trypanosoma* spp. in BSK-M medium inoculated with blood samples of striped leaf-nosed bats (*H. vittatus*). BSK medium and its derivatives are generally used for the isolation of *Borrelia* spp., however, Mafie and colleagues also incidentally isolated *T. dionisii* from an Eastern bent-winged bat (*Miniopterus fuliginosus*) using the BSK medium (Mafie et al., 2018). To the best of our knowledge, this study is the first record of trypanosome infection in *H. vittatus* bats and the first molecular study of bat trypanosomes in Zambia.

The nucleotide sequences of both *gGAPDH* and 18S rRNA genes of the samples ZB17-105, –111, and –115 (sequence type 1) showed high similarity with that of a trypanosome member of the *T. conorhini* clade. In the phylogenetic trees and PTP analysis based on the concatenated 18S rRNA and *gGAPDH* gene sequences, the detected trypanosome was a putative species forming a monophyletic group with *T. conorhini*. *T. conorhini*, a non-pathogenic rat trypanosome, is distributed all over the world (Hoare, 1972). The origin of *T. conorhini* is still unclear. Previously, *T. conorhini* was considered to originate in the Nearctic Ecozone and disseminated to the Afrotropic-Palaearctic Ecozones through rats in ships (Stevens et al., 1999). On the contrary, recent studies with the phylogenetic analysis of the vector bug *Triatoma rubrofasciata* suggested that *T. conorhini* might have originated in the Afrotropic-Palaearctic Ecozones (Hypsa et al., 2002; Patterson et al., 2010). Interestingly, the *Trypanosoma* sp. isolate NanDoum1, a member of the *T. conorhini* clade, was detected from civet in Cameroon, the Afrotropic Ecozones (Hamilton et al., 2009). The fact that *Trypanosoma* sp. sequence type 1 belonging to the *T. conorhini* clade was detected from *H. vittatus* bats in Zambia in this study supports the hypothesis that




Fig. 2. Phylogeny of *Trypanosoma cruzi* clade. Phylogenetic trees of *Trypanosoma cruzi* clade inferred using combined *gGAPDH* and 18S rRNA gene sequences (1,363 bp). The phylogenetic trees with outgroup (*Trypanosoma lewisi*) were generated using a maximum likelihood method (A), neighbor joining method (B), and minimum evolution method (C). All bootstrap values from 1,000 replications are shown on the interior branch nodes. Our isolated *Trypanosoma* spp. are highlighted in bold and under line. GenBank accession numbers used in this phylogeny are listed in Table S1.

the ancestor species of the *T. conorhini* clade might have originated in the Afrotropic Ecozones.

Both *gGAPDH* and 18S rRNA gene sequences of *Trypanosoma* sp. sequence type 2 (ZB17-108 and –109) showed 95.2% and 94.2%

identities with those of *T. dionisii*, respectively. In the phylogenetic trees based on the concatenated *gGAPDH* and 18S rRNA gene sequences, the detected trypanosome formed a monophyletic group with *T. dionisii* distributed in the European and American continents (Hoare, 1972;




Fig. 3. Species delimitation of *Trypanosoma cruzi* clade. Maximum likelihood phylogeny with outgroup (*Trypanosoma lewisi*) and with Bayesian support values presented 17 lineages recognized as species for the PTP analysis. Monophyletic groups in red indicated single putative species as well as terminal branches in blue. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Hamilton et al., 2012b). In addition, the PTP analysis suggested that *Trypanosoma* sp. sequence type 2 is a new putative trypanosome species in the *T. dionisii* clade. Recently, *T. dionisii* was isolated from Eastern bent-winged bats in Japan (Mafie et al., 2018), indicating that *T. dionisii* has a wide geographical distribution. Importantly, a human case of *T. dionisii* infection was reported in Brazil (Dario et al., 2016). Hence, it is particularly of interest to evaluate the pathogenicity and zoonotic potential of the *Trypanosoma* sp. sequence type 2.

Vectors of trypanosomes detected in the current study are not yet known. *T. dionisii* is mainly transmitted by bat bugs (cimicids) (Molyneux, 1991), although the contribution of other vectors such as triatomines cannot be ruled out (Dario et al., 2017). Furthermore, *Trypanosoma* spp. in Schizotrypanum clade were detected from a bat bug (*Stricticimex brevispinosus*) in Burundi, Africa (Van den Berghe et al., 1963). Thus, investigation of bat bugs and other blood-sucking insects in the cave is required to determine particular vector arthropods for the *Trypanosoma* spp. found in this study.

Our findings expanded the knowledge on genetic diversity of trypanosomes infesting African bats. Particularly, the first detection of the parasite in the *T. conorhini* clade in African bats provided important information to estimate the origin, dispersion, and speciation of the trypanosomes in the clade. The presence of the parasite in the Schizotrypanum clade is also of importance to study zoonotic potential of non-tsetse transmitted trypanosomes in the tested region. To better understanding of geographical distribution and individual speciation of trypanosomes in the *T. cruzi* clade, accumulation of the genetic information on trypanosomes in different locations is essential.

Conflicts of interest

The authors declare that they have no conflict of interest.

Acknowledgments

We appreciate Mr. Kenji Yokoi (Japan International Cooperation Agency) for his arrangement of the bat sampling activity. It was very kind of Dr. Keisuke Suganuma to advise us on the storage method of *Trypanosoma* spp. This study funded in part by the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID) [15FM0108008H0001], Japan Society for the Promotion of Science (JSPS) KAKENHI [16K19112], and the Agency for Cooperation Research and Development (AMED) and Japan International Cooperation Agency (JICA) within the framework of the Science and Technology Research Partnership for Sustainable Development (SATREPS) [15JM0110005H0004].

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijppaw.2019.04.009>.

References

- Bern, C., 2015. Chagas' disease. *N. Engl. J. Med.* 373, 456–466.
- Büscher, P., Cecchi, G., Jamonneau, V., Priotto, G., 2017. Human African trypanosomiasis. *Lancet* 390, 2397–2409.
- Calisher, C.H., Childs, J.E., Field, H.E., Holmes, K.V., Schountz, T., 2006. Bats: important reservoir hosts of emerging viruses. *Clin. Microbiol. Rev.* 19, 531–545.
- Cavazzana, M., Marcili, A., Lima, L., da Silva, F.M., Junqueira, A.C., Veludo, H.H., Viola, L.B., Campaner, M., Nunes, V.L., Paiva, F., Coura, J.R., Camargo, E.P., Teixeira, M.M., 2010. Phylogeographical, ecological and biological patterns shown by nuclear (ssrRNA and gGAPDH) and mitochondrial (Cyt b) genes of trypanosomes of the subgenus *Schizotrypanum* parasitic in Brazilian bats. *Int. J. Parasitol.* 40, 345–355.
- Changula, K., Kajihara, M., Mori-Kajihara, A., Eto, Y., Miyamoto, H., Yoshida, R., Shigeno, A., Hang'ombe, B.M., Qiu, Y., Mwizabi, D., Squarre, D., Ndebe, J., Ogawa, H., Harima, H., Simulundu, E., Moonga, L., Kapila, P., Furuyama, W., Kondoh, T., Sato, M., Takadate, Y., Kaneko, C., Nakao, R., Mukonka, V., Mweene, A., Takada, A., (in press). Seroprevalence of filovirus infection of *Rousettus aegyptiacus* bats in Zambia. *J. Infect. Dis.*
- Cottontail, V.M., Kalko, E.K.V., Cottontail, I., Wellinghausen, N., Tschapka, M., Perkins, S.L., Pinto, C.M., 2014. High local diversity of *Trypanosoma* in common bat species, and implications for the biogeography and taxonomy of the *T. cruzi* clade. *PLoS One* 9, e108603.
- da Silva, F.M., Noyes, H., Campaner, M., Junqueira, A.C., Coura, J.R., Añez, N., Shaw, J.J., Stevens, J.R., Teixeira, M.M., 2004. Phylogeny, taxonomy and grouping of *Trypanosoma rangeli* isolates from man, triatomines and sylvatic mammals from widespread geographical origin based on SSU and ITS ribosomal sequences. *Parasitology* 129, 549–561.
- Dario, M.A., Lisboa, C.V., Costa, L.M., Moratelli, R., Nascimento, M.P., Costa, L.P., Leite, Y.L.R., Llewellyn, M.S., Xavier, S.C.D.C., Roque, A.L.R., Jansen, A.M., 2017. High *Trypanosoma* spp. diversity is maintained by bats and triatomines in Espírito Santo state, Brazil. *PLoS One* 12, e0188412.
- Dario, M.A., Rodrigues, M.S., Barros, J.H., Xavier, S.C., D'Andrea, P.S., Roque, A.L., Jansen, A.M., 2016. Ecological scenario and *Trypanosoma cruzi* DTU characterization of a fatal acute Chagas disease case transmitted orally (Espírito Santo state, Brazil). *Parasites Vectors* 9, 477.
- Espinosa-Álvarez, O., Ortiz, P.A., Lima, L., Costa-Martins, A.G., Serrano, M.G., Herder, S., Buck, G.A., Camargo, E.P., Hamilton, P.B., Stevens, J.R., Teixeira, M.M.G., 2018. *Trypanosoma rangeli* is phylogenetically closer to Old World trypanosomes than to *Trypanosoma cruzi*. *Int. J. Parasitol.* 48, 569–584.
- Hamilton, P.B., Adams, E.R., Njiokou, F., Gibson, W.C., Cuny, G., Herder, S., 2009. Phylogenetic analysis reveals the presence of the *Trypanosoma cruzi* clade in African terrestrial mammals. *Infect. Genet. Evol.* 9, 81–86.
- Hamilton, P.B., Cruickshank, C., Stevens, J.R., Teixeira, M.M., Mathews, F., 2012a. Parasites reveal movement of bats between the new and old worlds. *Mol. Phylogenet. Evol.* 63, 521–526.
- Hamilton, P.B., Gibson, W.C., Stevens, J.R., 2007. Patterns of co-evolution between trypanosomes and their hosts deduced from ribosomal RNA and protein-coding gene phylogenies. *Mol. Phylogenet. Evol.* 44, 15–25.
- Hamilton, P.B., Stevens, J.R., Gaunt, M.W., Gidley, J., Gibson, W.C., 2004. Trypanosomes are monophyletic: evidence from genes for glyceraldehyde phosphate dehydrogenase and small subunit ribosomal RNA. *Int. J. Parasitol.* 34, 1393–1404.
- Hamilton, P.B., Teixeira, M.M., Stevens, J.R., 2012b. The evolution of *Trypanosoma cruzi*: the 'bat seeding' hypothesis. *Trends Parasitol.* 28, 136–141.
- Hypsa, V., Tietz, D.F., Zrzavý, J., Rego, R.O., Galvao, C., Jurberg, J., 2002. Phylogeny and biogeography of triatominae (Hemiptera: Reduviidae): molecular evidence of a new world origin of the Asiatic clade. *Mol. Phylogenet. Evol.* 23, 447–457.
- Hoare, C.A., 1972. *The Trypanosomes of Mammals*. Blackwell, Oxford.
- Keymer, I.F., 1971. Blood protozoa of insectivores, bats and primates in Central Africa. *J. Zool. Lond.* 163, 421–441.
- Lemos, M., Fermino, B.R., Simas-Rodrigues, C., Hoffmann, L., Silva, R., Camargo, E.P., Teixeira, M.M., Souto-Padrón, T., 2015. Phylogenetic and morphological characterization of trypanosomes from Brazilian armoured catfishes and leeches reveal high species diversity, mixed infections and a new fish trypanosome species. *Parasites Vectors* 8, 573.
- Lima, L., Espinosa-Álvarez, O., Hamilton, P.B., Neves, L., Takata, C.S., Campaner, M., Attias, M., de Souza, W., Camargo, E.P., Teixeira, M.M., 2013. *Trypanosoma livingstonei*: a new species from African bats supports the bat seeding hypothesis for the *Trypanosoma cruzi* clade. *Parasites Vectors* 6, 221.
- Lima, L., Silva, F.M., Neves, L., Attias, M., Takata, C.S., Campaner, M., de Souza, W., Hamilton, P.B., Teixeira, M.M., 2012. Evolutionary insights from bat trypanosomes: morphological, developmental and phylogenetic evidence of a new species, *Trypanosoma (Schizotrypanum) erneyi* sp. nov., in African bats closely related to *Trypanosoma (Schizotrypanum) cruzi* and allied species. *Protist* 163, 856–872.
- Mafie, E., Rupa, F.H., Takano, A., Suzuki, K., Maeda, K., Sato, H., 2018. First record of *Trypanosoma dionisii* of the *T. cruzi* clade from the Eastern bent-winged bat (*Miopterus fuliginosus*) in the Far East. *Parasitol. Res.* 117, 673–680.
- Molyneux, D.H., 1991. Trypanosomes of bats. In: Kreier, J.P., Baker, J.R. (Eds.), *Parasitic Protozoa*. Academic Press, New York, pp. 195–223.
- Morrison, L.J., Vezza, L., Rowan, T., Hope, J.C., 2016. Animal African trypanosomiasis: time to increase focus on clinically Relevant parasite and host species. *Trends Parasitol.* 32, 599–607.
- Patterson, J.S., Gaunt, M.W., 2010. Phylogenetic multi-locus codon models and molecular clocks reveal the monophyly of haematophagous reduviid bugs and their evolution at the formation of South America. *Mol. Phylogenet. Evol.* 56, 608–621.
- Qiu, Y., Nakao, R., Hang'ombe, B.M., Sato, K., Kajihara, M., Kanchela, S., Changula, K., Eto, Y., Ndebe, J., Sasaki, M., Thu, M.J., Takada, A., Sawa, H., Sugimoto, C., Kawabata, H., (in press). Human borreliosis caused by a new world relapsing fever *Borrelia*-like organism in the old world. *Clin. Infect. Dis.*
- Stevens, J., Noyes, H., Gibson, W., 1998. The evolution of trypanosomes infecting humans and primates. *Mem. Inst. Oswaldo Cruz* 93, 669–676.
- Stevens, J.R., Teixeira, M.M., Bingle, L.E., Gibson, W.C., 1999. The taxonomic position and evolutionary relationships of *Trypanosoma rangeli*. *Int. J. Parasitol.* 29, 749–757.
- Takano, A., Toyomane, K., Konnai, S., Ohashi, K., Nakao, M., Ito, T., Andoh, M., Maeda, K., Watarai, M., Sato, K., Kawabata, H., 2014. Tick surveillance for relapsing fever spirochete *Borrelia miyamotoi* in Hokkaido, Japan. *PLoS One* 8, e104532.
- Tamura, K., Stecher, G., Peterson, D., Filipiak, A., Kumar, S., 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729.
- Van den Berghe, L., Chardome, M., Peel, E., 1963. An African bat trypanosome in *Stricticimex brevispinosus* Usinger, 1959. *J. Protozool.* 10, 135–138.
- Zhang, J., Kapli, P., Pavlidis, P., Stamatakis, A., 2013. A general species delimitation method with applications to phylogenetic placements. *Bioinformatics* 29, 2869–2876.