

The Impact of Frailty on Artificial Urinary Sphincter Placement and Removal Procedures



Andrew R. Medendorp, Jennifer T. Anger, Chengshi Jin, Katherine A. Amin, Lindsay A. Hampson, Una J. Lee, and Anne M. Suskind

OBJECTIVE	To determine whether frailty is associated with increased odds of 30-day surgical complications among men undergoing both artificial urinary sphincter (AUS) placement and removal procedures and to determine whether frailty was associated with increased odds of having an AUS removal procedure.
METHODS	This is a retrospective cohort study of men undergoing AUS placement and removal procedures using data from the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) from 2006 to 2013. Frailty was quantified using the NSQIP-FI (frailty index) and was applied to logistic regression models predicting 30-day complications (overall, major, and minor) and the odds of having an AUS removal procedure (over an AUS placement procedure).
RESULTS	We identified a total of 624 and 147 men undergoing AUS placement and removal procedures, respectively. NSQIP-FI of ≥ 0.27 , but not age, was associated with major complications (aOR 3.5, 95% confidence interval 1.2-9.9), while age ≥ 85 years, but not NSQIP-FI, was associated with minor complications (aOR 7.9, 95% confidence interval 1.4-45.6). Men undergoing AUS removal procedures tended to be more frail compared to men undergoing AUS placement procedures (12.9% vs 6.1% had NSQIP-FI of ≥ 0.27 , $P < .01$).
CONCLUSION	Men undergoing AUS removal procedures are, on average, more frail compared to men undergoing AUS placement procedures. Frailty is associated with increased odds of major complications and with having an AUS removal procedure. These findings highlight the importance of incorporating measures of frailty, instead of age alone, into the perioperative decision-making process for adults considering these types of procedures. UROLOGY 129: 210–216, 2019. © 2019 Elsevier Inc.

Male stress urinary incontinence (SUI), defined as the involuntary loss of urine with effort or exertion,¹ is associated with significant detriments in quality of life² and costs of greater than \$8-10 billion per year in the United States alone.³ A history of prior prostate surgery, most commonly radical prostatectomy, is a major risk factor for the development of male SUI, and up to 20% of men undergoing these types of procedures ultimately undergo an intervention for SUI.³⁻⁶ While a variety of interventions exist, the artificial urinary

sphincter (AUS) is considered to be the gold standard definitive treatment for male SUI.³⁻⁸

Though overall success rates and satisfaction are high, the AUS is a procedure with many known potential complications.^{5,9} Among the most severe of these complications are device infection and erosion, necessitating removal of the device in its entirety.¹⁰ Known risk factors for device removal include diabetes, history of radiation or urethral stricture,¹¹ and higher American Society of Anesthesiologists (ASA) class.¹¹ However, there is reason to believe that frailty, a multifactorial syndrome associated with a reduction in physiologic reserve and an increased susceptibility to adverse events in response to stressors, may place men at heightened risk for device infection and erosion, leading to AUS removal. Some reasons why this may be the case include factors associated with frailty such as sarcopenia, changes in body composition, and malnutrition.¹² While frailty has been demonstrated to be associated with complications among men undergoing other types of urological procedures,^{13,14} its association specifically with AUS surgery has not been studied.

Financial Disclosure: Pepper Center Research Career Development Core (RCDC), NIH-NIDDK K12 DK83021-09; K12 Urologic Research (KUR) Career Development Program.

From the Department of Urology, University of California, Los Angeles (UCLA), Los Angeles, CA; the Division of Urology, Department of Surgery, Cedars Sinai Medical Center, Los Angeles, CA; the Department of Epidemiology and Biostatistics, University of California, San Francisco (UCSF), San Francisco, CA; the Section of Urology, Virginia Mason Medical Center, Seattle, WA; and the Department of Urology, University of California, San Francisco (UCSF), San Francisco, CA

Address correspondence to: Anne M. Suskind, M.D., M.S., F.A.C.S., Department of Urology, University of California, San Francisco (UCSF), 400 Parnassus Ave, Box 0738, San Francisco, CA 94123. E-mail: Anne.Suskind@ucsf.edu

Submitted: January 3, 2019, accepted (with revisions): April 9, 2019

In order to gain a better understanding of the role of frailty on outcomes related to AUS procedures, we performed a retrospective cohort study using data from the American College of Surgeons National Surgical Quality Improvement Project (ACS-NSQIP) from 2006 to 2013. The objectives of this study were (1) to explore the relationship between frailty and complications for both AUS placement and removal procedures and (2) to determine whether frailty is more likely to be associated with men undergoing AUS removal as opposed to AUS placement procedures. Findings from this study will serve to further our understanding of the role of frailty in older men undergoing both types of AUS procedures and will offer potential insights into the importance of incorporating measures of frailty into the preoperative decision-making process.

MATERIALS AND METHODS

Patients and Databases

We conducted a retrospective cohort study using ACS-NSQIP Participant Use Data Files from 2006 to 2013 to identify men undergoing AUS placement and AUS removal procedures. This study was deemed exempt by our institutional review board. The ACS-NSQIP Participant Use Data Files represent a prospectively maintained multicenter database composed of data extracted by trained and certified surgical clinical reviewers at each participating institution, with periodic audits performed to confirm inter-rater reliability. Extracted data include information on patient demographics, the surgical procedure, and 30-day postoperative outcomes.¹⁵ Cases were identified in the ACS-NSQIP database using the following current procedural terminology (CPT) codes: 53445 for AUS placement and 53446 for AUS removal.

Outcomes

We had 2 main outcomes in this study. The first outcome was to determine the incidence of 30-day complications among men undergoing AUS placement and removal procedures. Complications were evaluated overall and also stratified into major and minor complications, based existing categories in the literature.^{14,16-18} Major complications included hospital readmission, return to operating room, sepsis, deep vein thrombosis, cerebrovascular accident, reintubation, renal failure, myocardial infarction, pulmonary embolus, septic shock, wound dehiscence, cardiac arrest, deep wound infection, coma, and death. Minor complications included urinary tract infection, superficial wound infection, pneumonia, and blood transfusion. The second outcome of interest was to determine whether frailty was associated with men having an AUS removal procedure compared to an AUS placement procedure.

Covariates

We used the NSQIP frailty index (NSQIP-FI) as our measure of frailty, which was developed specifically for use with the NSQIP database. This frailty index was adapted from the Canadian Study of Health and Aging Functional Index which is a cumulative deficits model. The NSQIP-FI includes 11 factors—history of diabetes mellitus, impaired functional status, history of chronic obstructive pulmonary disease or pneumonia, congestive heart failure within 30 days preoperatively, history of myocardial infarction within 6 months prior to surgery, history of

percutaneous coronary intervention/angina/cardiac surgery, hypertension requiring medication, peripheral vascular disease or rest pain, impaired sensorium, history of transient ischemic attack, and history of cerebrovascular accident. The NSQIP-FI score is determined by summing the number of factors present and dividing by 11, yielding a number such as 0.09 for 1 out of 11 deficits, 0.18 for 2 out of 11 deficits, etc. For the purposes of this study, we categorized frailty into the following ascending categories: 0, 0.09, 0.18, or ≥ 0.27 consistent with the literature.¹⁹ The NSQIP-FI specifically assesses frailty and has been shown to correlate well with morbidity and mortality in surgical subspecialties including Urology.^{14,20}

Additional covariates obtained from NSQIP data included demographic information such as patient age in years (categorized as <65, 65-74, 75-84, ≥ 85), race (white and nonwhite), and body mass index (BMI) (underweight [<18.5], normal [$18.5-24.9$], overweight [$25-29.9$], and obese [≥ 30]).

Statistical Analysis

Continuous data were reported as means with standard deviations and compared with Mann-Whitney tests and categorical data were reported as numbers with percentages and were compared using the chi-square tests. The first study objective was to determine odds of complications, both (1) overall, and stratified by (2) major and (3) minor complications. Three separate logistic regression models were performed with each category of complications as the dependent variable (ie, overall, major, and minor complications) and with the type of procedure (AUS placement and AUS removal), age, race, BMI, and NSQIP-FI as the independent variables. The second study objective was to determine whether frailty was associated with having an AUS removed (vs having an AUS placed). In order to address this question, we performed a separate logistic regression where procedure type (AUS removal) was the dependent variable, and age, race, BMI, and NSQIP-FI were the independent variables. In order to determine whether there was an interaction between these 2 variables and found the interaction was not statistically significant (data not reported). Therefore, this term was left out of the final models. All analyses were performed using SAS, version 9.4.

RESULTS

We identified a total of 624 individuals undergoing AUS placement procedures and 147 individuals undergoing AUS removal procedures, demonstrated in Table 1. Individuals undergoing AUS removal procedures were significantly older than those having AUS placement procedures (75.2 ± 11.1 vs 69.8 ± 9.31 years, $P < .01$), with 20.4% and 3.7% over the age of 85 years, respectively ($P < .01$). There were no statistically significant differences in race or average BMI between men undergoing the 2 procedures; however, individual BMI categories did differ between groups. NSQIP-FI was significantly higher among individuals undergoing AUS removal, with 12.9% having a NSQIP-FI of ≥ 0.27 compared to only 6.1% among individuals undergoing AUS placement ($P < .01$).

Predictors of Postoperative Complications

Table 2 shows results of the multivariate logistic regression models predicting the odds of any type of complication, and both major and minor complications in 3 separate models.

Table 1. Demographic characteristics of study cohort

Variable	AUS Placement (N = 624)	AUS Removal (N = 147)	P Value
Age (mean ± SD)	69.8 ± 9.31	75.2 ± 11.1	<.01
Age category (N [%])			
<65	151 (24.2%)	21 (14.3%)	<.01
65-74	286 (45.8%)	34 (23.1%)	
75-84	164 (26.3%)	62 (42.2%)	
≥85	23 (3.7%)	30 (20.4%)	
Race (N (%))			
White	506 (90.0%)	122 (91.7%)	.75
Nonwhite	56 (10.0%)	11 (8.3%)	
BMI (mean ± SD)	29.5 ± 5.01	29 ± 7.06	.10
BMI category (N [%])			
Underweight (<18.5)	4 (0.6%)	5 (3.5%)	<.01
Normal (18.5-24.9)	95 (15.3%)	32 (22.5%)	
Overweight (25-29.9)	275 (44.2%)	53 (37.3%)	
Obese (≥30.0)	248 (39.9%)	52 (36.6%)	
NSQIP-FI score (mean ± SD (N))	0.10 ± 0.08 (N=624)	0.13 ± 0.10 (N=147)	<.01
NSQIP-FI categories (N (%))			
0	159 (25.5%)	27 (18.4%)	<.01
0.09	281 (45.0%)	55 (37.4%)	
0.18	146 (23.4%)	46 (31.3%)	
≥0.27	38 (6.1%)	19 (12.9%)	
Components of 11-item NSQIP FI (N (%))			
Diabetes mellitus	123 (19.7%)	42 (28.6%)	.01
Partially/totally dependent functional status	8 (1.3%)	16 (11.0%)	<.01
History of severe chronic obstructive pulmonary disease or pneumonia	24 (3.8%)	9 (6.1%)	.22
Congestive heart failure within 30 days preoperatively	3 (0.5%)	1 (0.7%)	.76
Myocardial infarction in 6 months prior to surgery	0 (0.0%)	2 (2.4%)	<.01
Any prior angina, percutaneous vascular intervention, or cardiac surgery	69 (18.8%)	28 (34.1%)	<.01
Hypertension requiring medication	437 (70.0%)	98 (66.7%)	.43
Peripheral vascular disease or rest pain	2 (0.5%)	1 (1.2%)	.50
Impaired sensorium	0 (0.0%)	2 (2.4%)	<.01
History of transient ischemic attack	21 (5.7%)	8 (9.8%)	.18
History of cerebral vascular accident	6 (1.6%)	4 (4.9%)	.07

The first model demonstrates that AUS removal procedures (compared to AUS placement procedures) were associated with increased odds of any complication (adjusted odds ratio [OR] 1.9, 95% confidence interval [CI] 1.1-3.5; $P = .03$). Age, race, BMI, and NSQIP-FI were not significant predictors of this outcome. The second model shows that NSQIP-FI of ≥ 0.27 was a significant predictor of major complications (adjusted OR 3.5, 95% CI 1.2-9.9; $P = .02$), while neither type of procedure (ie, AUS placement or removal) nor age were statistically significant. The third model shows that AUS removal procedures (adjusted OR 3.2, 95% CI 1.4-7.6; $P < .01$) and age ≥ 85 (adjusted OR 7.9, 95% CI 1.4-45.6; $P = .02$) were statistically significant predictors of minor complications, while frailty was not a significant predictor (NSQIP-FI ≥ 0.27 adjusted OR 1.0, 95% CI 0.2-6.5; $P = .97$).

Determinants of Having an AUS Removal Procedure

Table 3 depicts the multivariate logistic regression model predicting the odds of having an AUS removal procedure compared to having an AUS placement procedure. Age 75-84 years (adjusted OR 2.9, 95% CI 1.6-5.4; $P < .01$) and age ≥ 85 years (adjusted OR 9.8, 95% CI 4.3-22.2; $P < .01$), NSQIP-FI ≥ 0.27 (adjusted OR 2.4, 95% CI 1.1-5.5, $P < .04$) and underweight BMI (adjusted OR 6.0, 95% CI 1.4-26.8, $P = .02$) were associated with having an AUS removal procedure over an AUS placement procedure.

DISCUSSION

This nationwide study on AUS surgery demonstrated that individuals undergoing AUS removal procedures were, on average, older and more frail compared to those having AUS placement procedures. Additionally, frailty (and not age) was associated with major complications while age (and not frailty) was associated with minor complications. The type of procedure (having an AUS removal compared to an AUS placement procedure) was associated with increased odds of both major complications and with complications overall. Finally, increased frailty and age were both associated with increased odds of undergoing AUS removal procedures (compared to AUS placement procedures).

This is the first study to assess the impact of frailty specifically on AUS surgery, either removal or placement. Other studies have evaluated the association of several nonfrailty-related factors with complications following AUS surgery. One study of 155 patients undergoing primary AUS placement evaluated multiple factors including age, BMI, and comorbidities (ie, diabetes and hypertension) and found that none were significant predictors for repeat surgery on either univariate or multivariate analysis.¹⁰ A prospective study of 386 patients with a minimum of 3-month follow-up found that prior history of pelvic radiation, prior AUS

Table 2. Multivariate logistic regression model predicting odds of complications overall and by major and minor complications

Variable	Complications Overall				Major Complications				Minor Complications			
	N/total (%)	Odds ratio	95% CI	P Value	N/total (%)	Odds ratio	95% CI	P Value	N/total (%)	Odds ratio	95% CI	P Value
Procedure type												
AUS placement	43/560 (7.7%)	1.0			36/560 (6.4%)	1.0			14/560 (2.5%)	1.0		
AUS removal	24/130 (18.5%)	1.9	1.1-3.5	.03	17/130 (13.1%)	1.7	0.9-3.4	.11	14/130 (10.8%)	3.2	1.4-7.6	<.01
Age												
<65	11/157 (7.0%)	1.0			10/157 (6.4%)	1.0			2/157 (1.3%)	1.0		
65-74	21/287 (7.3%)	1.0	0.4-2.1	.92	19/287 (6.6%)	0.9	0.4-2.1	.84	8/287 (2.8%)	2.0	0.4-9.9	.39
75-84	26/201 (12.9%)	1.6	0.7-3.4	.28	20/201 (10.0%)	1.2	0.5-2.8	.68	11/201 (5.5%)	3.2	0.7-16.2	.14
≥85	9/45 (20.0%)	2.4	0.8-6.7	.11	4/45 (8.9%)	1.0	0.3-3.5	.94	7/45 (15.6%)	7.9	1.4-45.6	.02
Race												
White	58/623 (9.3%)	1.0			46/623 (7.4%)	1.0			25/623 (4.0%)	1.0		
Nonwhite	9/67 (13.4%)	1.6	0.7-3.6	.30	7/67 (10.4%)	1.2	0.5-3.1	.66	3/67 (4.5%)	2.0	0.6-7.4	.29
BMI												
Underweight (<18.5)	3/9 (33.3%)	3.1	0.6-16.0	.19	3/9 (33.3%)	3.6	0.7-18.9	.13	0/9 (0.0%)	<0.1		
Normal (18.5-24.9)	12/115 (10.4%)	1.0			10/115 (8.7%)	1.0			5/115 (4.3%)	1.0		
Overweight (25-29.9)	29/284 (10.2%)	1.1	0.5-2.3	.77	21/284 (7.4%)	0.8	0.4-1.9	.68	15/284 (5.3%)	1.6	0.5-4.7	.43
Obese (≥30.0)	23/282 (8.2%)	0.8	0.4-1.9	.67	19/282 (6.7%)	0.7	0.3-1.6	.40	8/282 (2.8%)	0.9	0.3-3.1	.85
NSQIP-FI score												
0	12/158 (7.6%)	1.0			9/158 (5.7%)	1.0			4/158 (2.5%)	1.0		
0.09	24/301 (8.0%)	1.0	0.5-2.2	.94	19/301 (6.3%)	1.2	0.5-2.7	.71	9/301 (3.0%)	1.0	0.3-3.3	.94
0.18	21/181 (11.6%)	1.4	0.7-3.1	.38	16/181 (8.8%)	1.6	0.7-3.8	.32	13/181 (7.2%)	2.2	0.7-7.5	.19
≥0.27	10/50 (20.0%)	2.6	1.0-6.9	.05	9/50 (18.0%)	3.5	1.2-9.9	.02	2/50 (4.0%)	1.0	0.2-6.5	.97

Table 3. Logistic regression model predicting odds of having an AUS removal procedure over an AUS placement procedure

Variable	N/total (%)	Odds Ratio	Lower 95% CI	Upper 95% CI	P Value
Age					
<65	17/157 (10.8%)	1.0			
65-74	30/287 (10.5%)	0.9	0.5	1.7	.67
75-84	58/201 (28.9%)	2.9	1.6	5.4	<.01
≥85	25/45 (55.6%)	9.8	4.3	22.2	<.01
Race					
White	119/623 (19.1%)	1.0			
Nonwhite	11/67 (16.4%)	0.9	0.4	1.9	.73
BMI					
Underweight (<18.5)	5/9 (55.6%)	6.0	1.4	26.8	.02
Normal (18.5-24.9)	27/115 (23.5%)	1.0			
Overweight (25-29.9)	49/284 (17.3%)	0.9	0.5	1.5	.57
Obese (≥30.0)	49/282 (17.4%)	0.9	0.5	1.7	.80
NSQIP-FI score					
0	22/158 (13.9%)	1.0			
0.09	48/301 (15.9%)	1.0	0.6	1.9	.94
0.18	44/181 (24.3%)	1.8	1.0	3.4	.07
≥0.27	16/50 (32.0%)	2.4	1.1	5.5	.04

erosion, and history of urethral stent placement were associated with increased risk of AUS explantation.²¹ Finally, a retrospective study of men undergoing AUS placement procedures found that pelvic irradiation, history of urethral stricture, and higher ASA classification (reflecting increased comorbidities) increased the risk of subsequent explantation.¹¹

Our prior work explored the impact of frailty on outcomes related to other types of urologic surgery. Our studies of multiple urologic procedures using the ACS-NSQIP database reported an increase in postoperative complications (both major and minor) and an increase in discharge to a skilled or assisted living facility associated with increasing NSQIP-FI score.^{14,18} We also found an association between increased frailty and higher complications in pelvic organ prolapse surgery.¹⁹

This study is novel in that it investigates the role of frailty specifically with regard to AUS surgery. We found that increased frailty (and not age) was associated with major complications, while age (and not frailty) was associated with minor complications. This finding highlights the point that frailty may be a more valuable and dynamic predictor of worse outcomes/complications than age alone and that frailty may not be as problematic for relatively minor complications.

We also found that the type of procedure made a difference in the odds of complications, whereby AUS removal procedures (compared to AUS placement procedures) were associated with increased odds of both major complications and complications overall, but not with minor complications. This is interesting, as AUS removal procedures are more likely to be performed in the setting of infection or device erosion and it would make sense that poor tissue quality associated with these procedures may lend themselves to increased rates of minor complications such as wound and urinary tract infections. However, this was not the case in our study, but rather, we found that AUS removal procedures were associated with heightened

risk of major complications such as hospital readmission, reintubation, renal failure, cerebrovascular accident, and cardiac arrest.

Interestingly, both age and frailty were significantly associated with device removal procedures (compared to device placement procedures). It makes sense that men undergoing device removal procedures were older than men undergoing device placement procedures, as one must first have a device placed in order to have it removed, making these men (by definition) older than those having the device placed. However, while frailty does tend to increase with increasing age, frailty is highly variable in its association with age, particularly among older age groups. Additionally, in our model, frailty was adjusted for age and there was no significant interaction between age and frailty when tested in a separate model. Therefore, the finding that frailty is significantly associated with device removal is meaningful and important.

There are many reasons why frailty may be associated with adverse outcomes related to AUS surgery. Possible contributing factors in frail older adults could include suboptimal tissue quality, issues with poor wound healing, and poor nutrition—all contributing to overall decreased resiliency and susceptibility to poor postoperative outcomes. These factors may ultimately result in increased likelihood of experiencing AUS erosion or infection, leading to subsequent device removal.

There are 2 important considerations in order to translate findings from this study into clinical practice: (1) how to best measure frailty in the clinical setting, and (2) once you have determined that a patient is frail, how to operationalize this information to improve patient outcomes. While the NSQIP-FI is well suited for frailty measurement within the NSQIP database, other frailty indices are more easily adapted for use in the clinical setting. We have previously published on our use of the Timed Up and Go Tests (TUGT) as a parsimonious measure for frailty in an academic nononcologic urology practice.²² Other

measurement tools include the frailty phenotype, consisting of 5 domains (unintentional weight loss, self-reported exhaustion, weakness, slow walking speed, and low physical activity),²³ and the Edmonton frail scale consisting of 9 domains (cognition, general health status, functional independence, social support, medication use, nutrition, mood, continence, and functional performance).²⁴

Once it is determined that a patient is frail, this information is useful for both risk stratification and risk optimization. For risk stratification, knowledge of a frail phenotype is useful to determine whether an individual is considered to be high- or low-risk for a procedure. This is helpful for patient counseling and in discussion of risks, benefits, and potential alternatives to surgical treatment. In frail patients in whom the decision has already been made to proceed with surgery or in whom surgery is not elective (ie, if they have an infected AUS device that needs removal), knowledge of a frail phenotype can be used to potentially optimize surgical prehabilitation, if there is time, in addition to peri- and postoperative care.

The findings of this study should be considered with certain limitations in mind. Due to the deidentified nature of the data source, we are unable to follow individual patients over time. Furthermore, based on these data, we cannot tell whether or not some patients included in the AUS placement cohort are also included in the AUS removal cohort. However, we think that the potential number of patients included in both groups is likely to be low and would have a minimal impact on our study findings in light of the strengths of the database as a large heterogeneous national sample of procedures. Additionally, we cannot determine whether AUS procedures were primary procedures or repeat procedures, which are known to have higher odd of complications.²⁵ Another limitation of our study is that the ACS-NSQIP database itself is limited to specific predefined variables and does not contain certain other factors likely to influence surgical outcomes such as history of radiation therapy, testosterone levels, prior AUS erosions, or complications beyond 30 days postoperatively. However, with an increasing emphasis on quality-based care, the 30-day complications have become an important metric with high visibility. Finally, our use of composite variables for major and minor complications puts equal statistical weight on adverse events within these groups, though clearly some complications have a greater clinical impact than others. Again, the role of complications in general as a metric for evaluating quality of care can mitigate this limitation and provide value to our findings.

CONCLUSION

This analysis of a nationwide database reveals the impact of frailty on postoperative complications from AUS

surgery. These findings suggest that relying on age alone may be inadequate for risk stratifying patients prior to AUS surgery and that incorporating frailty into the perioperative decision-making process may be beneficial.

References

1. Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology in lower urinary tract function: report from the standardisation sub-committee of the International Continence Society. *Urology*. 2003;61:37–49.
2. Namiki S, Saito S, Tochigi T, Numata I, Ioritani N, Arai Y. Psychological distress in Japanese men with localized prostate cancer. *Int J Urol*. 2007;14:924–929.
3. Brown JA, Elliott DS, Barrett DM. Postprostatectomy urinary incontinence: a comparison of the cost of conservative versus surgical management. *Urology*. 1998;51:715–720.
4. MacDonald S, Colaco M, Terlecki R. Waves of change: national trends in surgical management of male stress incontinence. *Urology*. 2017;108:175–179.
5. Chughtai B, Sedrakyan A, Isaacs AJ, et al. National study of utilization of male incontinence procedures. *Neurourol Urodyn*. 2016;35:74–80.
6. Constable L, Cotterill N, Cooper D, et al. Male synthetic sling versus artificial urinary sphincter trial for men with urodynamic stress incontinence after prostate surgery (MASTER): study protocol for a randomised controlled trial. *Trials*. 2018;19:131.
7. Yafi FA, Powers MK, Zurawin J, Hellstrom WJ. Contemporary review of artificial urinary sphincters for male stress urinary incontinence. *Sex Med Rev*. 2016;4:157–166.
8. Lucas MG, Bosch RJ, Burkhard FC, et al. EAU guidelines on surgical treatment of urinary incontinence. *Eur Urol*. 2012;62:1118–1129.
9. Alwaal A, Harris CR, Awad MA, Allen IE, Breyer BN. Comparison of complication rates related to male urethral slings and artificial urinary sphincters for urinary incontinence: national multi-institutional analysis of ACS-NSQIP database. *Int Urol Nephrol*. 2016;48:1571–1576.
10. Suh YS, Ko KJ, Kim TH, Sung HH, Lee KS. Long-term outcomes of primary implantation and revisions of artificial urinary sphincter in men with stress urinary incontinence. *Neurourol Urodyn*. 2017;36:1930–1937.
11. Husch T, Kretschmer A, Thomsen F, et al. Risk factors for failure of male slings and artificial urinary sphincters: results from a large middle European cohort study. *Urol Int*. 2017;99:14–21.
12. Bergman H, Ferrucci L, Guralnik J, et al. Frailty: an emerging research and clinical paradigm—issues and controversies. *J Gerontol A Biol Sci Med Sci*. 2007;62:731–737.
13. Tan HJ, Shirk JD, Chamie K, Litwin MS, Hu JC. Patient function and the value of surgical care for kidney cancer. *J Urol*. 2017;197:1200–1207.
14. Suskind AM, Walter LC, Jin C, et al. Impact of frailty on complications in patients undergoing common urological procedures: a study from the American College of Surgeons National Surgical Quality Improvement database. *BJU Int*. 2016;117:836–842.
15. American College of Surgeons National Quality Improvement Program (ACS NSQIP). Semiannual Report. 2014.
16. Etzioni DA, Wasif N, Dueck AC, et al. Association of hospital participation in a surgical outcomes monitoring program with inpatient complications and mortality. *JAMA*. 2015;313:505–511.
17. Molina CS, Thakore RV, Blumer A, Obrensky WT, Sethi MK. Use of the national surgical quality improvement program in orthopaedic surgery. *Clin Orthop Relat Res*. 2015;473:1574–1581.
18. Suskind AM, Jin C, Cooperberg MR, et al. Preoperative frailty is associated with discharge to skilled or assisted living facilities after urologic procedures of varying complexity. *Urology*. 2016;97:25–32.
19. Suskind AM, Jin C, Walter LC, Finlayson E. Frailty and the role of obliterative versus reconstructive surgery for pelvic organ prolapse: a national study. *J Urol*. 2017;197:1502–1506.

20. Velanovich V, Antoine H, Swartz A, Peters D, Rubinfeld I. Accumulating deficits model of frailty and postoperative mortality and morbidity: its application to a national database. *J Surg Res*. 2013;183:104–110.
21. Brant WO, Erickson BA, Elliott SP, et al. Risk factors for erosion of artificial urinary sphincters: a multicenter prospective study. *Urology*. 2014;84:934–938.
22. Pangilinan J, Quanstrom K, Bridge M, Walter LC, Finlayson E, Suskind AM. The Timed Up and Go Test as a measure of frailty in urologic practice. *Urology*. 2017;106:32–38.
23. Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56:M146–M156.
24. Rolfson DB, Majumdar SR, Tsuyuki RT, Tahir A, Rockwood K. Validity and reliability of the Edmonton Frail Scale. *Age Ageing*. 2006;35:526–529.
25. Tutolo M, Cornu JN, Bauer RM, et al. Efficacy and safety of artificial urinary sphincter (AUS): results of a large multi-institutional cohort of patients with mid-term follow-up. *Neurourol Urodyn*. 2019;38:710–718.