

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

# Resuscitation

journal homepage: [www.elsevier.com/locate/resuscitation](http://www.elsevier.com/locate/resuscitation)

## Clinical paper

# Chest compression release velocity factors during out-of-hospital cardiac resuscitation



Samuel Beger<sup>a,\*</sup>, John Sutter<sup>a</sup>, Tyler Vadeboncoeur<sup>b,1</sup>, Annemarie Silver<sup>c,2</sup>, Chengcheng Hu<sup>a,d,3</sup>, Daniel W. Spaite<sup>e,4</sup>, Bentley Bobrow<sup>a,e,f,5</sup>

<sup>a</sup> The University of Arizona College of Medicine — Phoenix, Phoenix, AZ, United States

<sup>b</sup> Mayo Clinic, Jacksonville, FL, United States

<sup>c</sup> ZOLL Medical Corporation, Chelmsford, MA, United States

<sup>d</sup> Mel & Enid Zuckerman College of Public Health (MEZCOPH), The University of Arizona, Tucson, AZ, United States

<sup>e</sup> Arizona Department of Health Services, Phoenix, AZ, United States

<sup>f</sup> Arizona Emergency Medicine Research Center, University of Arizona College of Medicine, United States

## Abstract

**Background:** Higher chest compression release velocity (CCRV) has been associated with better outcomes after out-of-hospital cardiac arrest (OHCA), and patient factors have been associated with variations in chest wall compliance and compressibility. We evaluated whether patient sex, age, weight, and time in resuscitation were associated with CCRV during pre-hospital resuscitation from OHCA.

**Methods:** Observational study of prospectively collected OHCA quality improvement data in two suburban EMS agencies in Arizona between 10/1/2008 and 12/31/2016. Subject-level mean CCRV during the first 10 min of compressions was correlated with categorical variables by the Wilcoxon rank-sum test and with continuous variables by the Spearman's rank correlation coefficient. Generalized estimating equation and linear mixed-effect models were used to study the trend of CCRV over time.

**Results:** During the study period, 2535 adult OHCA cases were treated. After exclusion criteria, 1140 cases remained for analysis. Median duration of recorded compressions was 8.70 min during the first 10 min of CPR. An overall decline in CCRV was observed even after adjusting for compression depth. The subject-level mean CCRV was higher for minutes 0–5 than for minutes 5–10 (mean 347.9 mm/s vs. 339.0 mm/s, 95% CI of the difference –12.4 to –5.4,  $p < 0.0001$ ). Males exhibited a greater mean CCRV compared to females [344.4 mm/s (IQR 307.3–384.6) vs. 331.5 mm/s (IQR 285.3–385.5),  $p = 0.013$ ]. Mean CCRV was negatively correlated with age and positively correlated with patient weight.

**Conclusion:** CCRV declines significantly over the course of resuscitation. Patient characteristics including male sex, younger age, and increased weight were associated with a higher CCRV.

**Keywords:** OHCA, Cardiac arrest, Haemodynamics, CPR, Resuscitation, CCRV, Chest compression release velocity, Compressions, Out of hospital cardiac arrest, EMS, Emergency medicine, Feedback, Chest compressions

\* Corresponding author.

E-mail addresses: [sbeger@email.arizona.edu](mailto:sbeger@email.arizona.edu) (S. Beger), [johnsutter@email.arizona.edu](mailto:johnsutter@email.arizona.edu) (J. Sutter), [vadeboncoeur.tyler@mayo.edu](mailto:vadeboncoeur.tyler@mayo.edu) (T. Vadeboncoeur), [asilver@zoll.com](mailto:asilver@zoll.com) (A. Silver), [hucc@email.arizona.edu](mailto:hucc@email.arizona.edu) (C. Hu), [dan@aemrc.arizona.edu](mailto:dan@aemrc.arizona.edu) (D.W. Spaite), [Bentley.Bobrow@azdhs.gov](mailto:Bentley.Bobrow@azdhs.gov) (B. Bobrow).

<sup>1</sup> Department of Emergency Medicine, Mayo Clinic, United States.

<sup>2</sup> ZOLL Medical Corporation, 269 Mill Rd., Chelmsford, MA 01824, United States.

<sup>3</sup> The University of Arizona Health Sciences, 1295 North Martin Avenue, Tucson, AZ 85721-0202, United States.

<sup>4</sup> Arizona Emergency Medicine Research Center, University of Arizona College of Medicine, United States.

<sup>5</sup> Arizona Emergency Medicine Research Center, University of Arizona College of Medicine, Phoenix, AZ, United States.

<https://doi.org/10.1016/j.resuscitation.2019.09.024>

Received 8 April 2019; Received in revised form 12 September 2019; Accepted 16 September 2019

0300-9572/© 2019 Published by Elsevier B.V.

## Introduction

Out-of-hospital cardiac arrest (OHCA) continues to be one of the leading public health concerns in the United States with over 359,800 EMS-assessed cases each year.<sup>1</sup> Several CPR quality measures have been independently associated with improved survival and functional outcomes including chest compression rate, compression depth, compression fraction and pre-shock pause.<sup>1–4</sup> The CPR measure termed chest compression release velocity (CCRV) is defined as the maximum velocity of chest release in the posterior to anterior direction.<sup>5</sup> Kovacs et al. and Indik et al. both demonstrate independent associations between higher CCRV and improved outcomes.<sup>5,6</sup> While CCRV has been associated with improved outcomes, there is wide variability in mean CCRV during OHCA resuscitations.<sup>5</sup> The sources of variability in CCRV in resuscitation are not well studied.

Multiple studies have shown that chest wall compliance increases significantly with repeated compressions, suggesting that compression dynamics are not constant during the performance of CPR.<sup>7,8</sup> Other studies have demonstrated that the amount of force required to achieve equal compression depth varies significantly between patients.<sup>7,9</sup> These findings suggest that patient-related factors influence compression dynamics and therefore may affect the ability to achieve higher rates of CCRV. Our aim was to evaluate how CCRV changes over the course of resuscitation and to assess associations between CCRV and patient sex, age, and weight.

## Methods

### Study design and population

This was a retrospective, observational, cohort study of consecutive adult OHCA patients treated by two EMS agencies that participate in the Save Hearts in Arizona Registry and Education (SHARE) Program and used minimally interrupted cardiac resuscitation as their adult CPR protocol between 10/1/2008 and 12/31/2016.<sup>10–14</sup> Cases were included for analysis if EMS attempted resuscitation and the case duration was at least one minute long. Cases were excluded if the patient had a do-not-resuscitate order, the cardiac arrest occurred in a

healthcare facility, the arrest was witnessed by EMS providers, the arrest was presumed to be non-cardiac in etiology (e.g., respiratory, trauma, or drowning), or if CPR quality data were not available including a minimum of 20 compressions during the first 10 min of the resuscitation (Fig. 1). Primary outcomes were the rate of decline of mean CCRV over time and a comparison of mean CCRV between minutes 0–10, 0–5, and 5–10. Our secondary analysis assessed associations between mean CCRV and patient sex, age, and weight.


### Data collection

Chest compression quality information, including CCRV, was obtained from defibrillators (E Series and X Series with ZOLL CPR-D Padz; ZOLL Medical, Chelmsford, MA) equipped with accelerometer-based compression sensing technology placed according to package instructions which align the CPR sensor with the victim's sternal notch.

Data were collected from the SHARE program OHCA database which has been described previously.<sup>10</sup> As an Arizona Department of Health Services (ADHS)-sponsored public health initiative, Arizona's Attorney General has determined that the SHARE Program is exempt from the requirements of the Health Insurance Portability and Accountability Act (HIPAA), allowing linkage of EMS and hospital data, tracking of OHCA events, and evaluation of efforts to improve resuscitation care. As a public health initiative, the ADHS Human Subjects Review Board and the University of Arizona Institutional Review Board (IRB) have determined that neither the interventions nor their evaluation constitute Human Subjects Research and have approved the publication of de-identified data. The SHARE Program is registered at clinical Trials.gov#NCT01999036.

### Statistical methods

The subject-level mean of CCRV was calculated for the time interval of the first 10 min, then separately for the first five minutes and the second five minutes. The subject-level mean CCRV was compared between males and females and across quartiles of age and weight using the Wilcoxon rank-sum test, and was also correlated with age and weight as continuous variables using the Spearman's rank correlation coefficient (Spearman's rho), with the rho parameter tested against 0. Mean CCRV levels in the first and second 5-min intervals



**Fig. 1 – Exclusion criteria and study population.**

were compared using the robust standard error of the difference obtained from generalized estimating equations (GEE) to account for intra-person correlation. Mixed effect models were applied to compression-level data to study the trend of CCRV over time, with or without adjustment for the depth of each compression. A non-linear function of CCRV over time was estimated non-parametrically using penalized thin plate regression splines through the generalized additive model.<sup>15</sup> A random intercept for each subject was included in the model to account for potential correlation of multiple measurements over time from the same subject. All tests were two-sided with significance levels of  $p < 0.05$ . The statistical environment R was used for all analyses.<sup>16</sup>

## Results


A total of 2535 adult OHCA cases were treated during the study period. After exclusion criteria were applied, 1140 cases remained for analysis (Fig. 1). The analysis focused on the first 10 min of CPR. The 1140 cases received a total of 852,963 compressions with a median of 790 compressions per subject (IQR 689.5, 858.0). The median duration of recorded compressions was 8.70 min (IQR 8.0, 9.1) per subject. The mean CCRV was 343.3 mm/s for the first ten minutes of

CPR. As shown in Fig. 2, the mean CCRV was higher for minutes 0–5 than for minutes 5–10 (mean 347.9 mm/s vs. 339.0 mm/s; decrease in the mean = 8.9 with 95% CI 5.4–12.4 from GEE with  $p < 0.0001$ ). CCRV declined over time with a rapid decrease in the first two minutes of resuscitation, even after adjusting for compression depth (Fig. 2).

Male patients exhibited a greater overall subject-level mean CCRV compared to female patients (Table 1). The mean CCRV was negatively correlated with age (older subjects having lower mean CCRV) and positively correlated with weight (heavier subjects having higher mean CCRV). All  $p$ -values were  $< 0.0001$ . Both sex and age independently were still strongly correlated with CCRV after adjusting for compression depth (both  $p$ -values  $< 0.0001$ ; Table 2). The association between patient weight and CCRV, however, disappeared after controlling for compression depth ( $p > 0.99$ ).

## Discussion

Cardiopulmonary resuscitation science is complex and our knowledge of factors that influence patient outcomes continues to evolve. While the assessment of CPR quality measures such as compression rate, depth, and fraction are traditionally rescuer-centered, patient and situation factors (e.g. duration of resuscitation) likely play a role. One



**Fig. 2 – Chest compression release velocity over time. Mean chest compression release velocity over time, without adjustment for compression depth (left side); mean chest compression release velocity over time adjusted for depth (right side). Overall mean release velocity variability 95% confidence bands are shown by dotted curves.**

**Table 1 – Associations between subject-level mean chest compression release velocity and victim sex, age and weight.**

Variable	Summary <sup>a</sup>	Subject-level Mean CCRV (mm/s) <sup>b</sup>	p-Value <sup>c</sup>
Sex	Female	385 (33.8%)	331.5 (285.3, 385.5)
	Male	755 (66.2%)	344.4 (307.3, 384.6)
Age (year)	65 (54, 76)		–0.226
Age quartile	[18,54]	300 (26.3%)	364.4 (323.1, 407.5)
	[54,65]	286 (25.1%)	341.4 (300.4, 388.6)
	[65,76]	272 (23.9%)	340.3 (298.2, 379.1)
	[76,104]	282 (24.7%)	315.8 (282.5, 362.6)
Weight (kg)	89.5 (73.0, 105.0)		0.239
Weight quartile	[32,73]	234 (20.5%)	316.4 (272.6, 367.7)
	[73,89.5]	188 (16.5%)	342.8 (308.4, 388.2)
	[89.5,105]	213 (18.7%)	354.2 (312.8, 395.1)
	[105,273]	209 (18.3%)	360.2 (317.3, 409.8)
	Unknown	296 (26%)	334 (295.4, 376.8)

<sup>a</sup> Count (percentage) for categorical variables, and median (IQR) for continuous variables.  
<sup>b</sup> Median (IQR) of subject-level CCRV was shown for categorical variables, and estimated Spearman's rank correlation coefficient (rho) was shown for continuous variables.  
<sup>c</sup> For each categorical variable, subject-level mean CCRV was compared across all categories by Wilcoxon rank-sum test; for each continuous variable, it's correlation with subject-level mean CCRV (Spearman's rho) was tested against 0.

**Table 2 – Association of release velocity and victim sex, age and weight after adjusting for compression depth and time in CPR.**

Variable	Levels	Effect (mm/s)	95% CI	p-Value
Compression depth (mm)		5.8	(5.78, 5.81)	<0.0001
Age (year) quartile	[18,54]	referent	–	<0.0001
	[54,65]	–8.19	(–13.98, –2.40)	
	[65,76]	–15.72	(–21.59, –9.86)	
	[76,104]	–20.29	(–26.18, –14.40)	
Weight (kg) quartile	[32,73]	Referent	–	0.9973
	[73,89.5]	–0.84	(–7.83, 6.15)	
	[89.5,105]	–0.33	(–7.09, 6.43)	
	[105,273]	–0.96	(–7.88, 5.96)	
Sex	Unknown	0.06	(–6.15, 6.27)	
	Female	Referent	–	<0.0001
	Male	11.07	(6.57, 15.58)	
Time after CPR started (minute)	Nonparametric function			<0.0001

Note: Linear mixed effect model applied to compression-level data.

of the most recently identified CPR measures is CCRV, which describes the maximum velocity at which the chest wall rises after compression.<sup>4–6</sup> Little is known about how arrest victim characteristics may affect CCRV and how CCRV evolves over the course of resuscitation. To our knowledge, we were the first study to evaluate CCRV over time and to assess any associations between CCRV and the patient factors of sex, age, and weight.

### CCRV over time

Studies to date suggest that higher CCRV is associated with better outcomes; however, the impact that repeated compressions have upon CCRV and how CCRV changes over the course of resuscitation is not documented. In animal models, the generation of intrathoracic vascular pressure during CPR is largely dependent on the initial thoracic anatomy and the change of the initial anatomy as a result of deformities produced during compression.<sup>17</sup> Tomlinson et al. demonstrated that a decreased amount of force is required to achieve an equal level of compression depth over time during CPR, suggesting that repeated compressions soften the chest wall.<sup>7</sup> Similarly, Segal

et al. showed that the compliance of the chest wall increases significantly over time during CPR.<sup>8</sup> Based upon these studies, we hypothesized, and our results suggest, that release velocity declines with repeated compressions. The mean CCRV in our sample was 8.9 mm/s higher for minutes 0–5 than for minutes 5–10 (Fig. 2). Additionally, our results demonstrate a sharp decline early in resuscitation even when controlled for depth (Fig. 2). While the decline in CCRV could be influenced by provider factors such as fatigue and increased incomplete rescuer release during recoil, this finding, in context of prior literature, suggests that loss of chest elasticity may impact CCRV over the course of resuscitation. Potential solutions which may stabilize CCRV over the course of CPR include implementation of resuscitation protocols designed to minimize fatigue through rescuer switching, defibrillator-rescuer feedback and active decompression devices.

### CCRV and patient characteristics

Our study found several patient factors that appear to impact CCRV, including patient sex, age, and weight. Variation in chest wall

compressibility has been demonstrated between male and female patients. Tomlinson et al. described how less force is required to achieve the same level of compression depth in female patients compared to male patients.<sup>7</sup> It was suggested that differences in chest wall anatomy contributed to increased chest wall compliance in female patients. In our study, mean CCRV values were statistically higher in male patients compared to female patients (344.4 mm/s and 331.5 mm/s, respectively  $p=0.01$ ) (Table 1), suggesting that anatomical differences between men and women may also contribute to the ability of the chest wall to achieve higher release velocities.

In addition to patient sex, there was significant variation in mean CCRV between age groups in our results. After adjusting for depth, we observed the CCRV to be 20.29 mm/s slower in the oldest age quartile compared to the youngest age quartile (Table 2). Several studies have demonstrated that increased age is associated with a decline in hyaline cartilage elasticity, a key component in chest wall recoil.<sup>18</sup> Porcine models have shown significant age-related chest morphology differences during compressions with subsequent differences in the intrathoracic pressures that are generated.<sup>17</sup> In humans, chest compliance has been shown to decline with age and rib cage deformation.<sup>19–21</sup> Our results, in context with the studies above, suggest that differences in chest wall physiology between older and younger victims of OHCA could be a factor contributing to the difference in CCRV observed between these groups.

Our study also suggested that CCRV tends to increase at greater patient weights. The heaviest quartile of victims demonstrated 13.8% higher CCRV than the lowest quartile [316.4 mm/s (272.6, 367.7) vs. 360.2 mm/s (317.3, 409.8)]. Increased chest wall resistance and atypical diaphragmatic position are anatomical differences that have been described in obese individuals.<sup>22</sup> As shown in Table 3, statistical differences between sex and age groups were maintained after adjusting for depth. However, the differences among weight quartiles were not significantly different after the adjustment. While it is possible that these anatomical differences in individuals with larger weights impact CCRV, our results suggest that deeper compressions in heavier patients may be impacting CCRV.

### **CCRV: victim vs. rescuer dependent factors**

Current guidelines recommend that the rescuer allow for complete chest wall recoil between compressions during CPR.<sup>23</sup> The ability of the chest wall to recoil is clearly impacted by the presence of rescuer leaning over the patient's chest, impeding full chest expansion.<sup>24</sup> In animal studies, leaning has been shown to decrease coronary perfusion pressure, cardiac output, and myocardial blood flow.<sup>25</sup> As CCRV is the maximum speed of recoil of the chest, CCRV may be impacted by incomplete recoil due to rescuer leaning. While rescuer-fatigue and leaning are described as likely causes of CCRV variance in previous studies, our results are the first to suggest that CCRV may be impacted by patient factors and duration of resuscitation in addition to rescuer-dependent elements.<sup>4–6</sup>

## **Conclusion**

Chest compression release velocity declines over time during prehospital cardiopulmonary resuscitation and is associated with patient sex, age, and weight. Male sex, younger age, and increased weight were associated with higher CCRV. Further research is necessary to elucidate how active decompression devices,

defibrillator-rescuer feedback, appropriate rescuer switching, and other CPR practices can optimize CCRV over the course of resuscitation.

## **Limitations**

The EMS systems studied in this analysis were involved in a dedicated quality assurance system making the applicability of our findings to other EMS systems without similar optimization efforts uncertain. The measurement of CCRV by the accelerometer may be affected by the release of other compressible surfaces (beds or EMS stretchers) beneath the patient. However, we believe the vast majority of the CPR provided by our EMS agencies occurred on a firm surface (backboard or ground) minimizing the potential impact on our CCRV data. Accurate placement of the accelerometer is another potential limitation. Although it cannot be guaranteed that the sensor was always placed in the recommended location, there is no reason to believe that it would be placed differently in one population vs. another. CPR quality data were missing for 286 cases and we excluded these cases from the main analyses. Variations in rescuer performance may have contributed to the observed differences in CCRV, however, the rescuers in this study were equipped with defibrillators with real-time feedback, minimizing inter-rescuer variability. Lastly, our study is observational and, as such, causality cannot be inferred from our study's findings.

## **Conflict of interest**

Annemarie Silver, PhD, is an employee of ZOLL Medical Corporation. The other authors certify that they have NO affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

## **REFERENCES**

1. Mozaffarian D, Benjamin EJ, Go AS, et al. Executive summary: heart disease and stroke statistics—2016 update: a report from the American heart association. *Circulation* 2016;133:447–54, doi:<http://dx.doi.org/10.1161/CIR.0000000000000366>.
2. Vaillancourt C, Everson-Stewart S, Christenson J, et al. The impact of increased chest compression fraction on return of spontaneous circulation for out-of-hospital cardiac arrest patients not in ventricular fibrillation. *Resuscitation* 2011;82:1501–7, doi:<http://dx.doi.org/10.1016/j.resuscitation.2011.07.011>.
3. Vadeboncoeur T, Stolz U, Panchal A, et al. Chest compression depth and survival in out-of-hospital cardiac arrest. *Resuscitation* 2014;85:182–8, doi:<http://dx.doi.org/10.1016/j.resuscitation.2013.10.002>.
4. Cheskes S, Common MR, Byers AP, Zhan C, Silver A, Morrison LJ. The association between chest compression release velocity and outcomes from out-of-hospital cardiac arrest. *Resuscitation* 2015;86:38–43, doi:<http://dx.doi.org/10.1016/j.resuscitation.2014.10.020>.
5. Kovacs A, Vadeboncoeur TF, Stolz U, et al. Chest compression release velocity: association with survival and favorable neurologic outcome after out-of-hospital cardiac arrest. *Resuscitation*

- 2015;92:107–14, doi:<http://dx.doi.org/10.1016/j.resuscitation.2015.04.026>.
6. Indik JH, Conover Z, McGovern M, et al. Amplitude-spectral area and chest compression release velocity independently predict hospital discharge and good neurological outcome in ventricular fibrillation out-of-hospital cardiac arrest. *Resuscitation* 2015;92:122–8, doi:<http://dx.doi.org/10.1016/j.resuscitation.2015.05.002>.
  7. Tomlinson AE, Nysaether J, Kramer-Johansen J, Steen PA, Dorph E. Compression force–depth relationship during out-of-hospital cardiopulmonary resuscitation. *Resuscitation* 2007;72:364–70, doi:<http://dx.doi.org/10.1016/j.resuscitation.2006.07.017>.
  8. Segal N, Robinson AE, Berger PS, et al. Chest compliance is altered by static compression and decompression as revealed by changes in anteroposterior chest height during CPR using the ResQPUMP in a human cadaver model. *Resuscitation* 2017;116:56–9, doi:<http://dx.doi.org/10.1016/j.resuscitation.2017.04.032>.
  9. Gruben KG, Guerci A, Halperin HR, Popel AS, Tsitlik J. Sternal force-displacement relationship during cardiopulmonary resuscitation. *J Biomech Eng* 1993;115:195–201.
  10. Vadeboncoeur T, Bobrow BJ, Clark L, et al. The save hearts in arizona registry and education (SHARE) program: who is performing CPR and where are they doing it? *Resuscitation* 2007;75:68–75.
  11. Bobrow BJ, Clark LL, Ewy GA, et al. Minimally interrupted cardiac resuscitation by emergency medical services for out-of-hospital cardiac arrest. *JAMA* 2008;299:1158–65.
  12. Ewy GA. Cardiocerebral resuscitation. *Circulation* 2005;111:2134–42, doi:<http://dx.doi.org/10.1161/01.CIR.0000162503.57657.FA>.
  13. Kern KB, Valenzuela TD, Clark LL, et al. An alternative approach to advancing resuscitation science. *Resuscitation* 2005;64:261–8, doi:<http://dx.doi.org/10.1016/j.resuscitation.2004.08.009>.
  14. Kellum MJ, Kennedy KW, Ewy GA. Cardiocerebral resuscitation improves survival of patients with out-of-hospital cardiac arrest. *Am J Med* 2006;119:335–40, doi:<http://dx.doi.org/10.1016/j.amjmed.2005.11.014>.
  15. Wood SN. Generalized additive models: an introduction with R. Chapman and Hall/CRC; 2017.
  16. R Core Team. A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2015 2015.
  17. Dean JM, Koehler RC, Schleien CL, et al. Age-related changes in chest geometry during cardiopulmonary resuscitation. *J Appl Physiol* (1985) 1987;62:2212–9, doi:<http://dx.doi.org/10.1152/jappl.1987.62.6.2212>.
  18. Lau A, Oyen ML, Kent RW, Murakami D, Torigaki T. Indentation stiffness of aging human costal cartilage. *Acta Biomater* 2008;4:97–103, doi:<http://dx.doi.org/10.1016/j.actbio.2007.06.008>.
  19. Mittman C, Edelman NH, Norris AH, Shock NW. Relationship between chest wall and pulmonary compliance and age. *J Appl Physiol* 1965;20:1211–6.
  20. Sharma G, Goodwin J. Effect of aging on respiratory system physiology and immunology. *Clin Interv Aging* 2006;1:253–60.
  21. Kent R, Patrie J. Chest deflection tolerance to blunt anterior loading is sensitive to age but not load distribution. *Forensic Sci Int* 2005;149:121–8.
  22. Brunette DD. Resuscitation of the morbidly obese patient. *Am J Emerg Med* 2004;22:40–7.
  23. Kleinman ME, Brennan EE, Goldberger ZD, et al. Part 5: adult basic life support and cardiopulmonary resuscitation quality: 2015 American heart association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2015;132:S414–35, doi:<http://dx.doi.org/10.1161/CIR.0000000000000259>.
  24. Aufderheide TP, Pirralo RG, Yannopoulos D, et al. Incomplete chest wall decompression: a clinical evaluation of CPR performance by EMS personnel and assessment of alternative manual chest compression–decompression techniques. *Resuscitation* 2005;64:353–62, doi:<http://dx.doi.org/10.1016/j.resuscitation.2004.10.007>.
  25. Niles DE, Sutton RM, Nadkarni VM, et al. Prevalence and hemodynamic effects of leaning during CPR. *Resuscitation* 2011;82: S23–6.