



ELSEVIER

Contents lists available at ScienceDirect

The Foot

journal homepage: www.elsevier.com/locate/foot

Original Article

Weight-bearing in ankle fractures: An audit of UK practice

BONE Collaborative¹

British Orthopaedic Network Environment — The British Orthopaedic Trainees Association, British Orthopaedic Association, London, UK



ARTICLE INFO

Keywords:

Fractures
Ankle
Weight-bearing
Early
Delayed
NICE

ABSTRACT

Introduction: The purpose of this national study was to audit the weight-bearing practice of orthopaedic services in the National Health Service (NHS) in the treatment of operatively and non-operatively treated ankle fractures. **Methods:** A multicentre prospective two-week audit of all adult ankle fractures was conducted between July 3rd 2017 and July 17th 2017. Fractures were classified using the AO/OTA classification. Fractures fixed with syndesmosis screws or unstable fractures (> 1 malleolus fractured or talar shift present) treated conservatively were excluded. No outcome data were collected. In line with NICE (The National Institute for Health and Care Excellence) criteria, “early” weight-bearing was defined as unrestricted weight-bearing on the affected leg within 3 weeks of injury or surgery and “delayed” weight-bearing as unrestricted weight-bearing permitted after 3 weeks. **Results:** 251 collaborators from 81 NHS hospitals collected data: 531 patients were managed non-operatively and 276 operatively. The mean age was 52.6 years and 50.5 respectively. 81% of non-operatively managed patients were instructed for early weight-bearing as recommended by NICE. In contrast, only 21% of operatively managed patients were instructed for early weight-bearing. **Discussion:** The majority of patients with uni-malleolar ankle fractures which are managed non-operatively are treated in accordance with NICE guidance. There is notable variability amongst and within NHS hospitals in the weight-bearing instructions given to patients with operatively managed ankle fractures. **Conclusion:** This study demonstrates community equipoise and suggests that the randomized study to determine the most effective strategy for postoperative weight-bearing in ankle fractures described in the NICE research recommendation is feasible.

1. Introduction

Each day approximately 200 patients sustain an ankle fracture in the United Kingdom (UK) [1]. The National Institute for Health and Care Excellence (NICE) guideline (NG38) [2] recommends that patients with uni-malleolar ankle fractures that are managed non-operatively should be advised to weight-bear as tolerated immediately or return within 2 weeks for further assessment if stability is uncertain. The most effective strategy for weight-bearing for patients with operatively managed ankle fractures remains uncertain and is one of the NICE priority recommendations for research in trauma. Current national practice is unknown. The purpose of this multicentre, prospective audit is to determine the weight-bearing practice for both conservative and operatively managed ankle fractures within orthopaedic services in the setting of the UK National Health Service (NHS).

Methods

A multicentre, prospective two-week audit of all adult ankle fractures was conducted between July 3rd 2017 and July 17th 2017. Collaborators were recruited and coordinated via the online British Orthopaedic Network Environment (BONE) on behalf of the **British Orthopaedic Trainees**

Association [3]. Research and Ethics Committee approval was not required under assessment of the Health Research Authority decision tool (<http://www.hra-decisiontools.org.uk/research/>). All collaborators obtained local approval from their respective institutions.

In line with NICE definitions, “early” weight-bearing was defined as unrestricted weight-bearing on the affected leg within 3 weeks of injury or surgery and “delayed” weight-bearing as unrestricted weight-bearing permitted after 3 weeks [4]. Restricted weight-bearing was defined as partial weight-bearing, touch weight-bearing, non-weight-bearing or protected weight-bearing. Unrestricted weight-bearing was defined as full weight-bearing or weight-bearing as tolerated [4]. These definitions applied whether or not the patient was in a splint or form of immobilisation.

Fractures were classified by AO/OTA classification (Fig. 1) [5] Both non-operatively and operatively managed cases were included. Within the operative group, only fractures treated with open reduction and internal fixation (ORIF) were included. Fractures fixed with syndesmosis screws or unstable ankle fractures (> 1 malleolus fractured or talar shift present) treated conservatively were excluded. Participants exited the study when they were allowed to weight-bear without restriction or a future date to do so was indicated in the clinical record.

Data were collected on patient demographics, method of treatment, time to surgery, time to unrestricted weight-bearing, form of immobilisation, time

¹ See Appendix A.




Fig 1. AO/OTA 44 classification.

and method of repeat radiographic assessment if stability was uncertain. Time to unrestricted weight-bearing was recorded both from date of injury and from date from first review by orthopedic services. Patient factors including documented diabetes, osteoporosis, peripheral neuropathy or mobility concerns (e.g. previously requiring mobility aids or assistance, or sustaining multiple injuries limiting mobility) were recorded. In addition, injury factors including open fracture, gross comminution or soft tissue compromise documented on the operation note were recorded. No outcome data were collected.

Statistical analysis

Count data were summarised as absolute numbers and proportions; continuous data as means and ranges. Data were analysed using IBM SPSS Statistics v.21 (IBM, Armonk, New York). No inferential statistical analyses were performed.

Results

251 collaborators from 81 NHS hospitals provided data from 996 adults with 996 ankle fractures before application of the exclusion criteria. A participant flow diagram is presented at Fig. 2.

Non-operatively treated fractures

Data from 531 patients with a mean age of 52.6 [16–97]. The mean time restricted weight-bearing from date of injury was 12.4 days [10–63] and from date of first assessment by orthopaedic services was 8.4 days [10–63]. 103 (19.4%) patients were advised delayed weight-bearing and 428 (80.6%) patients were allowed to weight-bear without restriction less than 3 weeks from injury. Table 1 demonstrates weight-bearing practice by fracture classification. Fig. 3 demonstrates the weight-bearing practice across participating hospitals. The proportion of patients permitted to weight-bear early with diabetes, osteoporosis,




Fig. 2. Participant flow diagram.

Table 1
Weight-bearing by fracture classification for patients managed conservatively.

Time till unrestricted weight-bearing	44A1	44B1	Isolated posterior malleolus	Isolated medial malleolus	Total
< 3 weeks	225 (94%)	181 (69%)	4 (80%)	18 (72%)	428 (81%)
> 3 weeks	15 (6%)	80 (31%)	1 (20%)	7 (28%)	103 (19%)

peripheral neuropathy or mobility concerns was 13/17 (77%), 21/26 (81%), 9/9 (100%) and 30/41 (73%) respectively. For those allowed to weight-bear early, the method of immobilisation was a boot in 284 (66%) cases, plaster in 72 (17%), nothing in 39 (9%), aircast stirrup in 22 (5%), tubigrip in 6 (1%), and other in 5 (1%).

311 (59%) patients had a check radiograph within 2 weeks to confirm stability, of which 131 (42%) were weight-bearing radiographs. 80 (61%) of

those that had a weight-bearing radiograph had them whilst in their respective form of immobilisation; 51 (39%) had no immobilisation whilst having their radiograph taken.

Operatively treated fractures

Data from 276 patients. The mean age was 50.2 [16–92], and the mean




Fig. 3. Weight-bearing by hospital for patients managed conservatively.

Table 2
Weight-bearing by fracture classification for operatively managed patients.

Time till unrestricted weight-bearing	44A1	44A2	44A3	44B1	44B2	44B3	44C1	44C2	44C3	Isolated medial malleolus	Posterior & medial malleolus	Total
< 3 weeks	1 (14%)	6 (33%)	1 (20%)	10 (24%)	19 (23%)	14 (22%)	4 (16%)	2 (12%)	1 (14%)	1 (11%)	0 (0%)	59 (21%)
> 3 weeks	6 (86%)	12 (67%)	4 (80%)	31 (76%)	63 (77%)	50 (78%)	21 (84%)	15 (88%)	6 (86%)	8 (89%)	1 (100%)	217 (79%)

days from injury to definitive fixation was 5.1 [1–24]. The mean time restricted weight-bearing postoperatively was 35.9 days [1–92]. 217 (79%) patients were advised delayed weight-bearing, with 59 (21%) allowed to weight-bear without restriction less than 3 weeks postoperatively. Table 2 demonstrates weight-bearing practice for post-operative patients by fracture classification. Fig. 4 demonstrates the weight-bearing practice across participating hospitals. Table 3 demonstrates how patient comorbidity or fracture factors influenced weight-bearing decisions. For those allowed to weight-bear early, 35 (59%) did so in a boot, 20 (34%) in plaster, and 4 (7%) other.

Discussion

The clinical and economic impacts of delayed weight-bearing are not fully understood [2]. NICE identified no randomised controlled trials (RCTs) evaluating weight-bearing practice in conservatively managed adult ankle fractures. In the absence of robust evidence, the recommendation was that patients with non-operatively managed uni-malleolar fractures should be allowed to weight-bear without restriction immediately [2].

A variety of factors may influence a clinician’s weight-bearing decision when treating fractures conservatively: clinical examination, medial tenderness, gravity stress and supination-external rotation stress views have all been used [6–8]. According to Egol et al. [6], medial tenderness and swelling have a low predictive correlation with positive stress tests, with sensitivity of 56% and 55%, and specificity of 80% and 71% respectively. Positive stress views do not predict deltoid ligament insufficiency, with Koval et al. [7] finding only 2 of 21 patients with positive stress tests having complete deltoid rupture on MRI scanning. Gill et al. [8] found that gravity stress views are equivalent to stress tests, but Weber et al. [9] proposed that these tests overestimate the need for surgery. They used

weight-bearing radiographs to determine stability, producing excellent clinical results with a low risk of complications. Dawe et al. [10] replicated these results, finding that a shift in protocol from using stress tests, to using weight-bearing radiographs greatly reduced the proportion of patients undergoing surgery and the rate of complications overall. Using the same method for determining stability, Hastie et al. [11] found that bracing and encouraging an early range of motion has a low risk of complications and good clinical results. Our study suggests that there is a greater inclination for clinicians to allow early weight-bearing for conservatively managed uni-malleolar fractures, but there is still variety in thinking for the best form of bracing in the early setting. Furthermore, whilst Hoshino et al. [12] demonstrated that a weight-bearing radiograph without immobilisation after 1 week is a robust method of predicting the success of conservative treatment for fractures of uncertain stability, our study has shown there is no uniformity for radiographically assessing stability within clinical practice at present.

A 2012 Cochrane review [13] examined rehabilitation for ankle fractures and NICE [2] has set a priority to investigate postoperative weight-bearing; they identified 6 and 8 RCTs respectively. Only Ahl et al. [14], Ahl et al. [15] and Finsen et al. [16] had a common form of immobilisation (plaster cast) in each group. In these studies, no difference in adverse events were found; early weight-bearing favoured improved dorsiflexion at 6 months but no difference in activity limitation at 1 year. Ahl et al. [17] compared a weight-bearing orthosis versus a non-weight-bearing dorsal splint and found more minor adverse events (e.g. skin irritation, superficial infection) in the early weight-bearing group. The trend of early functional improvements but an increase in minor wound healing adverse events was mirrored in the RCTs that focused on early-exercise postoperatively [13]. This would seem to reflect the findings of our study suggesting that 79% of clinicians may be anxious about the risk of adverse events compared to the




Fig. 4. Weight-bearing by hospital for patients managed operatively.

Table 3

Patients permitted unrestricted weight-bearing < 3 weeks post-operatively by comorbidity/fracture factors.

Diabetes	Osteoporosis	Peripheral neuropathy	Mobility concerns	Open fracture	Soft tissue compromise	Gross comminution	Total
3/10 (30%)	5/14 (36%)	1/8 (13%)	6/20 (30%)	1/11 (9%)	1/14 (7%)	4/29 (14%)	59/279 (21%)

minimal clinical gains of early weight-bearing. Overall NICE [2] and the Cochrane review [13] concluded that the studies were underpowered and had high risk of bias, serious imprecision and heterogeneity in outcomes measures, recommending further high-quality studies on the topic.

Kheir et al. [18] found the treatment of unstable ankle fractures required an average length of hospital stay of nine days and an associated mean cost of £4491 per patient. In patients over sixty, Keene et al. [19] found this increased to a mean NHS cost of £6648 and mean societal cost of £7684 per patient. No studies have identified the economic implications of prolonged restricted weight-bearing, but it is likely to have direct and indirect effects. It can diminish independence and, particularly in the elderly, comorbid patient, may lead to prolonged hospital or institutional care [20,21]. Younger patients can often mobilise with walking aids but it can affect their return to work, with Simanski et al. [21] and Gul et al. [22] suggesting a trend towards earlier return to work with early weight-bearing post-operative regimes.

Recent qualitative studies have shed light on the psychosocial effects of ankle fractures with patients having to “endure” a period of non-weight bearing. While some patients adjust, many find the experience “terrifying” and put their life on hold, even avoiding showering or bathing altogether until allowed to weight-bear [23,19].

This study has a number of limitations. It reports only the decisions of the treating clinicians; it did not collect data on patient outcomes, complications or cross over from the conservative into the operative group. It is not known if patients adhered to their weight-bearing instructions or if they were allowed to exercise their ankle. We used the AO/OTA classification as it has better intraobserver and interobserver reliability than the Lauge-Hansen classification [24]. However, its reliability and reproducibility was not independently assessed throughout all 81 participating centres. While patient comorbidity and injury factors were analysed, information on other factors such as whether some components of the fracture were left unfixed (e.g. posterior malleolus) was not, and could have influenced weight-bearing decisions.

This study is the first to prospectively look at weight-bearing decisions. A previous American survey of clinicians views towards postoperative ankle fractures found a preferred average time non-weight-bearing of 4.9 weeks in young healthy patients, increasing to 7.6 weeks in older, comorbid patients with complex fractures [25]. We wanted to examine what clinicians did in practice, faced with real patients and audit this against NICE guidelines [2]. A developing trainee collaborative provided the infrastructure to enable this

[3]. Nationally, this study demonstrates 81% adherence to NICE guidelines for non-operatively treated uni-malleolar ankle fractures though found considerable variability in the early radiographic reassessment. It establishes a national baseline with 21% of clinicians allowing early weight-bearing postoperatively. We found no evidence that fracture classification or patient factors such as diabetes or osteoporosis deterred clinicians from weight-bearing patients early. Fracture specific factors, particularly soft tissue compromise, were associated with a greater reluctance to allow early weight-bearing. Though numbers were too small to infer causation, these insights are useful if considering when and which patients' clinicians will be willing to randomise in future RCTs.

Conclusion

This study demonstrates the willingness of clinicians to allow early weight-bearing in conservatively managed AO44A1 fractures, the relative uncertainty in AO44B1 fractures and the reluctance to weight-bear early in operatively treated adult malleolar ankle fractures.

It is unclear if early or delayed weight-bearing influences late functional outcome. The economic and social cost may emerge as the key argument in the weight-bearing debate.

Summary

- The standard of care for weight-bearing after ankle fracture was previously unknown.
- 81% of non-operatively managed patients with uni-malleolar fractures are advised early weight-bearing.
- 21% of operatively managed patients are advised early weight-bearing.
- There is community equipoise and a randomized study examining weight-bearing in ankle fractures is feasible.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Appendix A

Collaborators

Hospital	First name	Middle initial	Surname	Email
Addenbrooke's Hospital, Cambridge, UK	James	R	Gill	james.ritchie.gill@gmail.com
Addenbrooke's Hospital, Cambridge, UK	Emilie		Lostis	el405@cam.ac.uk
Alexandra Hospital, Redditch, UK	Sabri	S	Bleibleh	sabri_bleibleh@yahoo.com
Arrowe Park Hospital, Wirral, UK	Michael	S	Hennessy	mhennessy@nhs.net
Arrowe Park Hospital, Wirral, UK	Femi		Thondickal	femijoy@nhs.net
Barnsley Hospital, Barnsley, UK	Richard	J	Gadd	richgadd@doctors.net.uk
Barnsley Hospital, Barnsley, UK	Rachael		Gruppung	rachael.gruppung@btinternet.com
Blackburn Royal Infirmary, Blackburn, UK	Albert		Chikate	albert.chikate@elht.nhs.uk
Broomfield Hospital, Chelmsford, UK	Suan		Khor	suan.khor@nhs.net
Broomfield Hospital, Chelmsford, UK	Sangam		Malani	sangam.malani@meht.nhs.uk
Broomfield Hospital, Chelmsford, UK	William		Nabulyato	willnabulyato@googlemail.com
Broomfield Hospital, Chelmsford, UK	Praveen		Rai	praveen.raimeht.nhs.uk
Broomfield Hospital, Chelmsford, UK	Kalpesh		Vaghela	kalpeshrvaghela@gmail.com
Cheltenham General Hospital, Cheltenham, UK	Shashank		Pillai	shan90.pillai@gmail.com
Colchester General Hospital, Colchester, UK	Dinnish		Baskaran	dinnish.baskaran04@alumni.imperial.ac.uk
Colchester General Hospital, Colchester, UK	Raviraj		Kugapiriyam	Raviraj.Kugapiriyam@colchesterhospital.nhs.uk

Colchester General Hospital, Colchester, UK			
Colchester General Hospital, Colchester, UK			
Conquest Hospital, Hastings, UK			
Conquest Hospital, Hastings, UK			
Conquest Hospital, Hastings, UK			
Conquest Hospital, Hastings, UK			
Croydon University Hospital, Croydon, UK			
Croydon University Hospital, Croydon, UK			
Croydon University Hospital, Croydon, UK			
Croydon University Hospital, Croydon, UK			
Croydon University Hospital, Croydon, UK			
Cumberland Infirmary, Carlisle, UK			
Cumberland Infirmary, Carlisle, UK			
Darent Valley Hospital, Dartford, UK			
Darent Valley Hospital, Dartford, UK			
Darlington Memorial Hospital, Darlington, UK			
Darlington Memorial Hospital, Darlington, UK			
Doncaster Royal Infirmary Doncaster, UK			
Dorset County Hospital, Dorchester, UK			
Forth Valley Royal Hospital, Larbert, UK			
Forth Valley Royal Hospital, Larbert, UK			
Forth Valley Royal Hospital, Larbert, UK			
Frimley Park Hospital, Frimley, UK			
Frimley Park Hospital, Frimley, UK			
Frimley Park Hospital, Frimley, UK			
Glasgow Royal Infirmary, Glasgow, UK			
Glasgow Royal Infirmary, Glasgow, UK			
Glasgow Royal Infirmary, Glasgow, UK			
Harrogate District Hospital, Harrogate, UK			
Hillingdon Hospital, Uxbridge, UK			
Hillingdon Hospital, Uxbridge, UK			
Hillingdon Hospital, Uxbridge, UK			
Hillingdon Hospital, Uxbridge, UK			
Hillingdon Hospital, Uxbridge, UK			
Hillingdon Hospital, Uxbridge, UK			
Hillingdon Hospital, Uxbridge, UK			
Hillingdon Hospital, Uxbridge, UK			
Horton General Hospital, Banbury, UK			
Horton General Hospital, Banbury, UK			
Horton General Hospital, Banbury, UK			
Horton General Hospital, Banbury, UK			
Horton General Hospital, Banbury, UK			
Ipswich Hospital, Ipswich, UK			
Ipswich Hospital, Ipswich, UK			
Ipswich Hospital, Ipswich, UK			
James Cook University Hospital, Middlesborough, UK			
James Cook University Hospital, Middlesborough, UK			
James Cook University Hospital, Middlesborough, UK			
John Radcliffe Hospital, Oxford, UK			
John Radcliffe Hospital, Oxford, UK			
John Radcliffe Hospital, Oxford, UK			
John Radcliffe Hospital, Oxford, UK			
Leeds General Infirmary, Leeds, UK			
Leeds General Infirmary, Leeds, UK			
Leicester Royal Infirmary, Leicester, UK			
Leicester Royal Infirmary, Leicester, UK			
Leicester Royal Infirmary, Leicester, UK			
Leicester Royal Infirmary, Leicester, UK			
Leighton Hospital, Crewe, UK			
Leighton Hospital, Crewe, UK			
Leighton Hospital, Crewe, UK			
Lincoln County Hospital, Lincoln, UK			
Lincoln County Hospital, Lincoln, UK			
Lincoln County Hospital, Lincoln, UK			
Lincoln County Hospital, Lincoln, UK			
Lincoln County Hospital, Lincoln, UK			
Milton Keynes General Hospital, Milton Keynes, UK			
Milton Keynes General Hospital, Milton Keynes, UK			
Morrison Hospital, Swansea, UK			
Morrison Hospital, Swansea, UK			
Morrison Hospital, Swansea, UK			
New Cross Hospital, Wolverhampton, UK			
Newham General Hospital, London, UK			
Newham General Hospital, London, UK			
Newham General Hospital, London, UK			
North Hampshire Hospital, Basingstoke, UK			
North Hampshire Hospital, Basingstoke, UK			
North Middlesex Hospital, London, UK			
North Middlesex Hospital, London, UK			
North Middlesex Hospital, London, UK			
North Middlesex Hospital, London, UK			
Northern General Hospital, Sheffield, UK			
Stanley			
Jeremy			
Fiona			
Tom			
Barry			
Roosbeh			
Thomas			
Piyush			
Ravikanth			
Reza			
Luke			
Ashley			
Lucy			
Syed			
Poonam			
Benjamin			
Patrick			
Andrew			
Zara			
Thomas			
Thomas	H		
Christopher			
Purnajyoti			
Vinoth			
Victor			
James			
Andraay			
David			
Mark			
Ahsan			
Khalid			
Emma			
Anna			
Edmund			
Sheraz			
Stephen			
Winster			
Shahid			
Gavin			
Mariam			
Ady			
Alexander			
Albert			
Alina			
Matt			
Antonia			
Henry	A		
Xavier	L		
Navraj	S		
Mustafa			
Vishal			
Ross			
Roberto	JR		
Thomas			
Sheweidin			
Ganapathy			
Bakir			
Marieta			
Purwa			
Mallikarjun			
Latif			
Rizwan			
Singapura			
Yiteng			
Ghazal			
John			
Simon			
Julia			
Tariq			
Jasprit			
Richard	L		
Steve			
Lee			
Hayleigh			
Sushmith			
Barbara			
Ravikiran			
Abhinav			
Sunil			
Paul			
Masunda			
Jeremy.Parker@colchesterhospital.nhs.uk			
Bintcliffe			
Murphy			
Rose			
Shafafy			
Holme			
Mahapatra			
Mallina			
Mansouri			
Render			
Scrimshire			
Walker			
Ahmed			
Kandel			
Drake			
Williams			
Hannah			
Hayat			
Aitken			
Carter			
Crome			
Banerjee			
Ravi			
West			
Bennett			
Leung			
Skipsey			
Blomfield			
Akhtar			
Al-Dadah			
Davies			
Kropelnicki			
Leong			
Malik			
Ng Man Sun			
D'Souza			
Mehmood			
Reynolds			
Sattar			
Abdelhaq			
Durst			
Ngu			
Budacan			
Gray			
Myatt			
Claireaux			
Griffin			
Nagra			
Rashid			
Borse			
Muir			
Alho			
Ankers			
Aziz			
Perianayagam			
Al-Dulaimy			
Franklin			
Wilson			
Chandrappa			
Mubasher			
Shahid			
Shashidhara			
Xu			
Hodhody			
Jeffery			
Humphry			
Street			
Yasin			
Kaur			
Donovan			
Kahane			
Parker			
Byrne			
Gowda			
Frank			
Shenoy			
Singh			
Trakru			
Brewer			

Northern General Hospital, Sheffield, UK	Mike	Petrie	mike.petrie@doctors.org.uk
Northwick Park Hospital, Harrow, UK	Ghias	Bhattee	g.bhattee@nhs.net
Northwick Park Hospital, Harrow, UK	Michael	Rafferty	michael.rafferty@nhs.net
Northwick Park Hospital, Harrow, UK	Quen	Tang	quentang@gmail.com
Peterborough City Hospital, Peterborough, UK	Mohamed	Elbashir	elbashirmb@hotmail.com
Peterborough City Hospital, Peterborough, UK	Vladislav	Kutuzov	vlad.kutuzov@yahoo.co.uk
Peterborough City Hospital, Peterborough, UK	Kate	Spacey	kate.spacey@gmail.com
Princess of Wales Hospital, Bridgend, UK	Darren	Ebreo	darren.ebreo@gmail.com
Princess of Wales Hospital, Bridgend, UK	Shreekant	Gupta	shreekgpt@aol.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Usman	Ahmed	ozman1984@gmail.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Mohammed	Al Mustafa	m.almustafa@nhs.net
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Seyed	Ali	seyed.ali@uhb.nhs.uk
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Govind Singh	Chauhan	gov.chauhan@gmail.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Saad	Elashry	saad_elashry@yahoo.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Naeil	Lotfi	naeilotfi@nhs.net
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Shahbaz	Malik	shahb.malik@gmail.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Alastair	Marsh	alastair.marsh@uhb.nhs.uk
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Iosif	Pagkalos	iosif.pagkalos@nhs.net
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Raghunaathan	Rangan	drrraghu@gmail.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Parag	Raval	praval@doctors.org.uk
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Islam	Sarhan	isarhan30@gmail.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Yousufuddin	Shaik	skyoufufs@yahoo.com
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Rajkumar	Thangaraj	rajt129@yahoo.co.uk
Queen Elizabeth Hospital, Birmingham, UK (Birmingham Orthopaedic Network)	Feiran	Wu	feiran.wu@gmail.com
Queen Elizabeth University Hospital, Glasgow, UK	Zainab	Kazmi	zainab.kazmi@nhs.net
Queen Elizabeth University Hospital, Glasgow, UK	Chia Yew	Kong	chiayew.kong@nhs.net
Queen's Hospital, Romford, UK	Moritz	Lebe	moritz.lebe@bhrhospitals.nhs.uk
Queen's Hospital, Romford, UK	Surjit	Lidder	surjitlidder@me.com
Queen's Hospital, Romford, UK	Sean	Masterson	seanmast@gmail.com
Queen's Hospital, Romford, UK	Farzad	Shabani	farzad.shabani@bhrhospitals.nhs.uk
Queens Medical Centre, Nottingham, UK	Adeel	Ikram	adeelikram0@gmail.com
Queens Medical Centre, Nottingham, UK	Ben	Marson	ben.marson@nottingham.ac.uk
Queens Medical Centre, Nottingham, UK	Paul	Matthews	Paul.Matthews1@nottingham.ac.uk
Queens Medical Centre, Nottingham, UK	Ganesh	Prasad	ganesh.prasad@nhs.net
Royal Albert Edward Infirmary, Wigan	James	Davenport	James.Davenport@wwl.nhs.uk
Royal Albert Edward Infirmary, Wigan	Mike	Karski	Mike.Karski@wwl.nhs.uk
Royal Albert Edward Infirmary, Wigan	Kevin	Syam	vinky_scorpio@yahoo.com
Royal Albert Edward Infirmary, Wigan	Nithin	Unnikrishnan	nithinunni@gmail.com
Royal Alexandra Hospital, Paisley, UK	Rashid	Abu-Rajab	Rashid.Abu-Rajab@ggc.scot.nhs.uk
Royal Alexandra Hospital, Paisley, UK	Ignatius	Liew	ignatiusliew@gmail.com
Royal Alexandra Hospital, Paisley, UK	Niak	Puei Koh	kpnoweretogo@gmail.com
Royal Blackburn Teaching Hospital, Blackburn, UK	Jim	Barrie	jim.barrie@elht.nhs.uk
Royal Cornwall Hospital, Truro, UK	Alex	Goubran	alexgoubran@nhs.net
Royal Cornwall Hospital, Truro, UK	Lucy	Maling	lucy.maling@nhs.net
Royal Derby Hospital, Derby, UK	Edward	Karam	edward.karam@doctors.org.uk
Royal Derby Hospital, Derby, UK	Richard	Limb	r.limb@icloud.com
Royal Derby Hospital, Derby, UK	George	Matheron	georgematheron@gmail.com
Royal Devon & Exeter Hospital, Exeter, UK	Fady	Awad	fady.awad@nhs.net
Royal Devon & Exeter Hospital, Exeter, UK	Kimberely	Shuttlewood	kshuttlewood@doctors.org.uk
Royal Gwent Hospital, Newport, UK	Jack	Pearce	jack.pearce@nhs.net
Royal Gwent Hospital, Newport, UK	Kar	Teoh	karhao@gmail.com
Royal Hampshire County Hospital, Winchester, UK	Jo	Higgins	jo_higgins@hotmail.com
Royal Hampshire County Hospital, Winchester, UK	Togay	Koc	togay.koc@doctors.org.uk
Royal Hampshire County Hospital, Winchester, UK	John	Mcfall	mcfalljohn@hotmail.com
Royal Infirmary of Edinburgh, Edinburgh, UK	George	Holland	gdbholland@gmail.com
Royal Infirmary of Edinburgh, Edinburgh, UK	Julian	Maempel	julian.maempel@nhs.net
Royal Infirmary of Edinburgh, Edinburgh, UK	Jamie	Nicholson	jami nicholson@nhs.net
Royal Infirmary of Edinburgh, Edinburgh, UK	Neil	Wickramasinghe	neilrwickcs@gmail.com
Royal Lancaster Infirmary, Lancaster, UK	Andrew	Berg	a.j.berg@doctors.org.uk
Royal Lancaster Infirmary, Lancaster, UK	David	Knowles	david.knowles@mbht.nhs.uk
Royal London Hospital, London, UK	Naem	Dowlut	m.dowlut@smd12.qmul.ac.uk
Royal London Hospital, London, UK	Mehvish	Hayat	mawish.hayat@gmail.com
Royal Preston Hospital, Preston, UK	James	Bourne	drjtbourne@gmail.com
Royal Preston Hospital, Preston, UK	Leanne	Dupley	leannedupley@doctors.org.uk
Royal Preston Hospital, Preston, UK	Min Soo	Kim	min.kim-2@student.manchester.ac.uk
Royal Preston Hospital, Preston, UK	Mark	Poustie	markpoustie1@gmail.com
Royal Stoke University Hospital, Stoke-on-Trent, UK	Laura	Bowen	lura@hotmail.co.uk
Royal Stoke University Hospital, Stoke-on-Trent, UK	Ross	Fawdington	fawdy@doctors.org.uk
Royal Stoke University Hospital, Stoke-on-Trent, UK	Lawrence	Moulton	lsmoulton@gmail.com
Royal Stoke University Hospital, Stoke-on-Trent, UK	Helen	Ribee	helenribee@googlemail.com
Royal Stoke University Hospital, Stoke-on-Trent, UK	Charlotte	Tunstall	lottietunstall@doctors.org.uk
Royal Surrey County Hospital, Guildford, UK	Robert	Boyd	robertboyd@doctors.org.uk
Royal Surrey County Hospital, Guildford, UK	Jonathan	Quayle	jonathan.quayle@gmail.com
Royal Sussex County Hospital, Brighton, UK	Felicity	Auld	felicity.auld@bsuh.nhs.uk
Royal Sussex County Hospital, Brighton, UK	Simon	Bellringer	Drbellringer@me.com
Royal Sussex County Hospital, Brighton, UK	Lucy	Cassidy	lucy.cassidy@bsuh.nhs.uk
Royal Sussex County Hospital, Brighton, UK	Arunray	Ray	arunray@doctors.org.uk
Royal Sussex County Hospital, Brighton, UK	Benedict	Rogers	benedict.rogers@bsuh.nhs.uk
Royal United Hospital, Bath, UK	Richard	Kapur	richard.kapur@btinternet.com
Royal United Hospital, Bath, UK	Katerina	Kyprianou	katerina.kyprianou@nhs.net

Royal United Hospital, Bath, UK	Payam	Tarassoli	Payam.Tarassoli@nhs.net
Royal Victoria Infirmary, Newcastle, UK	John	Ferns	j.fern@doctors.org.uk
Royal Victoria Infirmary, Newcastle, UK	Maire-Clare	Killen	maire-clare.killen@doctors.org.uk
Sandwell District General Hospital, West Bromwich	Hartej	Sur	hartej.sur@nhs.net
Sandwell District General Hospital, West Bromwich	Yin	Tam	ytam@nhs.net
South Tyneside General Hospital, South Shields, UK	Sathish	Bala	drskbala@doctors.org.uk
Southampton General Hospital, Southampton, UK	Rishi	Das	rishi.das@doctors.org.uk
Southampton General Hospital, Southampton, UK	Ryan	Higgin	rpchiggin@gmail.com
Southampton General Hospital, Southampton, UK	Ronathan	Topping	jetopping@outlook.com
Southmead Hospital, Bristol, UK	Sean	Duffy	seandxduffy@gmail.com
Southmead Hospital, Bristol, UK	Peter	Robinson	pwrobinson@doctors.org.uk
Southmead Hospital, Bristol, UK	Mike	Stoddart	michael.stoddart@nbt.nhs.uk
St Mary's Hospital, London, UK	Ali	ALTaweel	ali.altaweel@outlook.com
St Mary's Hospital, London, UK	Ernest	Chew	ernestchew@doctors.org.uk
St Mary's Hospital, London, UK	Ravi	Popat	dr.ravi.popat@googlemail.com
St Mary's Hospital, London, UK	Aadhar	Sharma	aadharsharma0@gmail.com
St Mary's Hospital, London, UK	Rupert	Wharton	rupertwharton@hotmail.co.uk
St Richard's Hospital, Chichester, UK	Samantha	Hook	samantha.hook@wsht.nhs.uk
St Richard's Hospital, Chichester, UK	Taran	Khangura	taran.khangura@wsht.nhs.uk
St Richard's Hospital, Chichester, UK	Sheena	Seewoonarin	sseewoonarin@gmail.com
St Richard's Hospital, Chichester, UK	Sian	Sokota	sian.sokota@wsht.nhs.uk
Stepping Hill Hospital, Stockport, UK	Charlotte	Cross	cecross@me.com
Stepping Hill Hospital, Stockport, UK	Kathryn	Mcloughlin	katmcloughlin@doctors.org.uk
Stoke Mandeville Hospital, Aylesbury, UK	Frances	Andrews	Frances.Andrews@buckshealthcare.nhs.uk
Stoke Mandeville Hospital, Aylesbury, UK	Chris	Bretherton	chris.bretherton@nhs.net
Stoke Mandeville Hospital, Aylesbury, UK	Jennie	Simpson	Jennie.Simpson@buckshealthcare.nhs.uk
Sunderland Royal Hospital, Sunderland, UK	Andreea	Lupu	dr.lupu.andreea@gmail.com
Sunderland Royal Hospital, Sunderland, UK	Charlotte	Thomas	charlottefonathomas@hotmail.com
Sunderland Royal Hospital, Sunderland, UK	Blair	Twieedie	blairtwieedie@gmail.com
Tameside General Hospital, Ashton-under-Lyne, UK	Kedar	Chirputka	kedar.chirputkar@tgh.nhs.uk
Tameside General Hospital, Ashton-under-Lyne, UK	Bernard	Ferns	bernard.fern@tgh.nhs.uk
Tameside General Hospital, Ashton-under-Lyne, UK	Steven	Ross	sajross@yahoo.com
Tameside General Hospital, Ashton-under-Lyne, UK	Yuhao	Zhang	yuhao.zhang@doctors.org.uk
Ulster Hospital, Dundonald, UK	Christopher	Goodland	cgoodland01@gmail.com
University Hospital Lewisham, London, UK	Benjamin	Barkham	benjamin.barkham@nhs.net
University Hospital Lewisham, London, UK	Jagmeet	Bhamra	j_s_bhamra@hotmail.com
University Hospital Lewisham, London, UK	Rory	Cuthbert	rory.cuthbert@nhs.net
University Hospital of Coventry & Warwickshire, Coventry, UK	Arun	Nair	arunnair@nhs.net
University Hospital of Coventry & Warwickshire, Coventry, UK	Robert	Payne	robert.payne@nhs.net
University Hospital of Coventry & Warwickshire, Coventry, UK	Matthew	Weston	matweston@doctors.org.uk
University Hospital of Coventry & Warwickshire, Coventry, UK	Ryan	Wong	ryanwong9@outlook.com
University Hospital of Coventry & Warwickshire, Coventry, UK	Tom	Wood	tomwood@doctors.org.uk
University Hospital of North Durham, Durham, UK	Philip	Thomas	pswthomas@doctors.org.uk
University Hospital of North Durham, Durham, UK	Hannah	Wilkinson	hannah.wilkinson14@nhs.net
University Hospital of North Tees, Stockton on Tees, UK	Muhammad	Adeelakhtar	muhammadadeelakhtar@me.com
University Hospital of South Manchester NHS Foundation Trust, Manchester, UK	Jo-Ann	Bate	joannbate@gmail.com
University Hospital of South Manchester NHS Foundation Trust, Manchester, UK	Bernardo	Duarte Brandao	bernardo.brandao@student.manchester.ac.uk
University Hospital of South Manchester NHS Foundation Trust, Manchester, UK	Martin	Lovell	martin.lovell@mft.nhs.uk
University Hospital of South Manchester NHS Foundation Trust, Manchester, UK	Richard	Unsworth	richard.unsworth@nhs.net
Victoria Hospital, Kirkcaldy, UK	Robert	Clayton	robertclayton@nhs.net
Victoria Hospital, Kirkcaldy, UK	Liam	Yapp	liam.yapp@gmail.com
Warrington Hospital, Warrington, UK	Ravi	Badgetu	ravi.badgetu@whh.nhs.uk
Warrington Hospital, Warrington, UK	Mike	Hughes	mwjhughes@hotmail.co.uk
Watford General Hospital, Watford, UK	Sophia	Burns	sophiaburns@doctors.org.uk
Watford General Hospital, Watford, UK	Ling	Kho	lingkho.9@gmail.com
Watford General Hospital, Watford, UK	Lily	Li	lily.xiaonuan.li@gmail.com
West Middlesex University Hospital, Isleworth, UK	Douglas	Evans	douglas.evans@doctors.org.uk
West Middlesex University Hospital, Isleworth, UK	Christopher	Fenner	christopher.oxton@gmail.com
West Middlesex University Hospital, Isleworth, UK	Kartik	Logishetty	k.logishetty@ic.ac.uk
West Middlesex University Hospital, Isleworth, UK	Sam	Nahas	sam.nahas@nhs.net
West Middlesex University Hospital, Isleworth, UK	Subra	Nathan	subra.nathan@chelwest.nhs.uk
West Suffolk Hospital, Bury Saint Edmunds, UK	Karadi	Kumar	drkhskumar@yahoo.com
West Suffolk Hospital, Bury Saint Edmunds, UK	Oliver	Negus	olovernegus@doctors.org.uk
West Suffolk Hospital, Bury Saint Edmunds, UK	Gemma	Salt	saltgemma@sky.com
West Suffolk Hospital, Bury Saint Edmunds, UK	Shahrier	Sarker	shahrier.fazal@gmail.com
Whittington Hospital, London, UK	Nicola	Blucher	nicblucher@doctors.org.uk
Whittington Hospital, London, UK	Shirley	Lyle	shirleylyle@doctors.org.uk
William Harvey Hospital (Ashford)	Christopher	Buckle	chrisbuckle@doctors.org.uk
William Harvey Hospital, Ashford, UK	Manafa	Chibuzo	chibuzomanafa@gmail.com
William Harvey Hospital, Ashford, UK	Mark	Jones	markdjones1984@gmail.com
William Harvey Hospital, Ashford, UK	Krasimir	Nachev	dr.nachev@gmail.com
William Harvey Hospital, Ashford, UK	Junaid	Sayani	junsayani@gmail.com
William Harvey Hospital, Ashford, UK	Olatoyosi	Williams	olatosi.williams@nhs.net
York District Hospital, York, UK	James	Cruikshank	jamescruikshank@doctors.org.uk
York District Hospital, York, UK	Nicholas	Edwards	nicholas.edwards5@nhs.net

References

- [1] Court-Brown CM, McBirnie J, Wilson G. Adult ankle fractures—an increasing problem? *Acta Orthop Scand* 1998;69(1):43–7.
- [2] Fractures (non – complex): Assessment and management. Nice Guidelines. 2016 <https://www.nice.org.uk/guidance/ng38> (date last Accessed 20 December 2018).
- [3] Rangan A, Jefferson L, Baker P, Cook L. Clinical trial networks in orthopaedic surgery. *Bone Joint Res* 2014;3:169–74.
- [4] Fractures (non – complex): Assessment and management, Appendices A-F. Nice Guidelines. 2016 <https://www.nice.org.uk/guidance/ng38/evidence/appendices-af-pdf-2358460766> (date last Accessed 20 December 2018).
- [5] Orthopaedic Trauma Association. Committee for Coding and Classification: fracture and dislocation compendium. *Orthop Trauma* 1996;10(1 Suppl):S1–54.
- [6] Egol KA, Amirtharajah M, Tejwani NC, Capla EL, Koval KJ. Ankle stress test for predicting the need for surgical fixation of isolated fibular fractures. *J Bone Joint Surg Am.* 2004;86-A(November (11)):2393–8.
- [7] Koval KJ, Egol KA, Cheung Y, Goodwin DW, Spratt KF. Does a positive ankle stress test indicate the need for operative treatment after lateral malleolus fracture? A preliminary report. *J Orthop Trauma* 2007;21(August (7)):449–55.
- [8] Gill JB, Risko T, Raducan V, Grimes JS, Schutt Jr. RC. Comparison of manual and gravity stress radiographs for the evaluation of supination-external rotation fibular fractures. *Bone Joint Surg Am* 2007;89(May (5)):994–9.
- [9] Weber M, Burmeister H, Flueckiger G, Krause FG. The use of weightbearing radiographs to assess the stability of supination-external rotation fractures of the ankle. *Arch Orthop Trauma Surg.* 2010;130(May (5)):693–8.
- [10] Dawe EJ, Shafafy R, Quayle J, Gougoulias N, Wee A, Sakellariou A. The effect of different methods of stability assessment on fixation rate and complications in supination external rotation (SER) 2/4 ankle fractures. *Foot Ankle Surg* 2015;21(June (2)):86–90.
- [11] Hastie GR, Akhtar S, Butt U, Baumann A, Barrie JL. Weightbearing radiographs facilitate functional treatment of ankle fractures of uncertain stability. *J Foot Ankle Surg* 2015;54(November–December (6)):1042–6.
- [12] Hoshino CM, Nomoto EK, Norheim EP, Harris TG. Correlation of weightbearing radiographs and stability of stress positive ankle fractures. *Foot Ankle Int.* 2012;33(February (2)):92–8.
- [13] Lin CW, Donkers NAJ, Refshauge KM, Beckenkamp PR, Khara K, Moseley AM. Rehabilitation for ankle fractures in adults. *Cochrane Database Syst Rev* 2012. Issue 11. Art. No.: CD005595.
- [14] Ahl T, Dalén N, Holmberg S, Selvik G. Early weight bearing of malleolar fractures. *Acta Orthop Scand* 1986 Dec;57(6):526–9.
- [15] Ahl T, Dalén N, Holmberg S, Selvik G. Early weight bearing of displaced ankle fractures. *Acta Orthop Scand* 1987 Oct;58(5):535–8.
- [16] Finsen V, Saetermo R, Kibsgaard L, Farran K, Engebretsen L, Bolz KD, et al. Early postoperative weight-bearing and muscle activity in patients who have a fracture of the ankle. *J Bone Joint Surg Am* 1989;71(1):23–7.
- [17] Ahl T, Dalén N, Lundberg A, Bylund C. Early mobilization of operated on ankle fractures. Prospective, controlled study of 40 bimalleolar cases. *Acta Orthop Scand* 1993;64(February (1)):95–9.
- [18] Khair E, Charopoulos I, Dimitriou R, Ghoo A, Dahabreh Z, Giannoudis P. The health economics of ankle fracture fixation. *Bulletin* 2012;94(April (4)):1–5.
- [19] Keene DJ, Mistry D, Nam J, Tutton E, Handley R, Morgan L, et al. The Ankle Injury Management (AIM) trial: a pragmatic, multicentre, equivalence randomised controlled trial and economic evaluation comparing close contact casting with open surgical reduction and internal fixation in the treatment of unstable ankle fractures in patients aged over 60 years. *Health Technol Assess* 2016;20(75).
- [20] McDonald MR, Sathiyakumar V, Apfeld JC, Hooe B, Ehrenfeld J, Obrensky WT, et al. Predictive factors of hospital length of stay in patients with operatively treated ankle fractures. *J Orthop Traumatol* 2014;15(4):255–8.
- [21] Simanski CJ, Maegele MG, Lefering R, Lehnen DM, Kawel N, Riess P, et al. Functional treatment and early weightbearing after an ankle fracture: a prospective study. *J Orthop Trauma* 2006;20(February (2)):108–14.
- [22] Gul A, Batra S, Mehmood S, Gillham N. Immediate unprotected weight-bearing of operatively treated ankle fractures. *Acta Orthop Belg* 2007;73(June (3)):360–5.
- [23] McPhail SM, Dunstan J, Canning J, Haines TP. Life impact of ankle fractures: qualitative analysis of patient and clinician experiences. *BMC Musculoskeletal Disord* 2012;21(13):224.
- [24] Yin MC, Yuan XF, Ma JM, Xia Y, Wang T, Xu XL, et al. Evaluating the reliability and reproducibility of the ao and lauge-hansen classification systems for ankle injuries. *Orthopedics* 2015;38(July (7)):e626–30.
- [25] Swart E, Bezhani H, Greisberg J, Vosseller JT. How long should patients be kept non-weight bearing after ankle fracture fixation? A survey of OTA and AOFAS members. *Injury* 2015;46(6):1127–30.