

## Clinical Reviews in Emergency Medicine

### CAN SYSTEMIC THROMBOLYSIS IMPROVE PROGNOSIS OF CARDIAC ARREST PATIENTS DURING CARDIOPULMONARY RESUSCITATION? A SYSTEMATIC REVIEW AND META-ANALYSIS

Yiwei Wang, MD, Maoyun Wang, MD, Yuenan Ni, PHD, Binmiao Liang, PHD, and Zongan Liang, MD

Department of Respiratory and Critical Care Medicine, West China School of Medicine and West China Hospital, Sichuan University, Sichuan, China

Reprint Address: Zongan Liang, MD, Department of Respiratory and Critical Care Medicine, West China School of Medicine and West China Hospital, Sichuan University, No. 37 Guoxue Alley, Chengdu 610041, China

**Abstract—Background:** Cardiac arrests are caused in most cases by thromboembolic diseases, such as acute myocardial infarction (AMI) and pulmonary embolism (PE). **Objective:** We aimed to ascertain the associations of thrombolytic therapy with potential benefits among cardiac arrest patients during cardiopulmonary resuscitation (CPR). **Methods:** We searched PubMed, Embase, and Cochrane databases for studies that evaluated systemic thrombolysis in cardiac arrest patients. The primary outcome was survival to hospital discharge, and secondary outcomes included return of spontaneous circulation (ROSC), 24-h survival rate, hospital admission rate, and bleeding complications. **Results:** Nine studies with a total of 4384 cardiac arrest patients were pooled in the meta-analysis, including 1084 patients receiving systemic thrombolysis and 3300 patients receiving traditional treatments. Compared with conventional therapies, the use of systemic thrombolysis did not significantly improve survival to hospital discharge (13.5% vs. 10.8%; risk ratio [RR] 1.13; 95% confidence interval [CI] 0.92–1.39;  $p = 0.24$ ,

$I^2 = 35\%$ ), ROSC (50.9% vs. 44.3%; RR 1.29; 95% CI 1.00–1.66;  $p = 0.05$ ,  $I^2 = 73\%$ ), and 24-h survival (28.1% vs. 25.6%; RR 1.25; 95% CI 0.88–1.77;  $p = 0.22$ ,  $I^2 = 63\%$ ). We observed higher hospital admission rates for patients receiving systemic thrombolysis (43.4% vs. 30.6%; RR 1.53; 95% CI 1.04–2.24;  $p = 0.03$ ,  $I^2 = 87\%$ ). In addition, higher risk of bleeding was observed in the thrombolysis group (8.8% vs. 5.0%; RR 1.65; 95% CI 1.16–2.35;  $p = 0.005$ ,  $I^2 = 7\%$ ). **Conclusions:** Systemic thrombolysis during CPR did not improve hospital discharge rate, ROSC, and 24-h survival for cardiac arrest patients. Patients receiving thrombolytic therapy have a higher risk of bleeding. More high-quality studies are needed to confirm our results. © 2019 Elsevier Inc. All rights reserved.

**Keywords—**thrombolysis; cardiac arrest; cardiopulmonary resuscitation; pulmonary embolism; acute myocardial infarction

Yiwei Wang and Maoyun Wang contributed equally to this work.

Funded by the National Key Research and Development Program of China, China (2016YFC1304303).

Each clinical trial included was approved by the corresponding institution's ethical committee.

Written informed consent was provided by participants of each study.

## INTRODUCTION

Cardiac arrest is a life-threatening condition and a major cause of sudden death. Each year, it causes more than 3.7 million deaths worldwide (1). Studies show that 75–85% of out-of-hospital cardiac arrest cases have a primary

cardiac cause, and respiratory diseases account for 8% of cardiac arrest (2,3).

Thromboembolic diseases, such as acute myocardial infarction (AMI) and pulmonary embolism (PE), are the most common causes of cardiac arrest (4–6). Thrombolytic therapy can reduce all-cause mortality and is recommended in guidelines for AMI and high-risk PE patients (presenting with shock or hypotension) (7–10). Dissolving blood emboli by administering fibrinolytic substances to re-establish circulation in cardiac arrest patients might work from a pathophysiological point of view. However, thrombolysis can increase bleeding risk and is one of the relative contraindications in patients receiving cardiopulmonary resuscitation (CPR) (11). Guidelines only suggest thrombolytic therapy for cardiac arrests secondary to PE, considering the life-threatening situation, and it is not routinely recommended for other etiologies (12,13).

Numerous studies have been published about thrombolytic therapy for cardiac arrest patients with inconsistent results. A retrospective study reported by Lederer et al. found that patients receiving recombinant tissue plasminogen activator (rt-PA) during CPR could improve return of spontaneous circulation (ROSC) and survival to hospital discharge (14). Bozeman et al. conducted a prospective cohort study in emergency departments and reported higher ROSC and hospital admission rates, but no difference was found for hospital discharge rate (15). The largest randomized controlled trial (RCT), conducted by Böttiger et al., did not detect any improvement in outcomes (16). Considering the clinical value and the current lack of evidence, we performed this systematic review and meta-analysis to identify the latest evidence on whether systemic thrombolysis has been shown to improve patient prognosis after cardiac arrest.

## METHODS

We conducted this review in adherence to the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analysis) statement according to current guidelines (17). We registered the protocol of this systematic review with PROSPERO (International Prospective Register of Systematic Reviews; registration number: CRD42017069703).

### Search Strategy

We searched PubMed, Embase, and Cochrane databases from inception to January 2019, and search strategies were adapted for each database, including medical subject headings and keywords for *heart arrest* or *cardiopulmonary resuscitation* and *thrombolysis* or *tissue*

*plasminogen activator* without limitations on publication type or language.

### Eligibility Criteria

We identified eligible comparative studies according to the following inclusion criteria: 1) exported data from original research studies, 2) human adults ( $\geq 16$  years of age) suffering from sudden cardiac arrest, 3) studies focusing on systemic thrombolysis therapy during CPR, and 4) studies including at least one of the predetermined outcomes. We excluded studies published as conference abstracts in our meta-analysis.

### Data Extraction and Quality Assessment

After removing duplications, two reviewers (YW-W and MY-W) independently assessed citation titles and abstracts using a “relevant,” “irrelevant,” or “unsure” designation. Disagreements were resolved by discussing with a third investigator (YN-N). The quality of the included studies was assessed by the Cochrane Risk of Bias Tool for RCTs and Newcastle-Ottawa Scale for observational studies (18,19).

### Outcomes


The primary outcome was survival to hospital discharge. The secondary outcomes included ROSC, hospital admission, 24-h survival, and bleeding complications.

### Heterogeneity and Sensitivity Analysis

We reported the overall effect size as well as the separate effect sizes for RCTs and observational studies. Studies were tested for heterogeneity using the  $I^2$  statistic. An  $I^2$  value  $> 50\%$  suggested substantial heterogeneity (20). We used fixed-effects models to pool data with insignificant heterogeneity and the random-effects models for data with significant heterogeneity. We performed a sensitivity analysis by excluding one trial in each round to test the influence of a single study on the overall pooled estimate.

### Statistical Analysis

We undertook this meta-analysis in Review Manager 5.3 software (Cochrane Collaboration, Oxford, UK) using a random-effects model to incorporate covariate adjustment to mitigate overspecification. Meta-analysis results were presented in terms of risk ratio (RR) and 95% confidence interval (CI) for dichotomous outcomes. All statistical tests were two-sided, and statistical significance was defined as  $p < 0.05$ .



**Figure 1. Flow diagram of primary study search process. CPR = cardiopulmonary resuscitation.**

## RESULTS

### Study Selection

A total of 2971 records were identified through database searching. After removing duplicates, 1935 records were removed. The 140 remaining records were identified as being potentially relevant, and abstracts were assessed for eligibility. A total of 19 articles discussed thrombolytic therapy in cardiac arrest patients, and full articles were read carefully. Finally, nine studies discussing thrombolysis during CPR were identified and included in the meta-analysis (Figure 1) (14–16,21–26).

As being potentially relevant, and abstracts were assessed for eligibility. A total of 19 articles discussed thrombolytic therapy in cardiac arrest patients, and full articles were read carefully. Finally, nine studies discussing thrombolysis during CPR were identified and included in the meta-analysis (Figure 1) (14–16,21–26).

**Table 1. Characteristics of Studies Included in the Meta-Analysis**


| First Author (Reference) | Year | Study Design       | Etiology                | Thrombolytic Agent      | Patients, n, Thrombolysis/Control | Outcomes                                            |
|--------------------------|------|--------------------|-------------------------|-------------------------|-----------------------------------|-----------------------------------------------------|
| Abu-Laban (22)           | 2002 | RCT                | Non-traumatic           | Alteplase               | 117/116                           | Discharge, 24-h survival, admission, ROSC, bleeding |
| Böttiger (21)            | 2001 | Prospective cohort | Non-traumatic           | Alteplase               | 40/50                             | Discharge, 24-h survival, admission, ROSC, bleeding |
| Böttiger (16)            | 2008 | RCT                | Presumed cardiac origin | Tenecteplase            | 525/525                           | Discharge, 24-h survival, admission, ROSC, bleeding |
| Bozeman (15)             | 2006 | Prospective cohort | Non-traumatic           | Tenecteplase            | 50/113                            | Discharge, 24-h survival, admission, ROSC, bleeding |
| Fatovich (23)            | 2004 | RCT                | Presumed cardiac or PE  | Alteplase               | 19/16                             | Discharge, admission, ROSC, bleeding                |
| Lederer (14)             | 2001 | Retrospect cohort  | Non-traumatic           | Alteplase               | 108/216                           | Discharge, 24-h survival, ROSC, bleeding            |
| Renard (25)              | 2011 | Retrospect cohort  | Non-traumatic           | Alteplase, tenecteplase | 107/1154                          | Admission                                           |
| Stadlbauer (24)          | 2006 | Post-hoc analysis  | Non-traumatic           | Tenecteplase, reteplase | 99/1087                           | Discharge, admission                                |
| Yousuf (26)              | 2016 | Retrospect cohort  | PE                      | Alteplase               | 19/23                             | Discharge, 24-h survival, bleeding                  |

AMI = acute myocardial infarction; PE = pulmonary embolism; RCT = randomized controlled trial; ROSC = return of spontaneous circulation.

**Table 2. Quality Assessment of Included Studies**

| First Author, Year<br>(Reference)                                           | Selection                               |                                       |                                                                                      |                                  |                                                        | Outcome                  |                          |                          |   | Total<br>Stars, n |
|-----------------------------------------------------------------------------|-----------------------------------------|---------------------------------------|--------------------------------------------------------------------------------------|----------------------------------|--------------------------------------------------------|--------------------------|--------------------------|--------------------------|---|-------------------|
|                                                                             | Representativeness<br>of Exposed Cohort | Selection of<br>Non-Exposed<br>Cohort | Demonstration<br>That Outcome of<br>Interest Was Not<br>Present at Start<br>of Study | Ascertainment<br>of Exposure     | Comparability<br>on Basis of<br>Design and<br>Analysis | Assessment<br>of Outcome | Follow-Up<br>Long Enough | Adequacy of<br>Follow-Up |   |                   |
| Newcastle-Ottawa Scale for Assessing the Quality of Included Cohort Studies |                                         |                                       |                                                                                      |                                  |                                                        |                          |                          |                          |   |                   |
| Böttiger, 2001 (21)                                                         | *                                       | *                                     | *                                                                                    | *                                | **                                                     | *                        | *                        | *                        | 9 |                   |
| Bozeman, 2006 (15)                                                          | *                                       | *                                     | *                                                                                    | *                                | —                                                      | *                        | *                        | *                        | 7 |                   |
| Lederer, 2001 (14)                                                          | *                                       | *                                     | *                                                                                    | *                                | *                                                      | *                        | *                        | *                        | 8 |                   |
| Renard, 2011 (25)                                                           | *                                       | *                                     | *                                                                                    | *                                | —                                                      | *                        | *                        | *                        | 7 |                   |
| Stadlbauer, 2006 (24)                                                       | *                                       | *                                     | *                                                                                    | *                                | *                                                      | *                        | *                        | *                        | 8 |                   |
| Yousuf, 2016 (26)                                                           | *                                       | *                                     | *                                                                                    | *                                | **                                                     | *                        | *                        | *                        | 9 |                   |
| Author, Year<br>(Reference)                                                 | Random<br>Assignment                    | Allocation<br>Concealment             | Blinding of<br>Participants                                                          | Blind Evaluation<br>For Outcomes | Incomplete<br>Outcome Data                             | Selective<br>Reporting   | Other Bias               |                          |   |                   |
| Quality Assessment of Included RCTs                                         |                                         |                                       |                                                                                      |                                  |                                                        |                          |                          |                          |   |                   |
| Abu-Laban, 2002 (22)                                                        | Low risk                                | Low risk                              | Low risk                                                                             | Low risk                         | Low risk                                               | Low risk                 | Low risk                 | Low risk                 |   |                   |
| Fatovich, 2004 (23)                                                         | Low risk                                | Low risk                              | Low risk                                                                             | Low risk                         | Low risk                                               | Low risk                 | Low risk                 | Low risk                 |   |                   |
| Böttiger, 2008 (16)                                                         | Low risk                                | Low risk                              | Low risk                                                                             | Low risk                         | Low risk                                               | Low risk                 | Low risk                 | Low risk                 |   |                   |

RCT = randomized controlled trial.



**Figure 2. Effect of systemic thrombolysis on survival to hospital discharge.** CI = confidence interval; M-H = Mantel-Haenszel.

### Study Characteristics

Nine studies with a total of 4384 cardiac arrest patients were pooled in the meta-analysis, including 1084 patients receiving systemic thrombolysis and 3300 patients receiving traditional treatments. Of the nine studies included, three were RCTs, three were retrospective cohort studies, two were prospective cohort studies, and one was a post-hoc analysis. Details of each study included in our analysis are summarized in Table 1.

### Quality Assessment

The Cochrane Risk of Bias Tool was used for RCTs, and the risk of bias was rated as “low,” “unclear,” or “high.” The Newcastle-Ottawa Scale was used to assess the quality of observational studies with a maximum of 9 stars, and a study with final stars  $\geq 6$  was regarded as high quality. Overall, all of the included trials were considered high quality. The quality assessment results are presented in Table 2.

### Survival to Hospital Discharge

Eight studies were pooled in this comparison, including 969 patients receiving thrombolysis and 2135 patients in the control group. The results showed that patients in both groups had survival rates similar to hospital

discharge rates (13.5% vs. 10.8%; RR 1.08; 95% CI 0.92–1.39;  $p = 0.24$ ,  $I^2 = 35\%$ ) (Figure 2).

### ROSC


We compared the rate of ROSC between the thrombolytic group and the control group. Six studies with a total of 1871 patients were enrolled, and we chose the random-effects model because of high heterogeneity. There was a trend for higher ROSC in the thrombolytic group, but the difference was not statistically significant (50.9% vs. 44.3%; RR 1.29; 95% CI 1.00–1.66;  $p = 0.05$ ,  $I^2 = 73\%$ ) (Figure 3).

### Hospital Admission

Seven studies with a total of 4018 patients were pooled in this analysis. No study reported the admission time between emergency department and hospital ward, and we used the random-effects models for high heterogeneity. The result showed that patients in the thrombolytic group had higher hospital admission rates (43.4% vs. 30.6%; RR 1.53; 95% CI 1.04–2.24;  $p = 0.03$ ,  $I^2 = 87\%$ ) (Figure 4).

### 24-Hour Survival Rate

Six studies with 1883 patients analyzed the 24-h survival rate. No significant statistical difference was found



**Figure 3. Effect of systemic thrombolysis on return of spontaneous circulation.** CI = confidence interval; M-H = Mantel-Haenszel.




Figure 4. Effect of systemic thrombolysis on hospital admission. CI = confidence interval; M-H = Mantel-Haenszel.

between the thrombolytic group and the control group (28.1% vs. 25.6%; RR 1.25; 95% CI 0.88–1.77;  $p = 0.22$ ,  $I^2 = 63\%$ ) (Figure 5).

### Bleeding Complications

Seven studies with a total of 1686 patients reported bleeding complications. Compared with non-thrombolytic therapy, patients receiving systemic thrombolysis showed a significantly higher risk of bleeding (8.8% vs. 5.0%; RR 1.65; 95% CI 1.16–2.35;  $p = 0.005$ ,  $I^2 = 7\%$ ) (Figure 6).

### Publication Bias

No obvious publication bias was observed by a visual inspection of the funnel plots in our meta-analysis (Figure 7).

## DISCUSSION

Our study suggested that, compared with conventional therapies for cardiac arrests, patients receiving systemic thrombolysis during CPR did not show any improvements in survival to hospital discharge, ROSC, and 24-h survival. A higher hospital admission rate was observed. In addition, more bleeding events were reported for patients receiving thrombolytic therapy.

Thromboembolic diseases, such as AMI and PE, are the main causes of non-traumatic cardiac arrests.

Obstruction of the coronary artery and pulmonary trunk can affect systolic function of the heart and causes hemodynamic instability or even cardiac arrest. Based on the pharmacologic properties of thrombolytic agents, systemic thrombolysis can dissolve blood clots, thus helping reperfusion of important organs. However, our meta-analysis did not find any improvement in hospital discharge, ROSC, and 24-h survival rates. The cause of these results may be related to the restriction of organ perfusion during chest compression, which may hamper the delivery of thrombolytic drugs to blood clots.

ROSC is the sustained perfusing cardiac activity and respiratory activity after cardiac arrest. It is regarded as an earlier achievement of CPR that may indirectly decrease mortality. In our meta-analysis, higher ROSC was observed in patients receiving thrombolysis, but the difference was not statistically significant. This finding might suggest that thrombolytic drugs have the potential to improve circulation, or it may be related to the small sample size of our study. Further increasing the number of articles may make this result more obvious. Approximately half of the cardiac arrest patients achieved ROSC in our meta-analysis, but only slightly more than 10% of patients survived to hospital discharge. According to some large studies, approximately 70% of ROSC patients died of complications (27,28). The survival rate remains poor, even after achieving ROSC, and post-cardiac arrest syndrome (PCAS) is the most common




Figure 5. Effect of systemic thrombolysis on 24-hour survival. CI = confidence interval; M-H = Mantel-Haenszel.




Figure 6. Effect of systemic thrombolysis on bleeding complications. CI = confidence interval; M-H = Mantel-Haenszel.

cause (29). PCAS refers to a series of abnormalities that develop after resuscitation, including post-cardiac arrest brain injury, post-cardiac arrest myocardial dysfunction, systemic ischemia/reperfusion response, and persistent precipitating pathology. The most common causes of death during the first 24 h after ROSC are refractory shock and multi-organ system failure, while later deaths result from neurologic injury (30).

In our meta-analysis, patients receiving systemic thrombolysis had higher hospital admission rates. None of the articles included mentioned the admission criteria, and very high heterogeneity was observed. It is difficult to assess whether this heterogeneity was related to the potential role of thrombolytics. Compared with the 24-h survival rate, we hypothesized that hospital admission was not representative to assess short-term survival. However, a higher admission rate also means more opportunities for treatment, and we should not ignore the potential benefits for patients.

Bleeding is one of the complications of thrombolytic therapy, resulting from systemic activation of plasmin outside the thrombus that leads to systemic fibrinolysis (31). Some studies we examined reported the total number of bleeding events, while other studies only reported major bleeding, or divided bleeding events into minor bleeding and major bleeding. Given the inconsistent definition and inclusion criteria, we analyzed all types of

bleeding events without classification. In our meta-analysis, an increase in bleeding complications was observed, but there was not sufficient evidence to prove that thrombolysis would increase bleeding-related mortality from the studies examined.

Systemic thrombolytic therapy is widely used in the treatment of coronary and pulmonary thrombosis. Current approaches for AMI patients aim to restore the blood flow to myocardial cells, and guidelines recommended thrombolysis for patients who have no access to percutaneous coronary intervention within 120 min (32,33). For PE, systemic thrombolytic therapy is recommended for patients with hemodynamic instability, and greatly improves their prognosis (10). For patients with clear-onset cardiac arrest, thrombolysis might improve outcomes according to some observational studies, but the results have yet to be verified by high-quality RCTs (34,35). For sudden cardiac arrest, physicians often do not have sufficient time to determine the cause of cardiac arrest. The results of our meta-analysis indicate that there is no survival benefit for systemic thrombolysis during CPR in cardiac arrest patients with unclear etiologies.

It is controversial whether thrombolysis could improve prognosis of cardiac arrest patients. Prior to our study, a meta-analysis with eight studies on cardiac arrest patients receiving thrombolytic therapy was published in 2004 (36). The meta-analysis showed that thrombolysis could improve survival to hospital discharge, ROSC, hospital admission, and 24-h survival, and was associated with better long-term neurologic function. After carefully reading all of the articles, only two studies met our inclusion criteria. Other studies were excluded for enrolling subjects receiving thrombolysis after CPR. All of the studies included in our review focused on subjects receiving systemic thrombolysis during CPR. The difference between our study and the conclusions of previous articles may be attributed to the research object itself. Despite the inconsistent results, systemic thrombolysis is still used for the rescue of patients with cardiac arrest. To ascertain the associations of thrombolytic therapy with potential benefits among




Figure 7. Funnel plots for publication bias. RR = risk ratio; SE = standard error.

cardiac arrest patients, we conducted this systematic review and meta-analysis by focusing on systemic thrombolysis during CPR.

### Limitations

This meta-analysis summarizes the most recent evidence available on systemic thrombolysis therapy in cardiac arrests, but some limitations should be noted. First, our research consists of nine articles. Of these studies, only three are RCTs, and the others are post-hoc analysis and observational studies. More high-quality studies are needed to evaluate the efficacy of systemic thrombolysis in the future. Second, heterogeneity was inevitable in many aspects, such as different etiologies for cardiac arrest, inconsistent baseline levels, and various supportive treatments. In addition, varying doses and types of thrombolytic drugs were used, and different anticoagulant therapies were employed in the studies. These differences potentially affect the evaluation of the thrombolytic treatment.

### CONCLUSIONS

Systemic thrombolysis does not improve survival to hospital discharge, ROSC, and 24-h survival rates, and is associated with more bleeding events for cardiac arrest patients. More large RCTs are needed to confirm our results.

**Acknowledgments**—Authors' contributions: YW-W designed the study, conducted the literature search, performed data analysis, and drafted the manuscript. MY-W conducted the literature search and revised the manuscript. YN-N helped in the design of the study and data analysis. BM-L helped in data analysis and revised the manuscript. ZA-L participated in modifying the important knowledge content and provided the final version for publication. All authors read and approved the final manuscript.

### REFERENCES

1. Kudenchuk PJ, Sandroni C, Drinhaus HR, et al. Breakthrough in cardiac arrest: reports from the 4th Paris International Conference. *Ann Intensive Care* 2015;5:22.
2. Berdowski J, Berg RA, Tijssen JG, Koster RW. Global incidences of out-of-hospital cardiac arrest and survival rates: systematic review of 67 prospective studies. *Resuscitation* 2010;81:1479–87.
3. Perkins GD, Lall R, Quinn T, et al., PARAMEDIC Trial Collaborators. Mechanical versus manual chest compression for out-of-hospital cardiac arrest (PARAMEDIC): a pragmatic, cluster randomized controlled trial. *Lancet* 2015;385(9972):947–55.
4. Hess EP, Campbell RL, White RD. Epidemiology, trends, and outcome of out-of-hospital cardiac arrest of non-cardiac origin. *Resuscitation* 2007;72:200–6.
5. Deasy C, Bray JE, Smith K, Harriss LR, Bernard SA, Cameron P. Out-of-hospital cardiac arrest in young adults in Melbourne, Australia—adding coronial data to cardiac arrest registry. *Resuscitation* 2011;82:1302–6.
6. Spaulding CM, Joly LM, Rosenberg A, et al. Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med* 1997;336:1629–33.
7. O'Gara PT, Kushner FG, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 2013;127:e362–e425.
8. Steg PG, James SK, Atar D, et al. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2012;33:2569–619.
9. Konstantinides SV, Torbicki A, Agnelli G, et al. 2014 ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J* 2014;35:3033–69. 69a–k.
10. Chatterjee S, Chakraborty A, Weinberg I, et al. Thrombolysis for pulmonary embolism and risk of all-cause mortality, major bleeding, and intracranial hemorrhage: a meta-analysis. *JAMA* 2014;311:2414–21.
11. Brunton LL, Parker KL, Lazo JS, eds. *Goodman & Gilman's The Pharmacological Basis of Therapeutics*. 11th ed. New York: McGraw Hill; 2011.
12. Truhlar A, Deakin CD, Soar J, et al. European Resuscitation Council Guidelines for Resuscitation 2015: Section 4. Cardiac arrest in special circumstances. *Resuscitation* 2015;95:148–201.
13. Lavonas EJ, Drennan IR, Gabrielli A, et al. Part 10: Special Circumstances of Resuscitation 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2015;132(suppl 2):S501–18.
14. Lederer W, Lichtenberger C, Pechlaner C, Kroesen G, Baubin M. Recombinant tissue plasminogen activator during cardiopulmonary resuscitation in 108 patients with out-of-hospital cardiac arrest. *Resuscitation* 2001;50:71–6.
15. Bozeman WP, Kleiner DM, Ferguson KL. Empiric tenecteplase is associated with increased return of spontaneous circulation and short term survival in cardiac arrest patients unresponsive to standard interventions. *Resuscitation* 2006;69:399–406.
16. Böttiger BW, Arntz HR, Chamberlain DA, et al. Thrombolysis during resuscitation for out-of-hospital cardiac arrest. *N Engl J Med* 2008;359:2651–62.
17. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med* 2009;151:W65–94.
18. Higgins JPT, Altman DG, Sterne JAC. Assessing risk of bias in included studies. In: Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*, version 5.1.0. 2011. <http://handbook.cochrane.org>. Accessed June 27, 2017.
19. Stang A. Critical evaluation of the Newcastle-Ottawa Scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010;25:603–5.
20. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
21. Böttiger BW, Bode C, Kern S, et al. Efficacy and safety of thrombolytic therapy after initially unsuccessful cardiopulmonary resuscitation: a prospective clinical trial. *Lancet* 2001;357(9268):1583–5.
22. Abu-Laban RB, Christenson JM, Innes GD, et al. Tissue plasminogen activator in cardiac arrest with pulseless electrical activity. *N Engl J Med* 2002;346:1522–8.
23. Fatovich DM, Dobb GJ, Clugston RA. A pilot randomised trial of thrombolysis in cardiac arrest (The TICA trial). *Resuscitation* 2004;61:309–13.
24. Stadlbauer KH, Krismer AC, Arntz HR, et al. Effects of thrombolysis during out-of-hospital cardiopulmonary resuscitation. *Am J Cardiol* 2006;97:305–8.
25. Renard A, Verret C, Jost D, et al. Impact of fibrinolysis on immediate prognosis of patients with out-of-hospital cardiac arrest. *J Thromb Thrombolysis* 2011;32:405–9.
26. Yousuf T, Brinton T, Ahmed K, et al. Tissue plasminogen activator use in cardiac arrest secondary to fulminant pulmonary embolism. *J Clin Med Res* 2016;8:190–5.

27. Nadkarni VM, Larkin GL, Peberdy MA, et al. First documented rhythm and clinical outcome from in-hospital cardiac arrest among children and adults. *JAMA* 2006;295:50–7.
28. Nolan JP, Laver SR, Welch CA, Harrison DA, Gupta V, Rowan K. Outcome following admission to UK intensive care units after cardiac arrest: a secondary analysis of the ICNARC Case Mix Programme Database. *Anaesthesia* 2007;62:1207–16.
29. Neumar RW, Nolan JP, Adrie C, et al. Post-Cardiac Arrest Syndrome: Epidemiology, Pathophysiology, Treatment, and Prognostication A Consensus Statement From the International Liaison Committee on Resuscitation (American Heart Association, Australian and New Zealand Council on Resuscitation, European Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Asia, and the Resuscitation Council of Southern Africa); the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; and the Stroke Council. *Circulation* 2008; 118:2452–83.
30. Jentzer JC, Chonde MD, DeZfulian C. Myocardial dysfunction and shock after cardiac arrest. *Biomed Res Int* 2015;2015(11):1–14.
31. Gurman P, Miranda OR, Nathan A, Washington C, Rosen Y, Elman NM. Recombinant tissue plasminogen activators (rtPA): a review. *Clin Pharmacol Ther* 2015;97:274–85.
32. Kireyev D, Tan HC, Poh KK. Management of acute ST-elevation myocardial infarction: Reperfusion options. *Ann Acad Med Singapore* 2010;39:927–33.
33. Pinto DS, Frederick PD, Chakrabarti AK, et al. Benefit of transferring ST-segment-elevation myocardial infarction patients for percutaneous coronary intervention compared with administration of onsite fibrinolytic declines as delays increase. *Circulation* 2011; 124:2512–21.
34. Janata K, Holzer M, Kırkcıyan I, et al. Major bleeding complications in cardiopulmonary resuscitation: the place of thrombolytic therapy in cardiac arrest due to massive pulmonary embolism. *Resuscitation* 2003;57:49–55.
35. van Campen LC, van Leeuwen GR, Verheugt FW. Safety and efficacy of thrombolysis for acute myocardial infarction in patients with prolonged out-of-hospital cardiopulmonary resuscitation. *Am J Cardiol* 1994;73:953–5.
36. Li X, Fu QL, Jing XL, et al. A meta-analysis of cardiopulmonary resuscitation with and without the administration of thrombolytic agents. *Resuscitation* 2006;70:31–6.

### ARTICLE SUMMARY

**1. Why is this topic important?**

The mortality rate of cardiac arrest patients is very high and limited measures can be taken in addition to cardiopulmonary resuscitation (CPR). Many studies reported systemic thrombolytic therapy during CPR with inconsistent results.

**2. What does this review attempt to show?**

We aim to ascertain associations of thrombolytic therapy with potential benefits among cardiac arrest patients during CPR with the latest evidence.

**3. What are the key findings?**

Systemic thrombolysis during CPR did not significantly improve survival to hospital discharge, return of spontaneous circulation, and 24-h survival rate. A higher hospital admission rate and more bleeding events were reported for patients receiving thrombolytic therapy.

**4. How is patient care impacted?**

Systemic thrombolysis does not improve hospital survival during CPR and can cause more bleeding events. Clinicians should be cautious in choosing thrombolytic therapy for cardiac arrest patients with unknown etiology.