




Contents lists available at ScienceDirect

Progress in Cardiovascular Diseases

journal homepage: www.onlinepcd.com



The role of cardiorespiratory fitness on the risk of sudden cardiac death at the population level: A systematic review and meta-analysis of the available evidence[☆]

David Jiménez-Pavón^{a,b,*}, Carl J. Lavie^c, Steven N. Blair^d

^a MOVE-IT Research Group and Department of Physical Education, Faculty of Education Sciences University of Cádiz, Cádiz, Spain

^b Biomedical Research and Innovation Institute of Cádiz (INIBICA) Research Unit, Puerta del Mar University Hospital University of Cádiz, Spain

^c Department of Cardiovascular Diseases, John Ochsner Heart and Vascular Institute, Ochsner Clinical School-The University of Queensland's School of Medicine, New Orleans, LA

^d Department of Exercise Sciences, University of South Carolina, Columbia, SC

ARTICLE INFO

Article history:

4 May 2019

4 May 2019

Keywords:

Cardiorespiratory fitness

Sudden cardiac death

Large cohorts

Coronary heart disease

Risk factors

ABSTRACT

Cardiorespiratory fitness (CRF) has been widely studied as a powerful and independent predictor of all-cause and disease-specific mortality. Sudden cardiac death (SCD) is recognized as a significant cause of mortality among the general population, including the general population without previous symptoms of any coronary heart disease (CHD). Consequently, SCD is an important public health problem, which constitutes a clinical challenge. Thus, prevention of SCD by detecting early risk factors could be a useful tool, contributing to the American Heart Association's goal of decreasing the incidence of SCD at the population level. The identification of these risk factors for CVD would facilitate the large-scale screening of those participants at higher risk of SCD. This systematic review collects information about the role of CRF on the risk of SCD at the available evidence, and analyzes the long-term influence of CRF as a risk factor and independent predictor of SCD.

© 2019 Elsevier Inc. All rights reserved.

Contents

Definition and relevance of sudden cardiac death	280
Definition and importance of assessing cardiorespiratory fitness	280
Methodology and search strategy	280
Search strategy	280
Selection criteria	280
Data extraction	281
Meta-analysis	281
Results	281
Overall results	281
The two population cohorts followed to assess the independent role of maximal CRF on SCD	281
Meta-analysis of the influence of CRF on SCD in two different cohorts	282
The combined role of maximal CRF or sub-maximal CRF on the risk of SCD	284
Discussion	285
Main findings	285
Recommendations for CRF assessment in the screening of risk for SCD	285
Limitations and strengths	286

Abbreviations and acronyms: ACLS, Aerobics Center Longitudinal Study; AHA, American Heart Association; BMI, body mass index; BP, blood pressure; CHD, coronary heart disease; CRF, cardiorespiratory fitness; CV, cardiovascular; CVD, cardiovascular disease; ECP, exercise cardiac power; eCRF, estimated-CRF; HTN, hypertension; ICD, the International Classification of Diseases; KIHd, Kuopio Ischemic Heart Disease; LQTS, long QT syndrome; LVEF, left ventricular ejection fraction; MET, metabolic equivalent; peO_{2p}, peak exercise oxygen pulse; PRSIMA, Preferred Reporting Items for Systematic Reviews and Meta-Analysis; SCD, sudden cardiac death; VAs, ventricular arrhythmias; VO_{2max}, maximal oxygen consumption.

[☆] Statement of conflict of interest: see page 287.

* Address reprint requests to David Jiménez-Pavón, PhD, Ramón y Cajal Senior Researcher, MOVE-IT Research Group, University of Cádiz, Av. República Saharaui s/n, University Campus of Puerto Real, Puerto Real, CP.11519 Cádiz, Spain.

E-mail address: david.jimenez@uca.es (D. Jiménez-Pavón).

Conclusion	286
Funding	287
Statement of conflict of interest	287
References	287

Definition and relevance of sudden cardiac death

Sudden cardiac death (SCD) is recognized as an important cause of mortality, with a significant incidence among the general population, including the general population without previous symptoms of any coronary heart disease (CHD).^{1–3} The concept of SCD is generally defined as death within 1 h of symptom onset or during sleep in a patient who was previously stable.⁴ Although most of the total incidence of SCD has been reported to occur in men (70–90%), this cause of mortality has been documented in both sexes in individuals with a history of CHD or other major cardiovascular (CV) disease (CVD) risk factors and in those without a history of CHD or CVD.^{2,5} Consequently, SCD is an important public health problem, which constitutes a clinical challenge.

One of the American Heart Association's (AHA) 2020 Impact Goals⁶ is to decrease CVD mortality. Thus, prevention of SCD by detecting early risk factors⁷ could be a useful tool contributing to the AHA's goal through decreasing the incidence of SCD. The identification of risk factors for CVD would facilitate the large-scale screening of those participants at higher risk of SCD. In line with this, the best-known predictor of death is depressed left ventricular ejection fraction (LVEF), which has been associated with a dramatically increased risk of SCD.⁸ Nonetheless, LVEF is limited by low sensitivity, as the majority of people who suffer from SCD have a preserved LVEF.⁹ Moreover, several studies analyzed the effect of specific risk factors, such as QRS duration, QT interval and dispersion, signal-averaged electrocardiography, short- and long-term heart rate variability,⁹ systemic blood pressure (BP) and hypertension (HTN)¹⁰ and asymptomatic ST-segment depression¹¹ in men or a cluster of lifestyle factors in women,¹² on the risk of SCD. However, although numerous studies^{13–15} have attempted to identify those factors associated with higher SCD risk for the general population, conclusive evidence on this point remains somewhat elusive.

On the other hand, genetic background is known to partially explain SCD events.¹⁶ Particularly, recent studies of rare genetic arrhythmic disorders provided information on mechanisms and risk factors associated with the risk of SCD in patients without structural heart disease.¹⁶ Although data on the risk associated with QT prolongation were derived from long-term genotype–phenotype studies of patients with the congenital long QT syndrome (LQTS), importantly, these studies of LQTS have also increased the understanding of SCD in the general population. In addition, the association between QT interval prolongation and SCD has also been reported in individuals without known genetic predisposition,¹⁷ and highlight the fact that the focus for risk assessment for SCD at the population level should be not only focused on the overall QT interval but also on its individual components.¹⁶

Definition and importance of assessing cardiorespiratory fitness

Cardiorespiratory fitness (CRF) can be defined, briefly, as the body's ability to capture and utilize oxygen to perform any movement, physical activity, or exercise. But in particular, the level of CRF requires the involvement of numerous organs and systems among which are the respiratory, circulatory, CV, muscular, and nervous systems. Thus, CRF represents the integrated functional capacity of the human body in response to a stressful physiological stimulus, such as physical exercise. Consequently, the quantification of this capacity (CRF) can be an indicator of total body health.¹⁸ Moreover, it is well-known that roughly half of the variance in CRF is attributable to heritable factors¹⁹; on the other hand, the rest of the variance depends on the contribution of

behavioral patterns. Therefore, having the possibility to influence half of the variance in CRF by modifiable factors makes it very interesting to investigate the potential role of CRF on health status. In fact, since the 1950s the relationship of CRF with several health outcomes has been studied, but particularly during the past 20 years there has been an exponential increase in the number of studies published analyzing the role of CRF on mortality and health.^{18,20–24} In summary, it has been reported¹⁸ that CRF is a strong and independent indicator of risk for CVD- and all-cause mortality, even to be a more powerful predictor of mortality risk than traditional CVD risk factors, such as HTN, smoking, obesity, hyperlipidemia, and type 2 diabetes. Therefore, CRF has been shown to be a more powerful predictor of risk than other exercise testing variables, including ST-segment depression, symptoms, and hemodynamic responses.¹⁸ Most of these studies express CRF in the context of survival benefit per metabolic equivalent (MET), showing that each 1-MET increase in CRF was associated with considerable improvement in survival, ranging from 10% to 25%.¹⁸

Despite the considerable evidence that CRF is associated with many health outcomes and types of mortality, few investigations have been done on how CRF is related with just one particular type of mortality, such as SCD.^{25,26} There is no clear consensus on the role of CRF in SCD, neither on its power as a predictor of SCD nor compared with other risk factors or exercise test variables used to assess CVD. In addition, an assessment of the potential influence that each 1-MET increase in CRF has on the risk of SCD is needed, similar to the relationship between CRF and CVD events, CVD-mortality, and all-cause mortality.¹⁸

On the other hand, although CRF is now recognized as an important marker of CV health and all-cause mortality, it has been highlighted that it is currently the only major risk factor not routinely assessed in clinical practice.¹⁸ Thus, summarizing the evidence in relation to CRF and SCD at the population level would also help to support this international call for assessing CRF in clinical practice.

Methodology and search strategy

This systematic review was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement²⁷ and was registered through the International Prospective Register of Systematic Reviews (PROSPERO registration number: CRD42019130259).

Search strategy

A literature search was conducted in PubMed, EBSCOhost (Sportdiscus) and Scopus from their inception to March 11th, 2019. The search terms used to perform the present study were the following: “cardiorespiratory fitness” OR “aerobic capacity” OR “maximal oxygen consumption (VO_{2max})” AND “sudden cardiac death”. The reference lists of the retrieved studies were reviewed to identify additional studies.

Selection criteria

The inclusion criteria were: (i) type of study (longitudinal and prospective studies; at least 5 years follow-up), (ii) the use of CRF or proxy measure as independent or combined predictor of SCD, (iii) endpoint including SCD events or direct risk factor for SCD. There were no restrictions on age, gender, or health status. The exclusion criteria

were (i) not longitudinal studies or follow-up shorter than 5 years, (ii) studies written in languages other than English or Spanish, (iii) studies published in non-peer-reviewed journals, (iv) reviews/letters, and, (v) studies in animal models and (vi) not achieving inclusion criteria.

Data extraction

The following data were collected from each included study: 1) study/year; 2) cohort and design (cohort, follow-up, analysis); 3) participants [sample size, location, health condition, age, sex and BMI]; 4) CRF or proxy measure [test type and protocol, test maximal parameters (maximal oxygen consumption [VO_{2max}], METs, total time, workload, heart rate)]; 5) Combined factors with CRF (systolic BP, ST depression, sauna bathing, etc.); 6) SCD information (Sources, Codes, Number of events, proxy measure of risk SCD); 7) Role of CRF on SCD.

Meta-analysis

In order to quantify the independent role of CRF on SCD, we performed a meta-analysis using the two prospective cohorts analyzed in this review. For this purpose, we extracted the hazard ratio and its 95% CI from each included study. Therefore, pooled hazard ratio estimates were obtained for the risk of SCD using a random model due to the greater heterogeneity evaluated by the I^2 statistics.

Results

Overall results

Fig 1 presents the PRISMA consort diagram for the search strategy. The initial search retrieved 245 articles, and 11 manuscripts were finally

included after applying the inclusion and exclusion criteria. The first study regarding the role of CRF on SCD was in 2010.²⁵ After that, 9 studies were published between 2015 and 2018,^{26,28–35} whereas 1 additional study was published in 2019³⁶ focusing on serious ventricular arrhythmias (VAs) as risk factor for SCD.

The predictive role of CRF has been analyzed by two different cohorts^{25,26} (Table 1) accumulating a total of 57,824 participants (96% United States). The other 9 studies^{28–36} analyzed the role of both the independent influence of CRF or proxy and combined with other physiological CV parameters on the risk of SCD (Table 2).

The two population cohorts followed to assess the independent role of maximal CRF on SCD

The characteristics of the only 2 population cohorts^{25,26} available analyzed in the systematic review are shown in Table 1. The first prospective cohort analyzing the relevance of CRF as a predictive marker of SCD was the Kuopio Ischemic Heart Disease (KIHD) risk factor study in 2010²⁵ with a sample of 2368 Caucasian men from Europe followed for 17.6 years. Six years later, Jiménez-Pavón et al.²⁶ reported the influence of CRF on the risk of SCD in 55,456 American men and women from the Aerobics Center Longitudinal Study (ACLS) who were followed for 14.7 years. These are the only two published epidemiologic cohorts analyzing this relationship; thus, their similarities and differences need to be considered in order to better understand the final findings. In this sense, the first cohort studied accounts for only 4% of the total number of participants and was focused in European men, while the second cohort analyzed a mixed population of men and women from the United States accounting for the remaining 96% of total participants. However, the mean difference for age among the cohorts was <9 years and only approximately 1 kg/m² for BMI.




Fig 1. Flow diagram of the literature search and studies selection according PRISMA.

Table 1
Characteristics of the two population cohorts followed to assess the independent role of CRF on SCD.

Study/Year	Cohort and Design	Participants	Maximal CRF Assessment	SCD Events	Role of CRF on SCD
Laukkanen et al. ²⁵	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 17.6 years Analysis: Cox proportional hazards regression (hazard ratios-HRs)	Sample: n = 2368 men; Location: Caucasian-Europe; Health condition: General population Age: 52.9 years BMI: 26.9 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). CRF: mean VO _{2max} 30.2 ml/kg/min (8.6 METs). Mean Maximal Workload = 181 W.	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 146	-Inverse association -Risk of SCD decrease by 22% per 1 MET increase. -CRF ≥12 METs no SCD events.
Jiménez-Pavón et al. ²⁶	Prospective cohort: Aerobics Center Longitudinal Study (ACLS). Follow-up: 14.7 years Analysis: Cox proportional hazards regression (hazard ratios-HRs)	Sample: n = 55,456 (13,507 women) Location: United States of America Health condition: General population and sub-analyses for hypertensive, overweight and unhealthy individuals. Age: 44.2 years BMI: 25.7 kg/m ²	Test type: Maximal treadmill test. Protocol: Modified balke protocol using maximal heart rate (220-age). Mean CRF: 11.1 METs. L.CRF/total time: 8.2 METs/10.5 min M.CRF/total time: 10.2 METs/14.9 min H.CRF/total time: 12.9 METs/20.5 min	Source: National death index and death certificates. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 109.	-Inverse association. -M.CRF-H.CRF provide 44–48% lower risk. -In hypertensive, overweight and unhealthy individuals M.CRF-H.CRF provide 58–72% lower risk. -Risk of SCD decrease by 14% per 1 MET increase.

Abbreviations: BMI, body mass index; CRF, cardiorespiratory fitness; H.CRF, high CRF; ICD, International Classification of Diseases; L.CRF, Low CRF; M.CRF, moderate CRF; MET, metabolic equivalent (1 MET = 3.5 ml/kg/min); SCD, sudden cardiac death; VO_{2max}, maximal oxygen uptake; W, watts.

Another very important difference among these two studies was the methodology used to assess or estimate CRF level. In fact, the European cohort²⁵ used a maximal cycle ergometer test with direct analyses of respiratory gases providing a precise information of VO_{2max}, while the ACLS cohort²⁶ used a maximal treadmill test (modified Balke protocol) to estimate the maximal metabolic equivalents (METs; 1 MET = 3.5 ml oxygen uptake/kg per minute) from exercise duration and treadmill speed and incline, using standard algorithms. Using these measurements, the difference between the mean CRF levels of both cohorts was 2.5 METs. These are two well-known and valid aerobic modalities for incremental testing, with different strengths and limitations. The use of cycle ergometer is the most controlled way to perform exercise testing in relation to physiological response of variables, low risk of falls and being safe during the final steps. However, the treadmill test is the most widely used means of assessing exercise capacity in the United States and also recreates the natural skill if walking/running is used as a tool for population-based screenings.

On the other hand, in relation to SCD, the sources used for identifying the events and for definition were similar, being based on the International Classification of Diseases (ICD) codes in both studies. The number of SCD events were 146²⁵ and 109²⁶ for the European and American cohorts, respectively, which in relative terms implicates a much higher incidence among the European population.

Regarding the main finding about the role of CRF on SCD, both studies consistently reported an inverse and independent association after controlling for other CVD risk factors.^{25,26} Moreover, the European cohort found that the risk of SCD decreased by 22% per 1 MET increase, while the American Cohort showed a risk decrease of 14% per 1 MET increase. The difference in the estimated risk decrease per 1 MET increase among studies could be partially affected by the higher incidence of SCD events reported in the European population.²⁵ However, other differences between studies, such as sample size, gender, CRF test protocols or location, and associated behaviors could hamper a weighted or equilibrated comparison and interpretation of the real effect of CRF level on SCD.

The two studies provided additional findings specific for their population cohorts. In this line, the study of Laukkanen et al.²⁵ analyzed the role of reclassification as a measure of CRF. The numbers of men

reclassified because of assessment of CRF was an important additional measure with an almost equal number of case subjects had reclassification upward rather than downward by the addition of CRF to the risk model (11 vs. 12), although more control subjects had their risk reclassified upward than downward (103 vs. 73). Moreover, Laukkanen et al.²⁵ studied the CRF levels according to health status and reported mean VO_{2max} values of 32.5 ml/kg/min and 27.3 ml/kg/min for healthy and unhealthy men, respectively. They found that an increase in VO_{2max} of 1 MET was related to a 20% risk reduction for healthy men and a 25% risk reduction for men with previously known disease.

In the case of the study performed by Jiménez-Pavón et al.,²⁶ the risk of SCD across CRF levels by the presence or absence of HTN, normal weight or overweight or obese, and healthy or unhealthy groups was reported. They found inverse trends in the risk of SCD across CRF levels in all subgroups, but only reaching statistical significance in HTN (and the normotensive group with high CRF levels), overweight or obese, and healthy or unhealthy (moderate CRF levels) groups. Therefore, those participants identified with HTN and with moderate or high CRF levels had 65% and 72% lower risk of SCD, respectively, than did those with low CRF levels. Similarly, the risk of SCD was 58% and 64% lower in overweight or obese participants with moderate or high CRF levels, respectively. Finally, the authors reported in the unhealthy group that those participants with moderate CRF levels had 69% lower risk of SCD, whereas participants classified in the healthy group and with moderate or high CRF levels had 47 and 65%, respectively, lower risk of SCD than did those with low CRF levels.

Meta-analysis of the influence of CRF on SCD in two different cohorts

To provide more evidence about the direct influence of CRF on SCD, and despite the availability of the only 2 cohorts^{25,26} reporting these data, we performed a meta-analysis of these studies. The results of the meta-analysis for the independent effect of CRF on SCD showed a significant difference in favour of CRF on the risk of SCD (Hazard ratio = 0.813, 95% CI: 0.740, 0.894, p < 0.001, I² = 47.5%, total N = 57,824) (Fig 2). The meta-analysis suggests that despite the methodological and design disparities among the studies, the

Table 2

The combined role of maximal CRF or sub-maximal CRF on the risk of SCD.

Study/Year	Cohort and Design	Participants	Independent/Combined Predictors	SCD Events	Role of CRF on SCD
Kurl et al. ²⁸	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 20 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	Sample: n = 2358 men; Location: Caucasian-Europe Health condition: General population Age: 52.8 years BMI: 26.6 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). Combined predictors: ECP: mean ECP 11.9 ml/mmHg. LQ.ECP: <8.2 ml/mmHg HQ.ECP: >16.1 ml/mmHg	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 205	- Inverse independent and combined association - 1 SD increase in ECP and VO _{2max} (3.6 ml/mmHg and 635.9 ml/min) decreased risks of SCD by 36% and 33%. - Risk of SCD was 4.6 fold higher in participants in LQ.ECP compared with those with HQ.ECP
Hagnäs et al. ²⁹	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 20.9 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	Sample: n = 2328 men; Location: Caucasian-Europe Health condition: General population Age: 52.8 years BMI: 26.8 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w + step-by-step increase of 20 W/min (maximal heart rate; 220-age). Independent/combined predictors: CRF: mean 8.6 METs L.CRF: ≤ 8 METs H.CRF: ≥ 8 METs Exercise-induced ST depression: 10.3% (n = 239)	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 156	- Inverse independent and combined association - Risk of SCD was 1.6 fold higher in participants with L.CRF compared with those with H.CRF - Risk of SCD was 2.3 fold higher in participants with exercise-induced ST segment depression compared with those without. -Men with L.CRF and ST depression had 4.8 fold higher risk of SCD than men with H.CRF and without ST depression. - Inverse association
Kunutsor et al. ³⁰	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 25.6 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	Sample: n = 1663 men Location: Caucasian-Europe Health condition: General population Age: 52.2 years BMI: 27 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). CRF: mean VO _{2max} 2496 ml/min. Independent predictor: oxygen uptake at AT.	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 138	- 1 SD increase in oxygen uptake at AT (ml/min) decreased risks of SCD by 30%. -Higher quartile of oxygen uptake at AT (ml/min) provide 52% lower risk of SCD.
Kunutsor et al. ³¹	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 25.6 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	Sample: n = 1639 men Location: Caucasian-Europe Health condition: General population Age: 52.2 years BMI: 26.9 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w + step-by-step increase of 20 W/min (maximal heart rate; 220-age). Independent predictor: Mean VO ₂ at VT 23.72 ml/kg/min (6.78 METs).	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 121	- Inverse association - Risk of SCD decrease by 19% per 1 MET increase. - Higher quartile of oxygen uptake at VT (ml/min) provide 63% lower risk of SCD.
Laukkanen et al. ³²	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 26.1 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	Sample: n = 2291 men Location: Caucasian-Europe Health condition: General population Age: 53 years BMI: 26.9 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). CRF: mean VO _{2max} 30.3 ml/kg/min Combined Predictors: L.CRF: ≤ 30.02 ml/kg/min H.CRF: ≥ 30.03 ml/kg/min L.FSB: ≤ 2 sessions/week H.FSB: 3–7 sessions/week	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 226	- Inverse independent and combined association - H.CRF provide 52% lower risk of SCD. -H.FSB provide 33% lower risk of SCD. -Men with H.CRF and H.FSB had 69% lower risk of SCD than men with L.CRF and L.FSB.
Laukkanen	Prospective cohort:	Sample: n =	Test type: Maximal cycle	Source: Hospital documents, wards	- Inverse independent and combined

(continued on next page)

Table 2 (continued)

Study/Year	Cohort and Design	Participants	Independent/Combined Predictors	SCD Events	Role of CRF on SCD
et al. ³³	Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 26.1 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	2227 men; Location: Caucasian-Europe Health condition: General population Age: 53 years BMI: 26.9 kg/m ²	Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). Combined predictors: Peak exercise oxygen pulse(peO2p)**: 15.5 ml/beats	of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 220	association - 1 SD increase in relative peak exercise oxygen pulse was associated with decreased risks of SCD by 24%. - Higher quartile of relative peak exercise oxygen pulse provides 46% lower risk of SCD.
Jae et al. ³⁴	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 22 years Analysis: Cox proportional hazards regression (hazard ratios-HRs)	Sample: n = 2357 men Location: Caucasian-Europe Health condition: General population Age: 52.7 years BMI: 26.7 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). CRF: mean VO _{2max} 30.3 ml/kg/min Combined predictors: FIT.NW: ≥8.6 METs, <26.5 kg/m ² UFIT.OW/OB: <8.6 METs, ≥26.5 kg/m ²	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 253	- Inverse independent and combined association. - Risk of SCD was 1.80-fold higher in participants in UFIT.OW/OB compared with those in FIT.NW.
Hagnäs et al. ³⁵	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 19 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	Sample: n = 2656 men Location: Caucasian-Europe Health condition: General population Age: 53 years BMI: 26.9 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). CRF: mean VO _{2max} 30.1 ml/kg/min (8.6 METs). Combined predictors: H.CRF-H.LTPA: 9.9 METs L.CRF-L.LTPA: 6.4 METs	Source: Hospital documents, wards of health centers, death certificates, autopsy and medicolegal reports. Codes and definition: ICD with codes 978.1 and 146 from 9th and 10th revision. Number of SCD events: 205	- Inverse independent and combined association. - Risk of SCD decrease by 18% per 1 MET increase. - Risk of SCD was 1.6 and 1.4 folds higher in participants with L.CRF and L.LTPA (respectively) compared with those with H.CRF and H.LTPA. - Risk of SCD was 2.2 fold higher in participants in L.CRF-L.LTPA compared with those with H.CRF-H.LTPA.
Laukkanen et al. ³⁶	Prospective cohort: Kuopio Ischemic Heart Disease (KIHD) risk factor study. Follow-up: 25.3 years Analysis: Cox proportional hazards regression (Hazard ratios-HRs)	Sample: n = 2299 men Location: Caucasian-Europe Health condition: General population Age: 53 years BMI: 26.9 kg/m ²	Test type: Maximal cycle ergometer test. Protocol: 3 min warm-up 50w+ step-by-step increase of 20 W/min (maximal heart rate; 220-age). Independent predictor: CRF: mean VO _{2max} 30.3 ml/kg/min.	Source: National Hospital Discharge Register. Codes and definition: VAs defined by ICD with codes 427.41 and 147.2–149.0 from 9th and 10th revision, respectively. Number of VAs events: 73 serious VAs.	- 1 SD increase in CRF was associated with decreased risks of VAs by 36%. - High CRF (top tertile) provide 71% lower risk of VAs.

Abbreviations: AT, aerobic threshold; VT, ventilatory threshold; BMI, body mass index; CRF, cardiorespiratory fitness; ECP, Exercise cardiac power; H.CRF, high CRF; H.FSB, high frequency of sauna bathing; HQ.ECP, high quartile ECP; H.LTPA, high leisure time physical activity; ICD, International Classification of Diseases; FIT.NW, Fit and normal weight; L.CRF, Low CRF; L.FSB, low frequency of sauna bathing; LQ.ECP, Low Quartile ECP; L.LTPA, low leisure time physical activity; MET, metabolic equivalent (1 MET = 3.5 ml/kg/min); SCD, sudden cardiac death; SD, standard deviation; UFIT.OW/OB, unfit and overweight-obese; VAs, ventricular arrhythmias; VO_{2max}, maximal oxygen uptake; W, watts. * ECP calculated as ratio of measured VO_{2max} with peak SBP; ** Peak exercise oxygen pulse calculated by dividing the measured peak VO₂ by the maximum exercise heart rate.

influence of CRF is constant and relevant across at least two continents.

The combined role of maximal CRF or sub-maximal CRF on the risk of SCD

Since 2010, 10 additional studies have been identified that meet the criteria of this systematic review, and 9 out of 10 (90%) belong to the first cohort of European men^{28–36} (Table 2). Most of these studies analyzed the role of maximal CRF combined with several physiological

parameters to predict the risk of SCD,^{28,29,32–35} while 2 studies described the predictive validity of sub-maximal CRF values (corresponding to an aerobic threshold and ventilatory threshold) as independent predictors of SCD risk.^{30,31} Moreover, 1 recent work focused on the direct association of maximal CRF with VAs as a closely related risk factor of SCD.³⁶

The 6 studies^{28,29,32–35} from the European cohort showing the combined role of CRF on the risk of SCD had follow-ups ranging from 20 to 26 years. The principal extra value of these works is related to the new quantification of the preventive character of CRF when combined

with several physiological and CV variables. In this regard, CRF level was studied in combination with maximal systolic BP (Exercise cardiac power-ECP, calculated as ratio of measured VO_{2max} with peak SBP),²⁸ exercise-induced ST segment depression,²⁹ frequency of sauna bathing,³² maximum exercise heart rate (peak exercise oxygen pulse-peO2p),³³ body weight categories³⁴ and leisure time physical activity.³⁵ The combined influence of these parameters reported a change in the risk of SCD ranging from 46% to 380% depending on the parameters included (Fig 3), with the highest change in the risk of SCD attributed to the combination of CRF with the presence or absence of exercise-induced ST segment depression.²⁹

Sub-maximal CRF level indicators, such as the oxygen uptake at aerobic threshold and ventilatory threshold (both ml/min) have been also studied.^{30,31} The authors found that both are sub-maximal indicators of reduction in the risk of SCD, with 52% (aerobic threshold)³⁰ and 63% (ventilatory threshold)³¹ differences in the risk of SCD when comparing the highest and lowest quartiles. These findings regarding submaximal CRF indicators are quite similar to those reported for the same cohort with maximal CRF when comparing low CRF versus high CRF (risk increase of 1.6-fold).²⁹

Finally, Table 2 includes a recent study on the European cohort³⁶ showing the inverse association of maximal CRF with the incidence of VAs which is considered a principal risk factor of SCD. Particularly, 1 standard deviation increase in CRF was associated with decreased risk of VAs by 36%, and being in the highest CRF group (top tertile) predicted a 71% lower risk of VAs.

Discussion

Main findings

The studies included in this systematic review were based on the only 2 cohorts available reporting this relationship. We found 11 studies^{25,26,28–36} showing association between CRF or its proxy and the risk of SCD, but only 2 large prospective studies^{25,26} reported the independent and direct role of maximal CRF level on risk of SCD, each one from one continent (Europe and North America). Consequently, our additional meta-analysis, based only on these 2 different cohorts, showed a significant difference in favour of higher CRF in the risk of SCD, suggesting that despite the methodological and design disparities among studies, the influence of CRF is constant and relevant across continents, and their findings support and complement each other. In fact, our meta-analysis reported an overall decreased risk of SCD of ~19% per each MET increase, which is in line with that suggested in the original studies from both cohorts (14% and 22%, respectively)^{25,26} and among the range of improvement in survival for mortality previously reported.¹⁸

In addition, we found that the combination of maximal CRF with several physiological parameters derived from an exercise test (e.g., systolic BP and maximal test heart rate) or some behaviors (e.g., sauna bathing or leisure time physical activity) also constituted very good predictors of the risk of SCD, in some cases being even better

predictors than CRF alone. On the other hand, the use of only CRF at sub-maximal intensity provides similar estimation of the risk decrease for SCD.

Recommendations for CRF assessment in the screening of risk for SCD

The overall importance to periodically obtain a directly measured VO_{2max} has been recently highlighted,¹⁸ although CRF estimated from the peak work rate achieved on a treadmill or a cycle ergometer or from non-exercise algorithms was also recognized as relevant, necessary and possibly more cost-effective. From the findings of the current systematic review, the capacity of both measured VO_{2max} and estimated CRF from peak workload showed good sensitivity as risk factor of SCD using cycle ergometer²⁵ and treadmill,²⁶ respectively. Particularly, one of the cohorts²⁵ reported that the ability of CRF to improve discrimination and risk classification of SCD did not significantly change when using directly or estimated CRF.³⁷ Thus, the main recommendation for assessing CRF would be to use one of these methods (estimated from workload or even non-exercise equations) as directly measured VO_{2max} may not be feasible or cost-effective in large populations or in routine clinical practices. In fact, assessing estimated-CRF (eCRF) from readily available clinical or research data base information has been suggested as a good alternative to objective CRF measurement,^{18,38} as it can be quickly and easily collected in epidemiological studies with good correlations with measured CRF and favorable predictive value for CVD events and mortality. Unfortunately, specific data on the ability of eCRF to discriminate the risk of SCD have not yet been published, therefore such data could not be included in our systematic review. However, based on the currently reviewed data and on the utility previously noted for eCRF in relation to other health outcomes and mortality types,^{38–40} it would be expected that future studies will also corroborate its utility in relation to SCD.

Additionally, when maximal exercise testing is not possible for reasons of safety, health, or cost, the utility of sub-maximal parameters, such as aerobic or ventilatory thresholds, have been analyzed in relation to SCD^{30,31}; and these parameters could be an alternative to full exercise testing with overall similar results. However, to highlight the fact that the parameters used as indicators of aerobic threshold and ventilatory threshold were oxygen uptake, gas exchange analysis would still be required. Future studies should add the estimation of hazard-ratio not only for the direct value of oxygen uptake at aerobic or ventilatory thresholds, but also the equivalent of percentage of heart rate reserve, exercise time or workload in order to provide evidence regarding other parameters that can be assessed without the necessity of gas exchange analysis.

Another important recommendation is to consider, when possible, the combination of CRF with other predictors which have been shown to provide extra or additional predictive value (Fig 3). Particularly, the combination of CRF with exercise-induced ST depression²⁹ or with maximal systolic BP (ECP)²⁸ reported even better assessment of the risk of SCD. These will be of great interest for those projects in which electrocardiogram or BP could be properly recorded during the maximal




Fig 2. Forest plot of the risk of SCD by CRF measured as 1-MET increase. CI: confidence intervals.




Fig 3. Combined role of CRF on decrease/increase of the risk of SCD where blue bars indicates decreased risk and red bars indicates increased risk (Abbreviations: H.CRF, high CRF; H.FSB, high frequency of sauna bathing; HQ.ECP, high quartile exercise cardiac power; HQ.Peo2p, high quartile peak exercise oxygen pulse; H.LTPA, high leisure time physical activity; FIT.NW, Fit and normal weight; L.CRF, Low cardiorespiratory fitness; L.FSB, low frequency of sauna bathing; LQ.ECP, Low Quartile exercise cardiac power; LQ.Peo2p, low quartile peak exercise oxygen pulse; LLTPA, low leisure time physical activity; UFIT.OW/OB, unfit and overweight-obese).

exercise test. However, an alternative combination for large epidemiological studies not able to include more precise technology would be to add the physical activity level as behavior combined with CRF that substantially increase the prediction level of the risk of SCD.³⁵ In general, it seems possible that a combined panel of measurements, including CRF and physiological/behavioral measures will have substantially improved degrees of discrimination and reclassification.

Limitations and strengths

This study presents several limitations. First, although two major cohorts were identified with consistent data about the topic of interest, the lack of availability of other cohorts or population studies addressing this relationship hampers the extrapolation, partially, of the current findings. Second, the natural differences among the 2 cohorts^{25,26} included in this systematic review, such as number of participants, gender, cycle ergometer versus treadmill or direct assessment versus MET estimation, certainly would require more studies to clarify some issues. Third, the fact that many of the results and conclusions are derived from studies based on the same prospective cohort could imply a methodological bias; therefore, more studies from different populations are needed to confirm part of the findings demonstrated in this review. Additionally, the longitudinal design of the included studies does not allow one to establish a direct causality of increasing CRF on decreasing risk of

SCD versus merely an association, and this would require intervention studies for further clarification.

Despite these limitations, to the best of our knowledge, this is the first systematic review with additional meta-analysis which focused on the role of CRF level and the risk of SCD, as well as showing data from 2 representative continents. The additional meta-analysis performed should be acknowledged as, despite the inclusion of only 2 studies, the output confirms a similar trend within these quite different prospective cohorts. Moreover, the constellation of data that we reviewed suggests a powerful association and potential role of higher CRF to markedly reduce the risk of SCD.

Conclusion

Our systematic review provides evidence, on the long-term influence of CRF as a risk factor of SCD. Few studies have assessed the relevance of CRF as an independent predictor of SCD at the population level, but the message from these studies is consistent. The independent ability of higher CRF to predict lower risk of SCD has been confirmed, but the ability of CRF to predict SCD is improved further when combined with some other behavioral or physiological parameters (most significantly, ST-segment depression on exercise testing).

Collectively, from a clinical perspective, our review supports the use of CRF as part of the early and initial screening in adult life for assessing

the risk of SCD, in line with recommendations for prediction of CVD- and all-cause mortality. Moreover, from a public health point of view, promoting the increase of CRF early in life and throughout adolescence and adult life to enhance long-term survival is important. However, additional research is needed to add novel insights on the specific role of CRF in women, different population cohorts, and lower-upper threshold values in order to improve the early risk stratification for SCD.

Funding

Current research activities of Dr. Jiménez-Pavón is supported by a grant from the Spanish Ministry of Science and Innovation - MINECO (RYC-2014-16938) and the Spanish Ministry of Economy and Competitiveness - MINECO/FEDER (DEP2016-76123-R); the Government of Andalusian, Integrated Territorial Initiative 2014-2020 for the province of Cádiz (PI-0002-2017); the European Union's ERASMUS+ SPORT programme (grant agreement: 603121-EPP-1-2018-1-ES-SPO-SCP); and the EXERNET Research Network on Exercise and Health in Special Populations (DEP2005-00046/ACTI).

Statement of conflict of interest

None of the authors have any conflicts of interests with regard to this publication.

References

- Chugh SS, Reinier K, Teodorescu C, et al. Epidemiology of sudden cardiac death: clinical and research implications. *Prog Cardiovasc Dis* 2008;51(3):213-228.
- Roger VL, Go AS, Lloyd-Jones DM, et al. American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics 2012 update: a report from the American Heart Association. *Circulation* 2012;125(1):e2-e220.
- Mahmoud KD, de Smet BJ, Zijlstra F, Rihal CS, Holmes DR Jr. Sudden cardiac death: epidemiology, circadian variation, and triggers. *Curr Probl Cardiol* 2011;36(2):56-80.
- Wilhelm M, Bolliger SA, Bartsch C, et al. Sudden cardiac death in forensic medicine - Swiss recommendations for a multidisciplinary approach. *Swiss Med Wkly* 2015;145, w14129.
- Fox CS, Evans JC, Larson MG, Kannel WB, Levy D. Temporal trends in coronary heart disease mortality and sudden cardiac death from 1950 to 1999: the Framingham Heart Study. *Circulation* 2004;110(5):522-527.
- Lloyd-Jones DM, Hong Y, Labarthe D, et al. American Heart Association Strategic Planning Task Force and Statistics Committee. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation* 2010;121(4):586-613.
- Adabag AS, Luepker RV, Roger VL, Gersh BJ. Sudden cardiac death: epidemiology and risk factors. *Nat Rev Cardiol* 2010;7(4):216-225.
- Bailey JJ, Berson AS, Handelsman H, Hodges M. Utility of current risk stratification tests for predicting major arrhythmic events after myocardial infarction. *J Am Coll Cardiol* 2001;38:1902-1911.
- Goldberger JJ, Cain ME, Hohnloser SH, et al. American Heart Association/American College of Cardiology Foundation/Heart Rhythm Society scientific statement on noninvasive risk stratification techniques for identifying patients at risk for sudden cardiac death: a scientific statement from the American Heart Association Council on Clinical Cardiology Committee on Electrocardiography and Arrhythmias and Council on Epidemiology and Prevention. *J Am Coll Cardiol* 2008;52:1179-1199.
- Laukkanen JA, Jennings JR, Kauhanen J, Mäkilä TH, Ronkainen K, Kurl S. Relation of systemic blood pressure to sudden cardiac death. *Am J Cardiol* 2012;110(3):378-382.
- Laukkanen JA, Mäkilä TH, Rauramaa R, Kurl S. Asymptomatic ST-segment depression during exercise testing and the risk of sudden cardiac death in middle-aged men: a population-based follow-up study. *Eur Heart J* 2009;30(5):558-565.
- Chiuvè SE, Fung TT, Rexrode KM, et al. Adherence to a low risk, healthy lifestyle and risk of sudden cardiac death among women. *JAMA* 2011;306(1):62-69.
- Greenland P, Smith Jr SC, Grundy SM. Improving coronary heart disease risk assessment in asymptomatic people: role of traditional risk factors and noninvasive cardiovascular tests. *Circulation* 2001;104(15):1863-1867.
- Müller D, Agrawal R, Arntz HR. How sudden is sudden cardiac death? *Circulation* 2006;114(11):1146-1150.
- Huikuri HV, Castellanos A, Myerburg RJ. Sudden death due to cardiac arrhythmias. *N Engl J Med* 2001;345(20):1473-1482.
- Goldenberg I, Moss AJ. Risk stratification for sudden cardiac death in individuals without structural disease: implications from studies of rare genetic arrhythmic disorders. *Circ Arrhythm Electrophysiol* 2017;10(10). pii: e005774. doi: <https://doi.org/10.1161/CIRCEP.117.005774>.
- Algra A, Tijssen JG, Roelandt JR, Pool J, Lubsen J. QTc prolongation measured by standard 12-lead electrocardiography is an independent risk factor for sudden death due to cardiac arrest. *Circulation* 1991;83:1888-1894.
- Ross R, Blair SN, Arena R, et al. Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American Heart Association. *Circulation* 2016;134:e653-e699.
- Bouchard C, Daw EW, Rice T, et al. Familial resemblance for VO_{2max} in the sedentary state: the HERITAGE family study. *Med Sci Sports Exerc* 1998;30:252-258.
- Myers J, McAuley P, Lavie CJ, Despres JP, Arena R, Kokkinos P. Physical activity and cardiorespiratory fitness as major markers of cardiovascular risk: their independent and interwoven importance to health status. *Prog Cardiovasc Dis* 2015;57:306-314. <https://doi.org/10.1016/j.pcad.2014.09.011>.
- Blair SN. Physical inactivity: the biggest public health problem of the 21st century. *Br J Sports Med* 2009;43:1-2.
- Kokkinos P, Myers J. Exercise and physical activity: clinical outcomes and applications. *Circulation* 2010;122:1637-1648. <https://doi.org/10.1161/CIRCULATIONAHA.110.948349>.
- Swift DL, Lavie CJ, Johannsen NM, et al. Physical activity, cardiorespiratory fitness, and exercise training in primary and secondary coronary prevention. *Circ J* 2013;77:281-292.
- Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA* 2009;301:2024-2035. <https://doi.org/10.1001/jama.2009.681>.
- Laukkanen JA, Mäkilä TH, Rauramaa R, Kiviniemi V, Ronkainen K, Kurl S. Cardiorespiratory fitness is related to the risk of sudden cardiac death: a population-based follow-up study. *J Am Coll Cardiol* 2010;56(18):1476-1483.
- Jiménez-Pavón D, Artero EG, Lee D-C, et al. Cardiorespiratory fitness and risk of sudden cardiac death in men and women in the United States: a prospective evaluation from the aerobics center longitudinal study. *Mayo Clin Proc* 2016;91(7):849-857.
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 2009;62(10):e1-34. <https://doi.org/10.1016/j.jclinepi.2009.06.006>.
- Kurl S, Jae SY, Kauhanen J, Ronkainen K, Rauramaa R, Laukkanen JA. Exercise cardiac power and the risk of sudden cardiac death in a long-term prospective study. *Int J Cardiol* 2015;181:155-159.
- Hagnas MJ, Lakka TA, Kurl S, et al. Cardiorespiratory fitness and exercise-induced ST segment depression in assessing the risk of sudden cardiac death in men. *Heart* 2017;103(5):383-389.
- Kunutsor SK, Kurl S, Khan H, Zaccardi F, Rauramaa R, Laukkanen JA. Oxygen uptake at aerobic threshold is inversely associated with fatal cardiovascular and all-cause mortality events. *Ann Med* 2017;49(8):698-709.
- Kunutsor SK, Kurl S, Khan H, Zaccardi F, Laukkanen JA. Associations of cardiovascular and all-cause mortality events with oxygen uptake at ventilatory threshold. *Int J Cardiol* 2017;236:444-450.
- Laukkanen JA, Laukkanen T, Khan H, Babar M, Kunutsor SK. Combined effect of sauna bathing and cardiorespiratory fitness on the risk of sudden cardiac deaths in Caucasian men: a long-term prospective cohort study. *Prog Cardiovasc Dis* 2018;60(6):635-641.
- Laukkanen JA, Araújo CGS, Kurl S, et al. Relative peak exercise oxygen pulse is related to sudden cardiac death, cardiovascular and all-cause mortality in middle-aged men. *Eur J Prev Cardiol* 2018;25(7):772-782.
- Jae SY, Franklin BA, Kurl S, et al. Effect of cardiorespiratory fitness on risk of sudden cardiac death in overweight/obese men aged 42 to 60 years. *Am J Cardiol* 2018;122(5):775-779.
- Hagnas MJ, Lakka TA, Mäkilä TH, et al. High leisure-time physical activity is associated with reduced risk of sudden cardiac death among men with low cardiorespiratory fitness. *Can J Cardiol* 2018;34(3):288-294.
- Laukkanen JA, Lavie CJ, Khan H, Kurl S, Kunutsor SK. Cardiorespiratory fitness and the risk of serious ventricular arrhythmias: a prospective cohort study. *Mayo Clin Proc* 2019;94(5):833-841. <https://doi.org/10.1016/j.mayocp.2018.11.027>.
- Lauer MS. Risk stratification for sudden cardiac death: a puzzle beyond p values. *J Am Coll Cardiol* 2010;56(18):1484-5. doi: <https://doi.org/10.1016/j.jacc.2010.05.047>.
- Gander JC, Sui X, Hébert JR, et al. Addition of estimated cardiorespiratory fitness to the clinical assessment of 10-year coronary heart disease risk in asymptomatic men. *Prev Med Rep* 2017;7:30-37.
- Nauman J, Tauschek LC, Kaminsky LA, Nes BM, Wisloff U. Global fitness levels: findings from a Web-based surveillance report. *Prog Cardiovasc Dis* 2017;60:78-88.
- Artero EG, Jackson AS, Sui X, et al. Longitudinal algorithms to estimate cardiorespiratory fitness: associations with nonfatal cardiovascular disease and disease-specific mortality. *J Am Coll Cardiol* 2014;63:2289-2296.