



Oral health and dental status in people with epilepsy in rural China

Mengjie Wang^a, Ding Ding^{b,*}, Qing Zhang^{c,**}, Guoxing Zhu^{a,*}, Yan Ge^a, Bin Yang^d,
Taiping Wang^e, Patrick Kwan^f, Wenzhi Wang^g, Zhen Hong^b, Josemir W. Sander^{h,i,j}

^a Department of Neurology, Huashan Hospital, Fudan University, Shanghai, China

^b Institute of Neurology, WHO Collaborating Center for Research and Training in Neurosciences, Huashan Hospital, Fudan University, Shanghai, China

^c Department of Neurology, Ningxia Medical University Affiliated Hospital, Yinchuan, China

^d Jiaozuo People's Hospital, Jiaozuo, Henan, China

^e Jincheng Emergency Center, Shanxi Province, China

^f Department of Neuroscience, Monash University, Alfred Hospital, Melbourne, Victoria, Australia

^g Beijing Neurosurgical Institute, Capital Medical University, Beijing, China

^h NIHR University College London Hospitals Biomedical Research Centre, UCL Queen Square Institute of Neurology, Queen Square, London, WC1N 3BG, United Kingdom

ⁱ Chalfont Centre for Epilepsy, Chalfont St Peter, SL9 0RJ, United Kingdom

^j Stichting Epilepsie Instellingen Nederland (SEIN), Achterweg 5, Heemstede, 2103 SW, the Netherlands

ARTICLE INFO

Keywords:

Seizure frequency
Dentistry
Buccal hygiene
Oral sampling
Dental caries
Tooth loss

ABSTRACT

Purpose: We evaluated the oral health and dental status of people with epilepsy, and their relationship to seizure frequency, in a community cohort in rural China.

Methods: A cross-sectional study of people with epilepsy was carried out in areas in Henan, Shanxi and Ningxia provinces of China. All participants underwent a specially designed “Oral health and Dental status Questionnaire”. Univariate and multivariate analyses were used to explore the relationships between seizure frequency and oral health and dental status.

Results: A total of 875 people participated. Almost two thirds (65.8%) reported brushing their teeth at least once a day but 634 (73.0%) brushed their teeth for less than 3 min each time. Only 80 (9.1%) had visited a dental clinic in the previous year. Multivariate logistic regression analysis showed that having 1–4 seizures/month (OR 0.60, 95%CI 0.40–0.90) or > 4 seizures/month (OR 0.38, 95%CI 0.22–0.66) was associated with decreased odds of brushing teeth at least once a day, and higher seizure frequency was associated with increased odds of dental injury due to seizures (OR 2.07, 95%CI 1.22–3.50 for < 1 seizure/month; OR 2.12, 95%CI 1.25–3.58 for 1–4 seizures/month; OR 3.09, 95%CI 1.57–6.07 for > 4 seizures/month).

Conclusion: Seizure frequency was significantly associated with seizure-related dental injury, and with the lack of good oral health practice. Improvement in the oral health of people with epilepsy in resource-poor areas should be encouraged.

1. Introduction

Epilepsy is characterized by recurrent seizures [1]. It is the second most common neurological condition in China with a prevalence of around 0.7%; this translates into just over 9 million people with epilepsy among whom an estimated 6 million have active epilepsy [2]. People affected by epilepsy have a significantly worse quality of life as well as more compromised physical and psychological health levels than the general population [3,4].

Oral health, as defined by WHO, is a condition of being free from chronic oro-facial ache, oral sores, dental caries and tooth loss,

periodontal diseases, or other factors that affect the well-being of the oral cavity [5]. Oral health, influencing the whole body directly or indirectly, is one determinant factor for quality of life [6,7]. Oral diseases can destroy the hard tissues of the teeth and the supporting tissues around the teeth, thus impairing not only the functions of chewing, speech, and aesthetics, but also social interaction and psychological state [5]. A close interrelationship has been demonstrated between oral and general health [6,7]. Some microorganisms in the oral cavity may, over time, cause or exacerbate certain systemic diseases such as diabetes or coronary heart disease [7,8].

Most existing oral health-related studies in epilepsy focus on

* Corresponding authors at: Huashan Hospital, 12 Middle Wulumuqi Rd., Shanghai 200040, China.

** Corresponding author at: General Hospital of Ningxia Medical University, 804 Shengli N Street, Yinchuan, Ningxia Hui Autonomic Region, 750004, China.

E-mail addresses: dingding@huashan.org.cn (D. Ding), nxzhangqing@aliyun.com (Q. Zhang), zhuguoxing@huashan.org.cn (G. Zhu).

gingival hyperplasia induced by anti-epileptic drugs (AEDs) [9–11]. Previous reports showed that people with epilepsy have poorer oral health status and awareness than people without epilepsy [12,13]. There is a dearth of data concerning oral health and dental status of people with epilepsy in resource-poor areas.

During a study involving the acquisition of DNA samples from people with epilepsy in rural communities in China, we aimed to obtain oral samples using swabs or saliva. During a pilot study, it became clear that this would not yield usable samples, largely due to contamination from oral flora. This led us to evaluate the oral health and dental status of people with epilepsy, as well as their relationships to seizure frequency, in these communities.

2. Methods

2.1. Study site and participants

Permanent residents in the study sites were eligible to participate. People with epilepsy previously recruited within the study “Validation of Clinical Assessment Tools for Population Genetic Studies of Epilepsy (1R21NS069223-01, NIH/NINDS)” from rural communities in Henan, Shanxi, and Ningxia provinces in China were asked to participate. Subjects were excluded for the following reasons: 1) seizures resulting from progressive systemic or neurological disorders, 2) nonepileptic attack disorder, 3) seizures related to drug or alcohol abuse, 4) lack of adequate assessment of seizure control or of whether the individuals adhere to their drug regime, 5) pregnancy, 6) presence of severe cognition or motor impairments and 7) inability to provide consent (from individuals or their guardians).

2.2. Clinical assessment and diagnosis of epilepsy

Clinical assessment was done in two steps. In step 1, trained village physicians interviewed individuals who had history of seizures. Family members who had witnessed seizures or were familiar with their medical history were also interviewed. In step 2, neurologists from the designated provincial epilepsy centre reviewed individuals who had been identified in step 1 and confirmed or refuted the diagnosis of epilepsy. This was based on clinical symptoms and on a comprehensive history including details of any potential risk factors for epilepsy (such as history of status epilepticus, febrile seizures, history of central nervous system infections, brain injury, stroke, and brain tumour, and family history of epilepsy (in close relatives), abnormal birth history), and results of neuro-imaging and EEGs which were available during this step. A group of experienced neurologists with a special interest in epilepsy (PK, JWS, and ZH), reviewed the diagnoses and any conflicts were resolved by consensus. Data collection included: 1) demographics; 2) age of onset; 3) seizure classification (focal without secondary generalization, focal with secondary generalization, or generalized, based on ILAE guidelines [14]); 4) seizure frequency (categorized as seizure free, < 1 seizure/month, 1–4 seizures/month, or > 4 seizures/month); 5) putative aetiology of epilepsy (classified as structural/metabolic, genetic, or unknown) [14]; 6) treatment history (whether or not the individual had had more than one year of AED treatment).

2.3. Questionnaire of oral health and dental status

Village physicians, who had been provided with basic training in oral health by experienced dentists, interviewed participants or their relatives using a specifically designed “Oral health and Dental status Questionnaire (ODQ)” (see Box 1). Questions addressed tooth brushing habits, visits to dental clinics, oral symptoms and diagnosis, number of caries, history of dental injury, tooth loss, and dental restoration experience, based on experts’ opinions from dentists and epileptologists in China.

2.4. Standard protocol approvals, registrations, and informed consent

The initial study was approved by the joint Chinese University of Hong Kong-New Territories East Cluster Research Ethics Committee and by the Ethics Committee of the Beijing Neurosurgical Institute in China. The oral health survey was approved by the Medical Ethics Committee of Huashan Hospital Fudan University, Shanghai, China. All participants provided informed consent; where applicable participants’ legally acceptable guardians provided assent.

2.5. Statistical approach

Continuous variables were expressed as median (minimum, maximum), and categorical variables as counts and frequencies (%). The Chi-square test was used for comparing categorical variables. One-way ANOVA or the Kruskal-Wallis test were used to compare continuous variables among 4 groups (seizure-free, < 1 seizure/month, 1–4 seizures/month, > 4 seizures/month) according to whether or not the data were normally distributed. Unadjusted and adjusted odds ratios and 95% confidence intervals (95% CI) were calculated using univariate and multivariate logistic regression models in order to explore the association between seizure frequency (seizure-free group as the reference) and oral health and dental status. Multivariate logistic regression was fitted, adjusting for potential confounders based on previous literatures and experts’ clinical experience. The confounders included age (as a continuous variable), age of onset (as a continuous variable), sex, aetiology of epilepsy (genetic, structural/metabolic, and unknown), seizure type (focal without secondary generalization, focal with secondary generalization, and generalized), and more than one year’s treatment with antiepileptic drugs (AED) (yes, no).

All p-values and 95% CIs were estimated in a two-tailed fashion. Differences were considered to be statistically significant at $p < 0.05$. Data were analyzed using SPSS 22.0 (SPSS Inc., IL, USA).

3. Results

3.1. Demographics and clinical characteristics of participants

A total of 875 people with epilepsy were included and their demographics and clinical details are given in Table 1. The majority (805; 92.0%) had taken AEDs for more than a year. Almost half (402; 46.0%) had seizures at least once a month and only 206 (23.5%) were seizure-free.

3.2. Oral health and dental status by seizure frequency

Two thirds (576; 65.8%) reported brushing teeth at least once a day and the remainder did not brush regularly (Table 2). The majority (634; 73.0%) brushed their teeth for less than 3 min each time. Less than ten percent (80; 9.1%) had visited a dental clinic in the previous year. Just under half (395; 45.1%) had dental symptoms, of whom 16.7% had gingival hyperplasia. Fifteen people of the 130 taking phenytoin had gingival hyperplasia. Nearly forty percent of participants (344; 39.3%) reported having caries. Only sixty-two (7.1%) had ever been formally diagnosed with a dental disease. Overall, 225 (25.7%) had missing teeth due to caries, and almost one fifth (19.0%) had injured teeth as a result of seizures.

Table 2 suggests that frequency of brushing teeth ($p = 0.011$), whether caries were present ($p = 0.027$), missing teeth due to seizures ($p = 0.025$), and dental injury due to seizures ($p = 0.044$) was statistically significant among four groups with different severity of seizure frequency (seizure-free, < 1 seizure/month, 1–4 seizures/month, and > 4 seizures/month).

Box 1

Oral health and dental status questionnaire.

Oral Health and Dental Status QUESTIONNAIRE (ODQ).

1 Do you brush your teeth regularly?

 0. No 1. Yes, brush teeth ___ a day, about ___ minutes each time

2 How many times did you visit dental clinics due to oral problems in the recent year:

___ times.

3 Do you have the following symptoms frequently:

1) Toothache: 0.No 1.Yes 2) Bleeding gums: 0.No 1.Yes
3) Swollen gums: 0.No 1.Yes 4) Gingival hyperplasia: 0.No 1.Yes
5) Bad breath: 0.No 1.Yes 6) Others: 0.No 1.Yes ___

4 Do you have caries?

 0.No 1.Yes, ___ number

5 Do you have any tooth detached or extracted due to caries?

 0.No 1.Yes, ___ number

6 Have you ever suffered from dental injury due to seizure?

 0.No 1.Yes, ___ number, fracture ____, completely detached or extracted ___

7 Have you ever suffered from dental injury due to other reasons?

 0.No 1.Yes, ___ number, fracture ____, completely detached or extracted ___

8 Apart from the change of teeth as a child, have you ever had natural tooth loss?

 0.No 1.Yes, ___ number

9 Have you had your tooth repaired after dental injury or tooth loss?

 0.No 1.Yes 1) Dental crown 2) Dental prosthesis 3) Others ___

10 Have you ever been diagnosed by your dentists for:

1) Pulpal and Periapical Disease: 0.No 1.Yes 2) Gingivitis: 0.No 1.Yes
3) Periodontitis: 0.No 1.Yes 4) Others: 0.No 1.Yes ___**3.3. Unadjusted and adjusted odds ratios for factors related to seizure frequency using logistic regression models**

Table 3 shows that, after adjusting for the confounding factors (age, age of onset, sex, aetiology of epilepsy, seizure type, and whether individuals had had more than one year of AED treatment), frequent seizures [1–4 seizures/month (OR 0.60, 95%CI 0.40–0.90) and > 4 seizures/month (OR 0.38, 95%CI 0.22–0.66)] were associated with decreased odds of brushing teeth at least once a day. Higher seizure frequency was associated with increased odds of dental injury due to seizures (OR 2.07, 95%CI 1.22–3.50 for < 1 seizure/month; OR 2.12, 95%CI 1.25–3.58 for 1–4 seizures/month; OR 3.09, 95%CI 1.57–6.07 for > 4 seizures/month). The lowest seizure frequency (< 1 seizure/month) was associated with a lower probability of caries than those seizure free (OR 0.61, 95%CI 0.41–0.91).

4. Discussion

This is one of the first community-based attempts to evaluate the oral health and dental status of people with epilepsy in resource-poor areas. Our findings suggest that people with epilepsy in rural China have poor oral health and are vulnerable to dental injury. There seems to be a significant association between seizure frequency and regularity of brushing teeth and seizure-related dental injury using multivariate logistic regression analyses.

Oral health is a significant part of physical and mental health but it is often neglected, particularly in people with epilepsy. Our findings show that fewer than 1 in 10 had visited a clinic in the previous year, which is lower than the approximately 16% attendance at dental clinics reported in rural areas [15]. We also found that less than one third brushed their teeth for at least 3 min each time and around one third did not brush teeth regularly. Participants with higher seizure

Table 1
Demographics and clinical characteristics of people with epilepsy.

	People with epilepsy n = 875
Age (years), median (range)	41 (8–9)
Age of onset (years), median (range)	14 (0–77)
Gender	
Male, n (%)	529 (60.5)
Aetiology	
Genetic, n (%)	158 (18.1)
Structural/Metabolic, n (%)	251 (28.7)
Unknown, n (%)	466 (53.3)
Seizure type	
Focal without secondary generalization, n (%)	62 (7.1)
Focal with secondary generalization, n (%)	467 (53.4)
Generalized, n (%)	346 (39.5)
Having more than one year of AED treatment, n (%)	805 (92.0)
Seizure frequency	
Seizure free, n (%)	206 (23.5)
< 1 seizure/month, n (%)	267 (30.5)
1–4 seizures/month, n (%)	313 (35.8)
> 4 seizures/month, n (%)	89 (10.2)

frequency were less likely to brush their teeth regularly and this may be due to the fact that people with more severe epilepsy are less likely to self-care [16]. It is possible that oral health may be improved through public oral hygiene education and perhaps also by effective control of the individuals' seizures [13]. The present finding may imply a lack of

awareness of oral health. People with epilepsy may not deem oral issues a high priority, unless they have severe pain or discomfort. The lower socioeconomic status of people with epilepsy, especially those with recurrent seizures, may also partly account for the relatively lower frequency of dental visits.

About a fifth had experienced at least a seizure-related dental injury. Previous studies found a higher risk of dental trauma in people with epilepsy than in the general population [17–20]. Dental injury has been reported as the most common hard tissue injury resulting from seizures [21]. Previous studies have also related seizure frequency with increased dental injuries [3,20]. A Brazilian study showed that dental injury was more common in people with epilepsy, affecting up to a third, but was even more common if the person did not have a useful aura prior to seizures [22]. The incidence of dental injury in Nigeria was also much more common amongst people with epilepsy than in the general population [13,23]. The unpredictability of seizures may lead to falls and accidents causing serious physical injury [24,25]. In many settings, it is not uncommon for a person having a seizure to have a spoon or other object placed in the mouth; this may lead to dental injury [17,26]. The powerful contraction of the masticatory muscles during convulsions may also play a role in dental injuries [12]. Convulsive seizures often result in minor oral lesions, such as bitten tongues, but also frequently cause dental injuries and, under some conditions, traumatic oro-facial injuries [27].

All participants on AED treatment were using cheaper AEDs, such as phenobarbital or phenytoin; gingival hyperplasia is a common complication, particularly of phenytoin treatment. The incidence of

Table 2
Oral health and dental status of the study participants by severity of seizure frequency.

	Total (N = 875)	Seizure frequency				P value
		Seizure free (N = 206)	< 1 seizure/month (N = 267)	1–4 seizures/month (N = 313)	> 4 seizures/month (N = 89)	
Brush teeth regularly						
No, n (%)	299(34.2)	59(28.6)	87(32.6)	110(35.1)	43(48.3)	0.011
> =once, n (%)	576(65.8)	147(71.3)	180(67.5)	203(64.8)	46(51.7)	
Brushing time ^a						
< 3 min, n (%)	634(73.0)	144(70.6)	196(74.5)	226(72.4)	68(76.4)	0.685
> = 3 min, n (%)	234(27.0)	60(29.4)	67(25.5)	86(27.6)	21(23.6)	
Visit dental clinics in the previous year > =once, n (%)	80(9.1)	20(9.7)	25(9.4)	25(8.0)	10(11.2)	0.777
Dental symptoms, n (%)	395(45.1)	97(47.1)	120(45.0)	139(44.4)	39(43.8)	0.965
Toothache, n (%)	180(45.6)	43(44.3)	54(45.0)	63(45.3)	20(51.3)	0.898
Bleeding gums, n (%)	164(41.5)	34(35.1)	52(43.3)	60(43.2)	18(46.2)	0.506
Swollen gums, n (%)	124(31.4)	22(22.7)	45(37.5)	44(31.7)	13(33.3)	0.135
Gingival hyperplasia, n (%)	66(16.7)	13(13.4)	19(15.8)	24(17.3)	10(25.6)	0.377
Bad breath, n (%)	172(43.5)	45(46.4)	44(36.7)	62(44.6)	21(53.8)	0.224
Others, n (%)	11(2.8)	1(1.0)	4(3.3)	4(2.9)	2(5.1)	0.566
Caries, n (%)	344(39.3)	96(46.6)	95(35.6)	124(39.6)	28(31.5)	0.027
Median (range)	2(1–32)	2(1–28)	2(1–24)	2(1–32)	2(1–30)	0.795
Missing teeth due to caries, n (%)	225(25.7)	68(33.0)	66(24.7)	75(24.0)	16(18.0)	0.025
Median (range)	2(1–32)	2(1–28)	2(1–24)	2(1–32)	2(1–28)	0.687
Dental injury due to seizures, n (%)	166(19.0)	26(12.6)	56(21.0)	62(19.8)	22(24.7)	0.044
Tooth fracture due to seizures, median (range)	1(0–32)	1(0–9)	1(0–10)	1(0–32)	2(0–6)	0.220
Tooth detached/extracted due to seizures, median (range)	1(0–32)	2(0–27)	1(0–10)	1(0–32)	2(0–25)	0.108
Dental injury for other reasons, n (%)	63(7.2)	16(7.8)	25(9.4)	17(5.4)	5(5.6)	0.295
Tooth fracture for other reasons, median (range)	1(0–15)	1(0–10)	1(0–15)	0(0–8)	0(0–2)	0.969
Tooth detached/extracted for other reasons, median (range)	1(0–28)	2(0–10)	0(0–6)	1(0–28)	2(0–5)	0.163
Natural tooth loss except change to adult teeth, n (%)	105(12.0)	29(14.1)	40(15.0)	29(9.3)	7(7.9)	0.082
Natural tooth loss except change to adult teeth, median (range)	2(1–28)	3(1–28)	2(1–28)	2(1–28)	2(1–25)	0.793
Dental restorative treatment, n (%)	178(20.3)	54(26.2)	53(19.9)	56(17.9)	15(16.9)	0.113
Dental crown, n (%)	27(15.1)	6(12.5)	8(16.0)	10(18.9)	3(21.4)	
Dental prosthesis, n (%)	127(71.3)	40(83.3)	40(80.0)	37(69.8)	10(71.4)	
Others, n (%)	11(6.2)	2(4.2)	2(4.0)	6(11.3)	1(7.1)	
Dental diagnosis, n (%)	62(7.1)	14(6.8)	24(9.0)	17(5.4)	7(7.9)	0.413
Pulpal and Periapical disease, n (%)	26(41.9)	3(21.4)	12(50.0)	9(52.9)	2(28.6)	0.220
Gingivitis, n (%)	37(59.7)	9(64.3)	12(50.0)	9(52.9)	7(100.0)	0.106
Periodontitis, n (%)	21(33.9)	6(42.9)	9(37.5)	4(23.5)	2(28.6)	0.672
Others, n (%)	5(8.1)	3(21.4)	1(4.2)	1(5.9)	0(0.0)	0.205

^a The total number of participants having data of brushing time data was 868. Among them, 204 were seizure-free, 263 had < 1 seizure/month, 312 had 1–4 seizures/month, and 89 had > 4 seizures/month.

Table 3
Unadjusted and adjusted odds ratios for factors related to seizure frequency using logistic regression models.

Seizure frequency	Brushing teeth at least once a day		Dental injury due to seizures		Presence of caries		Missing teeth due to caries	
	OR (95%CI)	P value	OR (95%CI)	P value	OR (95%CI)	P value	OR (95%CI)	P value
Univariate analysis								
Seizure free (reference)	1		1		1		1	
< 1 seizure/month	0.83(0.56–1.23)	0.358	1.84(1.11–3.05)	0.019	0.62(0.43–0.90)	0.012	0.66(0.44–0.99)	0.044
1–4 seizures/month	0.74(0.51–1.08)	0.122	1.72(1.05–2.83)	0.032	0.74(0.52–1.06)	0.097	0.64(0.43–0.94)	0.024
> 4 seizures/month	0.43(0.26–0.72)	0.001	2.26(1.20–4.26)	0.012	0.51(0.30–0.87)	0.012	0.44(0.24–0.82)	0.009
Multivariate analysis*								
Seizure free (reference)	1		1		1		1	
< 1 seizure/month	0.72(0.48–1.09)	0.120	2.07(1.22–3.50)	0.007	0.61(0.41–0.91)	0.015	0.70(0.46–1.09)	0.115
1–4 seizures/month	0.60(0.40–0.90)	0.013	2.12(1.25–3.58)	0.005	0.84(0.57–1.24)	0.381	0.82(0.54–1.26)	0.375
> 4 seizures/month	0.38(0.22–0.66)	0.001	3.09(1.57–6.07)	0.001	0.65(0.37–1.15)	0.143	0.69(0.35–1.34)	0.267

Note: The P values in the table were derived from the Wald test.

* Multivariate analyses were fitted after controlling for age, onset age, sex, epilepsy aetiology, seizure type, and whether the individual had more than one-year of AED treatment.

hyperplasia in people taking phenytoin in our study was much lower than widely reported [28,29]. The variance between studies may be a result of mild gingival hyperplasia being easily missed by individuals without subjective symptoms and with inadequate oral health education, or missed without comprehensive oral examinations. Mild gingival hyperplasia is less likely to induce strong emotional reactions and may therefore be under-reported. The aetiopathogenesis of drug-induced gingival hyperplasia is still not clear but likely multifactorial [30]; poor oral hygiene has been suggested as an important determinant [31].

Physicians and dentists need to cooperate in the management of people with epilepsy. Oral health should not be ignored by physicians and those who require dental care should be promptly referred to dentistry, as dentists have an important role to play.

Our study has limitations. Firstly, most demographic, clinical and oral health data were self-reported by the participants, and recollection retrieved from the past; this may lead to reporting bias (such as under-reporting of stigmatizing behaviors or over-reporting of healthy behaviors) and recall bias. Secondly, confounders such as levels of education and income, literacy, household poverty indices and nutritional status would also be expected to be associated with adverse dental outcomes. We could not adjust for these confounders in our model due to the limitation of the data. Participants were, however, all rural residents with similar socio-economic backgrounds. Thirdly, we had no controls, but were able to compare our findings to previous reports. Lastly, it is possible that village physicians may not have properly identified oral symptoms or diseases so this may have led to underestimation of the problems. To avoid this, future studies should include full dental examinations, performed under standard dentistry conditions.

5. Conclusions

Oral hygiene routine in people with epilepsy in rural China seems precarious and this results in poor oral health which may impact quality of life. Seizure frequency was significantly associated with seizure-related dental injury, and with bad oral health habits in people with epilepsy in rural China. Therefore, control of seizures must take precedence in epilepsy. As a determinant factor of quality of life, every effort should be made to improve oral health and dental status, especially in people who are already detrimentally overwhelmed by their condition.

Conflict of interest

JWS has received research funding from Eisai, GSK and UCB, personal fees from Eisai, UCB, and Janssen, outside the submitted work. All other authors have no conflict of interest to report.

Disclosures

JWS has received research funding from Eisai, GSK and UCB, personal fees from Eisai, UCB, and Janssen outside the submitted work. All other authors have no disclosures to report.

Ethical publication statement

We confirm that we have read the journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

Acknowledgements

JWS is based at UCLH/UCL Comprehensive Biomedical Research Centre, which receives a proportion of funding from the UK Department of Health's National Institute for Health Research Biomedical Research Centers funding scheme. He receives support from the Dr. Marvin Weil Epilepsy Research Fund and UK Epilepsy Society. This study was funded by Key Research Project of the Chinese Ministry of Science and Technology (Grant No. 2016YFC0904400), a NIH/NINDS grant (1R21NS069223-01), and National Natural Science Foundation of China (81271443). We thank all the village physicians for their hard field work of follow-up, and all participants for their cooperation.

References

- [1] Beghi E, Berg A, Carpio A, Forsgren L, Hesdorffer DC, Hauser WA, et al. Comment on epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* 2005;46:1698–9. author reply 1701–1702.
- [2] Wang WZ, Wu JZ, Wang DS, Dai XY, Yang B, Wang TP, et al. The prevalence and treatment gap in epilepsy in China: an ILAE/IBE/WHO study. *Neurology* 2003;60:1544–5.
- [3] Adebayo PB, Akinoyemi RO, Ogun SA, Ogunniyi A. Seizure severity and health-related quality of life of adult Nigerian patients with epilepsy. *Acta Neurol Scand* 2014;129:102–8.
- [4] Blond BN, Detyniecki K, Hirsch LJ. Assessment of treatment side effects and quality of life in people with epilepsy. *Neurol Clin* 2016;34:395–410. viii.
- [5] Petersen PE. Global policy for improvement of oral health in the 21st century - implications to oral health research of World Health Assembly 2007, World Health Organization. *Community Dent Oral Epidemiol* 2009(37):1–8.
- [6] Petersen PE. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Community Dent Oral Epidemiol* 2003;31(Suppl 1):3–23.
- [7] Biazevic MGH, Rissotto RR, Michel-Crosato E, Mendes LA, Mendes MOA. Relationship between oral health and its impact on quality of life among adolescents. *Braz Oral Res* 2008;22:36–42.
- [8] Kitagawa M, Kurahashi T, Matsukubo T. Relationship between general health, lifestyle, oral health, and periodontal disease in adults: a large cross-sectional study in Japan. *Bull Tokyo Dent Coll* 2017;58:1–8.
- [9] Huang JS, Ho KY, Chen CC, Wu YM, Wang CC, Ho YP, et al. Collagen synthesis in idiopathic and dilantin-induced gingival fibromatosis. *Kaohsiung J Med Sci*

- 1997;13:141–8.
- [10] Zeng K, Wang X, Xi Z, Yan Y. Adverse effects of carbamazepine, phenytoin, valproate and lamotrigine monotherapy in epileptic adult Chinese patients. *Clin Neurol Neurosurg* 2010;112:291–5.
- [11] Brodie MJ, Kwan P. Current position of phenobarbital in epilepsy and its future. *Epilepsia* 2012;53(Suppl 8):40–6.
- [12] Károlyházy K, Kovács E, Kivovics P, Fejérdy P, Arányi Z. Dental status and oral health of patients with epilepsy: an epidemiologic study: oral health of patients with epilepsy. *Epilepsia* 2003;44:1103–8.
- [13] Ogumbode EO, Adamolekun B, Akintomide AO. Oral health and dental treatment needs in Nigerian patients with epilepsy. *Epilepsia* 1998;39:590–4.
- [14] Berg AT, Berkovic SF, Brodie MJ, Buchhalter J, Cross JH, van Emde Boas W, et al. Revised terminology and concepts for organization of seizures and epilepsies: report of the ILAE Commission on Classification and Terminology, 2005-2009. *Epilepsia* 2010;51:676–85.
- [15] Liu L, Zhang Y, Wu W, Cheng R. Characteristics of dental care-seeking behavior and related sociodemographic factors in a middle-aged and elderly population in northeast China. *BMC Oral Health* 2015;15:66.
- [16] Dash D, Sebastian TM, Aggarwal M, Tripathi M. Impact of health education on drug adherence and self-care in people with epilepsy with low education. *Epilepsy Behav* 2015;44:213–7.
- [17] Dantas-Neta NB, Carvalho e Souza CH, Mendes Alencar SM, Prado Júnior RR, Mendes RF. Dental findings in children with west syndrome: dental findings in children with west syndrome. *Spec Care Dent* 2014;34:291–4.
- [18] Aragon CE, Burneo JG, Helman J. Occult maxillofacial trauma in epilepsy. *J Contemp Dent Pract* 2001;2:26–32.
- [19] Lawn ND, Bamlet WR, Radhakrishnan K, O'Brien PC, So EL. Injuries due to seizures in persons with epilepsy: a population-based study. *Neurology* 2004;63:1565–70.
- [20] Costa ALF, Yasuda CL, Jr MCF, Morita ME, Cendes F. Refractory epilepsy is highly associated with severe dentoalveolar and maxillofacial injuries. *Epileptic Disord* 2011;61–4.
- [21] Gurbuz T, Tan H. Oral health status in epileptic children. *Pediatr Int* 2010;52:279–83.
- [22] Nonato ER, Borges MA. Oral and maxillofacial trauma in patients with epilepsy: prospective study based on an outpatient population. *Arquivos de Neuro-Psiquiatria* 2011;69:491–5.
- [23] Ghafoor PAF, Rafeeq M, Dubey A. Assessment of oral side effects of Antiepileptic drugs and traumatic oro-facial injuries encountered in Epileptic children n.d.:3.
- [24] Buck D, Baker GA, Jacoby A, Smith DF, Chadwick DW. Patients' experiences of injury as a result of epilepsy. *Epilepsia* 1997;38:439–44.
- [25] Saengsuwan J, Laohasiriwong W, Boonyaleepan S, Sawanyawisuth K, Tiamkao S, Talkul A. Seizure-related vehicular crashes and falls with injuries for people with epilepsy (PWE) in northeastern Thailand. *Epilepsy Behav* 2014;32:49–54.
- [26] Fong CG, Hung A. Public awareness, attitude, and understanding of epilepsy in Hong Kong Special Administrative Region, China. *Epilepsia* 2002;43:311–6.
- [27] Aragon CE. Understanding the patient with epilepsy and seizures in the dental practice. *J Can Dent Assoc* 2007;73:6.
- [28] Angelopoulos AP. A clinicopathological review. Diphenylhydantoin gingival hyperplasia: 2. Aetiology, pathogenesis, differential diagnosis and treatment. *Dent J* 1975;41(275–7):283.
- [29] Angelopoulos AP. Diphenylhydantoin gingival hyperplasia. A clinicopathological review. 1. Incidence, clinical features and histopathology. *Dent J* 1975;41:103–6.
- [30] Moffitt ML, Bencivenni D, Cohen RE. Drug-induced gingival enlargement: an overview. *Compend Contin Educ Dent* 2013;34:330–6.
- [31] Seymour RA, Ellis JS, Thomason JM. Drug-induced gingival overgrowth and its management. *J R Coll Surg Edinb* 1993;38:328–32.